

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts¹, L. R. Poole², A. Dörnbrack³, and L. W. Thomason¹

¹NASA Langley Research Center, Hampton, Virginia 23681, USA

²Science Systems and Applications, Incorporated, Hampton, Virginia 23666, USA

³DLR, Institut für Physik der Atmosphäre, 82230 Oberpfaffenhofen, Germany

Received: 17 August 2010 – Accepted: 28 September 2010 – Published: 18 October 2010

Correspondence to: M. C. Pitts (michael.c.pitts@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

Spaceborne lidar measurements from CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations) are used to provide a vortex-wide perspective of the 2009–2010 Arctic polar stratospheric cloud (PSC) season to complement more focused measurements from the European Union RECONCILE (reconciliation of essential process parameters for an enhanced predictability of Arctic stratospheric ozone loss and its climate interactions) field campaign. The 2009–2010 Arctic winter was unusually cold at stratospheric levels, especially from mid-December 2009 until the end of January 2010, and was one of only a few winters from the past 52 years with synoptic-scale regions of temperatures below the frost point. More PSCs were observed by CALIPSO during the 2009–2010 Arctic winter than in the previous three Arctic seasons combined. In particular, there were significantly more observations of high number density nitric acid trihydrate (NAT) mixtures (referred to as Mix 2-enh) and ice PSCs. We found that the 2009–2010 season could roughly be divided into four periods with distinctly different PSC optical characteristics. The early season (15–30 December 2009) was characterized by patchy, tenuous PSCs, primarily low number density liquid/NAT mixtures. The second phase of the season (31 December 2009–14 January 2010) was characterized by frequent mountain wave ice clouds that nucleated widespread NAT particles throughout the vortex, including Mix 2-enh. The third phase of the season (15–21 January 2010) was characterized by synoptic-scale temperatures below the frost point which led to a rare outbreak of widespread ice clouds. The fourth phase of the season (22–28 January) was characterized by a major stratospheric warming that distorted the vortex, displacing the cold pool from the vortex center. This final phase was dominated by supercooled ternary solution (STS) PSCs, although NAT particles may have been present in low number densities, but were masked by the more abundant STS droplets at colder temperatures. We also found distinct variations in the relative proportion of PSCs in each composition class with altitude over the course of the 2009–2010 Arctic season. Lower number density liquid/NAT mixtures were most

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



frequently observed in the lower altitude regions of the clouds (below ~18–20 km), which is consistent with CALIPSO observations in the Antarctic. Higher number density liquid/NAT mixtures, especially Mix 2-enh, were most frequently observed at altitudes above 18–20 km, primarily downstream of wave ice clouds. This pattern is consistent with the conceptual model whereby low number density, large NAT particles are precipitated from higher number density NAT clouds (i.e. mother clouds) that are nucleated downstream of mountain wave ice clouds.

1 Introduction

In spite of more than two decades of research, a detailed understanding of the processes leading to polar ozone depletion has not yet fully emerged. The role of polar stratospheric clouds (PSCs) in catalyzing heterogeneous chemical reactions that activate ozone-depleting halogens has become well established, but there are still open questions with regard to the formation of solid PSC particles, especially nitric acid trihydrate, NAT (e.g., Lowe and MacKenzie, 2008). These remaining uncertainties limit our ability to accurately represent the ozone depletion processes in global climate models and call into question the capability of these models to predict the evolution of the ozone layer in a future stratosphere whose composition differs from the current state. This is of particular concern in the Arctic, where winter temperatures hover near the PSC formation threshold and, hence, future stratospheric cooling could lead to enhanced cloud formation and substantially greater ozone losses (WMO, 2007).

The general climatology of Arctic PSCs, such as their large interannual variability and asymmetric longitudinal distribution, has primarily been established through analyses of long-term data records from the Stratospheric Aerosol Measurement (SAM) II (1978–1993) and Polar Ozone and Aerosol Measurement (POAM) II/III (1996–2006) solar occultation satellite instruments (e.g. Poole and Pitts, 1994; Fromm et al., 2003). While extremely valuable, these historical occultation data sets provide only limited information on PSC composition and, due to significant sampling limitations, their

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



climatologies are not necessarily representative of the polar regions as a whole (Pitts et al., 2007). Ground-based and airborne lidars have also contributed significantly to our understanding of Arctic PSC occurrence and composition, but these datasets are limited in spatial coverage and are strongly influenced by local meteorology. For instance, ice PSCs have been observed frequently by lidar near the Scandinavian mountains (e.g. Dörnbrack et al., 2002; Reichardt et al., 2004; Blum et al., 2005) and over Sodankylä, Finland (Müller et al., 2001), but only under exceptional meteorological conditions over Ny Alesund (Maturilli and Dörnbrack, 2006; Massoli et al., 2006) and Greenland (di Sarra et al., 2002; Buss et al., 2004). A more complete picture of PSC morphology and composition on vortex-wide scales is now being provided by spaceborne lidar measurements from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) (Winker et al., 2007, 2009) and passive infrared measurements from MIPAS (Michelson Interferometer for Passive Atmospheric Sounding) on Envisat (e.g. Höpfner et al., 2006, 2009).

The European Union has also implemented the new four-year research project RECONCILE (reconciliation of essential process parameters for an enhanced predictability of Arctic stratospheric ozone loss and its climate interactions) for more comprehensive and detailed investigations of key processes governing Arctic ozone depletion. A vital component of RECONCILE is an intensive field campaign of coordinated aircraft, balloon, and ground-based measurements related to PSCs and ozone chemistry that was conducted in the Arctic during January–March 2010 (von Hobe et al., 2010). The aircraft campaign was based in Kiruna, Sweden where the M55 Geophysica high-altitude research aircraft (Stefanutti et al., 1999) conducted a series of science flights carrying a full suite of in situ and remote particle and gas-phase constituent sensors. The flight campaign was implemented in two phases, with eight flights in the first phase from 17–28 January 2010 and four flights in the second phase from 27 February–5 March 2010. The RECONCILE Match balloon campaign took place in parallel with the aircraft campaign with more than 300 sondes launched from 26 stations across the Arctic.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Five flights during the first phase of the RECONCILE aircraft campaign specifically targeted PSC measurements. Although these aircraft measurements are invaluable for detailed studies of polar processes, they are by nature limited both temporally (spanning a ten-day period) and spatially (localized along Geophysica flight tracks, which themselves were limited by the range of the aircraft and airspace constraints). Hence, the overall goal of this paper is to use spaceborne lidar measurements from CALIPSO to examine the evolution of PSCs during the 2009–2010 Arctic winter on vortex-wide scales and provide context to the more focused measurements obtained during the RECONCILE campaign. In Sect. 2, we briefly describe the CALIPSO PSC detection and composition discrimination algorithm and provide an update on the optical characteristics of PSCs observed by CALIPSO from 2006–2010. In Sect. 3, we describe the evolution of the polar vortex during the 2009–2010 Arctic winter and put this season into historical context from a meteorological perspective using European Centre for Medium-Range Weather Forecasts (ECMWF) and National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) Reanalyses datasets. In Sect. 4, we examine the four-year CALIPSO Arctic PSC record to show how this winter compares with the previous three Arctic winters in terms of PSC characteristics. In Sect. 5, we divide the 2009–2010 winter into four periods with distinctly different PSC optical characteristics and then analyze each of these periods in detail to illustrate how PSC composition evolved during the winter. The summary and conclusions are provided in Sect. 6.

2 CALIPSO PSC detection and composition classification

The CALIPSO satellite was launched in April 2006 and is a component of the A-train satellite constellation (Stevens et al., 2002). CALIPSO has an orbit inclination of 98.2° which provides extensive daily measurement coverage over the polar regions of both hemispheres up to 82° in latitude. The primary instrument on CALIPSO is a lidar (CALIOP, or Cloud-Aerosol Lidar with Orthogonal Polarization) that measures

backscatter at wavelengths of 1064 nm and 532 nm, with the 532-nm signal separated into orthogonal polarization components parallel and perpendicular to the polarization plane of the outgoing laser beam. A description of CALIOP and its on-orbit performance can be found in Hunt et al. (2009), and details on calibration of the CALIOP data are provided by Powell et al. (2009).

CALIOP has proven to be an excellent system for observing PSCs (Pitts et al., 2007, 2009, hereafter referred to as P09). The second-generation CALIPSO PSC algorithm, described in detail in P09, detects clouds using both the CALIOP 532-nm scattering ratio (R_{532} , the ratio of total to molecular backscatter, e.g., Cairo et al. 1999) and the 532-nm perpendicular backscatter coefficient in a successive horizontal averaging procedure that enhances the detection of tenuous PSCs. The algorithm also includes a scheme for classifying PSCs by composition based on the measured CALIOP aerosol depolarization ratio (δ_{aerosol} , the ratio of perpendicular to parallel components of aerosol backscatter, e.g., Cairo et al., 1999) and inverse scattering ratio ($1/R_{532}$). P09 defined four CALIPSO PSC composition classes: supercooled ternary solution (STS), water ice, and two classes (Mix 1 and Mix 2) of liquid/NAT mixtures. Mix 1 denotes mixtures with very low NAT number densities (from about $3 \times 10^{-4} \text{ cm}^{-3}$ – the inferred CALIOP NAT sensitivity threshold – to 10^{-3} cm^{-3}), while Mix 2 denotes mixtures with higher ($> 10^{-3} \text{ cm}^{-3}$) NAT number densities. As noted in P09, the STS class may also include low number densities of NAT particles whose optical signature is masked by the more numerous liquid droplets at cold temperatures. At the date of this writing, CALIPSO has produced a PSC dataset covering four winters in both the Arctic and Antarctic. Figure 1 shows a composite 2-D histogram of these observations (totalling approximately 96 million 5-km horizontal by 180-m vertical cloud “pixels”) in the δ_{aerosol} vs. $1/R_{532}$ coordinate system, with the solid black lines denoting the boundaries of the four PSC composition domains defined by P09. The histogram bin size is 0.02×0.02 and the color scale indicates the total number of cloud observations falling within each bin.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



most of the ice PSC domain, the maximum number of observations occurs at color ratios from 0.75–1.0, indicating large particles. But for ice PSCs with $1/R_{532} < 0.02$, the maximum in the number of observations shifts abruptly to color ratios of 0.25–0.35, which is consistent with mountain wave PSCs having high ice particle number densities (~100% ice activation from the background aerosol) but relatively small particles (1–1.5 μm radius, e.g., Füeglistaler et al., 2003; Carslaw et al., 1998). Therefore, we will interpret CALIPSO ice cloud observations with $1/R_{532} < 0.02$ as likely mountain wave PSCs.

3 The 2009–2010 Arctic winter

The 2009–2010 Arctic winter was unusually cold at stratospheric levels, especially the six-week period from mid-December 2009 until the end of January 2010. Cooling of the Arctic polar vortex generally tracked the 20-year ECMWF climatological mean through late November 2009, when the minimum temperature (T_{MIN}) at the 30 hPa pressure level between 65° N–90° N had dropped to about 192 K (Fig. 3). However, the polar vortex split into two unequally strong lobes during the first ten days of December (Fig. 4, upper row) and the associated Canadian warming prevented a further decline of the minimum temperatures, resulting in T_{MIN} values above the climatological mean. The stronger and colder vortex lobe located over the Canadian sector of the Arctic survived this early warming event, recovered and cooled gradually through mid-January 2010. There was a significant drop in T_{MIN} during this period, to values as much as 7 K below the climatological mean (Fig. 3). Typical flow and temperature fields from this exceptionally cold period are shown in Fig. 4 (bottom row) depicting a coherent polar vortex centered near the North Pole. Another interesting feature during this period was the frequent occurrence of wave-like structures, discernable in the stratospheric temperature and geopotential height fields (Fig. 4, bottom row). These undulations resulted from orographic gravity waves excited by the flow across Greenland. Due to the adiabatic expansion of the air flowing through these mountain waves, the actual stratospheric temperatures could have been below the T_{MIN} values seen in the ECMWF analyses.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the November–March period. In each figure, the 52-year daily average area is depicted by the black dotted curves, the daily 95th percentile value is represented by the upper limit of the gray-shaded region, and the daily values for the 2009–2010 winter are denoted by the red curves. Consistent with the period of unusually low T_{MIN} seen in Fig. 3, there was an extended period of large $A_{T_{\text{NAT}}}$ in 2009–2010. Daily values of $A_{T_{\text{NAT}}}$ exceed the 52-year average from mid-December through late January and were above the 95th percentile (i.e., in the top three years) during much of January. The depth of the cold air reached a peak during 15–21 January 2010 when synoptic-scale areas of $A_{T_{\text{ice}}}$ were observed on a series of days. The $A_{T_{\text{ice}}}$ values during this week were larger than the values for all but a few days over the past 52 years, underscoring how unique the 2009–2010 winter was in this regard. Although the 2009–2010 PSC season was relatively short due to the early major warming, the extent and depth of the cold air during the RECONCILE field campaign in January 2010 has rarely been seen over the past half-century.

The strong coupling between the evolution of the vortex and its associated cold pool and the actual occurrence of PSCs is clearly illustrated in Fig. 7 which shows a series of five-day composite maps of CALIPSO PSC frequency (fraction of observations identified as PSCs) at an altitude of 22 km (approximately 30 hPa) for the 2009–2010 winter. Overlaid on the figures are contours of T_{NAT} and T_{ice} based on Goddard Earth Observing System (GEOS-5) gridded temperature data used in the CALIPSO Level 1 data processing. Values of T_{NAT} and T_{ice} were calculated using estimates of the total abundances of HNO_3 and H_2O derived from daily averages of Aura MLS observations from cloud-free regions of the vortex. During the season, the spatial extent of the CALIPSO PSC observations corresponds well with the region of temperatures below T_{NAT} . Scattered patches of clouds were first observed in mid-December over Greenland as the vortex initially cooled below T_{NAT} . As the vortex continued to strengthen and cool, PSCs became widespread by early January and then persisted until late January when the major stratospheric warming occurred and ended the PSC season. Of particular note is the five-day period ending 19 January (Fig. 7f) when the vortex cooled below T_{ice} at

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

22 km over a large region between roughly 70°–80° N latitude and 10° W–80° E longitude. Although mesoscale cooling events associated with mountain waves commonly produce localized temperature excursions below T_{ice} (e.g., Dörnbrack and Leutbecher, 2001), synoptic-scale regions below T_{ice} such as observed in mid-January 2010 are unusual in the Arctic (Pawson et al., 1995). As we discuss later, this particular event led to a rare outbreak of synoptic-scale ice PSCs.

4 Comparison of 2009–2010 CALIPSO PSC data with prior years

The 2009–2010 PSC season was the most intense of the four-year CALIPSO PSC data record. Figure 8 shows the areal coverage of PSCs observed by CALIPSO in the Arctic for each of the past four winters as a function of altitude and time. The significant year-to-year differences in the observed PSC areal coverage illustrate the large interannual variability associated with Arctic PSC activity where the seasonal evolution of the vortex each winter is quite different with no real “typical” year. Although the PSC season was longer in some earlier years (e.g., 2006–2007 and 2007–2008), the areal extent of PSCs during the 2009–2010 winter was significantly larger than observed in any of the previous three winters, which is consistent with the abnormally cold temperatures observed during this winter (Figs. 3 and 7). In fact, the season total volume of PSCs (areal extent integrated over time and altitude) observed during the 2009–2010 winter was nearly as large as the total from the three previous seasons combined.

The CALIPSO dataset also provides the unique opportunity to examine how the Arctic PSC composition characteristics in 2009–2010 compare with previous years. Figure 9 shows composite 2-D histograms of CALIPSO PSC observations from each of the past four Arctic seasons in the $\delta_{aerosol}$ vs. $1/R_{532}$ coordinate system, with the domains of the CALIPSO PSC composition classes overlaid. The total number of CALIPSO PSC observations in each year is indicated above the color bar in each panel. In addition to there being many more PSC observations during the 2009–2010 season, there are also unique composition differences relative to previous years. A

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particularly striking feature of the 2009–2010 winter is the abundance of ice clouds, with large numbers of observations spread throughout the ice cloud optical domain (Fig. 9d). These widespread ice cloud observations occurred primarily during the week of 15–21 January when the vortex cooled below T_{ice} over synoptic scales as noted in the previous section. In most previous winters (i.e., 2007–2008 and 2008–2009), CALIPSO ice cloud observations are limited to a small number of mountain wave induced events. Another notable feature of the 2009–2010 winter is the large number of Mix 2 and Mix 2-enh PSC observations. In the following section, we will explore these unique aspects of the 2009–2010 season in more detail in attempt to understand the underlying physical mechanisms.

5 2009–2010 PSC season in detail

Through our examination of the 2009–2010 season, we found that it can roughly be divided into four time periods with distinctly different PSC optical characteristics. Figure 10 shows composite 2-D normalized histograms of CALIPSO PSC observations from each of these four periods over the altitude range from 15–30 km in the δ_{aerosol} vs. $1/R_{532}$ coordinate system. The early season (15–30 December 2009), shown in Fig. 10a, was dominated by patchy, tenuous clouds that can be characterized as liquid/NAT mixtures with NAT particles occurring in low numbers/volumes, falling predominantly into the Mix 1 domain. These tenuous PSCs were first detected by CALIPSO on 15 December and at least a few patches of clouds were observed on each day of this period. No mountain wave ice PSCs were observed by CALIPSO during this initial period. Since CALIPSO does not sample the entire stratospheric volume, we cannot preclude the possibility that isolated, small-scale wave clouds occurred, but it is quite unlikely since meteorological conditions during this period were not conducive to mountain wave formation. The detection of NAT particles in December, prior to any observed mountain wave ice PSCs, supports the conclusions of Voigt et al. (2005) and Pagan et al. (2004) that ice nuclei are not a prerequisite for NAT formation. Examples of CALIPSO PSC observations from this period are shown in Fig. 11. The figure shows

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



images of CALIPSO PSC observations, color-coded by composition class, from a sequence of four orbits sweeping across the vortex on 24 December 2009. On the right of each composition image is a map of GEOS-5 temperature and geopotential height fields at 30 hPa, with the corresponding CALIPSO orbit track overlaid as the solid white line. The composition images shown in Fig. 11 illustrate the patchy nature of the clouds observed by CALIPSO during December when it was only marginally cold enough to support PSC formation. A complete set of CALIPSO composition images from the entire 2009–2010 winter is provided as supplemental material.

The nature of the season changed abruptly on 31 December with the first observations of mountain wave ice clouds. This initiates the second phase of the PSC season (31 December–14 January) that was characterized by numerous mountain wave ice cloud events along the east coast of Greenland. The 2-D histogram from this period is shown in Fig. 10b. The cluster of observations with very large scattering ratio values at the right edge of the ice domain corresponds to the mountain wave ice PSC observations. This second phase was also marked by a dramatic increase in observations of Mix 2 and Mix 2-enh clouds, as well as fully developed liquid STS clouds. Figure 12 shows examples of CALIPSO composition images from this period (4 January). A distinct mountain wave ice cloud event ($R > 50$ denoted by dark blue pixels) is visible in Fig. 12b at an altitude of 25 km near 70° N, 28° W along the east coast of Greenland. Along the orbit track west of the mountain wave cloud (Fig. 12a), PSC composition is predominately liquid STS. Along orbits east of the wave cloud (Fig. 12c–e), the PSCs consist mostly of liquid/NAT mixtures, including distinct regions of Mix 2-enh clouds. This pattern of predominantly liquid STS clouds occurring west (generally upstream) and liquid/NAT mixtures occurring east (generally downstream) of wave ice clouds was frequently observed during this period, suggesting that NAT nucleation by wave ice clouds was a major source of the widespread NAT particles observed during this period. This has been observed by others on more limited scales in the Arctic (e.g., Carslaw et al., 1998; Dörnbrack et al., 2002; Füeglistaler et al., 2002; Voigt et al., 2003).

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Figure 13a shows the locations of all CALIPSO observations identified as mountain wave ice clouds ($R_{532} > 50$) during the 31 December–14 January period. Figure 13b shows analyzed locations for stratospheric mountain wave activity for the same period based on the ECMWF horizontal divergence (DIV) at 30 hPa (see Plougonven et al., 2003 for the use of DIV as an indicator of mountain wave activity). CALIPSO observed numerous mountain wave ice clouds along Greenland's east coast, in agreement with the maximum in mountain wave activity indicated in the ECMWF analyses. CALIPSO also observed a few wave ice clouds over northern Scandinavia on 10 January, which corresponds well with the short-lived (< 24 h) mountain wave event on 10–11 January depicted in the ECMWF analyses over northern Scandinavia. The overall agreement between the locations of CALIPSO wave ice clouds and the ECMWF analyzed mountain waves provides confidence that the CALIPSO wave cloud discrimination approach is robust.

The third distinct phase in the PSC season occurred the week of 15–21 January when a region of the vortex cooled below the frost point (see Fig. 6 and Fig. 7f), resulting in an extensive region of synoptic-scale ice PSCs. The 2-D histogram of CALIPSO PSC observations from this phase is shown in Fig. 10c. The relative maximum in the frequency of PSC observations extending upward to the right in the ice domain is the distinctive signature of synoptic-scale ice clouds as commonly seen during the Antarctic winter (e.g. P09). Also note that the relative number of observations in the Mix 1, Mix 2, and especially Mix 2-enh domains decreased significantly during this phase compared with the previous phase (Fig. 10b). The increase in ice PSC observations and concomitant decrease in NAT mixture observations may be an indicator that heterogeneous nucleation on pre-existing NAT particles is an important mechanism for formation of synoptic-scale ice PSCs. Examples of CALIPSO composition images from this period (18 January) are shown in Fig. 14. As was common during this week, large regions of ice clouds were observed by CALIPSO near the center of the cold pool (roughly north of Scandinavia, east of Spitsbergen, and west of Novaya Zemlya) at altitudes between 20–25 km (Fig. 14b–d). To the west of this ice region, STS was

**The 2009–2010 Arctic
polar stratospheric
cloud season: a
CALIPSO perspective**

M. C. Pitts et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

widespread (Fig. 14a), while liquid/NAT mixtures, including Mix 2-enh, were predominant along the eastern end of the cold pool (Fig. 14e). Also note that wave ice clouds ($R > 50$, denoted by dark blue pixels) were occasionally embedded in the large-scale regions of ice, such as seen near Novaya Zemlya (Fig. 14d). This is consistent with ECMWF analyzed mountain wave occurrence during this period (not shown).

Widespread ice PSC observations disappeared abruptly after 21 January as the vortex warmed above the frost point, signalling the transition to the last phase of the PSC season that lasted from 22–28 January. With the exception of a few wave ice PSCs, no ice clouds were observed during this period. The 2-D histogram of PSC observations from this last period, shown in Fig. 10d, is strikingly different than those of any of the previous periods. STS observations dominate this final phase with essentially no ice PSCs and many fewer liquid/NAT mixtures than in the previous two periods. Examples of CALIPSO composition images from this period (23 January) are shown in Fig. 15. Extensive areas of STS clouds (Fig. 15a–c) were observed by CALIPSO in the cold pool, which was displaced well south and west of the vortex center. Only patchy regions of NAT mixtures were observed, primarily on the eastern side of the cold pool. There are several possible explanations for the dramatic reduction in NAT mixture observations. Wave ice PSCs, a primary source of NAT nuclei, were limited during this period and occurred on the eastern side of the cold pool. Secondly, the displacement of the cold pool from the vortex center limits NAT particle growth, which increases the likelihood that any NAT particles present were masked by STS at colder temperatures. After 28 January, the vortex broke down and no more PSCs were observed by CALIPSO.

P09 showed that there were distinct variations in PSC composition class with altitude over the course of the season in the Antarctic. Following a similar approach, Fig. 16 shows the relative proportion of PSCs observed in the different composition classes as a function of altitude and time during the 2009–2010 Arctic season. For reference, maximum temperature departures from T_{NAT} , based on Aura MLS HNO_3 and H_2O mixing ratio observations and GEOS-5 temperature analyses, are shown in Fig. 16a. Low

number density liquid/NAT mixtures (Fig. 16b) were the most frequently observed composition class early in the season and in the lower altitude regions of the clouds (below ~18–20 km) throughout the season. The predominance of Mix 1 in the lower regions of the clouds is consistent with CALIPSO observations in the Antarctic and corresponds well with the altitude region where the temperature departures below T_{NAT} are smallest (<4 K) (see Fig. 16a). Temperatures in this regime are above the threshold where STS particles begin to grow substantially and, hence, there is a better likelihood of CALIPSO detecting the presence of low numbers of relative large NAT particles characterized by the Mix 1 class. Higher number density liquid/NAT mixtures (Mix 2 and Mix 2-enh) become the majority composition observed above 20 km during roughly the first half of January (Fig. 16c, d) which corresponds well with the period of wave ice cloud activity from 31 December–14 January. The widespread synoptic-scale ice episode is depicted by the relative maximum in ice PSC observations during the third week of January in Fig. 16e. Immediately following this period of widespread ice PSCs, STS becomes the most frequent composition observed above about 18 km (Fig. 16f). The transition of the dominant composition class above 18 km from Mix 2 and Mix 2-enh before the widespread ice event to STS after the ice event is consistent with the change in the dynamical flow regime that occurred in mid-January, as discussed earlier.

The spatial patterns of PSC composition observed by CALIPSO during the active mountain wave period (31 December–January 14) are consistent with the NAT “mother cloud” conceptual model (e.g., Füeglistaler et al., 2002; Dhaniyala et al., 2002; Mann et al., 2005) whereby mountain wave ice clouds nucleate high number density NAT clouds (“mother clouds”) with widespread regions of lower number density NAT mixtures (Mix 1) below these clouds.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

6 Summary and conclusions

An intensive field campaign focused on measurements related to PSCs and ozone chemistry was conducted in the Arctic during January–March 2010 as part of the European Union RECONCILE project. To complement the more focused measurements from the RECONCILE field campaign, we have used spaceborne lidar measurements from CALIPSO to characterize PSC occurrence and composition during the 2009–2010 season on vortex-wide scales. We have expanded the CALIPSO PSC composition definitions from P09 to include two new sub-classes: Mix 2-enh, which have the characteristics of higher number density NAT mixtures similar to Type 1a-enhanced identified in earlier Arctic studies, and mountain wave ice clouds characterized by high particle number densities but relatively small particles.

Compared to winters over the past half-century, the 2009–2010 winter was unusually cold from mid-December through January and one of only a few winters with synoptic-scale temperatures below T_{ice} . The 2009–2010 winter was also exceptional from the CALIPSO perspective with more PSCs observed than in the three previous Arctic seasons combined. The distribution of PSCs among the various composition classes was also different during the 2009–2010 winter with significantly more observations of ice clouds and higher number density NAT mixtures (Mix 2 and Mix 2-enh).

We found that the 2009–2010 season can roughly be divided into four time periods displaying different PSC optical characteristics. The first period (15–30 December) was marked by patchy, tenuous PSCs that were primarily liquid/NAT mixtures with low NAT number densities. The second period (31 December–14 January) was characterized by numerous wave ice clouds and a dramatic increase in liquid/NAT mixtures with higher NAT number densities (especially Mix 2-enh) and fully-developed STS. From an overall vortex perspective, the STS PSCs occurred upstream of the wave ice events, whereas liquid/NAT mixtures were observed downstream of the wave clouds, including distinct contiguous regions of Mix 2-enh PSCs. During the third phase (15–21 January), synoptic-scale ice clouds were observed over regions of the vortex

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that had cooled below the frost point. With the appearance of synoptic-scale ice clouds there was a noticeable decrease in liquid/NAT mixtures, suggesting that heterogeneous nucleation on pre-existing NAT particles may be an important mechanism for formation of synoptic-scale ice PSCs. The fourth period (22–28 January) was dominated by STS PSCs, with essentially no ice clouds and distinctly fewer observations of liquid/NAT mixtures than in any of the three earlier periods. The apparent disappearance of liquid/NAT mixtures is a result of many fewer wave ice clouds – a primary source of NAT nuclei – as well as a displacement of the cold pool away from the vortex center, which limited NAT particle growth and increased the likelihood of NAT particles being masked by STS droplets at low temperatures.

We also found interesting variations in the relative proportion of PSCs in each composition class with altitude over the course of the 2009–2010 Arctic season. Lower number density liquid/NAT mixtures were most frequently observed in the lower altitude regions of the clouds (below ~18–20 km), which is consistent with CALIPSO observations in the Antarctic. Higher number density liquid/NAT mixtures, especially Mix 2-enh, were most frequently observed at altitudes above 18–20 km, primarily downstream of wave ice clouds. This pattern is consistent with the conceptual model whereby low number density, large NAT particles are precipitated from higher number density NAT clouds (i.e. mother clouds) that are nucleated downstream of mountain wave ice clouds.

Supplementary material related to this article is available online at:

<http://www.atmos-chem-phys-discuss.net/10/24205/2010/acpd-10-24205-2010-supplement.zip>.

Acknowledgements. The Aura MLS gas species data were provided courtesy of the MLS team and obtained through the Aura MLS website (<http://mls.jpl.nasa.gov/index-eos-mls.php>). The NCEP Reanalysis data were provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, and obtained from their Web site at <http://www.esrl.noaa.gov/psd/>. The ECMWF data were available through the special project “Effect of nonhydrostatic gravity waves on the stratosphere above Scandinavia” by one of the authors (A. D.). The field activities in Kiruna were funded by

the EC as part of the FP7 project RECONCILE (Grant number: RECONCILE-226365-FP7-ENV-2008-1). We would also like to thank Hal Maring, NASA Radiation Sciences Program manager, and David Considine, Program Scientist for the CALIPSO/CloudSat Missions, for continued support of this research. Support for L. Poole is provided under NASA contract
5 NNL07AA00C.

References

- Biele, J., Tsias, A., Luo, B. P., Carslaw, K. S., Neuber, R., Beyerle, G., and Peter, T.: Nonequilibrium coexistence of solid and liquid particles in Arctic stratospheric clouds, *J. Geophys. Res.*, 106, 22991–23007, 2001.
- 10 Blum, U., Fricke, K. H., Müller, K. P., Siebert, J., and Baumgarten, G.: Long-term lidar observations of polar stratospheric clouds at Esrange in northern Sweden, *Tellus*, 57B, 412–422, 2005.
- Buss, S., Hertzog, A., Hostettler, C., Bui, T. B., Lüthi, D., and Wernli, H.: Analysis of a jet stream induced gravity wave associated with an observed ice cloud over Greenland, *Atmos. Chem. Phys.*, 4, 1183–1200, doi:10.5194/acp-4-1183-2004, 2004.
- 15 Cairo, F., Di Donfrancesco, G., Adriani, A., Pulvirenti, L., and Fierli, F.: Comparison of various linear depolarization parameters measured by lidar, *Appl. Opt.*, 38, 4425–4432, 1999.
- Carslaw, K. S., Wirth, M., Tsias, A., Luo, B. P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J. T., and Peter, T.: Particle microphysics and chemistry in remotely observed mountain polar stratospheric clouds, *J. Geophys. Res.*, 103(D5), 5785–5796, 1998.
- 20 Dhaniyala, S., McKinney, K., and Wennberg, P.: Lee-wave clouds and denitrification of the polar stratosphere, *Geophys. Res. Lett.*, 9, 1322, doi:10.1029/2001GL013900, 2002.
- Dörnbrack, A. and Leutbecher, M.: Relevance of mountain waves for the formation of polar stratospheric clouds over Scandinavia: a 20 year climatology, *J. Geophys. Res.*, 106(D2), 1583–1593, 2001.
- 25 Dörnbrack, A., Birner, T., Fix, A., Flentje, H., Meister, A., Schmid, H., Browell, E. V., and Mahoney, M. J.: Evidence for inertia gravity waves forming polar stratospheric clouds over Scandinavia, *J. Geophys. Res.*, 107(D20), 8287, doi:10.1029/2001JD000452, 2002.
- 30 di Sarra, A., Cacciani, M., Fiocco, G., Fuà, D., and Jørgensen, T. S.: Lidar observations of polar

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

stratospheric clouds over northern Greenland in the period 1990–1997, *J. Geophys. Res.*, 107(D12), doi:10.1029/2001JD001074, 2002.

Fromm, M. D., Alfred, J. M., and Pitts, M.: A unified long-term, high-latitude stratospheric aerosol and cloud database using SAM II, SAGE II, and POAM II/III data: Algorithm description, database definition and climatology, *J. Geophys. Res.*, 108(D12), 4366, doi:10.1029/2002JD002772, 2003.

Füeglistaler, S., Luo, B. P., Buss, S., Wernli, H., Voigt, C., Müller, M., Neuber, R., Hostetler, C. A., Poole, L. R., Flentje, H., Fahey, D. W., Northway, M. J., and Peter, Th.: Large NAT particle formation by mother clouds: Analysis of SOLVE/THESEO-2000 observations, *Geophys. Res. Lett.*, 29, 1610, doi:10.1029/2001GL014548, 2002.

Füeglistaler, S., Buss, S., Luo, B. P., Wernli, H., Flentje, H., Hostetler, C. A., Poole, L. R., Carslaw, K. S., and Peter, T.: Detailed modelling of mountain wave PSCs, *Atmos. Chem. Phys. Discuss.*, in preparation, 2003.

Hansen, D. R. and Mauersberger, K.: Laboratory studies of the nitric acid trihydrate: Implications for the south polar stratosphere, *Geophys. Res. Lett.*, 15, 855–858, 1988.

Höpfner, M., Luo, B. P., Massoli, P., Cairo, F., Spang, R., Snels, M., Di Donfrancesco, G., Stiller, G., von Clarmann, T., Fischer, H., and Biermann, U.: Spectroscopic evidence for NAT, STS, and ice in MIPAS infrared limb emission measurements of polar stratospheric clouds, *Atmos. Chem. Phys.*, 6, 1201–1219, doi:10.5194/acp-6-1201-2006, 2006.

Höpfner, M., Pitts, M. C., and Poole, L. R.: Comparison between CALIPSO and MIPAS observations of polar stratospheric clouds, *J. Geophys. Res.*, 114, D00H05, doi:10.1029/2009JD012114, 2009.

Hu, R.-M., Carslaw, K. S., Hostetler, C., Poole, L. R., Luo, B., Peter, T., Füeglistaler, S., McGee, T. J., and Burris, J. F.: Microphysical properties of wave polar stratospheric clouds retrieved from lidar measurements during SOLVE/THESEO 2000, *J. Geophys. Res.*, 107(D20), 8294, doi:10.1029/2001JD001125, 2002.

Hunt, W. H., Winker, D. M., Vaughan, M. A., Powell, K. A., Lucker, P. L., and Weimer, C.: CALIPSO Lidar Description and Performance Assessment, *J. Atmos. Oceanic Technol.*, 26, 1214–1228, doi:10.1175/2009JTECHA1223.1, 2009.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteor. Soc.*, 77, 437–470, 1996.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- List, R. J.: Smithsonian Meteorological Tables, 6th edition, Smithsonian Institution Press, Washington, DC, 527 pp., 1984.
- Liu, Z., Vaughan, M. A., Winker, D. M., Hostetler, C. A., Poole, L. R., Hlavka, D., Hart, W., and McGill, M.: Use of probability distribution functions for discriminating between cloud and aerosol in lidar backscatter data, *J. Geophys. Res.*, 109, D15202, doi:10.1029/2004JD004732, 2004.
- Lowe, D. and MacKenzie, A. R.: Polar stratospheric cloud microphysics and chemistry, *J. Atmos. Solar-Terr. Phys.*, 70, 13–40, 2008.
- Mann, G. W., Carslaw, K. S., Chipperfield, M. P., and Davies, S.: Large nitric acid trihydrate particles and denitrification caused by mountain waves in the Arctic stratosphere, *J. Geophys. Res.*, 110, D08202, doi:10.1029/2004JD005271, 2005.
- Manney, G. L., Daffer, W. H., Zawodny, J. M., et al.: Solar occultation satellite data and derived meteorological products: Sampling issues and comparisons with Aura Microwave Limb Sounder, *J. Geophys. Res.*, 112, D24S50, doi:10.1029/2007JD008709, 2007.
- Massoli, P., Maturilli, M., and Neuber, R.: Climatology of Arctic polar stratospheric clouds as measured by lidar in Ny-Ålesund, Spitsbergen (79° N, 12° E), *J. Geophys. Res.*, 111, D09206, doi:10.1029/2005JD005840, 2006.
- Maturilli, M. and Dörnbrack, A.: Polar stratospheric ice cloud above Spitsbergen, *J. Geophys. Res.* 111, D18, D18210, doi:10.1029/2005JD006967, 2006.
- Müller, M., Neuber, R., and Beyerle, G.: Non-uniform PSC occurrence within the Arctic polar vortex, *Geophys. Res. Lett.*, 28, 4175–4178, 2001.
- Pagan, K. L., Tabazadeh, A., Drdla, K., Hervig, M. E., Eckermann, S. D., Browell, E. V., Legg, M. J., and Foschi, P. G.: Observational evidence against mountain-wave generation of ice nuclei as a prerequisite for the formation of three solid nitric acid polar stratospheric clouds observed in the Arctic in early December 1999, *J. Geophys. Res.*, 109, D04312, doi:1029/2003JD003846, 2004.
- Pawson, S. and Naujokat, B.: The cold winters of the middle 1990s in the northern lower stratosphere, *J. Geophys. Res.*, 104(D12), 14209–14222, 1999.
- Pawson, S., Naujokat, B., and Labitzke, K.: On the polar stratospheric cloud formation potential on the northern stratosphere, *J. Geophys. Res.*, 100(D11), 23215–23225, 1995.
- Pitts, M. C., Thomason, L. W., Poole, L. R., and Winker, D. M.: Characterization of Polar Stratospheric Clouds with spaceborne lidar: CALIPSO and the 2006 Antarctic season, *Atmos. Chem. Phys.*, 7, 5207–5228, doi:10.5194/acp-7-5207-2007, 2007.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, doi:10.5194/acp-9-7577-2009, 2009.
- Plougonven, R., Teitelbaum, H., and Zeitlin, V.: Inertia gravity wave generation by the tropospheric midlatitude jet as given by the Fronts and Atlantic Storm-Track Experiment radio soundings, *J. Geophys. Res.*, 108(D21), 4686, doi:10.1029/2003JD003535, 2003.
- Poole, L. R. and Pitts, M. C.: Polar stratospheric cloud climatology based on Stratospheric Aerosol Measurement II observations from 1978 to 1989, *J. Geophys. Res.*, 99, 13083–13089, 1994.
- Powell, K. A., Hostetler, C. A., Liu, Z., Vaughan, M. A., Kuehn, R. E., Hunt, W. H., Lee, K., Trepte, C. R., Rogers, R. R., Young, S. A., and Winker, D. M.: CALIPSO Lidar Calibration Algorithms: Part I – Nighttime 532 nm Parallel Channel and 532 nm Perpendicular Channel, *J. Atmos. Oceanic Technol.*, 26, 2015–2033, doi:10.1175/2009-JTECHA1242.1, 2009.
- Reichardt, J., Dörnbrack, A., Reichardt, S., Yang, P., and McGee, T. J.: Mountain wave PSC dynamics and microphysics from ground-based lidar measurements and meteorological modeling, *Atmos. Chem. Phys.*, 4, 1149–1165, doi:10.5194/acp-4-1149-2004, 2004.
- Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., et al.: The CloudSat mission and the A-Train: A new dimension of space-based observations of clouds and precipitation, *Bull. Am. Meteorol. Soc.*, 83, 1771–1790, 2002.
- Stefanutti, L., Sokolov, L., Balestri, S., MacKenzie, A. R., and Khattatov, V.: The M-55 Geophysica as a platform for the airborne polar experiment, *J. Atmos. Ocean. Tech.*, 16, 1303–1312, 1999.
- Tsias, A., Wirth, M., Carslaw, K. S., Biele, J., Mehrtens, H., Reichardt, J., et al.: Aircraft lidar observations of an enhanced type Ia polar stratospheric clouds during APE-POLECAT, *J. Geophys. Res.*, 104(D19), 23961–23969, 1999.
- Voigt, C., Larsen, N., Deshler, T., Kröger, C., Schreiner, J., Mauersberger, K., Luo, B., Adriani, A., Cairo, F., Di Donfrancesco, G., Ovarlez, J., Ovarlez, H., Dörnbrack, A., Knudsen, B., and Rosen, J.: In situ mountain-wave polar stratospheric cloud measurements: Implications for nitric acid trihydrate formation, *J. Geophys. Res.*, 108, 8331, doi:10.1029/2001JD001185, 2003.
- Voigt, C., Schlager, H., Luo, B. P., Dörnbrack, A., Roiger, A., Stock, P., Curtius, J., Vössing, H., Borrmann, S., Davies, S., Konopka, P., Schiller, C., Shur, G., and Peter, T.: Nitric Acid Trihydrate (NAT) formation at low NAT supersaturation in Polar Stratospheric Clouds (PSCs),

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Atmos. Chem. Phys., 5, 1371–1380, doi:10.5194/acp-5-1371-2005, 2005.

von Hobe, M., Schlager, H., Stroh, F., et al.: The RECONCILE Arctic field campaign 2009/2010: an overview, Atmos. Chem. Phys. Discuss., in preparation, 2010.

Winker, D. M., McGill, M., and Hunt, W. H. : Initial performance assessment of CALIOP, Geophys. Res. Lett., 34,L19803, doi:10.1029/2007GL030135, 2007.

Winker, D. M., Vaughan, M. A., Omar, A. H., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.; Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms, J. Atmos. Oceanic Technol., 26, 2310–2323, doi:10.1175/2009JTECHA1281.1, 2009.

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project-Report No. 50, Geneva, Switzerland, 2007.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 1. Composite 2-D histogram of all CALIPSO PSC observations from 2006–2010 in the δ_{aerosol} vs. $1/R_{532}$ coordinate system, with the solid black lines denoting the boundaries of the four PSC composition domains defined by P09. The histogram bin size is 0.02×0.02 and the color scale indicates the total number of cloud observations falling within each bin. The domains of two new composition sub-classes, Mix 2-enh and wave ice, are denoted by dashed boxes.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**The 2009–2010 Arctic
polar stratospheric
cloud season: a
CALIPSO perspective**

M. C. Pitts et al.




Fig. 2. Composite 2-D histogram of lidar color ratio vs. $1/R_{532}$ (with a bin size of 0.054×0.02) for CALIPSO PSC observations with $1/R_{532} < 0.2$ from June 2006 to March 2010. Dashed red circle indicates abrupt shift in color ratio associated with wave clouds at $1/R_{532} < 0.02$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**The 2009–2010 Arctic
polar stratospheric
cloud season: a
CALIPSO perspective**

M. C. Pitts et al.




Fig. 3. Minimum temperatures T_{MIN} (K) between 65° N to 90° N at the 30 hPa pressure surface. Black line: mean value from 1989–2009; red line the T_{MIN} evolution from August 2009 through April 2010; shaded area encompasses the minimum/maximum T_{MIN} between 1989 and 2009. Source: ECMWF reanalyses interim (ERA interim) data provided at 6 hourly temporal resolutions, see: <http://www.ecmwf.int>.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion


Temperature (K) and Geopotential Height (m)



Scaled Potential Vorticity ($10^{-4} \text{ Km}^2/\text{s kg}$)



Temperature (K) and Geopotential Height (m)



Scaled Potential Vorticity ($10^{-4} \text{ Km}^2/\text{s kg}$)






Fig. 4. Left: Absolute Temperature (K, colour shaded) and geopotential height (m; black contour lines) at the 30 hPa pressure surface. Right: Scaled or modified potential vorticity $PV \cdot (\Theta/420 \text{ K})^{-9/2}$ (PVU; colour shaded) at $\Theta = 475 \text{ K}$. Operational ECMWF T1279/L91 analyses interpolated at a regular $0.5^\circ \times 0.5^\circ$ latitude/longitude grid.

Temperature (K) and Geopotential Height (m)




ECMWF Pressure Level 030 hPa VT: 24.01.2010 18 UT

Scaled Potential Vorticity ($10^8 \text{ Km}^2/\text{s kg}$)




ECMWF Theta Level 475 K VT: 24.01.2010 12 UT

Temperature (K) and Geopotential Height (m)



ECMWF Pressure Level 030 hPa VT: 03.02.2010 12 UT

Scaled Potential Vorticity ($10^8 \text{ Km}^2/\text{s kg}$)



ECMWF Theta Level 475 K VT: 03.02.2010 12 UT

Fig. 5. Left: Absolute Temperature (K, colour shaded) and geopotential height (m; black contour lines) at the 30 hPa pressure surface. Right: Scaled or modified potential vorticity $PV \cdot (\Theta/420 \text{ K})^{-9/2}$ (PVU; colour shaded) at $\Theta = 475 \text{ K}$. Operational ECMWF T1279/L91 analyses interpolated at a regular $0.5^\circ \times 0.5^\circ$ latitude/longitude grid.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 6. $A_{T_{\text{NAT}}}$ (top panel) and $A_{T_{\text{ice}}}$ (bottom panel) statistics at 30 hPa for the November–March period based on the temperature record for the 52-year period 1958–2009 from the NCEP/NCAR Reanalyses 1 project. In each figure, the 52-year daily average area is depicted by the black dotted curves, the daily 95th percentile value is represented by the upper limit of the gray-shaded region, and the daily values for the 2009–2010 winter are denoted by the red curves.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 7. Five-day composites of CALIPSO PSC frequency (number of PSC observations/total number of observations) at an altitude of 22 km (approximately 30 hPa) during the 2009–2010 Arctic winter. Contours of T_{NAT} (red) and T_{ice} (black) are indicated on the figures.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 8. Daily areal coverage of PSCs as a function of altitude for the four Arctic winters in the CALIPSO data record to date.

24235

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 9. Composite 2-D histograms of CALIPSO PSC observations from each of the last four Arctic seasons in the δ_{aerosol} vs. $1/R_{532}$ coordinate system, with the domains of the CALIPSO PSC composition classes overlaid.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 10. Normalized composite 2-D histograms of CALIPSO PSC observations from each of four periods during the 2009–2010 Arctic winter over the altitude range from 15–30 km in the δ_{aerosol} vs. $1/R_{532}$ coordinate system.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 11. Examples of CALIPSO PSC composition observations from four orbits during 24 December 2009. GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC are shown on the right with the location of the corresponding CALIPSO orbit tracks indicated by the white lines.




Fig. 12. Examples of CALIPSO PSC composition observations from five orbits during 4 January 2010. GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC are shown on the right with the location of the corresponding CALIPSO orbit tracks indicated by the white lines.

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The 2009–2010 Arctic
polar stratospheric
cloud season: a
CALIPSO perspective**

M. C. Pitts et al.




Fig. 13. Locations of wave ice PSCs as observed by CALIPSO ($R > 50$) during 31 December 2009–14 January 2010 (left panel) and mountain wave occurrence (number of six-hourly events) as derived from the ECMWF horizontal divergence conditions $DIV > 2 \times 10^{-4} \text{ s}^{-1}$ (right panel) at 30 hPa for the same period. Operational ECMWF T799 analyses interpolated at a regular $0.5^\circ \times 0.5^\circ$ latitude/longitude grid.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 14. Examples of CALIPSO PSC observations from five orbits during 18 January 2010. GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC are shown on the right with the location of the corresponding CALIPSO orbit tracks indicated by the white lines.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 15. Examples of CALIPSO PSC observations from four orbits during 23 January 2010. GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC are shown on the right with the location of the corresponding CALIPSO orbit tracks indicated by the white lines.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective

M. C. Pitts et al.




Fig. 16. Relative proportion of PSCs observed in the **(b)** Mix 1, **(c)** Mix 2, **(d)** Mix 2-enh, **(e)** ice, and **(f)** STS composition classes as a function of altitude and time. For reference, the maximum temperature departure from T_{NAT} derived from GEOS-5 analyses is shown in panel **(a)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion