

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**The effect of
meteorological and
chemical factors**

M. Z. Markovic et al.

The effect of meteorological and chemical factors on the agreement between observations and predictions of fine aerosol composition in Southwestern Ontario during BAQS-Met

M. Z. Markovic¹, K. L. Hayden², J. G. Murphy¹, P. A. Makar², R. A. Ellis¹,
R. Y.-W. Chang¹, J. G. Slowik¹, C. Mihele², and J. Brook²

¹Department of Chemistry, University of Toronto, Toronto, Ontario, Canada

²Air Quality Research Division, Environment Canada, Toronto, Ontario, Canada

Received: 13 September 2010 – Accepted: 1 October 2010 – Published: 22 October 2010

Correspondence to: J. G. Murphy (jmurphy@chem.utoronto.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

The Border Air Quality and Meteorology Study (BAQS-Met) was an intensive, collaborative field campaign during the summer of 2007 that investigated the effects of transboundary pollution, local pollution, and local meteorology on regional air quality in Southwestern Ontario. This analysis focuses on the measurements of the inorganic constituents of particulate matter with diameter of less than $1\ \mu\text{m}$ (PM_{10}), with a specific emphasis on nitrate. We evaluate the ability of AURAMS, the Environment Canada's chemical transport model, to represent regional air pollution in SW Ontario by comparing modelled aerosol inorganic chemical composition with measurements from Aerosol Mass Spectrometers (AMS) onboard the National Research Council (NRC) of Canada Twin Otter aircraft and at a ground site in Harrow, ON. The agreement between modelled and measured $p\text{NO}_3^-$ at the ground site (observed mean (M_{obs}) = $0.50\ \mu\text{g m}^{-3}$; modelled mean (M_{mod}) = $0.58\ \mu\text{g m}^{-3}$; root mean square error (RSME) = $1.27\ \mu\text{g m}^{-3}$) was better than aloft (M_{obs} = $0.32\ \mu\text{g m}^{-3}$; M_{mod} = $0.09\ \mu\text{g m}^{-3}$; RSME = $0.48\ \mu\text{g m}^{-3}$). Possible reasons for discrepancies include errors in (i) emission inventories, (ii) atmospheric chemistry, (iii) predicted meteorological parameters, or (iv) gas/particle thermodynamics in the model framework. Using the inorganic thermodynamics model, ISORROPIA, in an offline mode, we find that the assumption of thermodynamic equilibrium is consistent with observations of gas and particle composition at Harrow. We develop a framework to assess the sensitivity of PM_{10} nitrate to meteorological and chemical parameters and find that errors in both the predictions of relative humidity and free ammonia ($\text{FA} \equiv \text{NH}_3(\text{g}) + \text{NH}_4^+ - \text{SO}_4^{2-}$) are responsible for the poor agreement between modelled and measured values.

1 Introduction

Atmospheric PM_{10} are liquids and solids suspended in air with aerodynamic diameter of less than $1\ \mu\text{m}$. They are ubiquitous in the atmosphere and are important because they

ACPD

10, 24781–24820, 2010

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



alter the radiative balance of the Earth either by scattering incoming solar radiation directly, or by serving as cloud condensation nuclei (CCN) in cloud formation. PM₁ also reduces visibility through the formation of haze, affects sensitive ecosystems through acid deposition, and provides surface area for several important atmospheric chemical reactions (Seinfeld, 1996; Jacob, 1999; Finlayson-Pitts, 2000; Pinder et al., 2007). Recent studies have linked PM₁ to pulmonary disease and reduced lung function (Oberdorster, 2001; Gauderman et al., 2004), cardiac arrest (Dockery, 2001; Peters et al., 2001; Dominici et al., 2006) and in some cases even premature death (Burnett et al., 2000; Pope et al., 2002). For these reasons understanding the processes that govern the sources, sinks and chemical composition of PM₁ is of utmost importance.

In Eastern North America, more than 50% of the composition of atmospheric fine particulate matter is inorganic in nature, containing mostly sulphate (SO₄²⁻), ammonium (NH₄⁺) and nitrate (NO₃⁻) (Jacob, 1999; Pinder et al., 2007). These constituents of inorganic PM₁ are secondary in nature and are formed by physical processes (e.g. condensation) or chemical reactions of precursor gases such as H₂SO_{4(g)}, HNO_{3(g)} and NH_{3(g)}. Sulphuric and nitric acid are atmospheric oxidation products of the gases SO₂ and NO_x (≡ NO + NO₂) respectively, and ammonia is directly emitted into the atmosphere mainly from agricultural sources, although in urban areas other sources such as motor vehicles and human waste are non-trivial (Seinfeld, 1996; Jacob, 1999; Finlayson-Pitts, 2000; Pinder et al., 2007).

The secondary formation of inorganic PM₁ is governed by thermodynamics. Once formed, H₂SO₄ will remain in the condensed (particle) phase due to its low vapour pressure. NH_{3(g)} will react preferentially with H₂SO_{4(s,aq)} to neutralize it even in the presence of significant HNO_{3(g)}. Up to 2 moles of NH_{3(g)} will react with H₂SO_{4(s,aq)} to form particulate ammonium sulphate (Stelson and Seinfeld, 1982; Seinfeld, 1996).

Depending on the ambient temperature (T), relative humidity (RH) and the amounts of each species, semi-volatile HNO₃ and NH₃ can partition between the gas and particle phases in order to establish a thermodynamic equilibrium (HNO_{3(g)} ↔ NO_{3_(aq,s)}⁻; NH_{3(g)} ↔ NH_{4_(aq,s)}⁺). The formation of NO_{3_(aq,s)}⁻ depends strongly on not only

RH and T , but also on the absolute amounts and availability of total ammonia (TA \equiv $\text{NH}_{3(\text{g})} + \text{NH}_4^+(\text{aq,s})$), total sulphate (TS \equiv $\text{H}_2\text{SO}_{4(\text{g})} + \text{HSO}_4^-(\text{aq,s}) + \text{SO}_4^{2-}(\text{aq,s}) \sim \text{HSO}_4^-(\text{aq,s}) + \text{SO}_4^{2-}(\text{aq,s})$) and total nitrate (TN \equiv $\text{HNO}_{3(\text{g})} + \text{NO}_3^-(\text{aq,s})$) (Seinfeld, 1996).

Only when sulphuric acid is completely neutralized, can the remaining $\text{NH}_{3(\text{g})}$, if available, react with $\text{HNO}_{3(\text{g})}$ to form $\text{NH}_4\text{NO}_{3(\text{aq,s})}$. Thermodynamically, the formation of ammonium nitrate, and hence particulate nitrate, is governed by the $\text{NH}_4\text{NO}_{3(\text{aq,s})}$ dissociation constant, K_p which is related to the product of partial pressures of $\text{NH}_{3(\text{g})}$ and $\text{HNO}_{3(\text{g})}$ by the following equation (Stelson and Seinfeld, 1982; Mozurkewich, 1993; Seinfeld, 1996):

$$\ln K_p = 84.6 - (24\,220/T) - 6.1 \cdot \ln(T/298.15), \text{ where } K_p = P_{\text{NH}_3} \cdot P_{\text{HNO}_3} \quad (1)$$

where T is in Kelvin, and P_x is the gas phase partial pressure in ppb. When the aerosol is aqueous, the equilibrium product (K_p^*) depends on the solution concentrations, the Henry's law constants (H_x) and activities (α_x) must be considered (Mozurkewich, 1993; Finlayson-Pitts, 2000):

$$K_p^* = \frac{\alpha_{\text{NH}_3} \cdot \alpha_{\text{HNO}_3}}{H_{\text{NH}_3} \cdot H_{\text{HNO}_3}} \quad (2)$$

In this manuscript, we refer to particulate nitrate as ρNO_3^- , whether in the solid or aqueous form, and similarly particulate ammonium is referred to as ρNH_4^+ . The term ρSO_4^{2-} refers to the sum of $\text{SO}_4^{2-}(\text{aq,s})$ and $\text{HSO}_4^-(\text{aq,s})$, which cannot be distinguished by our measurements.

When meteorological conditions in the atmosphere change, the distribution of the ammonium-sulphate-nitrate system between the gas and particle phases may also change. The change subsequently alters the atmospheric lifetime of some of the constituents. Compared to the gas phase, the dry deposition velocities of ammonium and nitrate in fine particles are much lower, so partitioning to the particle phase results in higher steady state concentrations of atmospheric reactive nitrogen. The compounds

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

can then be transported longer distances downwind of emission sources to remote regions where they can affect fragile ecosystems through acid and nutrient deposition (Seinfeld, 1996; Finlayson-Pitts, 2000). As SO_2 and NO_x emissions are decreasing in much of the developed world, it is expected that nitrate will become a more important constituent of inorganic aerosol, both in relative and possibly absolute terms. This will impact the effectiveness of PM precursor emission controls, the spatial distribution of nitrogen deposition, and the rates of many heterogeneous reactions, including e.g. the reactive uptake of N_2O_5 . To assess the ability of a model to examine these impacts and make predictions, it is important to test its ability to determine ρNO_3^- and its sensitivity to a range of input parameters.

In order to describe the processes that govern the formation of particulate nitrate in the atmosphere, numerous thermodynamic partitioning models have been developed. Some of the more recent models include: AIM (Wexler and Seinfeld, 1990; Clegg et al., 1998a, b; Wexler and Clegg, 2002), SCAPE (Kim et al., 1993), EQUISOLV (Jacobson et al., 1996) and ISORROPIA (Nenes et al., 1999). In a review by Zhang et al. (2000), the authors demonstrated that most of the models compared well in predictions of particulate composition under most atmospherically relevant conditions. Compared to other thermodynamic partitioning models, ISORROPIA uses a set of unique process-solving mechanisms with nested algorithms that allow it to be more robust and require much less computational time for thermodynamic calculations (Nenes et al., 1999). The thermodynamic system modelled by ISORROPIA includes ammonium, sulphate, nitrate and water, which are partitioned between gas, liquid and solid phases depending on the meteorological input parameters. During computations, ISORROPIA assumes atmospheric particles to be internally mixed; i.e. it assumes that all particles have identical chemical composition. Due to its robustness and rapid computations, the model can be embedded in chemical transport models or used offline to model thermodynamic partitioning of the inorganic ammonium-sulphate-nitrate-water system (Nenes et al., 1999; Makar et al., 2003).

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A Unified Regional Air Quality Modelling System (AURAMS) (Zhang et al., 2002b; Makar et al., 2003, 2007; Gong et al., 2006, 2008) is a size- and composition-resolved, regional particulate matter and gas modelling system developed by Environment Canada for forecasts of regional air pollution and policy scenario simulations. The model consists of three main components: an emission processor (SMOKE), a meteorological driver model (GEM), and a chemical transport model. The chemical transport model (CTM) makes use of HETV (Makar et al., 2003) for thermodynamic partitioning of chemical species; a vectorized equilibrium solver based on the algorithms of ISORROPIA. In order to make accurate predictions of inorganic aerosol composition, an air quality model must provide accurate representations of meteorology, precursor emissions, oxidant fields, and gas-particle partitioning. In the case of $p\text{NO}_3^-$ predictions, the model is especially sensitive to inputs because of the strong temperature and humidity dependences (Eqs. 1 and 2), and the critical dependence on the availability of ammonia. In a recent assessment of the Community Multiscale Air Quality (CMAQ) model, which also uses the ISORROPIA inorganic equilibrium module, Yu et al. (2005) found the performance of the model for particulate nitrate was strongly dependent on the ability of the model to accurately predict TN, TA, and $p\text{SO}_4^{2-}$. Furthermore, the bias in TA in the model was found to be more critical to the accurate prediction of $p\text{NO}_3^-$ than the bias in $p\text{SO}_4^{2-}$.

The Border Air Quality and Meteorology Study (BAQS-Met) was an intensive, collaborative field campaign during the summer of 2007 that investigated the effects of transboundary pollution, local pollution, and local meteorology on regional air quality in Southwestern Ontario. Air pollution formation in this region is quite complex because local lake- and thermally-forced meteorological circulation interacts with local emissions, and long-range transport of pollutants, to generate some of the highest levels of ozone and PM_1 in Canada. The composition of PM_1 during the campaign was monitored by a network of ground-based and airborne Aerosol Mass Spectrometers (AMS, Aerodyne Research Inc.) (Drewnick et al., 2005; DeCarlo et al., 2006). Instrument details along with supporting meteorological and gas phase measurements relevant to

our analysis are described in Sect. 2.

The goal of this paper is to evaluate the ability of AURAMS to predict the speciation and mass loadings of the inorganic fraction of atmospheric PM₁ during BAQS-Met. A special emphasis was placed on the agreement between ρNO_3^- predictions and measurements (Sect. 3.1). In order to investigate the thermodynamic partitioning as a potential source of large bias in nitrate predictions we performed a series of sensitivity tests by using the ISORROPIA thermodynamic partitioning model in an offline mode (Sect. 3.2). Following this comparison, we develop a framework to assess the sensitivity of ρNO_3^- mass loadings to variability in meteorological and chemical variables used by the model (Sect. 3.3). In Sect. 3.4, we examine the model bias in the availability of ammonia, and in Sect. 3.5, we carry out some model sensitivity tests by replacing the modelled meteorology with measured temperature and relative humidity.

2 Experimental

2.1 Research aircraft

Sixteen research flights (33 project hours) with the NRC's Twin Otter research aircraft were completed between 23 June and 8 July 2007. Research flights were carried out throughout Southwestern Ontario and included multiple passes over Lake Erie and Lake St. Clair to investigate the effects of lake breezes on processing of pollutants (Fig. 1). Vertical profile measurements of trace pollutants were made over land and over the lakes during several day and night flights. In this project, we include the analysis of aircraft data over all flights.

The inorganic composition of PM₁ was measured with a compact time-of-flight aerosol mass spectrometer (c-ToF-AMS, Aerodyne Research, Inc., Billerica, MA, USA) which has been described in detail in the literature (Drewnick et al., 2005; DeCarlo et al., 2006). Briefly, the AMS consists of an aerodynamic lens that focuses particles < 1 μm into a narrow beam, a chopper and time-of-flight chamber for size-selective

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factorsM. Z. Markovic et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

separation of particles, a thermal vaporization plate with an ionizer for Electron Im-
paction (EI) of the vapourized particles, and a time-of-flight mass spectrometer for the
analysis of ionized fragments. The AMS provided mass concentrations of sulphate,
ammonium and nitrate with a time resolution of 30 s. A pressure controlled inlet (PCI)
5 was used in front of the AMS to remove variations in particle sizing and transmission
due to pressure changes in the aerodynamic lens of the AMS (Bahreini et al., 2008;
Hayden et al., 2008; Hayden, 2010). Transmission efficiency experiments indicated
that particles were transmitted through the PCI with near 100% efficiency. The particle
collection efficiency (CE) for the AMS, i.e. the fraction of particles that hit the vaporizer
10 and are vaporized, has been described in detail in the literature (Drewnick et al., 2005;
Kleinman et al., 2008; Dunlea et al., 2009) and is typically evaluated through compar-
isons with other instruments. The CE in this study was determined to vary between
0.5 and 1 as a function of particle acidity and the mass loadings reported in this paper
have been adjusted as described in Hayden et al. (2010). Values below the detection
15 limit were substituted with the detection limits (determined as 3 times the standard de-
viation of mass concentrations for 30 s filtered samples), which were 0.073, 0.024, and
0.014 $\mu\text{g m}^{-3}$ for $p\text{NH}_4^+$, $p\text{SO}_4^{2-}$, and $p\text{NO}_3^-$ respectively. Uncertainty in the measure-
ments is estimated at $\pm 20\%$. For comparison to model output, the observations were
averaged to two minutes.

20 The ambient temperature was measured with a Rosemount temperature probe
mounted on the nose of the aircraft; measurement uncertainty is $\pm 1\%$. The rela-
tive humidity was calculated using the measured dewpoint and temperature and the
estimated error is $\pm 4\%$.

2.2 Harrow monitoring site

25 The Harrow supersite was located in Harrow, ON (42.033° N, 82.893° W), about 5 km
north of Lake Erie and about 30 km south-east of the Detroit-Windsor metropolitan area
(Fig. 1). Measurements were made from 17 June to 11 July 2007. The site was subject
to frequent inflows of transboundary pollution from Michigan to the west and Ohio to

the south. More locally, the atmospheric composition at the site was often influenced by afternoon lake breezes from Lake Erie to the south and less frequently from Lake St. Clair to the north. Harrow is located in a rural area dominated by agricultural activities, mainly the cultivation of corn, soybeans, and fruit.

A c-ToF-AMS similar to that onboard the research aircraft (see above) was deployed to measure the composition of atmospheric PM_1 at the ground site. Instrument collection efficiency was estimated by the acquisition of single particle mass spectra in conjunction with particle counting by an optical scattering module (Cross et al., 2009), which provides a light scattering pulse for particles larger than ~ 215 nm. The collection efficiency was estimated as the fraction of optically-detected particles yielding single particle mass spectra above background levels, and this ratio was assumed to apply across the entire detectable size range of the AMS (Slowik et al., 2010). The vaporization of sulphate and bisulphate species in the AMS produces $H_2SO_{4(g)}$, $SO_{3(g)}$, and $H_2O_{(g)}$, with the ratios of these species dependent on particle composition. $H_2SO_{4(g)}$ and $SO_{3(g)}$ are treated separately in the AMS spectral analysis, with the $H_2O_{(g)}$ formed in equal molar quantities with $SO_{3(g)}$, and the total reported sulphate + bisulphate concentration is the sum of $H_2SO_{4(g)}$, $SO_{3(g)}$ and calculated $H_2O_{(g)}$ (Allan et al., 2004). Detection limits at 5 min time resolution were determined to be 0.200, 0.014, and $0.006 \mu g m^{-3}$ for pNH_4^+ , pSO_4^{2-} and pNO_3^- respectively. Uncertainty in measurements was estimated to be 20%. Data were acquired at 5 min time resolution and averaged to 30 min time steps for our analysis of ground-based data.

Gas phase measurements of ammonia were made using a Quantum Cascade Tunable Infrared Laser Differential Absorption Spectrometer (QC-TILDAS), developed by Aerodyne Research Inc. (Billerica, Massachusetts). Data was acquired with a time resolution of 1 Hz using a specially designed heated quartz inlet (Ellis et al., 2010b) to minimize interferences and sampling biases. Further description of the QC-TILDAS along with a detailed analysis of ammonia during BAQS-Met can be found in Ellis et al. (2010a).

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A chemiluminescence-based NO/NO_x instrument (Thermo Scientific Model 42C) was modified to conduct measurements of NO, NO₂ and NO_y, where NO_y is defined as the sum of all nitrogen oxides (NO_y = NO + NO₂ + HNO₃ + PAN + HONO + NO₃ radical + 2 * N₂O₅ + gaseous organic nitrates + pNO₃⁻). A photolytic converter unit from Ecophysics (model PLC 760 MH) was used to selectively convert NO₂ to NO, and a molybdenum converter heated at 325 °C was used to reduce all NO_y species to NO. The photolytic converter was operated inside the building, while the molybdenum converter unit was housed outside to minimize the losses for the reactive NO_y components such as HNO₃, HONO and NO₃ radical. For this analysis, we used NO_z (≡ NO_y – NO_x) as a constraint on TN levels at Harrow. NO_z is calculated as the difference between measurement of NO following the molybdenum converter and NO following the photolytic converter, after correction for conversion efficiency. Data were available at 1 min time resolution and averaged to 30 min time steps for our analysis.

Relative humidity (RH) and temperature (*T*) measurements were made via Orion Weather Station™ from Columbia Weather Systems Inc. (Hillsboro, Oregon) with 1 min resolution. The uncertainties were ± 1.5% for *T* and ± 4% for RH. Measurements averaged to 30 min were used in our analysis.

2.3 AURAMS model

AURAMS version 1.4.0 was used for the simulations during BAQS-Met. The model was run in forecast mode with 15 km resolution during the campaign to guide the research aircraft and mobile ground-based platforms. After the campaign, a new high-resolution nested version of AURAMS was re-run for BAQS-Met with three levels of nesting: a 2.5 km grid-size high-resolution local domain nested within a 15 km regional domain, nested in turn within a 42 km North American domain. The model ran with a 2 min time step for the innermost local domain and a 15 min time step on the two larger domains. The two lower resolution simulations make use of 15 km grid spacing meteorological input from the Global Environmental Multiscale weather forecast model (GEM, v3.2.2; Cote et al., 1998), while the higher resolution AURAMS simulation made use of GEM

output in a 2.5 km grid spacing configuration. In this analysis, the highest simulation AURAMS output (2.5 km) was used. AURAMS makes use of HETV (Makar et al., 2003) for thermodynamic partitioning of chemical species; a vectorized equilibrium solver based on the algorithms of ISORROPIA. In this configuration, the model was ran with the “metastable state” option on, which treats all particles as exclusively aqueous and prevents crystalline solid formation in the model output. More information about the model can be found in Makar et al. (2010).

2.4 ISORROPIA

ISORROPIA v2.1 was used for all offline re-partitioning work including a series of sensitivity tests. All of the runs in ISORROPIA were performed in “forward mode” where TA, TS, TN, relative humidity and temperature were entered and the model was allowed to re-partition the species between gas and particle phase according to its thermodynamic algorithms. All runs were executed with the “metastable state” option on. Both observations and AURAMS output were separately entered into ISORROPIA and repartitioned in these studies. ISORROPIA output which included, among other things, mass loadings of $\text{NH}_{3(\text{g})}$, $\text{NH}_{4(\text{aq})}^+$, $\text{HSO}_{4(\text{aq})}^-$, $\text{SO}_{4(\text{aq})}^{2-}$, $\text{HNO}_{3(\text{g})}$, and $\text{NO}_{3(\text{aq})}^-$ was compared to original modelled and measured data. We focused our sensitivity studies on the ground-based data because the additional measurements of NH_3 and NO_z (not available on the aircraft) provided better constraints on the levels of TA and TN.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 The comparison between measured and modelled inorganic composition of PM₁

3.1.1 Harrow data

AURAMS-modelled PM₁ sulphate and AMS sulphate measurements in Harrow (Fig. 2a) were reasonably well correlated (Table 1) throughout the campaign with three major exceptions that occurred during 19–20 June, 24–29 June and 10–11 July periods. These periods were characterized by air masses originating from the high emission region of the Ohio Valley, which is a well-established contributor to SO₂, NO_x and PM₁ pollution in Southwestern Ontario (Canada – United States Transboundary Particulate Matter Science Assessment, 2004). During these periods of high PM₁, AURAMS was biased high for sulphate compared to measured data. In some instances the model predicted 2–3 times higher mass loadings. However, some of this bias may actually be attributable to the falloff in transmission efficiency of the AMS between 0.4–1.0 μm. Particle size distributions measured by the AMS at Harrow suggest that particularly at high mass loadings significant sulphate mass may be present above the nominal size cut. Overall at Harrow, AURAMS had a low mean bias for $p\text{SO}_4^{2-}$, but a substantial RMSE. Modelled $p\text{NH}_4^+$ were biased low (Fig. 2b), also with a large RMSE = 1.50 μg m⁻³ (Table 1).

AURAMS tended to predict more extreme values of PM₁ nitrate than those measured by the AMS (Fig. 2c). While AURAMS frequently predicted zero, AMS observations were rarely below the detection limit. On the other hand, AURAMS tended to predict much higher mass loadings when the observations indicated significant levels of nitrate. As a result, while the mean bias appears small (MB = 0.08 μg m⁻³), RMSE is very large (1.27 μg m⁻³). Because the model predicted peaks in $p\text{NO}_3^-$ at approximately the same times of day as the observations, the correlation coefficient ($R = 0.39$) is higher than it might otherwise be.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1.2 Aircraft data

Compared to the evaluation of the model predictions at the ground site, the evaluation of aircraft data poses the additional challenges of (i) the simultaneous evaluation of predictions in both time and space, and (ii) the precise nature of the relatively fast aircraft measurements vs. 2.5 km grid resolution constraining the evaluation. Both modelled sulphate (Fig. 3a) and ammonium (Fig. 3b) were biased high relative to observations for most of the flights and the bias was more significant for time periods characterized by high PM_1 levels. The positive bias in pSO_4^{2-} was much larger than that for pNH_4^+ , resulting in predictions of aerosol that were more acidic than indicated by the observations. The model seldom predicted significant nitrate mass loadings ($>1 \mu g m^{-3}$) aloft, but when it did, the predictions were also biased high relative to the AMS observations (Fig. 3c). Thus, a very low correlation coefficient ($R = 0.12$) was calculated from the correlation plot between modelled and measured aircraft pNO_3^- data.

3.1.3 The influence of plume effects on the model/measurement agreement

While the data in Table 1 and Figs. 2 and 3 demonstrate that AURAMS did accurately predict the overall magnitude of regional nitrate mass loadings during the BAQS-Met campaign, individual observations both at a specific ground site and in the lowest 3000 m aloft were not well-represented by the model. Because point sources (e.g. power plants) are expected to be the dominant source of NO_x and SO_2 in the region, one explanation of the large RMSE for pNO_3^- is inaccurate predictions of when plumes are influencing the measurements. The erroneous placement of a plume in space or time due to inaccurate representations of advection or diffusion could lead to poor correlations on short timescales. In addition, the spatial resolution of AURAMS is currently limited to a $2.5 km \times 2.5 km$ grid box which is larger than the spatial extent of most plumes. To examine the significance of these “hit-and-miss” instances on the agreement between measurements and predictions of PM_1 composition, we examine the data through a series of histograms. The histogram shown in Fig. 4 illustrates

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the range of measurements and predictions of ρNO_3^- . All modelled values lower than $0.01 \mu\text{g m}^{-3}$ were accumulated into the lowest bin in the figure. As mentioned previously, all measured values lower than the detection limit were assigned a value equal to the detection limit (aircraft = $0.014 \mu\text{g m}^{-3}$; Harrow = $0.006 \mu\text{g m}^{-3}$). The great majority of aircraft measurements were made within the boundary layer during daytime (Hayden et al., 2010). Most of the nitrate mass loadings measured by the aircraft AMS were between 0.1 and $1 \mu\text{g m}^{-3}$. In comparison to aircraft measurements, the mass loadings of ρNO_3^- at the Harrow site were slightly higher, as evident by greater frequency of points in 1 to $4 \mu\text{g m}^{-3}$ range.

While the measured average mass loading of ρNO_3^- at Harrow was higher than that aloft (Table 1), both datasets are well-represented by a log-normal distribution. In contrast, the model predicts a much wider range of particulate nitrate, with a significant fraction of the data predicted to be below the detection limit of the AMS. However, according to Fig. 4 only 1% of the observations at Harrow and 8% of the aircraft observations were in fact below the detection limit. If hit-and-miss errors were the significant driver of model bias, we would expect the overall distribution to be well-represented by the model, even if the point-to-point comparisons were not good. Because the distributions of nitrate mass loadings predicted by the model do not match the distributions observed from the aircraft or on the ground, we can conclude that inaccurate plume placement is not the main reason for disagreement between the model and the measurements. We now examine alternative explanations for the disagreement in the remainder of the paper.

3.2 The applicability of thermodynamic equilibrium at Harrow

In order to determine whether the ammonium-sulphate-nitrate system was in thermodynamic equilibrium, Harrow observation data were re-partitioned using ISORROPIA. At Harrow, TA values were calculated using the sum of $\text{NH}_{3(\text{g})}$ measurements by QC-TILDAS and ρNH_4^+ measurements by AMS. TS values were represented by AMS

ρSO_4^{2-} . AMS ρSO_4^{2-} is a good proxy for TS for three reasons: (i) the vapour pressure of $\text{H}_2\text{SO}_{4(\text{g})}$ is low and it is expected to be exclusively in condensed phase in the atmosphere, (ii) AMS ρSO_4^{2-} is actually the sum of particulate bisulphate (HSO_4^-), sulphate (SO_4^{2-}) and sulphuric acid (H_2SO_4), and (iii) most of the ρSO_4^{2-} is expected to be at particle sizes less than $1\ \mu\text{m}$ diameter. The lack of specific $\text{HNO}_{3(\text{g})}$ measurements at the site precluded an exact calculation of TN from direct observations. We estimated TN as a fraction of the NO_z measurements from the chemiluminescence analyzer. We re-partitioned the data assuming (i) a low TN case, $\text{TN} = \text{AMS}\ \rho\text{NO}_3^-$, (ii) a medium TN case, $\text{TN} = 0.5\cdot\text{NO}_z$, and (iii) a high TN case, $\text{TN} = \text{NO}_z$. The medium TN case provided the most reasonable TN concentrations because the re-partitioned ρNO_3^- agreed most closely with the ρNO_3^- observed by the AMS. Therefore, we calculated $\text{HNO}_{3(\text{g})}$ by subtracting AMS ρNO_3^- from TN obtained by $\text{TN} = 0.5\cdot\text{NO}_z$. While our approximation for TN is likely not consistently valid throughout the campaign, it provides a reasonable range of ρNO_3^- values and there is no justification for adopting a variable fraction. This analysis could not be carried out for the aircraft data because of the lack of gas phase NH_3 and NO_z measurements.

Observed Harrow TA, TS, TN, RH and T values were used as inputs to ISORROPIA and the model was allowed to re-partition the data between the gas and the particle phase assuming thermodynamic equilibrium conditions. The re-partitioned data was in very good agreement with the original measurements suggesting that the ammonium-sulphate-nitrate system was in thermodynamic equilibrium in Harrow. According to ISORROPIA, TS was almost exclusively in the form of SO_4^{2-} as opposed to HSO_4^- . In terms of neutralization of the ammonium-sulphate-nitrate system this means that PM_1 sulphate was completely neutralized and that thermodynamic formation of particulate ammonium nitrate is possible under such conditions. Re-partitioned TA is in good agreement with the original observations suggesting that most of the ρNH_4^+ is contained in particles with diameter $<1\ \mu\text{m}$, and that the observed species are sufficient to describe the system. The scatter plot of re-partitioned ρNH_4^+ versus AMS ρNH_4^+ has a

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



slope of 1.072 and $R = 0.982$ (Fig. 5a). The close agreement between observed and re-partitioned $p\text{NH}_4^+$ also suggests that the estimated values for TN were not totally unreasonable.

The correlation plot (Fig. 5b) between the re-partitioned and original $p\text{NO}_3^-$ has a slope of $m = 1.413$ and $R = 0.74$, suggesting that ISORROPIA predicted, on average, more $p\text{NO}_3^-$ than measured. According to the average percent difference calculation, ISORROPIA predicted, on average, about 1% more $p\text{NO}_3^-$. Most likely, $\text{TN} = 0.5 \cdot \text{NO}_z$ is, at least in some cases, an overestimate which led to the overestimate in $p\text{NO}_3^-$ by the model in conditions of high TA. This is also consistent with a slope of greater than one for the $p\text{NH}_4^+$ comparison (Fig. 5a). Predictions of high nitrate can be also attributed to increased $p\text{NO}_3^-$ formation due to dissolution of $\text{HNO}_{3(\text{g})}$ in water (Fig. 5b). Under high RH conditions, the amount of $p\text{NO}_3^-$ that will form from dissolution in water is directly proportional to TN. For this reason, the re-partitioned $p\text{NO}_3^-$ points that are furthest away from the 1:1 line in the positive direction correspond to high RH regimes as shown by colour scale. The fact that the re-partitioned TA data agrees well with the measured data and that re-partitioned TN data agrees reasonably well (considering the constraints in TN measurements) suggests that gas-particle partitioning can be described by thermodynamic equilibrium expressions at Harrow during BAQS-Met. Thermodynamic equilibrium was also observed in the field by Zhang et al. (2002a). In addition, Meng and Seinfeld (1996) showed that the submicron (PM_{10}) ammonium-sulphate-nitrate system will reach thermodynamic equilibrium on the order of minutes. Because the measurement site is not directly located next to a source that would strongly perturb the levels of $\text{NH}_{3(\text{g})}$ or acidic aerosols, and an adequate representation of $p\text{NO}_3^-$ is achieved by the re-partitioning using ISORROPIA, it is expected that the HETV algorithms in AURAMS should appropriately describe the gas-particle partitioning.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.3 Free ammonia (FA) chemical maps

In Sect. 3.2, we demonstrated that thermodynamic equilibrium expressions are appropriate to model the gas-particle partitioning of TA and TN. The accurate prediction of ρNO_3^- will depend not only on accurate predictions of TN, but also on RH, T , and the availability of $\text{NH}_{3(\text{g})}$, as a result of the equilibrium expressions in Eqs. (1) and (2). In order to assess the variables that control the ability of AURAMS to accurately predict PM_{10} nitrate, we examined the parameters governing the mass loading of particulate nitrate by using free ammonia chemical maps. Free ammonia is defined as $\text{FA} = \text{TA} - 2 \cdot \text{TS}$ (Seinfeld, 1996), where the factor of 2 accounts for the equivalents of sulphate. If FA of the ammonium-sulphate-nitrate system is positive, the system contains sufficient ammonia to form ammonium nitrate (particulate nitrate), depending on the RH and T conditions. If FA of the system is negative, the system does not contain sufficient ammonia and the formation of ammonium nitrate is thermodynamically unfavourable. At sufficiently high RH, $\text{HNO}_{3(\text{g})}$ may dissolve to form aqueous nitrate, even in the absence of high FA. In order to investigate the combined effects of absolute concentrations and meteorological variables on the formation of particulate nitrate, as driven by algorithms of the model, we generated a series of FA chemical maps, an example of which is shown in Fig. 6.

The map in Fig. 6 was created by running ISORROPIA to predict ρNO_3^- mass loadings over a range of temperatures and FA values at a constant relative humidity (0.65) and TN value ($100 \text{ nmol m}^{-3} \equiv 6.2 \mu\text{g m}^{-3}$). A range of FA values (-275 to 275 nmol m^{-3}) were generated by varying TS values while keeping TA constant at an atmospherically relevant level (275 nmol m^{-3}). The map space in Fig. 6 is coloured by the amount of nitrate predicted to be in particle phase for each combination of FA and T . By generating these maps, we can identify regimes in which the formation of particulate nitrate is favourable for a given pair of T and FA values. It suggests that in order to form $>2 \mu\text{g m}^{-3}$ of particulate nitrate (given $6.2 \mu\text{g m}^{-3}$ of TN) at $\text{RH} = 0.65$, temperatures must be lower than 295 K, and if they are above 270 K, FA must also

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be positive. The S-shaped transition region between where formation of nitrate is not favourable (light) and where the formation is favourable (dark) is where the predictions of nitrate are most sensitive to model input parameters. The map indicates that at low temperatures the role of FA in ρNO_3^- formation becomes less important. It should be noted that the position of the transition region depends on RH as the formation of particulate nitrate becomes more pronounced at high water concentrations in the atmosphere. At high RH, the amount of water in the atmosphere is sufficiently high for the dissolution of $\text{HNO}_{3(g)}$ to form $\rho\text{NO}_3^-(\text{aq})$ and become more important than chemically driven formation of nitrate through the reaction with $\text{NH}_{3(g)}$.

Figure 6 shows that temperature strongly impacts both predictions of particulate nitrate and their sensitivity to FA. Because the absolute amount of TN, the temperature and the relative humidity impact the predictions of particulate nitrate, we have generated a series of panels (Fig. 7) representing different scenarios relevant to the measurements during BAQS-Met. These partitioning maps were generated at three RH regimes (low RH = 0.40, medium RH = 0.65 and high RH = 0.90) and three TN regimes (low TN = 20 nmol m^{-3} , medium TN = 50 nmol m^{-3} and high TN = 100 nmol m^{-3}). TA was held constant at 275 nmol m^{-3} and TS was varied between 0 and 275 nmol m^{-3} in order to generate a range of FA values that were observed during the field campaign (-275 to 275 nmol m^{-3}). RH increases from left to right (a to c) and TN increases from top to bottom (a to g).

The AURAMS model predictions (red circles) and observations from Harrow (yellow triangles) are overlaid on the panels. AMS and AURAMS data were binned according to measured and modelled RH respectively in order to match three RH regimes (low RH \equiv 0–0.50, medium RH \equiv 0.50–0.80, and high RH \equiv 0.80–1.0) to account for the effect of RH on the formation of nitrate. For example, all measured and modelled data corresponding to the low RH bin are plotted on maps generated at 0.40 RH, because there is little variation in the characteristics of the maps at all RH < 0.50 in the temperature range observed during BAQS-Met. Data were not binned according to TN because the NO_2 measurements did not offer a strong constraint on TN, so the same

data appear on e.g. panels a, d, and g. At $RH = 0.65$, the average observed RH at Harrow, there is little variation in the position of the S-shaped transition between the three TN regimes (Fig. 7b, e, and h). At high RH there is sufficient water in the atmosphere such that the dissolution of $HNO_{3(g)}$ in water completely outcompetes chemical formation of ammonium nitrate. This is evident from the vertical shift of the S-shaped transition regime from Fig. 7d, e, and f.

As indicated by the colour scale, the absolute amount of particulate nitrate will significantly increase going from low to high TN concentrations. For an intermediate to high level of FA ($50\text{--}100\text{ nmol m}^{-3}$), the higher the level of TN in the atmosphere, the more of it will partition into particle phase, both in relative and absolute terms. At high TN levels, FA becomes the limiting factor in formation of particulate nitrate according to Eqs. (1) and (2). On these maps, both modelled and observed FA values are located close to the transition between the light region where the majority of TN is $HNO_{3(g)}$ and dark region where the majority of TN is ρNO_3^- . Most of the observations (yellow triangles) are located in the positive FA region of the map where significant ρNO_3^- formation is expected if temperatures are low. In contrast, the model points (red circles) are located across a wider range of FA and temperature values, producing a wider range of ρNO_3^- predictions. There is a pronounced bias in modelled FA in the low RH regime (Fig. 7a, d, and g), but the chemical maps indicate that this causes little bias in ρNO_3^- because its formation is not favoured under these warm, dry conditions, even at higher FA. The difference in the modelled and measured FA parameters on these partitioning maps can explain much of the discrepancy between modelled and measured nitrate mass loadings observed during BAQS-Met, especially at intermediate RH and assuming intermediate values of TN (Fig. 7e).

3.4 Measured and modelled FA

The bias in AURAMS FA values results either from overpredictions of TS, or underpredictions of TA. In Ellis et al. (2010a), it is shown that the TA was consistently

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



underpredicted at Harrow. To compare the range of measured and modelled FA values, we generated histograms of FA at Harrow (Fig. 8a) and from the aircraft data (Fig. 8b). Compared to FA values calculated from measurements, FA values predicted by AURAMS were distributed across a much wider range both at the ground and aloft.

5 In addition, a significant number of modelled FA values were negative while all of the ground-based and > 95% of the aircraft measurements of FA values were positive. One of the reasons why measured FA values were lower aloft is because we had no gas phase NH_3 instrument onboard the aircraft. In the absence of $\text{NH}_{3(\text{g})}$ data we calculated TA by $\text{TA} \equiv \rho\text{NH}_4^+$, so measurements of FA from the aircraft are an underestimate, which further exacerbates the differences with respect to the model, which is already
10 biased low for FA. We expect that the contribution of $\text{NH}_{3(\text{g})}$ to TA (and hence FA) is less significant for the aircraft data than for Harrow because ammonia is emitted at the ground from agricultural and traffic sources; thus, we expect TA, and hence FA, values to be higher at the ground site. According to Fig. 7, the impact on ρNO_3^- of a negative bias in FA ranges from negligible (under warm, dry conditions and low TN loadings) to
15 significant (under cool, moist conditions with moderate to high TN).

3.5 Measured and modelled meteorological parameters

Modelled values of relative humidity compared with measurements at Harrow were significantly lower across the entire spectrum of RH values (Fig. 9a). Only in a few
20 instances was the modelled RH higher than the measurements, and this generally occurred in higher RH regimes ($\text{RH} = 0.70\text{--}0.90$). The difference between modelled and measured RH for the most part ranged between -0.20 and -0.30 . The comparison between temperature data at Harrow (Fig. 9b) showed that the model was biased high at high temperatures and biased low at low temperatures. The difference between modelled and measured temperatures rarely exceeded 5 K. The chemical maps
25 in Fig. 7 show that the predictions of ρNO_3^- are much more sensitive to temperature between 280 K and 290 K than between 300 K and 310 K. The low bias at low tempera-

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tures likely contributes to the overpredictions of $p\text{NO}_3^-$ seen on several days at Harrow (Fig. 2), which occurred between 02:00 and 08:00 when the temperature was lowest.

Compared to measurements of relative humidity aboard the aircraft, modelled RH (Fig. 10a) values were significantly lower, much like at the ground site. The model was seldom biased high relative to the observations and the high bias was most pronounced at lower RH (< 0.60). The comparison between measured and modelled temperature data (Fig. 10b) showed that the model was biased low (by 1–5 K) over the entire study period. According to Fig. 7, the impact on $p\text{NO}_3^-$ of biases in RH and T depends on the levels of TN and FA and are most important in the transition region.

Sensitivity studies with meteorological variables

In order to further investigate the significance of meteorological parameters on AURAMS's bias in $p\text{NO}_3^-$ we performed a series of sensitivity tests in which we re-partitioned modelled PM_1 data using ISORROPIA offline with (i) modelled meteorological data (case I), (ii) measured RH only (case II), (iii) measured T only (case III), and (iv) measured T and RH (case IV), with the results compiled in Table 2. These sensitivity tests were carried out both for the Harrow data and the aircraft data. Because of missing temperature and relative humidity measurements (more significant at Harrow than on aircraft), the “AURAMS” column in Table 2 refers to model output only from periods when meteorological measurements were available for use in the other cases. For case I, any differences compared to the original AURAMS data should only reflect differences in the partitioning calculated by the algorithms used in HETV and those in ISORROPIA v2.1, which was used in the offline re-partitioning analysis.

In terms of the chemical maps (Fig. 7), vertical shifts in the location of data brought about by changes in T were more significant when the FA of the system was positive because regions of favourable and non-favourable nitrate formation were as little as 2–3 K apart for T values observed during BAQS-Met (282–308 K). However, RH affects the entire chemical map and caused a large upward shift in the location of S-shaped

The effect of meteorological and chemical factors

M. Z. Markovic et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

transition line – increasing RH from 0.65 to 0.90 is the equivalent of moving a point from Fig. 7e to Fig. 7f.

As shown in Table 2, re-partitioning of aircraft data with measured RH and T had little effect on predictions of ρNO_3^- . MB and RMSE values between measured and modelled ρNO_3^- showed very little variation between the cases. Re-partitioning with measured T and RH (case IV) led to a slight improvement in correlation ($R = 0.18$, compared to $R = 0.12$) between modelled and measured nitrate. The slight improvement in correlation was the net result of the opposing effects of measured RH (case II, $R = 0.20$) and T (case III, $R = 0.06$) on the modelled formation of ρNO_3^- . The limited effect of the substitution of measured RH and T on the re-partitioning of the aircraft data is the result of the low FA values predicted by AURAMS during the campaign. As evident from Fig. 8, approximately half of all modelled FA values were negative. In terms of the chemical maps, this means that modelled FA values were located in the upper left part of the map far from the transition line where RH and T have very little effect on formation of nitrate.

Compared to the aircraft data, re-partitioning of Harrow data with measured meteorological parameters had a more significant impact on predicted ρNO_3^- (Table 2). Re-partitioning with measured RH led to an increase in predicted nitrate ($\text{MB} = 0.12 \mu\text{g m}^{-3}$) and a better correlation with measured ρNO_3^- data. Re-partitioning with measured T values, led to an overall decrease in predicted nitrate ($\text{MB} = -0.12 \mu\text{g m}^{-3}$) and in the correlation with measurements. The combined effect (case IV) of re-partitioning was the same as for the aircraft data which was an overall increase in predicted nitrate ($\text{MB} = 0.08 \mu\text{g m}^{-3}$) and slight improvement in the correlation ($R = 0.30$). Even though the overall effect was an increase in predicted nitrate, for the most part, the increase occurred in instances when the FA of the system is positive and significant levels of nitrate already existed. This exacerbated the instances of ρNO_3^- overprediction and led to an overall increase in RMSE from 1.09 to $1.22 \mu\text{g m}^{-3}$.

These sensitivity studies demonstrate that the bias in modelled RH and T contributes to some of the errors in ρNO_3^- predicted by the AURAMS model during BAQS-Met.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, constraining RH and T without also removing the bias in FA prevents significant improvements in the predictions. This simple offline re-partitioning does not fully account for the impact of inaccuracies in the meteorological parameters driving AURAMS, because gas-particle partitioning of TN (and TA) determines their lifetimes, which feeds back to the predictions of ρNO_3^- . For example, a low bias in RH is likely to produce a low bias in $\rho\text{NO}_3^-/\text{HNO}_{3(\text{g})}$, which will lead to a low bias in TN (because $\text{HNO}_{3(\text{g})}$ deposits more quickly), further exacerbating the low bias in ρNO_3^- . From the chemical maps in Fig. 7, it is clear that in a sulphate-rich summertime atmosphere, the model output is most sensitive to temperature biases when $T < 290$ K.

4 Conclusions

During the BAQS-Met campaign, A Unified Regional Air-quality Modeling System (AURAMS) exhibited large biases in the prediction of PM_{10} nitrate. The agreement between model predictions and observations at the ground site ($M_{\text{obs}} = 0.50 \mu\text{g m}^{-3}$; $M_{\text{mod}} = 0.58 \mu\text{g m}^{-3}$; $\text{RMSE} = 1.27 \mu\text{g m}^{-3}$; and $R = 0.39$), was better than aloft ($M_{\text{obs}} = 0.32 \mu\text{g m}^{-3}$; $M_{\text{mod}} = 0.09 \mu\text{g m}^{-3}$; $\text{RMSE} = 0.48 \mu\text{g m}^{-3}$; $R = 0.12$). In order to better understand the factors that limited AURAMS' ability to accurately predict ρNO_3^- during BAQS-Met, we generated a set of FA chemical maps that illustrated the relationship between the absolute amounts of TS, TA, TN, and meteorological variables (RH, T) in the formation of ρNO_3^- . From the histogram in Fig. 8b, we determined that for the aircraft data, AURAMS was biased low for ρNO_3^- because it was making predictions in the chemical map space where ρNO_3^- formation is not allowed due to negative FA. At Harrow, ρNO_3^- measurements ranged between 0.1 and $3 \mu\text{g m}^{-3}$, whereas the model predicted a wide range of values up to $8 \mu\text{g m}^{-3}$, with the most frequent prediction being zero. The underpredictions appear to be the result of a low bias in FA, whereas the overpredictions occur when FA is positive, but the temperature predictions are too low by several K. AURAMS was biased low for FA during the campaign because it was biased high for TS and/or biased low for TA. From the chemical FA maps, we

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concluded that at low temperatures and high relative humidity, the role of FA in $p\text{NO}_3^-$ formation becomes less important. Sensitivity tests with ISORROPIA using measured RH and T showed that constraining meteorological inputs with measurements can only offer modest improvements in the predictions of $p\text{NO}_3^-$ if the FA values in the model remain biased. In this work, we developed a novel framework to examine the variables that affect the model's ability to accurately predict fine particulate nitrate. The results of our analysis are relevant to all regional air quality models and as such can help improve approaches to modelling regional PM pollution.

Acknowledgements. The authors would like to thank Ontario Ministry of Environment, Environment Canada, and Agriculture and Agri-Food Canada for support during BAQS-Met field campaign. The authors would also like to thank Jeffrey A. Geddes and Ilan Levy for their assistance with data analysis and G. J. Evans, C.-H. Jeong, and M. L. McGuire for providing us with meteorological data measurements from the Harrow site. We would also like to acknowledge the support of the pilots, technical crew and scientists from the National Research Council of Canada. M. Z. Markovic is grateful to Environment Canada for funding this analysis through Research Affiliate Position (RAP) programme.

References

- Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A. M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A generalised method for the extraction of chemically resolved mass spectra from aerodyne aerosol mass spectrometer data, *J. Aerosol Sci.*, 35, 909–922, 2004.
- Bahreini, R., Dunlea, E. J., Matthew, B. M., Simons, C., Docherty, K. S., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., and Middlebrook, A. M.: Design and operation of a pressure-controlled inlet for airborne sampling with an aerodynamic aerosol lens, *Aerosol Sci. Technol.*, 42, 465–471, doi:10.1080/02786820802178514, 2008.
- Burnett, R. T., Brook, J., Dann, T., Delocla, C., Phillips, O., Cakmak, S., Vincent, R., Goldberg, M. S., and Krewski, D.: Association between particulate- and gas-phase components of urban air pollution and daily mortality in eight Canadian cities, *Inhalation Toxicology*, 12, 15–39, 2000.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factorsM. Z. Markovic et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$ at 298.15 K, *J. Phys. Chem. A*, 102, 2155–2171, 1998a.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{H}_2\text{O}$ at tropospheric temperatures, *J. Phys. Chem. A*, 102, 2137–2154, 1998b.
- 5 Cote, J., Gravel, S., Methot, A., Patoine, A., Roch, M., and Staniforth, A.: The operational CMC-MRB Global Environmental Multiscale (GEM) model. Part I: Design considerations and formulation, *Mon. Weather Rev.*, 126, 1373–1395, 1998.
- Cross, E. S., Onasch, T. B., Canagaratna, M., Jayne, J. T., Kimmel, J., Yu, X.-Y., Alexander, M. L., Worsnop, D. R., and Davidovits, P.: Single particle characterization using a light scattering module coupled to a time-of-flight aerosol mass spectrometer, *Atmos. Chem. Phys.*, 9, 7769–7793, doi:10.5194/acp-9-7769-2009, 2009.
- 10 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.
- Dockery, D. W.: Epidemiologic evidence of cardiovascular effects of particulate air pollution, *Environ. Health Persp.*, 109, 483–486, 2001.
- Dominici, F., Peng, R. D., Bell, M. L., Pham, L., McDermott, A., Zeger, S. L., and Samet, J. M.: Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases, *Jama-Journal of the American Medical Association*, 295, 1127–1134, 2006.
- 20 Drennick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) – Instrument description and first field deployment, *Aerosol Sci. Technol.*, 39, 637–658, doi:10.1080/02786820500182040, 2005.
- 25 Dunlea, E. J., DeCarlo, P. F., Aiken, A. C., Kimmel, J. R., Peltier, R. E., Weber, R. J., Tomlinson, J., Collins, D. R., Shinozuka, Y., McNaughton, C. S., Howell, S. G., Clarke, A. D., Emmons, L. K., Apel, E. C., Pfister, G. G., van Donkelaar, A., Martin, R. V., Millet, D. B., Heald, C. L., and Jimenez, J. L.: Evolution of Asian aerosols during transpacific transport in INTEX-B, *Atmos. Chem. Phys.*, 9, 7257–7287, doi:10.5194/acp-9-7257-2009, 2009.
- 30 Ellis, R. A., Murphy, J. G., Markovic, M. Z., VandenBoer, T. C., Makar, P. A., Brook, J., and Mihele, C.: The influence of gas-particle partitioning and surface-atmosphere exchange on ammonia during BAQS-Met, *Atmos. Chem. Phys. Discuss.*, 10, 21895–21929,

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



doi:10.5194/acpd-10-21895-2010, 2010a.

Ellis, R. A., Murphy, J. G., Pattey, E., van Haarlem, R., O'Brien, J. M., and Herndon, S. C.: Characterizing a Quantum Cascade Tunable Infrared Laser Differential Absorption Spectrometer (QC-TILDAS) for measurements of atmospheric ammonia, *Atmos. Meas. Tech.*, 3, 397–406, doi:10.5194/amt-3-397-2010, 2010b.

Finlayson-Pitts, B. J. and Pitts Jr., J. N.: *Chemistry of the Upper and Lower Atmosphere*, Second ed., Academic Press, New York, 2000.

Gauderman, W. J., Avol, E., Gilliland, F., Vora, H., Thomas, D., Berhane, K., McConnell, R., Kuenzli, N., Lurmann, F., Rappaport, E., Margolis, H., Bates, D., and Peters, J.: The effect of air pollution on lung development from 10 to 18 years of age, *New England Journal of Medicine*, 351, 1057–1067, 2004.

Gong, W., Zhang, J., Moran, M. D., Makar, P. A., Gong, S. L., Stroud, C., Bouchet, V. S., Cousineau, S., Menard, S., Samaali, M., Sassi, M., Pabla, B., Leaitch, R., Macdonald, A. M., Anlauf, K., Hayden, K., Toom-Saunry, D., Leithead, A., and Strapp, J. W.: Modelling regional aerosols: Impact of cloud processing on gases and particles over eastern North America and in its outflow during ICARTT 2004, in: *Air Pollution Modeling and Its Application Xix*, edited by: Borrego, C. and Miranda, A. I., *Nato Science for Peace and Security Series C – Environmental Security*, 539–547, 2008.

Gong, W. M., Dastoor, A. P., Bouchet, V. S., Gong, S. L., Makar, P. A., Moran, M. D., Pabla, B., Menard, S., Crevier, L. P., Cousineau, S., and Venkatesh, S.: Cloud processing of gases and aerosols in a regional air quality model (AURAMS), *Atmos. Res.*, 82, 248–275, doi:10.1016/j.atmosres.2005.10.012, 2006.

Hayden, K. L., Macdonald, A. M., Gong, W., Toom-Saunry, D., Anlauf, K. G., Leithead, A., Li, S. M., Leaitch, W. R., and Noone, K.: Cloud processing of nitrate, *J. Geophys. Res.-Atmos.*, 113, D18201, doi:10.1029/2007JD009732, 2008.

Hayden, K. L., Sills, D., Li, S. M., Brook, J. R., Liu, P., Anlauf, K. G., Makar, P. A., O'Brien, J. M., Strawbridge, K., Firanski, B., and McLaren, R.: The impact of lake breezes on trace gases and particles during the Border Air Quality and Meteorology Study (BAQS-Met), to be submitted to *Atmos. Chem. Phys. Discuss.*, 2010.

Jacob, D. J.: *Introduction to Atmospheric Chemistry*, Princeton University Press, Princeton, NJ, 1999.

Jacobson, M. Z., Tabazadeh, A., and Turco, R. P.: Simulating equilibrium within aerosols and nonequilibrium between gases and aerosols, *J. Geophys. Res.-Atmos.*, 101, 9079–9091,

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sci. Technol., 41, 380–386, 2007.

Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *Jama-Journal of the American Medical Association*, 287, 1132–1141, 2002.

5 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, John Wiley & Sons, New York, 1996.

Slowik, J. G., Brook, J., Chang, R. Y.-W., Evans, G. J., Hayden, K. L., Jeong, C.-H., Li, S.-M., Liggio, J., Liu, P. S. K., McGuire, M., Mihele, C., Sjostedt, S., Vlasenko, A., and Abbatt, J. P. D.: Photochemical processing of organic aerosol at nearby continental sites: Contrast between urban plumes and regional aerosol, in preparation, 2010.

10 Stelson, A. W. and Seinfeld, J. H.: Thermodynamic Prediction of the Water Activity, NH_4NO_3 Dissociation-Constant, Density and Refractive-Index for the NH_4NO_3 - $(\text{NH}_4)_2\text{SO}_4$ - H_2O System at 25-Degrees-C, *Atmos. Environ.*, 16, 2507–2514, 1982.

Wexler, A. S. and Seinfeld, J. H.: The Distribution of Ammonium-Salts among a Size and Composition Dispersed Aerosol, *Atmos. Environ. Part a*, 24, 1231–1246, 1990.

15 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J. Geophys. Res.-Atmos.*, 107, 4207, doi:10.1029/2001jd000451, 2002.

20 Yu, S. C., Dennis, R., Roselle, S., Nenes, A., Walker, J., Eder, B., Schere, K., Swall, J., and Rorbarge, W.: An assessment of the ability of three-dimensional air quality models with current thermodynamic equilibrium models to predict aerosol NO_3 , *J. Geophys. Res.-Atmos.*, 110, D07s13, doi:10.1029/2004jd004718, 2005.

Zhang, J., Chameides, W. L., Weber, R., Cass, G., Orsini, D., Edgerton, E., Jongejan, P., and Slanina, J.: An evaluation of the thermodynamic equilibrium assumption for fine particulate composition: Nitrate and ammonium during the 1999 Atlanta Supersite Experiment, *J. Geophys. Res.-Atmos.*, 108(D7) 8414, doi:10.1029/2001JD001592, 2002a.

25 Zhang, L. M., Moran, M. D., Makar, P. A., Brook, J. R., and Gong, S. L.: Modelling gaseous dry deposition in AURAMS: a unified regional air-quality modelling system, *Atmos. Environ.*, 36, 537–560, 2002b.

30 Zhang, Y., Seigneur, C., Seinfeld, J. H., Jacobson, M., Clegg, S. L., and Binkowski, F. S.: A comparative review of inorganic aerosol thermodynamic equilibrium modules: similarities, differences, and their likely causes, *Atmos. Environ.*, 34, 117–137, 2000.

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Table 1. Statistical evaluation of modelled and measured inorganic PM₁ mass at the ground and aloft during BAQS-Met. M_{mod} = modelled mean, M_{obs} = observed mean, MB = mean bias, RMSE = root mean square error, all in $\mu\text{g m}^{-3}$ and R = correlation coefficient.

| | AIRCRAFT | | | HARROW | | |
|------------------|---------------------|------------------------|---------------------|---------------------|------------------------|---------------------|
| | ρNH_4^+ | ρSO_4^{2-} | ρNO_3^- | ρNH_4^+ | ρSO_4^{2-} | ρNO_3^- |
| M _{mod} | 1.58 | 6.71 | 0.09 | 1.40 | 3.90 | 0.58 |
| M _{obs} | 1.34 | 2.94 | 0.32 | 1.60 | 3.82 | 0.50 |
| MB | 0.24 | 3.77 | -0.23 | -0.20 | 0.08 | 0.08 |
| RMSE | 1.24 | 6.82 | 0.48 | 1.50 | 4.64 | 1.27 |
| R | 0.65 | 0.63 | 0.12 | 0.44 | 0.46 | 0.39 |

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The effect of meteorological and chemical factors

M. Z. Markovic et al.

Table 2. Comparison of measured ρNO_3^- mass loading with offline model runs using measured temperature and/or relative humidity at the ground and aloft during BAQS-Met. The column showing AURAMS data in this table is slightly different from Table 1, because it has been filtered to only include points for which measured T and RH are available, which were needed to generate cases I–IV. Re-partitioning of AURAMS data was performed: (i) with original (AURAMS) met data (case I), (ii) with measured RH only (case II), (iii) with measured T only (case III), and (iv) with measured RH and T (case IV). For each case, the model is compared to the AMS measurements. MB = mean bias, RMSE = root mean square error, all in $\mu\text{g m}^{-3}$ and R = correlation coefficient.

| | AIRCRAFT | | | | | HARROW | | | | |
|------|----------|--------|---------|----------|---------|--------|--------|---------|----------|---------|
| | AURAMS | case I | case II | case III | case IV | AURAMS | case I | case II | case III | case IV |
| MB | −0.22 | −0.25 | −0.22 | −0.28 | −0.25 | 0.01 | −0.05 | 0.12 | −0.12 | 0.08 |
| RMSE | 0.47 | 0.48 | 0.48 | 0.47 | 0.46 | 1.09 | 1.00 | 1.28 | 0.91 | 1.22 |
| R | 0.12 | 0.09 | 0.20 | 0.06 | 0.18 | 0.29 | 0.28 | 0.32 | 0.25 | 0.30 |

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 1. Map of the BAQS-Met region showing the location of Harrow and the flight tracks of the Twin Otter research aircraft coloured by altitude.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 2. Time series of measured and modelled (a) $p\text{SO}_4^{2-}$, (b) $p\text{NH}_4^+$ and (c) $p\text{NO}_3^-$ at the Harrow, ON site during BAQS-Met.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 3. Vertical distributions of measured and modelled (a) $p\text{SO}_4^{2-}$, (b) $p\text{NH}_4^+$ and (c) $p\text{NO}_3^-$ from the Twin Otter aircraft over all flights during BAQS-Met.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 4. Normalized histograms of the measured and modelled ρNO_3^- mass loadings **(a)** from the Twin Otter aircraft **(b)** at Harrow.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 5. Scatter plots between re-partitioned and measured **(a)** $p\text{NH}_4^+$ and **(b)** $p\text{NO}_3^-$ at Harrow. The black line is 1:1, and the red line is the least squares fit.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. FA chemical map illustrating the formation of $p\text{NO}_3^-$ at $\text{RH} = 0.65$ for $T = 255\text{--}310\text{ K}$. TA was held constant at 275 nmol m^{-3} and TS was varied between 0 and 275 nmol m^{-3} in order to generate a range of FA values that were observed during the field campaign (-275 to 275 nmol m^{-3}). TN was 100 nmol m^{-3} .

The effect of meteorological and chemical factors

M. Z. Markovic et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 7. A matrix of FA chemical maps depicting the correlation between FA, meteorological data and the concentration of nitrate in the particle phase. These partitioning maps were generated at three RH regimes (low RH = 0.40 **(a,d,g)**, medium RH = 0.65 **(b,e,h)** and high RH = 0.90 **(c,f,i)**) and three TN regimes (low TN = 20 nmol m^{-3} **(a,b,c)**, medium TN = 50 nmol m^{-3} **(d,e,f)** and high TN = 100 nmol m^{-3} **(g,h,i)**). TA was held constant at 275 nmol m^{-3} and TS was varied between 0 and 275 nmol m^{-3} in order to generate a range of FA values that were observed during the field campaign (-275 to 275 nmol m^{-3}). RH increases from left to right (a to c) and TN increases from top the bottom (a to g). The maps were coloured by the amount of nitrate in the particle phase in nmol m^{-3} and overlaid with measurements and model predictions from Harrow, averaged to 30 min time resolution.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**The effect of
meteorological and
chemical factors**

M. Z. Markovic et al.




Fig. 8. Histogram of FA values calculated from measured and modelled TS and TA for **(a)** 30 min averages from the ground and **(b)** 2 min averages from the aircraft. The aircraft FA was calculated with $\text{TA} = \text{NH}_4^+$ due to lack of $\text{NH}_{3(\text{g})}$ measurements aloft.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 9. Harrow modelled vs. measured **(a)** RH **(b)** T . The black line is 1:1 line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of meteorological and chemical factors

M. Z. Markovic et al.




Fig. 10. Aircraft modelled vs. measured **(a)** RH **(b)** T . The black line is 1:1 line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

