

⁸Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, CO, USA

⁹Atmospheric Chemistry Services, Okehampton, UK

¹⁰National Centre for Atmospheric Science, School of Chemistry, University of Leeds, UK

Received: 4 October 2010 – Accepted: 20 October 2010 – Published: 1 November 2010

Correspondence to: H. Coe (hugh.coe@manchester.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Isoprene is the most abundant non-methane biogenic volatile organic compound (BVOC), but the processes governing secondary organic aerosol (SOA) formation from isoprene oxidation are only beginning to become understood and selective quantification of the atmospheric particulate burden remains difficult. Organic aerosol above a tropical rainforest located in Danum Valley, Borneo, Malaysia, a high isoprene emission region, was studied during Summer 2008 using Aerosol Mass Spectrometry and offline detailed characterisation using comprehensive two dimensional gas chromatography. Observations indicate that a substantial fraction (up to 15% by mass) of atmospheric sub-micron organic aerosol was observed as methylfuran (MF) after thermal desorption. This observation was associated with the simultaneous measurements of established gas-phase isoprene oxidation products methylvinylketone (MVK) and methacrolein (MACR). Observations of MF were also made during experimental chamber oxidation of isoprene. Positive matrix factorisation of the AMS organic mass spectral time series produced a robust factor which accounts for an average of 23% ($0.18 \mu\text{g m}^{-3}$), reaching as much as 53% ($0.50 \mu\text{g m}^{-3}$) of the total oraganic loading, identified by (and highly correlated with) a strong MF signal. Assuming that this factor is generally representative of isoprene SOA, isoprene derived aerosol plays a significant role in the region. Comparisons with measurements from other studies suggest this type of isoprene SOA plays a role in other isoprene dominated environments, albeit with varying significance.

1 Introduction

In order to assess the regional and global impacts of anthropogenic aerosols, it is necessary to quantify the contribution of the natural background. Aerosols can interact directly with incoming radiation by scattering or absorption, or indirectly through their effect on cloud formation and lifetime. A large fraction of the background par-

ACPD

10, 25545–25576, 2010

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 ticulate matter in the atmosphere is composed of organics thought to be formed by secondary processes from biogenic precursors (Kanakidou et al., 2005). Isoprene (2-methyl-1,3-butadiene) is potentially a major source of secondary organic aerosol (SOA) (Henze and Seinfeld, 2006), but highly time resolved atmospheric identification of isoprene derived SOA have so far proven elusive (Hallquist et al., 2009). In Amazonia, 2-methyltetrols identified on ambient filter samples were hypothesised to arise from isoprene oxidation (Claeys et al., 2004), and subsequently found to be present in chamber photo-oxidation of isoprene under low NO_x conditions (<1 ppb of NO_x with isoprene:NO_x ~500:1; Surratt et al., 2006). Low NO_x isoprene chamber studies have identified reactive photo-oxidation products in the gaseous and particulate phases such as epoxides (Paulot et al., 2009; Surratt et al., 2010), tetrols (Surratt et al., 2006, 2010; Kleindienst et al., 2009) and organosulphates (Surratt et al., 2008). These isoprene derived SOA species have also been measured offline in field studies (Chan et al., 2010; Surratt et al., 2008).

15 While previous work has identified a number of potentially important mechanisms (Chan et al., 2010; Paulot et al., 2009), the ambient measurements are based on offline analysis of bulk samples. These measurements are limited by low time-resolution, which prevents detailed photochemical analysis and comparison to rapidly changing parameters, such as photolysis rates and oxidant concentrations. The Aerodyne Aerosol Mass Spectrometer (AMS; Canagaratna et al., 2007) allows for online measurements of organic aerosols with a much higher time resolution and has been used previously for studies concerning isoprene oxidation (e.g. Capes et al., 2009). While the AMS reports bulk organic matter, the identification of specific chemical markers is required for more comprehensive biogenic source apportionment.

25 Here we present results from intensive measurements in an isoprene-dominated tropical forest environment and attempt to interpret a previously unreported chemical marker within the measurements. Factor analysis and comparisons with other datasets are used to estimate its importance both locally and in a global context.

Evidence for a significant proportion of SOA from isopreneN. H. Robinson et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2 Measurements

Measurements were made above a South East Asian Rainforest at the Danum Valley Conservation Area Global Atmospheric Watch (GAW) station, Sabah, in Malaysian Borneo (4.981° N, 117.844° E) as part of both the Oxidant and Particulate Photochemical Processes Above a South East Asian Rainforest project (OP3) (Hewitt et al., 2009, 2010) and the Aerosol Coupling in the Earth's System (ACES) project, during June and July 2008. A wide suite of aerosol, gas, radical and meteorological instruments were deployed in order to study the local atmospheric chemical processes as comprehensively as possible. In addition to the ground-based measurements, airborne measurements were made using instruments aboard the Facility for Airborne Atmospheric Measurements (FAAM) BAe 146 aircraft.

At the ground site, air was sampled from a height of 33 m above ground level on the top of a ridge surrounded by rainforest. The inlet sampled at a rate of 1500 l min⁻¹ through a 30 m long, 15 cm i.d. diameter tube insulated against solar heating. Air was then sub-sampled isokinetically from the centre of this flow at 35 l min⁻¹ and dried using a 780 tube Nafion drier using a dry air counter flow, where the air was decelerated to the laminar flow regime. After drying the air was decelerated further before being sampled by the suite of online instruments. The aircraft missions consisted of two separate sets of stacked straight and level runs (at 100–250, 1500, 3000 and 6000 m) over regions of rainforest (centred around the ground site) and homogeneous agro-industrial region of oil palm agriculture (centred around 5.25° N, 118.25° E). Aerosol was sampled on board the FAAM through a Rosemount inlet (Foltescu et al., 1995) and approximately 0.7 m of stainless steel tubing with a total residence time of ~4 s in the inlet system.

Filter samples (Thermo Scientific Partisol) were collected at the ground site from a height of 10 m. Particles less than 2.5 µm were collected onto pre-fired quartz filters which were transported and stored at -20 °C until analysis. Organic aerosol material was thermally desorbed from the filter for off-line analysis by comprehensive two dimensional gas chromatography with detection by Time-of-Flight Mass Spectrometry

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(GC×GC/ToF-MS) (Phillips and Beens, 1999; Hamilton et al., 2004). A dual stage commercial thermal desorption injector was used incorporating a thermal desorption unit (TDU) connected to a programmable-temperature vaporisation (PTV) injector, CIS-4 plus (Gerstel, Mulheim an der Ruhr, Germany), using a heated transfer line. 10-20 mg of filter paper containing each different sample was loaded into cleaned thermodesorption tubes using tweezers to ensure no contamination of the sample. The GC×GC/ToF-MS system consisted of an Agilent 6890 (Agilent Technologies, Palo Alto, CA, USA) gas chromatograph and a Pegasus III TOF-MS (LECO, St. Joseph, MI, USA). The first column was a non-polar DB5 (29 m × 0.32 mm i.d. × 0.25 μm film thickness) and the second column a DB17 (1.5 m × 0.10 mm i.d. × 0.10 μm film thickness).

Ground and airborne VOC measurements were made using Proton Transfer Reaction Mass Spectrometry (PTRMS) (Lindinger et al., 1998). Further details can be found in Langford et al. (2010) and Murphy et al. (2010). The PTRMS can measure methylvinylketone (MVK) and methacrolein (MACR), both first generation isoprene oxidation products. These species have the same molecular mass and cannot be separated using this technique, so are reported as the sum of the two, referred to as MVK+MACR.

Online aerosol composition was measured using two Aerodyne time of flight Aerosol Mass Spectrometers; a high mass resolution version (HR-AMS) (DeCarlo et al., 2006) for measurements at the 33 m level of the ground site measurement tower, and a compact version (C-AMS) (Drewnick et al., 2005) for airborne measurements on the FAAM. The AMS provides quantitative online measurements of sub-micron non-refractory aerosol composition and size. Despite extensive fragmentation of particulate matter species, compositional information can be gained from inspection of ions in the mass spectra. For example, the ratio of m/z 44 (mainly CO_2^+) to m/z 43 (mainly $\text{C}_2\text{H}_3\text{O}^+$) can be used as a proxy for the oxidation of organic aerosols (Morgan et al., 2010; Ng et al., 2010) and m/z 60 can be used as a marker for fresh biomass burning (Alfarra et al., 2007; Capes et al., 2008). The high resolution mode of the HR-AMS can resolve ion mass at a higher resolution (~ 4300) than the C-AMS, allowing separation of ions

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

with different elemental composition detected at the same m/z .

Positive matrix factorisation (PMF) analysis was also performed on the organic HR-AMS unit mass resolution data. This multivariate technique attempts to explain the AMS ensemble organic mass spectral time series as the sum of time series of differing amounts of static “factor” spectra which can then be linked to distinct contributions to the total organic mass (Paatero and Tapper U., 1994; Paatero, 1997; Ulbrich et al., 2009). Details of this analysis are included in the supplementary material. In short, a variety of solutions were calculated exploring different starting parameters and rotational ambiguities, and the most satisfactory solution was obtained when four factors were used. A detailed discussion of the results of PMF analysis will be included in Robinson et al. (2010).

3 Results

Total sub-micron non-refractory organic aerosol mass was less in Borneo than typical measurements in the Northern mid latitudes, which have an average of $2.8 \mu\text{g m}^{-3}$ for remote sites (Zhang et al., 2007). A mean organic matter loading of $0.74 \mu\text{g m}^{-3}$ was measured from the ground site, which is comparable to a mean loading of $0.64 \mu\text{g m}^{-3}$ measured in Amazonia (Chen et al., 2009). A time series of organic and inorganic sub-micron non-refractory aerosol loading is shown in Fig. 1. The ambient organic spectra from both AMS instruments shows a distinctive peak at m/z 82 (Fig. 2(a) and 2(b)), a feature not seen in the mid-latitude environments that make up the majority of ambient AMS observations to date. Furthermore, the high mass resolution capability of the HR-AMS was able to attribute the majority of the m/z 82 peak of the AMS mass spectrum to the $\text{C}_5\text{H}_6\text{O}^+$ ion (Fig. 2(b)). One of the PMF factors exhibited strong signals at m/z 82 and 53, henceforth called 82Fac, which was found to be very robust when subjected to variations in the starting conditions (starting seed) or when rotational ambiguity in the solutions was explored (fpeak; see Supplementl and Fig. 1). It was closely correlated with the m/z 82 peak with an r^2 of 0.86. Also, the other factors

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



contained largely insignificant contributions from this m/z .

GC×GC chromatography was used to investigate the source of the m/z 82 ion, as shown in Fig. 3. The dominant peak was identified as methylfuran (MF) based on a standard solution retention time and electron ionisation mass spectrum, although under this column set, the 2- and 3-methylfuran isomers cannot be separated. It is clear that there are no other substantial contributions to $C_5H_6O^+$. The AMS m/z 82 peak was found to occur in conjunction with a prominent m/z 53 peak (Pearson's r^2 of 0.96) corresponding to $C_4H_5^+$, the most abundant electron ionisation fragment ion of MF (NIST mass spec database, Stein, 2009). Combined with the GC×GC/ToF-MS results, this suggests that the m/z 82 fragment in the AMS can be attributed to the MF molecular ion, and confirms that the filter-identified MF was particulate and was not attributable to a gaseous artifact.

An AMS spectrum of pure 3-methylfuran (3MF) (Fig. 2(d)) was obtained by nebulising commercially available 3MF (Fisher Scientific, UK; 98% pure) suspended in deionized water in the laboratory. A background spectrum of deionized water measured using the same equipment was subtracted to remove any influences from contamination sources. The measured sample spectrum compared well to the 3MF reference spectrum (NIST mass spec database, Stein, 2009) and the unusual features of the ambient organic spectrum, such as the prominent m/z 82 and 53 peaks. Inspection of the high resolution m/z 82 peak of the pure 3MF mass spectrum showed only $C_5H_6O^+$ ions to be present, comprising 13% of the total organic mass.

The average diurnal profile of the $C_5H_6O^+$ signal shows an increase throughout the day with a maximum in the early evening (Fig. 4(a)). This is also the case for the diurnal profile of the fraction of organic aerosol at $C_5H_6O^+$ (Fig. 4(b)). These elevated levels in the evening may be caused by partitioning of semi-volatile aerosol as the ambient temperature drops. They may also be transported from the isoprene rich oil palm plantations (Hewitt et al., 2009) close to the site (around 30 km), causing a time lag between production and measurement. The first-generation isoprene oxidation products MVK+MACR, also show a strong diurnal cycle linked to the peak in the emission

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

of isoprene at midday with the delay caused by the photochemical reaction rate. If it is assumed that the $C_5H_6O^+$ signal has a similar precursor, the fact it peaks later in the day could imply that it is produced by a longer or slower sequence of reactions.

Airborne PTRMS measurements of MVK+MACR, and AMS measurements of m/z 82 were also compared (Fig. 5). The median of the AMS m/z 82 and m/z 53 peaks measured in the boundary layer over each of eight individual flights correlated well with gas phase MVK+MACR, with respective Pearson's r^2 values of 0.92 and 0.87. This shows that both measurements exhibited systematically similar daily increases, which would be expected if they shared a common precursor. A substantial m/z 82 signal was observed throughout the surface mixed layer to a height of 3000 m, with little signal observed in the lower free troposphere above (Fig. 6(a)). That MF was measured throughout the boundary layer shows that it is of regional importance. Altitude profiles show the AMS aerosol signal at m/z 82 increases throughout the day, at all altitudes in the boundary layer, which is consistent with production from photochemical processing of precursors rather than direct emission from the surface. This increase in the afternoon is also observed in the altitude profile of gas phase MVK+MACR (Fig. 6(c)), although this increase is mostly below 2000 m. The fractional contribution of m/z 82 to the total organic mass is greater at the top of the boundary layer than at the surface (Fig. 6(b)), with a maximum fractional loading of 1.7 times the low altitude fractional loading in the morning, and 2.5 times in the afternoon. There is an increase in this fractional contribution throughout the day, with measurements at low altitude showing an average increase of $\sim 9\%$, which is similar to the change seen in the ground site diurnal profiles. The altitude profiles are also consistent with semi-volatile aerosol partitioning at the top of the atmosphere where it is cooler, with a change of temperature of 298 K at ~ 500 m to 283 K at ~ 3000 m. This may also explain the disparity between the MVK+MACR profile and the m/z 82 profile at the top of the boundary layer.

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Possible sources of MF

Previous studies have presented evidence of gas-phase methylfuran formation through oxidation of isoprene by OH radicals: MF has been measured in simulation chamber studies of isoprene oxidation (Atkinson et al., 1989; Ruppert and Becker, 2000; Sprengnether et al., 2002) and, in a study of a rural US forest, ambient measurements of gaseous MF were found to correlate with isoprene (Montzka et al., 1995). The likelihood of MF being present in the condensed phase was investigated using methods to estimate first the normal boiling point (Nannoolal et al., 2004), and then the vapour pressure (Nannoolal et al., 2008), evaluated as being the best available for atmospheric purposes (Barley and McFiggans, 2010). A literature boiling point was used (Burness, 1956) (338.65 K) with the above vapour pressure method to give an estimated value of 0.217 atm at 298.15 K. This value was then used to estimate the abundance of MF that would be expected to be present in secondary organic aerosol under equilibrium conditions at 298.15 K. Using a partitioning model (Barley et al., 2009) with a reasonable and representative set of assumptions ($2\mu\text{g m}^{-3}$ of completely involatile, perfectly organic core of molar mass 320 g mol^{-1} with total concentration of the considered molecule set to $2\mu\text{g m}^{-3}$), it was estimated that MF contributed $1.41 \times 10^{-9}\mu\text{g m}^{-3}$ to the condensed mass ($7.05 \times 10^{-8}\%$ by mass). Based upon an equilibrium partitioning consideration, MF should not be present in the condensed phase in detectable quantities under reasonable atmospheric conditions.

That MF is measured from condensed material may be explained if it is directly incorporated into the particles by some means other than absorptive partitioning, such as reversible reactive uptake. However, no such mechanism is known. Instead, we hypothesise that MF measurements are a result of decomposition of some, as yet unknown, condensed phase isoprene photo-oxidation product upon the thermal vaporisation used in the GC×GC and AMS analyses reported here. This is supported by the association of detected MF with the gas-phase isoprene photo-oxidation products MVK+MACR. The diurnal profiles shown in Fig. 4 also support this, being consistent

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Evidence for a significant proportion of SOA from isopreneN. H. Robinson et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

with comparisons of day and night filter samples from previous studies which show greater levels of isoprene SOA markers during the day when isoprene emissions and photochemistry are greatest (Kourtchev et al., 2008; Ion et al., 2005). Known potential sources of isoprene SOA include epoxides (Paulot et al., 2009; Surratt et al., 2010), tetrols (Surratt et al., 2006, 2010; Kleindienst et al., 2009), peroxides (Surratt et al., 2006) or oligomers (Surratt et al., 2010) (formed through accretion reactions) but at present it is unclear how they would produce MF upon volatilisation. It is conceivable that this relation of MF to MVK+MACR could be caused by terpene photo-oxidation products produced at a similar rate to isoprene products. However, there is no evidence in the literature for MF formation from terpene photo-oxidation, in contrast to various isoprene mechanistic studies. This suggests that the MF is a product of a form of isoprene SOA not previously identified.

The previously reported isoprene:NO_x ratio measured during OP3 had a typical value of 20:1 (Hewitt et al., 2009) which is comparable to ratios in NO_x-limited chamber oxidation studies of isoprene (e.g. Lane et al., 2008). The formation of gas-phase 3MF during isoprene photo-oxidation studies has previously been attributed to cyclisation/dehydration of unsaturated C₅ 1,4-hydroxycarbonyls formed (Sprengnether et al., 2002; Dibble, 2007), in a process that would also be possible during the heating involved in the analyses reported here. A C₅H₈O⁺ ion has been attributed to the fragmentation of 1,4-hydroxycarbonyls in previous isoprene photo-oxidation studies (Zhao et al., 2004). This signal was present in the ambient mass spectrum in Borneo and was covariant with the molecular ion of MF with a Pearson's *r*² of 0.57 but absent in the laboratory 3MF spectrum, consistent with the hypothesis that unsaturated 1,4-hydroxycarbonyls are the source of the ambient MF measurements. While these studies were performed at NO_x levels higher than in Borneo, they do provide a possible candidate for a mechanism that yields MF measurements. Two unsaturated C₅-hydroxycarbonyls (4-hydroxy-2-methyl-but-2-enal and 4-hydroxy-3-methyl-but-2-enal), are known to be formed as substantial first generation products of the OH-initiated degradation of isoprene, with their collective molar yield reported to be approximately

15–20% in a number of studies performed in the presence of NO_x, albeit at higher concentrations than in the atmosphere (Kwok et al., 1995; Zhao et al., 2004; Baker et al., 2005). Hydroxycarbonyls have also been observed in NO_x-free systems with a collective yield approaching 20% (Ruppert and Becker, 2000), with the dominant contribution estimated to be attributable to the above 1,4 isomers (Jenkin et al., 1998) and substantial formation may occur under the range of conditions of the OP3 field campaign.

Similar partitioning calculations as for MF were performed for C₅-hydroxycarbonyls. They were estimated to have a boiling point of 463.06 K and a vapour pressure at 298.15 K of 3.38 × 10⁻⁵ atm. The C₅-hydroxycarbonyls contributed 9.05 × 10⁻⁶ μg m⁻³ to the condensed mass (4.5 × 10⁻⁴% by mass). It is apparent from these calculations that, if either or both of the C₅-hydroxycarbonyls are degrading upon measurement to give MF, it must itself be a degradation product of another, less volatile compound. Again it is unclear how C₅-hydroxycarbonyls could be formed from established isoprene or terpene SOA species. One possibility is that some isoprene SOA could be in the form of peroxyhemiacetals which produce MF upon analysis. Hydroperoxide species (ROOH), such as isoprene derived hydroxyhydroperoxides, can be formed under (low NO_x) NO_x-limited conditions where RO₂+HO₂ reactions become important. Under such conditions peroxyhemiacetals can be formed from the (reversible) reactions of isoprene derived 1,4-hydroxycarbonyls with hydroperoxides and hydroperoxides with aldehydes. The formation of high molecular weight peroxyhemiacetals has been characterised in previous studies of condensed organic material formed from the ozonolysis of alkene and monoterpene systems (Tobias and Ziemann, 2000; Docherty et al., 2005). Future chamber studies investigating isoprene SOA should seek to identify the potential sources of C₅-hydroxycarbonyls and MF in analytical measurements.

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Mass estimates of MF related aerosol

The fraction of the ambient m/z 82 peak present as $C_5H_6O^+$ was calculated as a function of time using the high resolution AMS data. Averaged over the whole data set, these time series have a ratio of mean values of 80%. This high resolution time series was scaled to the unit mass resolution data to ensure consistency with calibrations, and used to derive a quantitative estimate of the $C_5H_6O^+$ signal (used in Fig. 1). By scaling the laboratory 3MF spectrum so its m/z 82 peak is the same magnitude as the ambient $C_5H_6O^+$ signal, it is possible to estimate an upper limit to the MF loading detected, MF_{amb} , using

$$MF_{amb} = \frac{3MF_s}{C_5H_6O_s^+} C_5H_6O_{amb}^+ \quad (1)$$

where $3MF_s$ is the total organic loading of the laboratory sample spectrum, $C_5H_6O_{amb}^+$ is the $C_5H_6O^+$ signal in the ambient spectrum and $C_5H_6O_s^+$ is the $C_5H_6O^+$ signal in the laboratory 3MF sample spectrum (m/z 82 loading). The mean ambient $C_5H_6O^+$ signal measured at the ground site during OP3 was 1.0% of the mean total organic aerosol loading, which gives a mean MF_{amb} loading of $0.06 \mu g m^{-3}$, or 8% of the organic aerosol. The maximum (95th percentile) MF_{amb} loading was $0.14 \mu g m^{-3}$, and the maximum fractional loading was 15% of the simultaneously measured total organic aerosol.

The mass of particulate SOA due to isoprene oxidation in the ambient will be greater than the mass of MF it produces upon analysis, meaning this approach underestimates the loading of isoprene derived SOA. An estimate of the total amount of organic aerosol due to isoprene oxidation can be made using the 82Fac from the PMF analysis of the HR-AMS unit mass resolution spectrum (see supporting material). It should be noted that the mass spectrum of 82Fac (Fig. 2(c)) is more complex than the 3MF laboratory mass spectrum (Fig. 2(d)) implying other organic components also contribute to the 82Fac PMF factor. However, the close correlation of 82Fac with m/z 82 ($r^2 = 0.86$)

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



identifies it as being derived from the same organic aerosol as MF, either directly as other components that are also formed on thermal degradation, or indirectly as isoprene SOA species that are produced simultaneously but through a different mechanism.

Note that it is conceivable that isoprene SOA produced at a different rate to the MF-precursor aerosol may also be contributing to other factors (such as OOA2), although we have no evidence to suggest this is the case. It is likely that, if MF is indeed indicative of isoprene SOA, the 82Fac PMF factor provides a good estimate of the amount of isoprene SOA in the region.

6 Global extent of MF

Such a prominent m/z 82 signal has not been reported in previously published studies over biogenic sources of VOCs. However, analysis of HR-AMS mass spectra from a study in the Amazon Basin, previously published as part of the Amazonian Aerosol Characterization Experiment (AMAZE-08) (Chen et al., 2009), also shows evidence for $C_5H_6O^+$ during periods dominated by in-Basin sources, though with a lower mean signal comprising 0.5% of the organic mass, compared to the mean $C_5H_6O^+$ signal of 1% measured in Borneo. Table 1 compares m/z 82 and $C_5H_6O^+$ signals from projects in various locations around the World showing the fractional contribution of m/z 82 to the total organic matter and its comparison with the adjacent peaks, which is an indicator of how distinct the signal is compared to other peaks in that part of the mass spectrum. Though these two metrics are useful for comparing projects, care should be taken in their interpretation as both are vulnerable to conflating circumstances: for example, the m/z 82 peak will appear as a low fraction of the total mass if MF yielding aerosol is present in low levels compared to other organic aerosol, and if large amounts of a species that fragment at masses around m/z 82 are measured then the m/z 82 peak may be a significant fraction of the total mass without being prominent. It should also be noted that it is impossible to assess the contribution of non-MF ions to m/z 82

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



peaks unless high mass resolution data is available.

The m/z 82 peak in the rainforest environments is more prominent, implying these measurements are from stable molecular ions, rather than background fragments resultant from the breakup of non-MF-precursor organic aerosol. Higher levels of $C_5H_6O^+$ yielding aerosol have been measured in these two isoprene dominated rainforest environments than in a North American temperate forest of mixed terpene and isoprene influence (BEARPEX campaign, see Appendix A), which in turn has higher levels than a terpene dominated environment (Kulmala et al., 2009). This is consistent with the detection of MF being a result of isoprene derived SOA.

HR-AMS data from low NO_x isoprene photo-oxidation experiments (isoprene: $NO_x > 200 : 1$) in the Harvard Environmental Chamber (King et al., 2010) also show an m/z 82 peak distinct from the adjacent peaks. While the m/z 82 peak is responsible for less signal than in either of the tropical field studies (0.4% of total organic mass), the prominence of the $C_5H_6O^+$ ion at m/z 82 strongly suggests that the same SOA formation mechanism is taking place and could be isolated through further laboratory work. Sulphate levels in Borneo are around four times greater than in the Amazon. As acidic sulphate had been shown to play an important role in isoprene SOA formation in previous studies, its presence may contribute to the greater significance of MF in Borneo, although chamber studies have not shown sulphate-isoprene SOA mechanisms likely to yield MF (Surratt et al., 2010).

7 Conclusions

AMS and GC \times GC measurements of organic aerosol composition in Borneo identified a MF signal at m/z 82 and 53, the parent ion and major fragment respectively. This was confirmed by laboratory measurements of a 3MF sample which showed the same m/z 82 and 53 peaks. The m/z 82 signal was present throughout the boundary layer with the signal increasing throughout the day, associated with the isoprene oxidation products MVK+MACR through correlation of aircraft data and diurnal profiles

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



unidentified isoprene product(s). As such, identification of the MF precursor aerosol may provide key insights into missing SOA, and the identification of the relationship of the AMS m/z 82 peak to isoprene SOA opens up the opportunity for high time resolution isoprene SOA measurements in other projects. The PMF analysis suggests that an average of 23% ($0.18 \mu\text{g m}^{-3}$), and as much as 53% ($0.5 \mu\text{g m}^{-3}$) of the organic aerosol may be produced from isoprene oxidation in Borneo. The natural balance of isoprene is being affected by deforestation for oil palm agriculture (McMorrow and Talip, 2001) which emits five times as much isoprene as the natural forest (Hewitt et al., 2009), meaning isoprene SOA will only become more important with further similar land use change.

Appendix A

Details of unpublished data from the BEARPEX campaign

The North American temperate HR-AMS measurements presented in Table 1 of the main text, took place during the BEARPEX (Biosphere Effects on AeRosols and Photochemistry EXperiment) campaign at Blodgett Forest Ameriflux site in 2007. The site is a mid-elevation ponderosa pine plantation owned by Sierra Pacific Industries, and located $\sim 75\text{km}$ northeast of Sacramento on the western slope of California's mid-elevation Sierra Nevada mountains (38.90° N , 120.63° W , 1315 m elevation). AMS measurements were taken between 19 August and 28 September 2007, at the top of an 18 m tower. The site has been previously described in detail Dreyfus et al. (2002). Wind patterns throughout the measurement period were consistent, with westerly to southwesterly wind during the day and northeasterly to easterly winds at night Murphy et al. (2006). This results in a consistent diurnal pattern of VOCs in which a mix of terpenes and other pine-derived biogenic VOCs are observed in the morning followed by isoprene and its oxidation products at mid-day as air is transported over a lower elevation band of oak forest Dreyfus et al. (2002). In the afternoon, a mix of biogenic

25561

ACPD

10, 25545–25576, 2010

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and anthropogenic VOCs are caused by transport of polluted air from the Sacramento area. The Blodgett forest canopy is dominated by *Pinus ponderosa* L., planted in 1990 with a mean height of 8 m, with few Douglas fir, white fir and incense cedar. The understory includes mountain whitehorn and manzanita. The data presented have been screened for the influence of biomass burning.

Supplement related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/10/25545/2010/
acpd-10-25545-2010-supplement.pdf](http://www.atmos-chem-phys-discuss.net/10/25545/2010/acpd-10-25545-2010-supplement.pdf).

Acknowledgements. Thanks to J. Kroll (Massachusetts Institute of Technology) and M. Alfara (NCAS, University of Manchester) for conversations during the preparation of this manuscript. This work was supported by the UK Natural Environment Research Council through the OP3 (grant NE/D002117/1) and ACES (grant NE/E011179/1) projects. QC and STM thank US NSF ATM-0723582. KSD, DKF, and JLJ thank US NSF ATM-0449815, ATM-0449815, and NOAA NA08OAR4310565. We thank the Malaysian and Sabah Governments for their permission to conduct research in Malaysia; the Malaysian Meteorological Department for access to the measurement site; Yayasan Sabah and the Royal Society's South East Asian Rain Forest Research Programme for logistical support; and the ground staff, engineers, scientists and flight crew of the FAAM aircraft. This is paper number 515 of the Royal Societys South East Asian Rainforest Research Programme.

References

- Alfara, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the Mass Spectral Signature of Organic Aerosols from Wood Burning Emissions, *Environ. Sci. Technol.*, 41, 5770–5777, doi:10.1021/es062289b, 2007. 25550
- Atkinson, R., Aschmann, S. M., Tuazon, E. C., Arey, J., and Zielinska, B.: Formation of 3-Methylfuran from the gas-phase reaction of OH radicals with isoprene and the rate constant for its reaction with the OH radical, *Int. J. Chem. Kin.*, 21, 593–604, 1989. 25554

25562

ACPD

10, 25545–25576, 2010

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Baker, J., Arey J., and Atkinson, R.: Formation and reaction of hydroxycarbonyls from the reaction of OH radicals with 1, 3-butadiene and isoprene, *Environ. Sci. Technol.*, 39, 4091–4099, 2005. 25556

Barley, M., Topping, D. O., Jenkin, M. E., and McFiggans, G.: Sensitivities of the absorptive partitioning model of secondary organic aerosol formation to the inclusion of water, *Atmos. Chem. Phys.*, 9, 2919–2932, 2009. 25554

Barley, M. H. and McFiggans, G.: The critical assessment of vapour pressure estimation methods for use in modelling the formation of atmospheric organic aerosol, *Atmos. Chem. Phys.*, 10, 749–767, doi:10.5194/acp-10-749-2010, 2010. 25554

Burness, D. M.: Beta-Keto acetals. II. Synthesis of 3-methyl- and 3-phenyl-furans., *J. Org. Chem.*, 21, 102–104, 1956. 25554

Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., and Others: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26, 185–222, 2007. 25548

Capes, G., Johnson, B., McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of biomass burning aerosols over West Africa: Aircraft measurements of chemical composition, microphysical properties, and emission ratios, *J. Geophys. Res.*, 113, D00C15, doi:10.1029/2008JD009845, available online at: <http://www.agu.org/pubs/crossref/2008/2008JD009845.shtml>, 2008. 25550

Capes, G., Murphy, J. G., Reeves, C. E., McQuaid, J. B., Hamilton, J. F., Hopkins, J. R., Crosier, J., Williams, P. I., and Coe, H.: Secondary organic aerosol from biogenic VOCs over West Africa during AMMA, *Atmos. Chem. Phys.*, 9, 3841–3850, doi:10.5194/acp-9-3841-2009, 2009. 25548, 25560

Carlton, A., Wiedinmyer, C., and Kroll, J.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987–5005, doi:10.5194/acp-9-4987-2009, 2009. 25560

Chan, M. N., Surratt, J. D., Claeys, M., Edgerton, E. S., Tanner, R. L., Shaw, S. L., Zheng, M., Knipping, E. M., Eddingsaas, N. C., Wennberg, P. O., and Seinfeld, J. H.: Characterization and Quantification of Isoprene-Derived Epoxydiols in Ambient Aerosol in the Southeastern United States, *Environ. Sci. Technol.*, 44, 4590–4596, available online at: <http://dx.doi.org/10.1021/es100596b>, 2010. 25548

Chen, Q., Farmer, D. K., Schneider, J., Zorn, S. R., Heald, C. L., Karl, T. G., Guenther, A., Allan, J. D., Robinson, N., Coe, H., Kimmel, J. R., Pauliquevis, T., Borrmann, S., Poschl, U.,

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Andreae, M. O., Artaxo, P., Jimenez, J. L., and Martin, S. T.: Mass spectral characterization of submicron biogenic organic particles in the Amazon Basin, *Geophys. Res. Lett.*, 36, L20806, doi:10.1029/2009GL039880, 2009. 25551, 25558, 25570

5 Claeys, M. et al.: Formation of Secondary Organic Aerosols Through Photooxidation of Isoprene, *Science*, 303, 1173–1176, 2004. 25548

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and J.L., J.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer., *Anal. Chem.*, 78, 8281–8289, 2006. 25550

10 Dibble, T. S.: Cyclization of 1,4-hydroxycarbonyls is not a homogenous gas phase process, *Chemical Physics Letters*, 447, 5–9, doi:10.1016/j.cplett.2007.08.088, available online at: <http://dx.doi.org/10.1016/j.cplett.2007.08.088>, 2007. 25555

Docherty, K. S., Wu, W., Lim, Y. B., and Ziemann, P. J.: Contributions of Organic Peroxides to Secondary Aerosol Formed from Reactions of Monoterpenes with O₃, *Environ. Sci. Technol.*, 39, 4049–4059, doi:10.1021/es050228s, available online at: <http://pubs.acs.org/doi/abs/10.1021/es050228s>, 2005. 25556

15 Drewnick, F., Hings, S. S., Decarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Borrmann, K. L. D. S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) – Instrument description and first field deployment, *Aerosol Sci. Technol.*, 39, 637–658, 2005. 25550

Dreyfus, G. B., Schade, G. W., and Goldstein, A.: Observational constraints on the contribution of isoprene oxidation to ozone production on the western slope of the Sierra Nevada, California, *J. Geophys. Res.*, 107, 4365, doi:10.1029/2001JD001490, 2002. 25561

20 Foltescu, V. L., Selin, E., and Below, M.: Corrections for particle losses and sizing errors during aircraft aerosol sampling using a rosemounts inlet and the PMS LAS-X, *Atmos. Environ.*, 29, 449–453, 1995. 25549

25 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A., Hamilton, J. F., Herrmann, H., Hudson, L. E., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prvt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5235, doi:10.5194/acp-9-5155-2009, 2009. 25548

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hamilton, J. F., Webb, P. J., Lewis, A. C., Hopkins, J. R., Smith, S., and Davy, P.: Partially oxidised organic components in urban aerosol using GC×GC-TOF/MS, *Atmos. Chem. Phys.*, 4, 1279–1290, doi:10.5194/acp-4-1279-2004, 2004. 25550
- 5 Heald, C., Jacob, D., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and Weber, R. J.: A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, 32, 1–4, 2005.
- Henze, D. and Seinfeld, J. H.: Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33, L09812, doi:10.1029/2006GL025976, 2006. 25548
- 10 Hewitt, C. N., MacKenzie, A. R., Carlo, P. D., Marco, C. F. D., Dorsey, J. R., Evans, M., Fowler, D., Gallagher, M. W., Hopkins, J. R., Jones, C. E., Langford, B., Lee, J. D., Lewis, A. C., Lim, S. F., McQuaide, J., Misztal, P., Moller, S. J., Monks, P. S., Nemitz, E., Oram, D. E., Owen, S. M., Phillips, G. J., Pugh, T. A. M., Pyle, J. A., Reeves, C. E., Ryder, J., Siong, J., Skiba, U., and Stewart, D. J.: Nitrogen management is essential to prevent tropical oil palm plantations from causing ground level ozone pollution, *Proc. Natl. Acad. Sci. USA*, 106, 18447–18451, doi:10.1073/pnas.0907541106, 2009. 25549, 25552, 25555, 25561
- 15 Hewitt, C. N., Lee, J. D., MacKenzie, A. R., Barkley, M. P., Carslaw, N., Carver, G. D., Chappell, N. A., Coe, H., Collier, C., Commane, R., Davies, F., Davison, B., DiCarlo, P., Di Marco, C. F., Dorsey, J. R., Edwards, P. M., Evans, M. J., Fowler, D., Furneaux, K. L., Gallagher, M., Guenther, A., Heard, D. E., Helfter, C., Hopkins, J., Ingham, T., Irwin, M., Jones, C., Karunaharan, A., Langford, B., Lewis, A. C., Lim, S. F., MacDonald, S. M., Mahajan, A. S., Malpass, S., McFiggans, G., Mills, G., Misztal, P., Moller, S., Monks, P. S., Nemitz, E., Nicolas-Perea, V., Oetjen, H., Oram, D. E., Palmer, P. I., Phillips, G. J., Pike, R., Plane, J. M. C., Pugh, T., Pyle, J. A., Reeves, C. E., Robinson, N. H., Stewart, D., Stone, D., Whalley, L. K., and Yin, X.: Overview: oxidant and particle photochemical processes above a south-east Asian tropical rainforest (the OP3 project): introduction, rationale, location characteristics and tools, *Atmos. Chem. Phys.*, 10, 169–199, doi:10.5194/acp-10-169-2010, 2010. 25549
- 20 lon, A. C., Vermeylen, R., Kourtchev, I., Cafmeyer, J., Chi, X., Gelencsr, A., Maenhaut, W., and Claeys, M.: Polar organic compounds in rural PM_{2.5} aerosols from K-pusztá, Hungary, during a 2003 summer field campaign: Sources and diel variations, *Atmos. Chem. Phys.*, 5, 1805–1814, doi:10.5194/acp-5-1805-2005, 2005. 25555
- 25 Jenkin, M. E., Boyd, A. A., and Lesclaux, R.: Peroxy radical kinetics resulting from the OH-initiated oxidation of 1, 3-butadiene, 2, 3-dimethyl-1, 3-butadiene and isoprene, *J. Atmos. Chem.*, 29, 267–298, 1998. 25556
- 30

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Kanakidou, M., Seinfeld, J., Pandis, S. N., Barnes, I., Dentener, F., Facchini, M., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C., and Others: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005. 25548

5 King, S. M., Rosenoern, T., Shilling, J. E., Chen, Q., Wang, Z., Biskos, G., McKinney, K. A., Poeschl, U., and Martin, S. T.: Cloud droplet activation of mixed organic-sulfate particles produced by the photooxidation of isoprene, *Atmos. Chem. Phys.*, 10, 3953–3964, doi:10.5194/acp-10-3953-2010, 2010. 25559

10 Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Jaoui, M., and Edney, E. O.: The formation of secondary organic aerosol from the isoprene+OH reaction in the absence on NO_x, *Atmos. Chem. Phys.*, 9, 6541–6558, doi:10.5194/acp-9-6541-2009, 2009. 25548, 25555

15 Kourtchev, I., Ruuskanen, T. M., Keronen, P., Sogacheva, L., Dal Maso, M., Reissell, A., Chi, X., Vermeylen, R., Kulmala, M., Maenhaut, W., and Claeys, M.: Determination of isoprene and alpha-/beta-pinene oxidation products in boreal forest aerosols from Hyytiälä, Finland: diel variations and possible link with particle formation events., *Plant biology (Stuttgart, Germany)*, 10, 138–49, doi:10.1055/s-2007-964945, available online at: <http://www.ncbi.nlm.nih.gov/pubmed/18211553>, 2008. 25555

20 Kulmala, M., Asmi, A., Lappalainen, H. K., Carslaw, K. S., Pschl, U., Baltensperger, U., Hov, ., Brenquier, J.-L., Pandis, S. N., Facchini, M. C., Hansson, H.-C., Wiedensohler, A., and O'Dowd, C. D.: Introduction: European Integrated Project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) - integrating aerosol research from nano to global scales, *Atmos. Chem. Phys.*, 9, 2825–2841, doi:10.5194/acp-9-2825-2009, 2009. 25559, 25570

Kwok, E. S. C., Atkinson, R., and Arey, J.: Observation of hydroxycarbonyls from the OH radical-initiated reaction of isoprene, *Environ. Sci. Technol.*, 29, 2467–2469, 1995. 25556

25 Lane, T. E., Donahue, N. M., and Pandis, S. N.: Effect of NO_x on Secondary Organic Aerosol Concentrations, *Environ. Sci. Technol.*, 42, 6022–6027, doi:10.1021/es703225a, available online at: <http://pubs.acs.org/doi/abs/10.1021/es703225a>, 2008. 25555

Langford, B., Misztal, P. K., Nemitz, E., Davison, B., Helfter, C., Pugh, T. A. M., MacKenzie, A. R., Lim, S. F. and Hewitt, C. N.: Fluxes and concentrations of volatile organic compounds from a South-East Asian tropical rainforest, *Atmos. Chem. Phys.*, 10(17), 8319–8412, doi:10.5194/acp-10-8391-2010, 2010 25550

30 Lim, Y. B. and Ziemann, P. J.: Kinetics of the heterogeneous conversion of 1,4-hydroxycarbonyls to cyclic hemiacetals and dihydrofurans on organic aerosol particles.,

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Physical Chemistry Chemical Physics, 11, 8029–39, doi:10.1039/b904333k, available online at: <http://www.ncbi.nlm.nih.gov/pubmed/19727510> 2009. 25560

Lindinger, W., Hansel, A., and Jordan, A.: Proton-transfer-reaction mass spectrometry (PTR-MS): On-line monitoring of volatile organic compounds at pptv levels, *Chemical Society reviews*, 27, 347–354, 1998. 25550

McMorrow, J. and Talip, M. A.: Decline of forest area in Sabah, Malaysia: Relationship to state policies, land code and land capability, *Global Environ. Change*, 11, 217–230, doi:10.1016/S0959-3780(00)00059-5, available online at: [http://dx.doi.org/10.1016/S0959-3780\(00\)00059-5](http://dx.doi.org/10.1016/S0959-3780(00)00059-5) 2001. 25561

Montzka, S. A., Trainer, M., Angevine, W. M., and Fehsenfeld, F. C.: Measurements of 3-methylfuran, methyl vinyl ketone, and methacrolein at a rural forested site in the southeastern United States, *J. Geophys. Res.*, 100(11), 11393–11401, <http://www.agu.org/journals/jd/v100/iD06/95JD01132/95JD01132.pdf>, 1995. 25554

Morgan, W. T., Allan, J. D., Bower, K. N., Highwood, E. J., Liu, D., McMeeking, G. R., Northway, M. J., Williams, P. I., Krejci, R., and Coe, H.: Airborne measurements of the spatial distribution of aerosol chemical composition across Europe and evolution of the organic fraction, *Atmos. Chem. Phys.*, 10, 4065–4083, doi:10.5194/acp-10-4065-2010, 2010. 25550

Murphy, J. G., Day, D. A., Cleary, P. A., Wooldridge, P. J., and Cohen, R. C.: Observations of the diurnal and seasonal trends in nitrogen oxides in the western Sierra Nevada, *Atmos. Chem. Phys.*, 6, 5321–5338, doi:10.5194/acp-6-5321-2006, 2006. 25561

Murphy, J. G., Oram, D. E. and Reeves, C. E.: Measurements of volatile organic compounds over West Africa, *Atmos. Chem. Phys.*, 10, 5281–5294, doi:10.5194/acp-10-5281-2010, 2010. 25550

Nannoolal, Y., Rarey, J., Ramjugernath, D., and Cordes, W.: Estimation of pure component properties: Part 1. Estimation of the normal boiling point of non-electrolyte organic compounds via group contributions and group interactions, *Fluid Phase Equilibria*, 226, 45–63, doi:10.1016/j.fluid.2004.09.001, <http://dx.doi.org/10.1016/j.fluid.2004.09.001>, 2004. 25554

Nannoolal, Y., Rarey, J., and Ramjugernath, D.: Estimation of pure component properties: Part 3. Estimation of the vapor pressure of non-electrolyte organic compounds via group contributions and group interactions, *Fluid Phase Equilibria*, 269, 117–133, doi:10.1016/j.fluid.2008.04.020, <http://dx.doi.org/10.1016/j.fluid.2008.04.020>, 2008. 25554

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt,

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, doi:10.5194/acp-10-4625-2010, 2010. 25550

5 Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kurten, A., St. Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325, 730–733, doi:10.1126/science.1172910, 2009. 25548, 25555

Phillips, J. B. and Beens, J.: Comprehensive two-dimensional gas chromatography: a hyphenated method with strong coupling between the two dimensions, *J. Chromatogr. A*, 856, 331–347, doi:10.1016/S0021-9673(99)00815-8, <http://www.sciencemag.org/cgi/content/abstract/325/5941/730> 1999. 25550

Ruppert, L. and Becker, H. H.: A product study of the OH radical-initiated oxidation of isoprene: formation of C5-unsaturated diols, *Atmos. Environ.*, 34, 1529–1542, 2000. 25554, 25556

Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometr. Intell. Lab. Syst.*, 37, 23–35, 1997. 25551

Paatero, P. and Tapper U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111–126, 1994. 25551

Robinson, N. H., Newton, H., Allan, J. D., Irwin, M., Hamilton, J. F., Chen, Q., Martin, S. T., McFiggans, G., and Coe, H.: Source attribution during the OP3 project using backwards air mass trajectories, in preparation, *Atmos. Chem. Phys.*, 2010. 25551

20 Sprengnether, M., Demerjian, K. L., Donahue, N. M., and Anderson, J. G.: Product analysis of the OH oxidation of isoprene and 1,3-butadiene in the presence of NO, *J. Geophys. Res.*, 107, 4268, doi:10.1029/2001JD000716, 2002. 25554, 25555

Stein, S. E.: “Mass Spectra” in NIST Chemistry WebBook, NIST Standard Reference Database Number 69, <http://webbook.nist.gov>, 2009. 25552

25 Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of Secondary Organic Aerosol Formed from the Photooxidation of Isoprene, *J. Phys. Chem. A*, 110, 9665–9690, doi.org/10.1021/jp061734m, 2006. 25548, 25555

30 Surratt, J. D., Gomez-Gonzalez, Y., Chan, A. W. H., Vermeylen, R., Shahgholi, M., Kleindienst, T. E., Edney, E. O., Offenberg, J. H., Lewandowski, M., Jaoui, M., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Organosulfate Formation in Biogenic Secondary Organic Aerosol, *The Journal of Physical Chemistry A*, 112, 8345–8378, available online at: <http://>

//dx.doi.org/10.1021/jp061734m, 2008. 25548

Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Atmospheric Chemistry Special Feature: Reactive intermediates revealed in secondary organic aerosol formation from isoprene., *Proc. Natl. Acad. Sci. USA*, 15, 6640–6645, doi:10.1073/pnas.0911114107, , 2010. 25548, 25555, 25559, 25560

Tobias, H. J. and Ziemann, P. J.: Thermal Desorption Mass Spectrometric Analysis of Organic Aerosol Formed from Reactions of 1-Tetradecene and O₃ in the Presence of Alcohols and Carboxylic Acids, *Environ. Sci. Technol.*, 34, 2105–2115, doi:10.1021/es9907156, available online at: <http://pubs.acs.org/doi/abs/10.1021/es9907156> , 2000. 25556

Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009. 25551

Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, *Environ. Sci. Technol.*, 39, 4938–4952, 2005.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T. B., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K. N., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, available online at: <http://www.agu.org/pubs/crossref/2007/2007GL029979.shtml>, 2007. 25551

Zhao, J., Zhang, R., Fortner, E. C., and North, S. W.: Quantification of Hydroxycarbonyls from OH-Isoprene Reactions, *J. Am. Chem. Soc.*, 126, 2686–2687, doi:10.1021/ja0386391, available online at: <http://dx.doi.org/10.1021/ja0386391>, 2004. 25555, 25556

ACPD

10, 25545–25576, 2010

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.




Fig. 1. Top, time series of inorganic and, below that, organic aerosol loading as measured by the AMS. The organic time series also details the *m/z* 82 containing PMF factor (82Fac). On the bottom are bars showing: the mean aerosol composition (with same colour scheme as in the time series legend); an expansion of the organic component to show the individual PMF factors; and an expansion of the 82Fac to show the contribution from MF which is calculated from the $C_5H_6O^+$ signal using Eq. (1).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.




Fig. 2. Various AMS mass spectra showing a MF peak at m/z 82, including (a) a C-AMS mass spectrum measured aboard the FAAM aircraft during a period of high m/z 82 signal, (b) a high resolution mass spectrum, with inset detailing the m/z 82 peak fits and indicating major peaks, measured from the ground site during a period of high MF signal, (c) the 82Fac organic aerosol factor from PMF analysis of the ground site mass spectra and (d) a mass spectrum of 3MF suspended in water and measured in the laboratory.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. GC×GC chromatogram of m/z 82. The dominant peak is based on methylfuran.

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.




Fig. 5. AMS m/z 82 aerosol loading (in $\mu\text{g sm}^{-3}$, defined as $\mu\text{g m}^{-3}$ at 273 K and 1013 hPa with no condensation or evaporation) vs. PTRMS gas phase MVK+MACR as measured in the boundary layer from the FAAM research aircraft. Points are median values of all boundary layer data from one flight and bars are 25th and 75th percentiles.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evidence for a significant proportion of SOA from isoprene

N. H. Robinson et al.




Fig. 6. Average altitude profiles of **(a)** m/z 82 signal, **(b)** m/z 82 signal as a percentage of total organic mass, **(c)** gas phase MVK+MACR and **(d)** organic aerosol loading as measured from the FAAM research aircraft. Red lines from data before midday and blue points from data after midday, local time. Thick lines are median values and shaded areas (outlined with thin lines) denote interquartile ranges. Points are averages of all available data over 14 flights.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
