

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

# A 6-year global cloud climatology from the Atmospheric InfraRed Sounder AIRS and a statistical analysis in synergy with CALIPSO and CloudSat

C. J. Stubenrauch, S. Cros, A. Guignard, and N. Lamquin

Laboratoire de Météorologie Dynamique, IPSL/CNRS, UMR 8539, Ecole Polytechnique, France

Received: 15 February 2010 – Accepted: 20 March 2010 – Published: 30 March 2010

Correspondence to: C. J. Stubenrauch (stubenrauch@lmd.polytechnique.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

We present a six-year global climatology of cloud properties, obtained from observations of the Atmospheric Infrared Sounder (AIRS) onboard the NASA Aqua satellite. Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) combined with CloudSat observations, both missions launched as part of the A-Train in 2006, provide a unique opportunity to evaluate the retrieved AIRS cloud properties such as cloud amount and height as well as to explore the vertical structure of different cloud types. AIRS-LMD cloud detection agrees with CALIPSO about 85% over ocean and about 75% over land. Global cloud amount has been estimated as about 66% to 74%, depending on the weighting of not cloudy AIRS footprints by partial cloud cover (0 or 0.3). 40% of all clouds are high clouds, and about 44% of all clouds are single layer low-level clouds. The “radiative” cloud height determined by the AIRS-LMD retrieval corresponds well to the height of the maximum backscatter signal and of the “apparent middle” of the cloud. Whereas the real cloud thickness of high opaque clouds often fills the whole troposphere, their “apparent” cloud thickness (at which optical depth reaches about 5) is on average only 2.5 km. The real geometrical thickness of optically thin cirrus as identified by AIRS-LMD is identical to the “apparent” cloud thickness with an average of about 2.5 km in the tropics and midlatitudes. High clouds in the tropics have slightly more diffusive cloud tops than at higher latitudes. In general, the depth of the maximum backscatter signal increases nearly linearly with increasing “apparent” cloud thickness. For the same “apparent” cloud thickness optically thin cirrus show a maximum backscatter about 10% deeper inside the cloud than optically thicker clouds. We also show that only the geometrically thickest opaque clouds and (the probably surrounding anvil) cirrus penetrate the stratosphere in the tropics.

ACPD

10, 8247–8296, 2010

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# 1 Introduction

Clouds cover more than two thirds of the Earth's surface, and hence they play a dominant role in the energy and water cycle of our planet. Satellite observations offer a unique possibility to survey cloud properties on a global and continuous scale, and their record length exceeds now more than 25 years. Within the framework of the World Climate Research Programme (WCRP), the Radiation panel of the Global Energy and Water Experiment (GEWEX) has initiated a cloud assessment to evaluate the quality of climate records of cloud properties, with special emphasis on the GEWEX cloud products from the International Satellite Cloud Climatology Project (ISCCP, Rossow and Schiffer, 1999). To resolve the diurnal cycle of clouds, ISCCP uses VIS (day only) and IR atmospheric window radiance measurements from imagers on geostationary and polar orbiting weather satellites. Time sampling is three hourly, and the initial spatial resolution of about 7 km is sampled at about 30 km. To resolve the diurnal cycle of clouds, ISCCP uses VIS (day only) and IR atmospheric window radiance measurements from imagers on geostationary and polar orbiting weather satellites. Time sampling is three hourly, and the initial spatial resolution of about 7 km is sampled at about 30 km. Cloud detection is based on space and time variability of the IR and VIS radiances. First intercomparisons of about ten different cloud climatologies have highlighted the different sensitivities of various instruments and retrieval methods (Stubenrauch et al., 2009). Compared to other passive remote sensing instruments, the high spectral resolution of IR vertical sounders leads to especially reliable properties of cirrus with optical depth as low as 0.1, day and night (e.g. Wylie et al., 1994; Ackerman et al., 1995; Smith et al., 1998; Stubenrauch et al., 1999b, 2006; Wylie and Menzel, 1999; Chung et al., 2000; Kahn et al., 2007).

CO<sub>2</sub> sensitive channels of IR vertical sounders allow the determination of cloud height and cloud emissivity of a single cloud layer (the uppermost cloud layer in the case of multi-layer cloud scenes). Radiances measured from near the centre of a CO<sub>2</sub> absorption band are only sensitive to the upper atmosphere while radiances from the

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



wings of the band (away from the band centre) successively probe lower levels of the atmosphere. The TIROS-N Operational Vertical Sounders onboard the NOAA polar satellites provide data since 1979, the Atmospheric InfraRed Sounder (AIRS) onboard Aqua since 2002 and the IR Atmospheric Sounding Interferometer (IASI) onboard METOP since 2006. The A-Train mission (Stephens et al., 2002), consisting of several passive and two active remote sensing instruments in constellation with the Aqua satellite, provides a unique possibility to explore the geometrical depth and multi-layer structure of clouds. The Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) of the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) mission (Winker et al., 2007, 2009) is even sensitive to very thin cirrus (such as subvisible cirrus with optical depth down to 0.01) and provides information on multiple cloud layers as long as clouds are optically not too thick. In the latter case, the cloud profiling radar (CPR) of the CloudSat mission (Stephens et al., 2002; Mace et al., 2007) helps to complete the information on vertical cloud layer structure. For this purpose, the Cloudsat Geometrical Profiling Product (GEOPROF; Mace et al., 2007; Marchand et al., 2008) and the CALIPSO Vertical Feature Mask (VFM, Vaughan et al., 2004) have been merged into a combined Radar-Lidar Geometrical Profile Product (Radar-Lidar GEOPROF; Mace et al., 2009).

In this article we present cloud properties, retrieved from AIRS data by a weighted  $\chi^2$  method (Stubenrauch et al., 1999a). The AIRS-LMD cloud property retrieval, first developed for tropical and subtropical latitude bands (30° N to 30° S) and presented in Stubenrauch et al. (2008), has been refined and extended to the whole globe. The retrieval is applied to all data, and then a test based on the spectral coherence of cloud emissivities, determined at wavelengths between 9 and 12  $\mu\text{m}$  by assuming the retrieved cloud pressure, decides whether the AIRS footprint is cloudy or mostly clear. Thresholds have been established by comparing clear and cloudy scenes within the AIRS footprints, distinguished by coincident CALIOP measurements.

Section 2 describes the AIRS-LMD cloud property retrieval algorithm, which makes use of retrieved atmospheric temperature and water vapour profiles of the AIRS L2 data

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Susskind et al., 2003, 2006) and of atmospheric spectral transmissivity profiles which have been simulated for atmospheric profiles of the Thermodynamic Initial Guess Retrieval (TIGR) data base (Chédin et al., 1985; Chevallier et al., 1998). AIRS data have been collocated with CALIPSO data and then with the Radar-Lidar GEOPROF data.

5 The latter complete the information on vertical cloud layer structure, when the lidar cannot completely penetrate the whole cloud column. These data are used to choose tests to determine the AIRS cloud amount and to evaluate the AIRS cloud height. Section 3 gives an overview of average cloud properties of the 6-year AIRS-LMD cloud climatology. High clouds are studied in more detail in Sect. 4, in combination with CALIPSO and Radar-Lidar GEOPROF data. Conclusions are drawn in Sect. 5.

## 2 Data and methods

### 2.1 AIRS data

15 Launched in May 2002 onboard the Earth Observing System (EOS) platform Aqua, the AIRS instrument (Aumann et al., 2003; Chahine et al., 2006) provides very high spectral resolution measurements of Earth emitted radiation in three spectral bands (3.74–4.61  $\mu\text{m}$ , 6.20–8.22  $\mu\text{m}$  and 8.80–15.40  $\mu\text{m}$ ) using 2378 channels with a spectral resolution given by  $\Delta\lambda/\lambda=0.0008$ . The polar orbiting Aqua satellite provides observations at 01:30 and 13:30 local time (LT). The spatial resolution of these measurements is 13.5 km at nadir. Nine AIRS measurements (3×3) correspond to one footprint of the Advanced Microwave Sounder Unit (AMSU), also called a “golf ball”. AIRS L2 standard products include temperature at 28 pressure levels from 0.1 hPa to the surface and water vapour mixing ratios in 14 pressure layers from 50 hPa to the surface (Susskind et al., 2003, 2006). These atmospheric profiles were retrieved from cloud-cleared AIRS radiances (Chahine et al., 2006) within each AMSU footprint. Validations with radiosonde data from the NOAA-NESDIS operational meteorological database archive (Divakarla et al., 2006) and with Atmospheric Radiation Measurement (ARM)

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data (Tobin et al., 2006) have shown that the accuracy is close to 1 K in 1 km layers for temperature and better than 15% in 2 km layers for water vapour. For the cloud property retrieval we have collocated the AIRS L2 standard products (version 5) with a subset of AIRS L1B radiance measurements which have been downloaded from the NASA data pool (<http://mirador.gsfc.nasa.gov>).

## 2.2 AIRS, CALIPSO and L2 Radar-Lidar GEOPROF data and their collocation

The lidar CALIOP (Winker et al., 2007, 2009) of the CALIPSO mission provides backscatter profiles at 532 nm and at 1064 nm, at a vertical resolution of about 30 m to 60 m, respectively below and above an altitude of 8 km. The size of the lidar footprints is about 70 m×70 m. Horizontal sampling is 333 m along the track, and the distance between two orbits is about 1000 km. The CALIPSO L2 cloud data (version 2) at 5 km spatial resolution along the track provide the number of vertical cloud layers and their geometrical height at cloud top,  $z_{\text{top}}$ , and at the “apparent” cloud base,  $z_{\text{base}}^{\text{app}}$ . The “apparent” cloud base will be higher than the real cloud base in the case of optically thick clouds, because in that case the lidar signal penetrates the cloud only up to an optical depth,  $\tau_{\text{clid}}$ , of about 5 (Winker et al., 2003). Geometrical height is transformed into cloud top pressure,  $p_{\text{top}}$ , and “apparent” cloud base pressure,  $p_{\text{base}}^{\text{app}}$ , using the atmospheric profiles provided by the Global Modelling and Assimilation Office (GMAO) and available in the CALIPSO L1 data. The pressure of the “apparent middle” of the cloud is then:  $p_{\text{mid}}=0.5(p_{\text{top}} + p_{\text{base}}^{\text{app}})$ . CALIPSO L2 cloud data also provide cloud optical depth,  $\tau_{\text{clid}}$ , and a cloud feature flag, which indicates if the cloud is opaque. In that case we have set the cloud optical depth to 5 (after discussion with D. Winker). The CALIPSO L2 cloud data also indicate at which horizontal averaging the cloud was detected (1 km, 5 km, 20 km or 80 km), and we determine the height of the maximum backscattering signal,  $z_{\text{mbsc}}$ , at 532 nm from the backscatter profiles of the CALIPSO L1 data. For a close comparison with AIRS data, we use clouds which have been detected at horizontal averaging over 5 km or less, corresponding to a sensitivity in terms of minimum detectable particle backscatter coefficient of about

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

0.0008 km<sup>-1</sup> sr<sup>-1</sup> at night and about 0.0015 km<sup>-1</sup> sr<sup>-1</sup> during day, for a cirrus with an altitude of about 12 km (Fig. 4 of Winker et al., 2009). This corresponds to clouds with  $\tau_{\text{clid}}$  larger than about 0.05 to 0.1 (Winker et al., 2008). We only use  $\tau_{\text{clid}}$  in Sect. 4, being aware that this product is still not final and has large uncertainties also due to multiple scattering corrections (e.g. Lamquin et al., 2008). CALIPSO data were obtained through the NASA Atmospheric Sciences Data Center (ASDC) by the ICARE Thematic Center created by CNES (<http://www-icare.univ-lille1.fr/>) and its interface ClimServ at IPSL (<http://climserv.ipsl.polytechnique.fr/>).

The cloud profiling radar (CPR) of the CloudSat mission (Stephens et al., 2002; Mace et al., 2007) is capable to probe optically thick cloud layers and therefore provides also in this case the correct cloud base. Combined with the information on optically thin cloud layers from CALIOP, these two instruments provide a complete vertical profiling of all clouds. The CPR footprint is about 2.5 km × 1.4 km, and it provides measurements at a vertical resolution of about 250 m. The method to merge the geometrical profiling of CALIOP and CPR (Mace et al., 2009) was designed to extract maximum information on cloud layering from the combined radar and lidar sensors. The data (version 3) have been acquired from the CloudSat data processing center (<http://www.cloudsat.cira.colostate.edu>).

All satellites of the A-Train follow each other within a few minutes. First, CALIPSO cloud properties averaged over 5 km are combined with the corresponding AIRS footprints in such a way that for each AIRS golf ball (3 × 3 AIRS footprints) three CALIPSO samples are kept, each close to the centre of an AIRS golf ball. For our comparisons, we have to keep in mind, however, that CALIPSO provides only a small sample (5 km × 90 m) of the AIRS footprint (14 km × 14 km).

In a second step we collocate these data with the complete vertical profiling of the L2 Radar-Lidar GEOPROF data. Therefore, we keep for each CPR footprint (1.4 km × 2.3 km) the information of the CALIPSO sample and of the corresponding AIRS footprint. We also add information on scene homogeneity, by using clear/cloudy information of the three CALIPSO samples and cloud type information of the nine AIRS

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

footprints within the AIRS golf ball. For the analysis of high clouds in Sect. 4 we only keep situations for which the cloud top height of the CALIPSO sample and cloud top height averaged over the CPR footprint lie within 200 m.

### 2.3 AIRS-LMD cloud property retrieval method

5 The cloud property retrieval scheme is based on a weighted  $\chi^2$  method using channels around the  $15\ \mu\text{m}$   $\text{CO}_2$  absorption band (Stubenrauch et al., 1999a), providing pressure and effective emissivity of a single cloud layer (of the uppermost cloud layer in the case of multi-layer clouds).  $\chi^2$  is computed by summation over  $N$  wavelengths  $\lambda_i$  of the  $\text{CO}_2$  absorption band around  $15\ \mu\text{m}$ .

$$10 \quad \chi^2(\rho_k) = \sum_{i=1}^N [(I_{\text{clid}}(\rho_k, \lambda_i) - I_{\text{clr}}(\lambda_i)) \cdot \varepsilon_{\text{clid}}(\rho_k) - (I_{\text{m}}(\lambda_i) - I_{\text{clr}}(\lambda_i))]^2 \cdot W^2(\rho_k, \lambda_i) \quad (1)$$

The measured radiance  $I_{\text{m}}$  is obtained from the AIRS L1B data. We have chosen AIRS channels corresponding closely to the five channels used in the TIROS-N Operational Vertical Sounder (TOVS) Path-B cloud retrieval, at wavelengths of 14.190, 14.002, 13.928, 13.279 and  $10.901\ \mu\text{m}$ , and three additional channels at 14.298, 14.094 and 15 13.239  $\mu\text{m}$  (AIRS channels 174, 193, 210, 226, 239, 355, 362 and 787). The weighting functions of these channels are shown in Fig. 1 as the derivatives of the transmission function with respect to pressure. They have been simulated by the Automatized Atmospheric Absorption Atlas (4A) radiative transfer model (Scott and Chédin, 1981; operational version available at <http://www.noveltis.net/4AOP>) for an average tropical atmosphere, taken from the TIGR data base (Chédin et al., 1985, 2003; Chevallier et al., 1998).

$I_{\text{clr}}$  is the radiance which would be measured by AIRS in the case of clear sky, and  $I_{\text{clid}}$  is the radiance emitted by a homogenous opaque single cloud layer, calculated for 25 39 assumed cloud pressure levels  $\rho_k$  above surface (984 hPa to 106 hPa). For their computation we need the AIRS L2 temperature profiles as well as atmospheric transmissivity profiles at the corresponding wavelengths for an atmospheric situation similar

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the one described by the AIRS L2 atmospheric temperature and water vapour profiles (Susskind et al., 2003). These atmospheric spectral transmissivity profiles have been simulated by the 4A radiative transfer model, separately for each satellite viewing zenith angle (up to 50°) and for about 2000 representative clear sky atmospheric temperature and humidity profiles of the TIGR data base. The proximity recognition between the AIRS L2 atmospheric profiles and the TIGR atmospheric profiles is described in detail in Stubenrauch et al. (2008). If no simultaneous AIRS L2 atmospheric profile of good quality is available (which may occur if the situation is too cloudy), a running mean average of atmospheric profiles with good quality (Tobin et al., 2006) over one week, at a spatial resolution of 1° latitude × 1° longitude, is used. The third choice is a monthly mean of atmospheric profiles with good quality, at 1° latitude × 1° longitude. For the computation of  $I_{\text{clr}}$  we also need spectral surface emissivities. These are provided for the latitude band 30° N–30° S as climatological monthly averages from three years of AIRS data (Péquignot et al., 2008), at a spatial resolution of 1° latitude × 1° longitude. For the rest of the globe we use climatological monthly averages from six years of MODIS data (Seemann et al., 2008), at a spatial resolution of 0.5° latitude × 0.5° longitude, which have then been spectrally interpolated to the AIRS channels.

By introducing empirical weights  $W(\rho_k, \lambda_j)$ , the method takes into account the vertical weighting of the different channels, the growing uncertainty in the computation of  $\varepsilon_{\text{cld}}$  with increasing  $\rho_k$  and uncertainties in atmospheric profiles. These weights are determined for each of the five TIGR air mass classes (tropical, midlatitude summer and winter, polar summer and winter) as in Eqs. (8) and (10) of (Stubenrauch et al., 1999a). Minimizing  $\chi^2$  in Eq. (1) is equivalent to  $d\chi^2/d\varepsilon_{\text{cld}}=0$ , from which one can extract  $\varepsilon_{\text{cld}}$  as:

$$\varepsilon_{\text{cld}}(\rho_k) = \frac{\sum_{i=1}^N [I_m(\lambda_i) - I_{\text{clr}}(\lambda_i)] \cdot [I_{\text{cld}}(\rho_k, \lambda_i) - I_{\text{clr}}(\lambda_i)] \cdot W^2(\rho_k, \lambda_i)}{\sum_{i=1}^N [I_{\text{cld}}(\rho_k, \lambda_i) - I_{\text{clr}}(\lambda_i)]^2 \cdot W^2(\rho_k, \lambda_i)} \quad (2)$$

It is important to allow values larger than 1, because at larger pressure levels  $I_{\text{clr}}$  and  $I_{\text{clid}}$  become very similar and their uncertainties can lead to values larger than 1 (Stubenrauch et al., 1999a). When the  $\chi^2$  method leads to a non-acceptable value of  $\varepsilon_{\text{clid}}$  (larger than 1.5), the scene is set to clear sky.

Cloud temperature  $T_{\text{clid}}$  is determined from  $p_{\text{clid}}$ , using the AIRS L2 temperature profile. Cloud types are distinguished according to  $p_{\text{clid}}$  and  $\varepsilon_{\text{clid}}$ . High clouds are defined by  $p_{\text{clid}} < 440$  hPa, midlevel clouds by  $440 \text{ hPa} < p_{\text{clid}} < 680$  hPa and low clouds by  $p_{\text{clid}} > 680$  hPa. High clouds may be further distinguished into opaque ( $\varepsilon_{\text{clid}} > 0.95$ ), cirrus ( $0.95 > \varepsilon_{\text{clid}} > 0.50$ ) and thin cirrus ( $\varepsilon_{\text{clid}} < 0.50$ ). The transformation of  $p_{\text{clid}}$  into cloud altitude  $z_{\text{clid}}$  makes use of the virtual temperature profile determined from the AIRS L2 temperature and water vapour profiles.

The retrieval is applied to all AIRS footprints. Then a test based on the spectral coherence of retrieved cloud emissivities decides whether the AIRS footprint is cloudy (overcast or mostly cloudy) or clear (or not cloudy enough to determine reliably the cloud properties). Thresholds have been established by comparing clear and cloudy scenes within the AIRS footprints, distinguished by coincident CALIOP measurements (see next section).

When extending the cloud property retrieval to the whole globe, we have revised the algorithm presented in Stubenrauch et al. (2008) in several ways:

- Instead of five channels along the  $\text{CO}_2$  absorption band we now use eight channels.
- Therefore, we have increased the vertical resolution of possible pressure levels for clouds (between 984 hPa and 106 hPa) from 29 to 39 levels.
- Since the bias corrections between observed and simulated brightness temperatures for the chosen channels are small (less than 0.5 K), we do not apply them anymore.
- When evaluating the cloud altitude of high clouds in the midlatitudes, using simultaneous CALIPSO data (see below), we have discovered that for few cases

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


the AIRS-LMD cloud altitude happened to be higher than the CALIPSO cloud altitude. This happened in cases where the effective cloud emissivity in Eq. (2) did only very slightly change from one pressure level to the next. The pressure level with minimum  $\chi^2$  and second minimum  $\chi^2$  were very close together. For these cases we now determine the cloud level as the average of all cloud levels for which the retrieved effective cloud emissivity lies within 0.1.

- Detection of cloudy scenes now depends on the spectral variability of the retrieved cloud emissivity within six wavelengths instead of on a cloud emissivity difference between two wavelengths (see next section).
- Instead of considering only clouds with  $T_{\text{cld}} - T_{\text{surf}}(\text{air}) < -4.5 \text{ K}$ , we consider all clouds over ocean and clouds with  $T_{\text{cld}} - T_{\text{surf}}(\text{air}) < -3 \text{ K}$  over land.

The impact of these changes, however, is small, as can be seen in the latitudinal averages of total, high, midlevel and low-level cloud amounts shown in Fi. 7 (described in Sect. 3).

## 2.4 Determination of AIRS-LMD clear and cloudy scenes

To distinguish cloudy from clear sky scenes in a way which does not depend on regionally and seasonally varying cloud detection thresholds, we have developed a method which is applied *after* the cloud property retrieval. It is based on the spectral coherence of retrieved cloud emissivities. Therefore, for each AIRS footprint cloud emissivities  $\varepsilon_{\text{cld}}$  are determined at six wavelengths  $\lambda_i = 11.85, 10.90, 10.69, 10.40, 10.16, 9.12 \mu\text{m}$  as:

$$\varepsilon_{\text{cld}}(\lambda_i) = \frac{I_m(\lambda_i) - I_{\text{clr}}(\lambda_i)}{I_{\text{cld}}(\rho_{\text{cld}}, \lambda_i) - I_{\text{clr}}(\lambda_i)} \quad (3)$$

where  $I_{\text{cld}}$  is now determined for  $\rho_{\text{cld}}$  which has been retrieved by the  $\chi^2$  method (see above). When  $\rho_{\text{cld}}$  is well determined, the cloud emissivities should only slightly differ between 9 and  $12 \mu\text{m}$ . The variability should be larger, when the footprint is partly

cloudy or clear and hence the cloud pressure could not be well determined. In this case, the footprint is declared as not cloudy. Figure 2 presents distributions of the standard deviation  $\sigma(\varepsilon_\lambda)$  over the six wavelengths divided by the retrieved  $\varepsilon_{\text{cloud}}$ , separately for cloudy scenes and for clear sky scenes as determined by CALIPSO. Cloudy/clear scenes are situations for which all three CALIPSO samples within the AIRS golf ball are cloudy/clear. Distributions are shown for clouds which have been determined by the  $\chi^2$  method as low or high clouds ( $p_{\text{cloud}} > 680$  hPa or  $p_{\text{cloud}} < 440$  hPa) and separately for clouds determined as midlevel clouds ( $440 \text{ hPa} < p_{\text{cloud}} < 680$  hPa) over surfaces not covered by snow or ice as well as for all clouds over snow and ice regions, using observations at 01:30 LT. A microwave flag giving information on snow and ice is provided by the AIRS L2 data. We observe that in general the distributions are narrower for cloudy scenes than for clear sky (or partly cloudy scenes), with a relatively good separation when using a threshold of 0.2 for low and high clouds and of 0.1 for midlevel clouds. However, the discrimination is much less pronounced when the surface is covered by snow or ice. In this case we have explored another variable: the brightness temperature difference between  $11 \mu\text{m}$  and an average of four channels around  $7 \mu\text{m}$  (at 7.180, 7.183, 7.223 and  $7.240 \mu\text{m}$ ). The first wavelength corresponds to an atmospheric window, whereas the latter correspond to the absorption band of water vapour. In general, one would expect positive differences, because the second brightness temperature is reduced by the absorption of water vapour in the atmosphere. In the case of cold temperatures and dry air, predominant in polar regions, atmospheric inversions would lead to a negative difference (Liu and Key, 2003). From Fig. 2 we discriminate cloudy from clear scenes by using a threshold of  $-5 \text{ K}$ . A similar test is applied in the MODIS cloud retrieval (Frey et al., 2008) during polar night. Indeed, when comparing the distributions for observations at 13:30 LT in Fig. 3, the brightness temperature difference is less pertinent, because temperature inversions occur mostly during night and early morning. When applying these thresholds, we observed that especially over land there were clouds with a temperature close to the surface air temperature. By exploring distributions of  $T_{\text{cloud}} - T_{\text{surf}}(\text{air})$  (not shown), we discovered that these scenes correspond

**6-year global cloud climatology from AIRS**

C. J. Stubenrauch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

mostly to CALIPSO clear sky scenes. Therefore, we apply over land an additional test which resets to clear sky all cloudy situations with  $T_{\text{cld}} - T_{\text{surf}}(\text{air}) > -3 \text{ K}$ .

In the following, we summarize the tests for the AIRS-LMD cloud detection.

The AIRS footprint is cloudy if the following conditions are fulfilled:

5  $\varepsilon_{\text{cld}} > 0.05$

for regions not covered by snow or sea ice:

$$\sigma(\varepsilon_{\lambda})/\varepsilon_{\text{cld}} < 0.2 \quad \text{if} \quad p_{\text{cld}} < 440 \text{ hPa} \quad \text{or} \quad p_{\text{cld}} > 680 \text{ hPa}$$

$$\sigma(\varepsilon_{\lambda})/\varepsilon_{\text{cld}} < 0.1 \quad \text{if} \quad 440 \text{ hPa} < p_{\text{cld}} < 680 \text{ hPa}$$

or for regions covered by snow or sea ice:

10  $TB(11 \mu\text{m}) - TB(7 \mu\text{m}) > -5 \text{ K}$

$$\sigma(\varepsilon_{\lambda})/\varepsilon_{\text{cld}} < 0.3$$

and over land or snow:

$$T_{\text{cld}} - T_{\text{surf}}(\text{air}) < -3 \text{ K}$$

15 This “a posteriori” cloud detection leads to an agreement with the CALIPSO cloud detection (at a horizontal averaging of 5 km or less) of about 82.7%/90.3%/93.8% over ocean and 76.2%/73.4%/84.2% over land, respectively in tropical latitudes, midlatitudes and polar latitudes and of about 83.0%/78.8% over sea ice and 71.8%/69.4% over snow, respectively in midlatitudes and polar latitudes for observations at 01:30 LT and of about 69.2%/87.1%/93.2% over ocean and 84.3%/82.4 %/83.3% over land,  
20 respectively in tropical latitudes, midlatitudes and polar latitudes and of about 84.9%/84.7% over sea ice and 77.4%/66.3% over snow, respectively in midlatitudes and polar latitudes for observations at 13:30 LT. In general these agreements are quite high, especially if we consider that CALIPSO only samples the AIRS footprint. They are slightly higher over ocean than over land.

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.5 Evaluation of AIRS-LMD cloud height

We have analyzed two years (2007–2008) of collocated AIRS CALIPSO data, separately for three latitude bands: tropical/subtropical latitudes (30° N–30° S), midlatitudes (30° N–60° N and 30° S–60° S) and polar latitudes (60° N–90° N and 60° S–90° S). Figure 4 presents normalized distributions of  $p_{\text{clid}}(\text{AIRS}) - p_{\text{mid}}(\text{CALIPSO})$ , using cloud layers detected by CALIPSO at horizontal averaging over 5 km or less (see Sect. 2.2). In the case of multiple cloud layers we choose the one which is closest in height between the highest and second highest, as in Stubenrauch et al. (2008). This is justified, because CALIPSO only samples sparsely the AIRS footprint, and AIRS could observe a mixture of both clouds. We compare the results of the AIRS-LMD cloud retrieval to those provided by the NASA AIRS L2 data. In general, all AIRS-LMD distributions peak around 0. The distribution in the polar latitudes is broader, most probably because the cloud height determination is less precise over snow and ice surfaces. Distributions for the NASA L2 cloud pressure also peak around 0, but they have large tails towards negative values. The range of retrieved  $p_{\text{clid}}$  is much smaller than for AIRS-LMD, with large negative biases for low clouds. These biases have already been shown in Kahn et al. (2008); Stubenrauch et al. (2008). In polar latitudes the peak is even shifted to –100 hPa and the tail is even larger.

In Fig. 5 we compare the AIRS-LMD cloud height with the height of the maximum backscatter signal within the cloud, determined by CALIPSO, separately for high clouds and for midlevel/low clouds. We observe that in all three latitude bands the difference distributions have again a peak around 0 and distributions for midlevel/low clouds are slightly narrower than for high clouds. This can be explained by the fact that for midlevel/low clouds  $z_{\text{mbsc}}$  is close to  $z_{\text{top}}$ , whereas high clouds can be much more diffuse, leading to much broader distributions of  $z_{\text{top}} - z_{\text{mbsc}}$ , as seen in the right panel of Fig. 5.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Average cloud properties from 2003 to 2008

In this section we give a short overview of physical cloud properties obtained from the AIRS-LMD cloud climatology. We concentrate on total cloud amount (CA) as well as on high cloud amount (HCA, clouds with  $p_{\text{clid}} < 440$  hPa), midlevel cloud amount (MCA,  $440 \text{ hPa} < p_{\text{clid}} < 680$  hPa) and low-level cloud amount (LCA,  $p_{\text{clid}} > 680$  hPa). These quantities are computed as the ratio of the number of AIRS footprints including clouds, or respectively only high clouds, midlevel clouds or low-level clouds, and the total number of AIRS measurements per  $1^\circ$  latitude  $\times$   $1^\circ$  longitude. Table 1 presents 6-year averages of these cloud amounts over the whole globe, over ocean and over land, in the midlatitudes ( $30^\circ \text{ N} - 60^\circ \text{ N}$  and  $30^\circ \text{ S} - 60^\circ \text{ S}$ ) and in the inner tropics ( $15^\circ \text{ N} - 15^\circ \text{ S}$ ). An AIRS footprint is either cloudy (CA=100%) or not cloudy (CA=0%). Because of the relatively large size of the AIRS footprints, it is more likely that not cloudy AIRS footprints are on average partly covered by clouds instead of being completely clear. Therefore, we also present (second column) cloud amounts which have been calculated by adding the not cloudy AIRS footprints with a weight of 0.3 (corresponding to about 30% cloud amount on average). This leads to a rise of global cloud amount from 68% to 75% and to a larger proportion of low clouds. Globally, there are about 10% more clouds over ocean than over land. 40% of all clouds are high clouds, and about 44% of all clouds are single layer low-level clouds. The largest fraction of high clouds is situated in the tropics and the largest fraction of single layer low-level clouds in the Southern Hemisphere midlatitudes, since single layer low-level clouds are about 10% more frequent over ocean than over land. Only about 10% of all clouds in the tropics are single layer midlevel clouds.

For comparison, Table 1 also shows results from the TOVS Path-B cloud climatology (1987–1995; Scott et al., 1999; Stubenrauch et al., 2006) as well as from an analysis of CALIPSO L2 cloud data (2007–2008) in which only the uppermost cloud layers were counted, once only those detected at horizontal averaging over 5 km or less and once all detected layers, including subvisible cirrus. Results from TOVS Path-B are very

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



similar, with slightly more high clouds than AIRS-LMD. The lidar of CALIPSO is globally 10% more sensitive to very thin cirrus (subvisible cirrus). When comparing to CALIPSO clouds detected at horizontal averaging over 5 km or less, the cloud amounts become very close to the AIRS-LMD values. Differences between summer and winter in the midlatitudes are shown in Table 2. They are stronger in the Northern Hemisphere due to more land, with about 10% more clouds in winter, appearing as midlevel and low-level clouds. This difference, however, is less pronounced (about 5%) in the CALIPSO observations.

Figure 6 presents geographical maps of CA, HCA, MCA and LCA, for January and for July. Apparent features are the InterTropical Convergence Zone (ITCZ), jet streams and winter storm tracks in the midlatitudes, as well as continental cirrus in summer. We also observe large single layer low-level cloud amount in the stratocumulus regions off the West coasts of the continents, even if this type of cloud is easier to detect by using instruments including visible channels (during daytime) or active instruments.

For a more detailed comparison with other cloud climatologies, we present in Fig. 7 latitudinal averages of these cloud amounts, again for January and July. Averages of six years AIRS-LMD (2003–2008) are compared to those of two years CALIPSO (2007–2008; all uppermost cloud layers and only those detected at horizontal averaging over 5 km or less), as well as to averages of eight years TOVS Path-B (1987–1995). Cloud detection of TOVS Path-B is based on spectral IR brightness temperature differences, also in comparison with those from the Microwave Sounding Unit (MSU). In addition, we present cloud amount averages of ISCCP (Rossow and Schiffer, 1999) for the period 1984–2004. Data and documentation are available at <http://isccp.giss.nasa.gov>. We have analyzed the monthly D2 statistics (Rossow et al., 1996). CA is given in the D2 data set after interpolation between daytime and nighttime observations. Cloud type determination is more reliable during daytime than during night, because by using “IR only” measurements semi-transparent cirrus are classified as midlevel or low-level clouds (Jin and Rossow, 1997; Stubenrauch et al., 1999b). Therefore, we present in Fig. 7 for ISCCP HCA, MCA and LCA daytime results as well as results obtained

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from averages over all observations. The latter are obtained by weighting the ISCCP daytime cloud type amounts with those from “IR only” measurements. The sum of all three gives CA for day and night. It is slightly smaller than the one which is directly given in the D2 data set.

5 Even if sampling (CALIPSO) and observation times (TOVS Path-B 07:30 and 19:30 LT; ISCCP three hourly) differ, the latitudinal behaviour of all data sets agree quite well, except the very high cloud amount from TOVS Path-B over Antarctica. Compared to CALIPSO data in this region, the TOVS Path-B cloud amount seems too high. AIRS-LMD underestimates CA in polar latitudes in winter. These regions are the most  
10 difficult for cloud detection, because of the small contrast between clouds and surface. ISCCP determines well CA in these regions because of the additional use of the 3.7  $\mu\text{m}$  channel of the Advanced Very High Resolution Radiometer (AVHRR). Between 60° N and 60° S CA of all climatologies agree very well clouds detected by CALIPSO at horizontal averaging over 5 km or less. The cloud type amount differences can be  
15 explained by instrument sensitivities: HCA of CALIPSO is about 10% larger than HCA of CALIPSO for clouds detected at horizontal averaging over 5 km or less. In the tropics, the difference can even attain 20%, suggesting that these regions are covered by more thin cirrus (e.g. Wang et al., 1996; Winker and Trepte, 1998). For all three cloud types AIRS-LMD and TOVS Path-B agree quite well with CALIPSO clouds detected at  
20 horizontal averaging over 5 km or less. Daytime cloud type amounts from ISCCP are missing in the winter hemisphere at latitudes higher than 60°, because of permanent lack of daylight. Compared to the results from the IR sounder cloud climatologies, ISCCP daytime HCA is about 15% lower in the tropics, due to the smaller sensitivity to thin cirrus. When combining ISCCP day and night observations the difference in HCA  
25 between IR sounders and ISCCP increases to 20% in the tropics. Especially if thin cirrus is overlying low clouds, this cloud type is misidentified as midlevel or low-level cloud (Jin and Rossow, 1997; Stubenrauch et al., 1999b). Therefore MCA of ISCCP appears 5 to 10% larger than that of CALIPSO. In the polar latitudes during winter, the ISCCP MCA is up to 20% larger than the one of CALIPSO.

---

**6-year global cloud  
climatology from  
AIRS**C. J. Stubenrauch et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

At last we present in Fig. 8 zonally averaged vertical distributions of relative cloud amount as obtained from AIRS-LMD, from CALIPSO (once for the uppermost cloud layer and once for all clouds detected at horizontal averaging of 5 km or less) and from Radar-Lidar GEOPROF, separately for boreal winter (left) and boreal summer (right). The CALIPSO cloud amount is presented at the height of maximum backscatter and the Radar-Lidar GEOPROF cloud amount at the height of the middle of the cloud. The data are normalized in such a way that the sum over all heights for each latitude interval corresponds to 100%. We have to keep in mind that AIRS-LMD provides only information on the uppermost cloud layers, whereas for CALIPSO and Radar-Lidar GEOPROF we have used the information on all cloud layers. Therefore the features from AIRS-LMD compare better to those from CALIPSO for the uppermost cloud layer. However, low-level clouds from the AIRS-LMD climatology seem to be situated slightly lower than those observed by CALIPSO, probably because of partly covered cloud fields. When considering all cloud layers, the fraction of low clouds increases slightly, even more with Radar-Lidar GEOPROF data. However, features look quite similar, which means that lowlevel clouds also appear as single layer clouds. All data sets show well the structure of the ITCZ with high clouds near the tropopause and only few low level clouds. The maximum of relative cloud amount from AIRS-LMD seems to be slightly lower than the one by CALIPSO or Radar-Lidar GEOPROF, because in the case of thin cirrus and thicker cirrus underneath, AIRS-LMD determines the cloud height of the cloud underneath, whereas the other data sets account for both cloud layers. Other interesting features in the midlatitudes are the winter storm tracks for which AIRS-LMD shows a full cloud column in contrast to cirrus in summer. In the polar latitudes (especially over Antarctica) in winter CALIPSO detects stratospheric clouds (as observed for example by Sassen et al., 2008) which are not determined by AIRS-LMD.

**6-year global cloud climatology from AIRS**

C. J. Stubenrauch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 4 Synergy of AIRS, CALIPSO and CloudSat: properties of high clouds

In this section we further explore the properties of high clouds, by distinguishing opaque clouds, cirrus and thin cirrus (see Sect. 2.3). Table 3 presents their amount over the whole globe, over ocean and over land, as well as in the midlatitudes and in the tropics. Whereas high opaque clouds cover only a very small area of about 3%, cirrus and thin cirrus have each about a four times larger coverage and have therefore more influence on the Earth's radiation budget. For comparison we also show results from the TOVS Path-B cloud climatology. Values are quite similar with slightly more high opaque clouds and slightly less cirrus for AIRS, especially in the tropics. This is certainly linked to the better spatial resolution of 13.5 km for the AIRS-LMD cloud retrieval compared to 100 km for TOVS Path-B.

In the following we use the collocated AIRS, CALIPSO and CloudSat data set (see Sect. 2.2), which provides, amongst other parameters, cloud type identification and “radiative” cloud height by AIRS, cloud top, height of maximum backscatter signal and “apparent” cloud base by CALIPSO as well as cloud top and “real” cloud base by CloudSat.

### 4.1 Position of maximum backscatter, geometrical and “apparent” thickness

Figure 9 presents distributions of normalized frequency of  $z_{\text{top}} - z_{\text{mbsc}}$  (left panel) and  $(z_{\text{top}} - z_{\text{mbsc}})/(z_{\text{top}} - z_{\text{base}})$  (right panel), separately for high opaque clouds, cirrus and thin cirrus in the tropics and subtropics ( $30^{\circ}\text{N}$ – $30^{\circ}\text{S}$ ), midlatitudes ( $30^{\circ}\text{N}$ – $60^{\circ}\text{N}$  and  $30^{\circ}\text{S}$ – $60^{\circ}\text{S}$ ) and polar latitudes ( $60^{\circ}\text{N}$ – $90^{\circ}\text{N}$  and  $60^{\circ}\text{S}$ – $90^{\circ}\text{S}$ ). Only AIRS scenes for which all three CALIPSO samples within the AIRS golf ball are declared cloudy and which are homogenous (same AIRS cloud type over nine AIRS footprints of the corresponding AIRS golf ball) are considered. The distributions of  $z_{\text{top}} - z_{\text{mbsc}}$  are quite similar for the three cloud types, with a peak of maximum backscatter near the cloud top. Nevertheless, high clouds seem to be slightly more diffuse in the tropics, where the distributions are broader than in the midlatitudes and in the polar latitudes. The

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



distributions of  $(z_{\text{top}} - z_{\text{mbsc}})/(z_{\text{top}} - z_{\text{base}})$  are quite different for high opaque clouds compared to thinner cirrus: the height of maximum backscatter is near the top, with very few cases of maximum backscatter height in the lower three quarters of the cloud, whereas the distributions of cirrus and thin cirrus are much broader. The difference between high opaque clouds and thinner cirrus can be explained by the much larger vertical extent of high opaque clouds (including Cumulonimbus) compared to the one of cirrus and thin cirrus, as shown in the left panel of Fig. 10. The distributions of vertical extent of high opaque clouds are quite broad (the tropical/subtropical distribution is even nearly flat), with a peak around 10 km in the midlatitudes and around 8 km in the polar latitudes. This means that these clouds may extend over the whole troposphere. Note, however, that this type of cloud only covers about 3% of the globe (Table 3). Cirrus vertical extent peaks around 4 km in the tropics, 5 km in the midlatitudes and around 8 km in the polar latitudes. The distributions of vertical extent of optically thin cirrus peak around 2 km, with a quite broad distribution in the polar latitudes. It is interesting to note that the cirrus distribution resembles more the distribution of thin cirrus in the tropics and shifts towards the one for opaque clouds towards higher latitudes. This is probably linked to the different formation processes. To highlight the importance of including the CloudSat cloud base, we present in the right panel of Fig. 10 distributions of the difference between cloud vertical extents determined by CALIPSO alone and by the CALIPSO-CloudSat GEOPROF data base. For high opaque clouds the difference may reach more than 10 km in the tropics, 7 km in the midlatitudes and 6 km in the polar latitudes, whereas for optically thin cirrus there is no difference. This result also indicates the quality of the AIRS-LMD cloud type determination.

Since for high opaque clouds the “apparent” geometrical cloud thickness (at which the cloud reaches an optical depth of 5) can be much smaller than the real geometrical cloud thickness, we present in Fig. 11 normalized distributions of this quantity separately for the three latitude bands. Indeed, the distributions lie between 1 and 7 km with a peak value around 2.5 km for all three latitude bands. It is the “apparent” geometrical cloud thickness which is more relevant for radiative effects, and therefore we also

---

**6-year global cloud  
climatology from  
AIRS**C. J. Stubenrauch et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

explore in Fig. 11 the normalized frequency distributions of the relative height of the maximum backscatter signal compared to the “apparent” cloud geometrical thickness, separately for the three different latitude bands. These distributions show again that in the tropics high opaque clouds are more diffusive, with a peak around 25% below cloud top in the tropics compared to 20% below cloud top in the other latitude bands.

## 4.2 Relationship between position of maximum backscatter, “radiative” height and “apparent” thickness

In Sect. 2.4 we have shown that in general the AIRS-LMD “radiative” cloud height compares quite well to the middle of the “apparent” geometrical cloud thickness and also to the height of the maximum backscatter, with a slightly larger dispersion. Two case studies (Holz et al., 2006) have shown that the cloud height retrieved from the Scanning High-Resolution Interferometer Sounder (S-HIS) corresponds to the level in the cloud where the Cloud Physics Lidar (CPL) integrated optical depth is approximately 1. The question in this case would be if  $\tau_{\text{cld}}=1$  is reached faster for an optically thick cloud than for an optically thin cloud which would then induce that  $z_{\text{cld}}(\text{AIRS})$  (further on  $z_{\text{AIRS}}$ ) would be closer to the cloud top for optically thick clouds than for optically thin clouds. Therefore we try to explore further the relation between the position of the maximum backscatter signal, the “radiative” height and the “apparent” cloud geometrical thickness. For a better separation between optically thick and thin clouds, we consider high opaque clouds with  $\varepsilon_{\text{cld}} \geq 0.95$ , cirrus with  $0.8 > \varepsilon_{\text{cld}} > 0.5$  and optically thin cirrus with  $0.4 > \varepsilon_{\text{cld}} > 0.05$ , and we demand  $\tau_{\text{cld}}=5$ ,  $\tau_{\text{cld}} > 0.5$  and  $\tau_{\text{cld}} < 1.5$ , respectively, to reduce AIRS misidentification due to heterogeneous scenes. For the determination of  $z_{\text{top}} - z_{\text{AIRS}}$ , we need atmospheric profiles of good quality. These are available only for optically thin cirrus. For the other cloud types, a running mean over a week or a monthly average of good quality profiles is used.

Figure 12 presents  $z_{\text{top}} - z_{\text{mbsc}}$  (left) and  $(z_{\text{top}} - z_{\text{mbsc}})/(z_{\text{top}} - z_{\text{base}}^{\text{app}})$  (right) as a function of “apparent” geometrical cloud thickness,  $z_{\text{top}} - z_{\text{base}}^{\text{app}}$ , again separately for the three

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cloud types and for the three latitude bands. We observe a nearly linear increase of the average depth of the signal of maximum backscatter with increasing “apparent” cloud geometrical thickness, reaching 2.5 km at an “apparent” geometrical cloud thickness of 6 km. High opaque clouds and cirrus show the same behaviour. There seems to be a slightly stronger increase for thin cirrus. The slope is slightly weaker in the polar latitudes. Considering the relative depth of the signal of maximum backscatter, we observe an increase of only about 10% over the range of “apparent” cloud thickness. At the same “apparent” cloud thickness the maximum backscatter of optically thin cirrus is about 10% deeper inside the cloud than for the other cloud types.

Figure 13 presents  $z_{\text{top}} - z_{\text{AIRS}}$  (left) and  $(z_{\text{top}} - z_{\text{AIRS}})/(z_{\text{top}} - z_{\text{base}}^{\text{app}})$  (right) as function of  $z_{\text{top}} - z_{\text{base}}^{\text{app}}$ , for the same cloud types and latitude bands as in Fig. 12. High opaque clouds and cirrus show an increase of  $z_{\text{top}} - z_{\text{AIRS}}$  as function of  $z_{\text{top}} - z_{\text{base}}^{\text{app}}$ , similar to the behaviour of  $z_{\text{top}} - z_{\text{mbsc}}$ . It is interesting to note that even for high opaque clouds the “radiative” cloud height depends on the “apparent” cloud thickness and can be as low as 3 km beneath the cloud top. Earlier case studies by Heymsfield et al. (1991) and Sherwood et al. (2004) have already drawn attention to this problem. On average, the “radiative” cloud height of high opaque clouds lies between 0.5 and 1.5 km below the cloud top. In comparison to Fig. 12, “radiative” cloud height seems to be about 10 to 20% lower than the position of maximum backscatter at the same “apparent” geometrical thickness. In general, the “radiative” cloud height seems to be close to the “apparent middle” of the cloud (or slightly below), independent of “apparent” geometrical cloud thickness. In polar latitudes, the spread of “radiative” height between cirrus and high opaque clouds is larger. This could perhaps be linked to the different vertical structure of the clouds and their different formation process. Further investigations are needed to draw conclusions, because in these regions the atmospheric profiles are more difficult to retrieve and the cloud height over ice has larger uncertainties.

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 4.3 Relationship between distance of cloud top to tropopause and geometrical thickness

Our collocated data set should also give some information on the height of clouds with respect to the tropopause. We are especially interested in clouds penetrating into the stratosphere. This phenomenon has been observed for deep convection in the tropics (e.g. Zhang, 1993; Gettelman et al., 2002; Hong et al., 2008). The tropics should also show widespread layers of cirrus near the tropopause. Figure 14 presents the difference between the height of the thermal tropopause, given by CALIPSO (from GMAO), and CALIPSO cloud top height as function of real geometrical cloud thickness, obtained from radar-lidar GEOPROF data, separately for high opaque clouds with  $\varepsilon_{\text{cld}} \geq 0.95$ , cirrus with  $0.8 > \varepsilon_{\text{cld}} > 0.5$  and optically thin cirrus with  $0.4 > \varepsilon_{\text{cld}} > 0.05$  identified by AIRS in the three latitude bands. In the tropics we observe that only the very geometrically thickest opaque clouds and (the probably surrounding anvil) cirrus penetrate the stratosphere. These vertically extending clouds correspond possibly to larger organized, mesoscale convective systems, which more often lead to penetrating convection than smaller, unorganized convective systems as has been shown by Rossow and Pearl (2007). Even if this is no cluster analysis as for example in Machado et al. (1993) we observe that at similar geometrical cloud thickness the cloud top of high opaque clouds (probably the convective part of a tropical cloud system) seems to be about 1 km lower than the cloud top of cirrus (probably the surrounding anvil). In the midlatitudes and polar latitudes, the cloud top height relative to the tropopause differs much less between high opaque clouds and cirrus, and there is a smooth increase with increasing geometrical cloud thickness.

## 5 Conclusions

We have presented properties of a six-year global cloud climatology from the Atmospheric InfraRed Sounder AIRS, using the AIRS LMD cloud retrieval. The retrieval is

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



based on a weighted  $\chi^2$  method on radiances around the  $15\ \mu\text{m}$   $\text{CO}_2$  absorption band. Auxiliary data are AIRS L2 atmospheric profiles and pre-computed spectral transmissivity profiles from the TIGR data set, as well as spectral surface emissivities from AIRS and MODIS. Cloud amount is essentially determined by testing the spectral coherence of retrieved cloud emissivities. In addition, clouds over land are restricted to a temperature 3 K less than the surface air temperature, and we check on atmospheric inversions over snow and ice covered surface.

This “a posteriori” cloud detection leads to an overall coincident agreement with CALIPSO of about 85% over ocean and of about 75% over land. Also the zonal averages of cloud amount agree very well with the one determined by CALIPSO, when taking into account only clouds which have been detected at a horizontal averaging of 5 km or less (and therefore excluding subvisible cirrus). Only at higher latitudes in winter, total cloud amount over snow and ice seems to be underestimated, whereas the total cloud amount of the TOVS Path-B climatology, using a multi-spectral cloud detection also based on the simultaneous use of microwave channels, and of the ISCCP climatology, using a  $3.7\ \mu\text{m}$  channel during night, compare well to the one of CALIPSO. Unfortunately, channel 7 of AMSU, with a maximum of the weighting function in the lower troposphere, was defect since the beginning of operation.

Global cloud amount has been estimated as about 66% to 74%, depending on the weighting of not cloudy AIRS footprints by partial cloud cover (0 or 0.3). There are about 10% more clouds over ocean than over land. 40% of all clouds are high clouds, and about 44% of all clouds are single layer low-level clouds. When considering also clouds detected after horizontal averaging over up to 80 km (including subvisible cirrus), global cloud amount increases to 80% and high clouds make out 50% of all clouds. The largest fraction of high clouds is situated in the tropics and the largest fraction of single layer low-level clouds in the Southern Hemisphere midlatitudes. Only about 10% of all clouds in the tropics are single layer midlevel clouds, in agreement also with earlier observations by Mace and Benson-Troth (2002). In summary, the fraction of detected cirrus depends on the instrument sensitivity: the active lidar is the most sensitive

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



instrument, followed by IR sounders. ISCCP underestimates high cloud amount, especially in the tropics where these clouds are most abundant. Thin cirrus overlying lowlevel clouds are falsely identified as midlevel clouds by ISCCP. When distinguishing high clouds by their IR emissivity, one observes that high opaque clouds only cover a very small fraction of our globe: about 3%, in agreement with ISCCP (Rossow and Schiffer, 1999).

Cloud height of the AIRS LMD cloud retrieval has been evaluated using the height of the maximum backscatter signal and of the “apparent middle” of the highest cloud layer detected by CALIPSO, at horizontal averaging over 5 km or less. All difference distributions (for high and low clouds as well as in the tropics, midlatitudes and polar latitudes) are Gaussian with a strong peak around 0. This means that the “radiative” height determined by the AIRS-LMD retrieval corresponds well to the height of the maximum backscatter signal and of the “apparent middle” of the cloud. A comparison of the cloud height provided by the AIRS L2 products with CALIPSO exhibits a strong underestimation of cloud pressure for low clouds, in agreement with a study of Kahn et al. (2008).

We used the unique constellation of the A-Train, with its two active instruments, to explore the “apparent” and the real geometrical cloud thickness, as well as the depth of maximum backscatter, indicating the diffusiveness of clouds. Whereas the real cloud thickness of high opaque clouds may fill the whole troposphere, their “apparent” cloud thickness (at which optical thickness reaches about 5) is on average only 3 km. We could also show that the real geometrical thickness of thin cirrus as identified by AIRS-LMD is identical to the “apparent” cloud thickness with an average of about 2.5 km in the tropics and midlatitudes.

In general, the depth of the maximum backscatter signal increases nearly linearly with increasing “apparent” cloud thickness. Even for high opaque clouds, the height of maximum backscatter lies on average about 35%/30% and 25% below cloud top relative to “apparent” geometrical thickness, respectively in the tropics, midlatitudes and polar latitudes. This indicates that high clouds in the tropics have slightly more

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



diffusive cloud tops than at higher latitudes. For the same “apparent” cloud thickness optically thin cirrus show a maximum backscatter about 10% deeper inside the cloud than optically thicker clouds.

The difference between cloud top and “radiative” height also increases with increasing “apparent” cloud thickness for high opaque clouds and cirrus. However, relatively speaking, the “radiative” cloud height seems to be close to the “apparent middle” of the cloud (or slightly below), independent of “apparent” geometrical cloud thickness. It also seems to be slightly lower than the position of maximum backscatter. It is interesting to note that even for high opaque clouds the “radiative” cloud height depends on the “apparent” cloud thickness and can be as low as 3 km beneath the cloud top.

When relating the distance between the tropopause and the cloud top to the real cloud thickness, we observe in the tropics that only the geometrically thickest opaque clouds and (the probably surrounding anvil) cirrus penetrate the stratosphere. These vertically extending clouds correspond possibly to larger organized, mesoscale convective systems, which lead more often to penetrating convection than smaller, unorganized convective systems as has been shown by Rossow and Pearl (2007). At similar geometrical cloud thickness the cloud top of high opaque clouds (probably the convective part of a tropical cloud system) seems to be about 1 km lower than the cloud top of cirrus (probably the surrounding anvil). In the midlatitudes and polar latitudes, the cloud top height relative to the tropopause differs much less between high opaque clouds and cirrus, and there is a smooth increase with increasing geometrical cloud thickness.

This 6-year global cloud climatology participates in the GEWEX cloud assessment (<http://ipsl.polytechnique.fr/gewexca>), and in this framework further analyses of average cloud properties as well as their regional, seasonal and interannual variations from all participating climatologies are in progress (Stubenrauch et al., 2009). The AIRS-LMD cloud climatology will be made available at <http://ara.lmd.polytechnique.fr>.

*Acknowledgements.* This work has been financially supported by CNRS and by CNES, and in part by the European Community under the contract FP6-516099. The authors thank the

**6-year global cloud climatology from AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



members of the AIRS, CALIPSO and CloudSat science teams for their efforts and cooperation in providing the data as well as the engineers and space agencies who control the quality of the data. We also want to thank our other colleagues from the Atmospheric Radiation Analysis (ARA) team at LMD for technical support. Special thanks to Dave Winker for fruitful discussions about analyzing the CALIPSO L2 data and to Gerald Mace and Donald Reinke for help with the Radar Lidar GEOPROF data.



The publication of this article is financed by CNRS-INSU.

## References

- Ackerman, S. A., Smith, W. L., Collard, A. D., Ma, X. L., Revercomb, H. E., and Knuteson, R. O.: Cirrus cloud properties derived from high-spectral resolution infrared spectrometry during FIRE II, Part II: Aircraft HIS results, *J. Atmos. Sci.*, 52, 4246–4263, 1995.
- Aumann, H. H., Chahine, M. T., Gautier, C., Goldberg, M. D., et al.: AIRS/AMSU/HSB on the Aqua mission: Design, science objectives, data products, and processing systems, *IEEE Trans. Geosci. Remote Sens.*, 41, 253–264, 2003.
- Chahine, M. T., Pagano, T. S., Aumann, H. H., Atlas, R., et al.: AIRS: Improving weather forecasting and providing new data on greenhouse gases, *B. Am. Meteorol. Soc.*, 87, 911–926, 2006.
- Chédin, A., Scott, N. A., Wahiche, C., and Moulinier, P.: The improved initialization inversion method: A high resolution physical method for temperature retrievals from satellites of the TIROS-N series, *J. Climate Appl. Meteor.*, 24, 128–143, 1985.
- Chédin, A., Serrar, S., Scott, N. A., Crevoisier, C., and Armante, R.: First global measurement of midtropospheric CO<sub>2</sub> from NOAA polar satellites: Tropical zone, *J. Geophys. Res.*, 108, D18, doi:101029/2003JD003439, 2003.

ACPD

10, 8247–8296, 2010

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chevallier, F., Cheruy, F., Scott, N. A., and Chédin, A.: A neural network approach for a fast and accurate computation of longwave radiative budget, *J. Appl. Meteor.*, 37, 1385–1397, 1998.
- Chung, S. G., Ackerman, S. A., Van Delst, P. F., and Menzel, W. P.: Model Calculations and Interferometer Measurements of Ice-Cloud Characteristics, *J. Appl. Meteor.*, 39, 634–644, 2000.
- 5 Divakarla, M. G., Barnet, C. D., Goldberg, M. D., McMillin, L. M., Maddy, E., Wolf, W., Zhou, L., and Liu, X.: Validation of Atmospheric Infrared Sounder temperature and water vapour retrievals with matched radiosonde measurements and forecasts, *J. Geophys. Res.*, 111, D9, doi:10.129/205JD006116, 2006.
- 10 Frey, R. A., Ackerman, S. A., Liu, Y., Strabala, K. I., Zhang, H., Key, J. R., and Wang, X.: Cloud Detection with MODIS. Part I: Improvements in the MODIS Cloud Mask for Collection 5, *J. Atmos. Ocean. Technol.*, 25, 1057–1072, 2008.
- Gettelman, A., Salby, M. L., and Sassi, F.: Distribution and influence of convection in the tropical tropopause region, *J. Geophys. Res.*, 107, D10, doi:10.1029/2001JD001048, 2002.
- 15 Heymsfield, G. M., Fulton, R., and Spinhirne, J. D.: Aircraft overflight measurements of Midwest severe storms: Implications on geosynchronous satellite interpretations, *Mon. Weather Rev.*, 119, 436–456, 1991.
- Holz, R. E., Ackerman, S., Antonelli, P., Nagle, F., Knuteson, R. O., McGill, M., Hlavka, D. L., and Hart, W. D.: An Improvement to the High-Spectral-Resolution CO<sub>2</sub>-Slicing Cloud-Top Altitude Retrieval, *J. Atmos. Oceanic Technol.*, 23, 653–670, 2006.
- 20 Hong, G., Heygster, G., Notholt, J., and Buehler, S. A.: Interannual to Diurnal Variations in Tropical and Subtropical Deep Convective Clouds and Convective Overshooting from Seven Years of AMSU-B Measurements, *J. Climate*, 21, 4168–4189, 2008.
- Jin, Y. and Rossow, W. B.: Detection of cirrus overlapping low-level clouds, *J. Geophys. Res.*, 25, 1727–1737, 1997.
- Kahn, B. H., Eldering, A., Braverman, A. J., Fetzer, E. J., Jiang, J. H., Fishbein, E., and Wu, D. L.: Toward the characterization of upper tropospheric clouds using Atmospheric Infrared Sounder and Microwave Limb Sounder observations, *J. Geophys. Res.*, 112, D05202, doi:10.1029/2006JD007336, 2007.
- 30 Kahn, B. H., Chahine, M. T., Stephens, G. L., Mace, G. G., Marchand, R. T., Wang, Z., Barnet, C. D., Eldering, A., Holz, R. E., Kuehn, R. E., and Vane, D. G.: Cloud type comparisons of AIRS, CloudSat, and CALIPSO cloud height and amount, *Atmos. Chem. Phys.*, 8, 1231–1248, 2008,

---

**6-year global cloud  
climatology from  
AIRS**C. J. Stubenrauch et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<http://www.atmos-chem-phys.net/8/1231/2008/>.

Lamquin, N., Stubenrauch, C. J., and Pelon, J.: Upper tropospheric humidity and cirrus thickness: a statistical analysis using one year of collocated AIRS and CALIPSO data, *J. Geophys. Res.*, 113, D00A10, doi:10.1029/2008JD009928, 2008.

Liao, X., Rossow, W. B., and Rind, D.: Comparison between SAGE II and ISCCP high-level clouds: 2. Locating cloud tops, *J. Geophys. Res.*, 100, 1137–1147, 1995.

Liu, Y. and Key, J. R.: Detection and analysis of clear-sky, low-level atmospheric temperature inversions with MODIS, *J. Atmos. Ocean. Technol.*, 20, 1727–1737, 2003.

Mace, G. G. and Benson-Troth, S.: Cloud overlap characteristics derived from long-term cloud radar data, *J. Climate*, 15, 2405–2515, 2002.

Mace, G. G., Marchand, R., Zhang, Q., and Stephens, G.: Global hydrometeor occurrence as observed by Cloudsat: Initial observations from summer 2006, *Geophys. Res. Lett.*, 34, L09808, doi:10.1029/2006GL029017, 2007.

Mace, G. G., Zhang, Q., Vaughan, M., Marchand, R., Stephens, G., Trepte, C., and Winker, D.: A description of hydrometeor layer occurrence statistics derived from the first year of merged Cloudsat and CALIPSO data, *J. Geophys. Res.*, 114, D00A26, doi:10.1029/2007JD009755, 2009.

Machado, L. A. T. and Rossow, W. B.: Structural characteristics and radiative properties of tropical cloud clusters, *Mon. Weather Rev.*, 121, 3234–3260, 1993.

Marchand, R. T., Mace, G. G., and Ackerman, T. P.: Hydrometeor detection using CloudSat: An Earth orbiting 94 GHz cloud radar, *J. Atmos. Oceanic. Technol.*, 25, 519–533, 2008.

Péquignot, E., Chédin, A., and Scott, N. A.: Infrared continental surface emissivity spectra retrieved from AIRS hyperspectral sensor, *J. Appl. Meteor. Climatol.*, 47, 1619–1633, 2008.

Rossow, W. B., Walker, A. W., Beuschel, D., and Roiter, M.: International Satellite Cloud Climatology Project (ISCCP): Description of New Cloud Datasets, WMO/TD-No.737, World Climate Research Programme (icsu and WMO), Geneva, February 1996, 115 pp., 1996.

Rossow, W. B. and Schiffer, R. A.: Advances in understanding clouds from ISCCP, *B. Am. Meteorol. Soc.*, 80, 2261–2287, 1999.

Rossow, W. B. and Pearl, C.: 22-Year survey of tropical convection penetrating into the lower stratosphere, *Geophys. Res. Lett.*, 34, L04803, doi:10.1029/2006GL028635, 2007.

Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) measurements, *J. Geophys. Res.*, 113, D00A12, doi:10.1029/2008JD009972, 2008.

**6-year global cloud climatology from AIRS**

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Scott, N. A. and Chédin, A.: A fast line-by-line method for atmospheric absorption computations: the 4A Automized Atmospheric Absorption Atlas, *J. Appl. Meteor.*, 20, 801–812, 1981.
- Scott, N. A., Chédin, A., Armante, R., Francis, J., Stubenrauch, C. J., Chaboureau, J.-P., Chevallier, F., Claud, C., and Chéruiy, F.: Characteristics of of the TO,VS Pathfinder Path-B Data set, *B. Am. Meteorol. Soc.*, 80, 2679–2701, 1999.
- Seemann, S. W., Borbas, E. E., Knuteson, R. O., Stephenson, G. R., and Huang, H. L.: Development of a Global Infrared Land Surface Emissivity Database for Application to Clear Sky Sounding Retrievals from Multispectral Satellite Radiance Measurements, *J. Appl. Meteor. Climatol.*, 47, 108, u2013123, doi:10.1175/2007JAMC1590.1, 2008.
- Smith, W. L., Ackerman, S. A., Revercomb, H., Huang, H., DeSlover, D. H., Feltz, W., Gumley, L., and Collard, A. D.: Infrared spectral absorption of nearly invisible cirrus clouds, *Geophys. Res. Lett.*, 25, 1137–1140, 1998.
- Stephens, G., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., et al.: The CloudSat mission and the A-train, *B. Am. Meteorol. Soc.*, 83, 1771–1790, 2002.
- Stubenrauch, C. J., Chédin, A., Armante, R., and Scott, N. A.: Clouds as Seen by Satellite Sounders (3I) and Imagers (ISCCP): II) A New Approach for Cloud Parameter Determination in the 3I Algorithms, *J. Climate*, 12, 2214–2223, 1999a.
- Stubenrauch, C. J., Rossow, W. B., Scott, N. A., and Chédin, A.: Clouds as seen by Infrared Sounders (3I) and Imagers (ISCCP): Part III) Spatial Heterogeneity and Radiative Effects, *J. Climate*, 12, 3419–3442, 1999b.
- Stubenrauch, C. J., Chédin, A., Rädcl, G., Scott, N. A., and Serrar, S.: Cloud properties and their seasonal and diurnal variability from TOVS Path-B, *J. Climate*, 19, 5531–5553, 2006.
- Stubenrauch, C. J., Cros, S., Lamquin, N., Armante, R., Chédin, A., Crevoisier, C., and Scott, N. A.: Cloud properties from AIRS and evaluation with CALIPSO, *J. Geophys. Res.*, 113, D00A10, doi:10.1029/2008JD009928, 2008.
- Stubenrauch, C. J., Kinne, S., and GEWEX Cloud Assessment Team: Evaluation of Global Cloud Data Products, *GEWEX news article*, 19(1), 6–7, 2009.
- Susskind, J., Barnet, C., and Blaisdell, J.: Retrieval of atmospheric and surface parameters from AIRS/AMSU/HSB data in the presence of clouds, *IEEE Trans. Geosci. Remote Sens.*, 41, 390–409, 2003.
- Susskind, J., Barnet, C., Blaisdell, J., Iredell, L., Keita, F., Kouvaris, L., Molnar, G., and Chahine, M.: Accuracy of geophysical parameters derived from AIRS/AMSU as a function of fractional cloud cover, *J. Geophys. Res.*, 111, D09S17, doi:10.1029/2005JD006272, 2006.

**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Tobin, D. C., Revercomb, H. E., Knuteson, R. O., Lesht, B. M., Strow, L. L., Hannon, S. E., Feltz, W. F., Moy, L. A., Fetzer, E. J., and Cress, T. S.: Atmospheric Radiation Measurement site atmospheric state best estimates for Atmospheric Infrared Sounder temperature and water vapour retrieval validation, *J. Geophys. Res.*, 111, D9, doi:10.1029/2005JD006103, 2006.
- 5 Vaughan, M., Young, S., Winker, D., Powell, K., Omar, A., Liu, Z., Hu, Y., and Hostetler, C.: Fully automated analysis of space-based lidar data: An overview of the CALIPSO retrieval algorithms and data products, *Proc. SPIE Int. Soc. Opt. Eng.*, 5575, 16–30, 2004.
- Wang, J., Rossow, W. B., and Zhang, Y.: Cloud Vertical Structure and its Variations from a 20-Yr Global Rawinsonde Dataset, *J. Climate*, 13, 3041–3056, 2000.
- 10 Wang, P.-H., Minnis, P., McCormick, M. P., Kent, G. S., and Skeens, K. M.: A 6-Year Climatology of Cloud Occurrence Frequency from Stratospheric Aerosol and Gas Experiment II Observations (1985–1990), *J. Geophys. Res.*, 101, 29407–29429, 1996.
- Winker, D. M. and Trepte, C. R.: Lamina cirrus observed near the tropical tropopause by LITE, *Geophys. Res. Lett.*, 25, 3351–3354, 1998.
- 15 Winker, D. M., Hunt, W. H., and McGill, M. J.: Initial performance assessment of CALIOP, *Geophys. Res. Lett.*, 34, L19803, doi:10.1029/2007GL030135, 2007.
- Winker, D., Getzewitch, B., and Vaughan, M.: Evaluation and Applications of Cloud Climatologies from CALIOP, *Proc. Int. Laser Radar Conference (ILRC)*, 2008.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., and Powell, K. A.: Overview of the CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Oceanic. Technol.*, 26, 2310–2323, 2009.
- 20 Wylie, D. P., Menzel, W. P., and Strabala, K. I.: Four years of global cirrus cloud statistics using HIRS, *J. Climate*, 7, 1972–1986, 1994.
- Wylie, D. P. and Menzel, W. P.: Eight years of cloud statistics using HIRS, *J. Climate*, 12, 170–184, 1999.
- 25 Zhang, C.: On the annual cycle in highest, coldest clouds in the tropics, *J. Climate*, 6, 1987–1990, 1993.

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

**Table 1.** Cloud amount from the AIRS-LMD cloud climatology, determined as the fraction of cloudy AIRS spots (not cloudy spots have a weight of 0). Results are also shown when in the computation not cloudy AIRS footprints are added with a weight of 0.3 (meaning that not cloudy footprints have on average 30% cloud). For further comparison are shown results from the TOVS Path-B cloud climatology (1987–1995), and results using CALIPSO considering only uppermost layers detected at horizontal averaging over 5 km or less and all uppermost layers, including subvisible cirrus. Averages are shown over the globe, separately over ocean and over land, and over NH midlatitudes (30° N–60° N), tropics (15° S–15° N) and SH midlatitudes (30° S–60° S): **(a)** total (CA), **(b)** high, **(c)** midlevel and **(d)** lowlevel.

**(a) CA (%)**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
Global	66	74	70	70	80
Global – ocean	71	76	74	74	85
Global – land	56	65	60	59	70
60° N–30° N	68	75	68	67	76
15° N–15° S	66	72	69	71	86
30° S–60° S	78	83	83	82	88

**(b) HCA/CA (%)**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
Global	40	36	43	42	50
Global – ocean	37	32	40	39	46
Global – land	46	43	50	52	60
60° N–30° N	39	34	41	45	51
15° N–15° S	58	55	62	57	66
30° S–60° S	28	26	30	34	39

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

**Table 1.** Continued.**(c) MCA/CA (%)**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
Global	18	16	20	16	12
Global – ocean	16	13	16	13	9
Global – land	24	22	28	25	19
60° N–30° N	22	19	24	19	15
15° N–15° S	11	9	10	10	5
30° S–60° S	21	18	19	16	13

**(d) LCA/CA**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
Global	42	47	44	42	38
Global – ocean	47	56	50	48	44
Global – land	31	34	30	24	22
60° N–30° N	39	45	39	36	34
15° N–15° S	31	36	34	34	28
30° S–60° S	51	56	52	51	46

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Table 2.** Cloud amount from AIRS-LMD, TOVS Path-B and CALIPSO as in Table 1, averaged over NH midlatitudes (30° N–60° N) and SH midlatitudes (30° S–60° S), separately in winter and in summer of the corresponding hemisphere: **(a)** total (CA), **(b)** high, **(c)** midlevel and **(d)** lowlevel.

**(a) CA**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
30° N–60° N win	71	79	72	70	78
30° N–60° N sum	63	70	64	66	74
30° S–60° S win	76	81	82	81	87
30° S–60° S sum	79	83	82	80	89

**(b) HCA/CA**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
30° N–60° N win	35	30	34	42	47
30° N–60° N sum	43	38	47	44	51
30° S–60° S win	26	25	32	32	37
30° S–60° S sum	30	27	29	35	39

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.

**Table 2.** Continued.**(c) MCA/CA**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
30° N–60° N win	25	22	32	19	16
30° N–60° N sum	19	17	18	18	15
30° S–60° S win	21	18	19	15	12
30° S–60° S sum	20	17	19	16	13

**(d) LCA/CA**

Latitude band	AIRS-LMD not cldy = 0	AIRS-LMD not cldy = 0.3	TOVS Path-B climatology	CALIPSO upper clds	CALIPSO upper clds + subvis Ci
30° N–60° N win	40	47	36	38	35
30° N–60° N sum	37	44	40	36	33
30° S–60° S win	52	57	50	53	50
30° S–60° S sum	51	56	54	48	47

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.


**Table 3.** Cloud amount (CA) of high opaque clouds ( $p_{\text{cld}} < 440$  hPa and  $\varepsilon_{\text{cld}} > 0.95$ ), cirrus ( $p_{\text{cld}} < 440$  hPa and  $0.5 > \varepsilon_{\text{cld}} > 0.95$ ), and thin cirrus ( $p_{\text{cld}} < 440$  hPa and  $\varepsilon_{\text{cld}} < 0.5$ ), from the AIRS-LMD cloud climatology. For comparison, results are shown in italic from the TOVS Path-B cloud climatology (1987–1995). Averages are presented over the globe, separately over ocean and over land, and over NH midlatitudes ( $30^\circ\text{N}$ – $60^\circ\text{N}$ ), tropics ( $15^\circ\text{S}$ – $15^\circ\text{N}$ ) and SH midlatitudes ( $30^\circ\text{S}$ – $60^\circ\text{S}$ ).

Latitude band	High opaque CA	Cirrus CA	Thin Ci CA
Global	3.1 <i>2.4</i>	12.2 <i>13.8</i>	12.6 <i>12.5</i>
Global – ocean	3.0 <i>1.9</i>	11.8 <i>13.7</i>	12.3 <i>12.8</i>
Global – land	3.3 <i>3.7</i>	13.2 <i>14.6</i>	13.4 <i>12.3</i>
$60^\circ\text{N}$ – $30^\circ\text{N}$	3.2 <i>2.9</i>	13.6 <i>14.0</i>	10.4 <i>10.2</i>
$15^\circ\text{N}$ – $15^\circ\text{S}$	4.5 <i>2.4</i>	16.7 <i>20.1</i>	20.9 <i>21.1</i>
$30^\circ\text{S}$ – $60^\circ\text{S}$	3.2 <i>2.3</i>	11.9 <i>14.1</i>	7.6 <i>7.2</i>

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.




**Fig. 1.** Weighting functions of eight AIRS channels, from near the centre towards the wing of the CO<sub>2</sub> absorption band around 15  $\mu\text{m}$ , used in the cloud property retrieval.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.



**Fig. 2.** Normalized distributions of spectral variability of effective cloud emissivity over six wavelengths between 9 and 12  $\mu\text{m}$  divided by cloud effective emissivity retrieved by the  $\chi^2$  method, separately for scenes declared as cloudy (full line) and as clear (broken line) by CALIPSO. Distributions are shown for observations at 01:30 LT, separately for low clouds, midlevel clouds over ocean/land and all clouds over ice/snow. Over ice/snow are also shown normalized distributions of the brightness temperature difference between 11 and 7  $\mu\text{m}$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.




**Fig. 3.** Same distributions as in lower panels of Fig. 2, but this time for observations at 13:30 LT.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.




**Fig. 4.** Normalized frequency distributions of the difference between AIRS cloud pressure and pressure of the “apparent middle” of the uppermost cloud layer detected by CALIPSO at horizontal averaging over 5 km or less. Observations at 01:30 LT. AIRS cloud retrievals from LMD are compared to those of NASA L2 products, separately for the latitude bands 30° N–30° S (upper panel), 30°–60° (middle panel) and 60°–90° (lower panel).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.



**Fig. 5.** Normalized frequency distributions of the difference between the cloud height of AIRS-LMD and the height of maximum backscatter of the uppermost cloud layer detected by CALIPSO at horizontal averaging over 5 km or less (left) and the difference between cloud top and height of maximum backscatter (right), separately for high clouds (full line) and mid-level/low clouds (broken line) as determined by CALIPSO, for the latitude bands  $30^{\circ}$  N– $30^{\circ}$  S (upper panels),  $30^{\circ}$ – $60^{\circ}$  (middle panels) and  $60^{\circ}$ – $90^{\circ}$  (lower panels).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.




**Fig. 6.** Geographical maps of total, high, midlevel and low cloud amount from AIRS-LMD, in January (left) and in July (right). Averages over the period from 2003 to 2008.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.



**Fig. 7.** Zonal averages of total, high, midlevel and low-level cloud amount from AIRS-LMD compared to results from various cloud climatologies, in January (left) and in July (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**6-year global cloud  
climatology from  
AIRS**

C. J. Stubenrauch et al.



**Fig. 8.** Zonally averaged vertical distributions of relative cloud amount as obtained from AIRS-LMD (top panel), from CALIPSO (once for the uppermost cloud layer and once for all clouds detected at horizontal averaging of 5 km or less) and from Radar-Lidar GEOPROF (bottom panel), separately for boreal winter (left) and boreal summer (right). The data are normalized in such a way that the sum over all heights for each latitude interval corresponds to 100%. Averages over the period from 2007 to 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.



**Fig. 9.** Normalized frequency distributions of the height of maximum backscatter in relation to the cloud top, separately for high opaque clouds (full line), cirrus (broken line) and thin cirrus (dotted line), as absolute difference  $z_{\text{top}} - z_{\text{mbsc}}$  (left panel) and as relative difference  $(z_{\text{top}} - z_{\text{mbsc}})/(z_{\text{top}} - z_{\text{base}})$  (right panel). Distributions are presented for three different latitude bands:  $30^{\circ}$  N– $30^{\circ}$  S (upper panels),  $30^{\circ}$ – $60^{\circ}$  (middle panels) and  $60^{\circ}$ – $90^{\circ}$  (lower panels).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.



**Fig. 10.** Normalized frequency distributions of the geometrical cloud thickness (left panel), separately for high opaque clouds (full line), cirrus (broken line) and thin cirrus (dotted line), obtained from Radar-Lidar GEOPROF data, and of the difference between geometrical cloud thickness from Radar-Lidar GEOPROF data and the one obtained from CALIPSO (right panel). Distributions are presented for three different latitude bands:  $30^{\circ}$  N– $30^{\circ}$  S (upper panels),  $30^{\circ}$ – $60^{\circ}$  (middle panels) and  $60^{\circ}$ – $90^{\circ}$  (lower panels).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## 6-year global cloud climatology from AIRS

C. J. Stubenrauch et al.



**Fig. 11.** Normalized frequency distributions of “apparent” cloud geometrical thickness (left) and of the relative height of the maximum backscatter signal compared to the “apparent” cloud geometrical thickness (right) for high opaque clouds, separately for three different latitude bands: 30° N–30° S (full line), 30°–60° (broken line) and 60°–90° (dotted line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.



**Fig. 12.** Average difference between cloud top and height of maximum backscatter (right panels) and average difference relative to “apparent” cloud geometrical thickness (left panels) as a function of “apparent” cloud geometrical thickness, separately for high opaque clouds (full circles), cirrus (open triangles) and thin cirrus (open circles) and for three different latitude bands: 30° N–30° S (upper panels), 30°–60° (middle panels) and 60°–90° (lower panels).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.



**Fig. 13.** Average difference between cloud top and “radiative” cloud height (left panels) and average difference relative to “apparent” cloud geometrical thickness (right panels) as a function of “apparent” cloud geometrical thickness, separately for high opaque clouds (full circles), cirrus (open triangles) and thin cirrus (open circles) and for three different latitude bands: 30° N–30° S (upper panels), 30°–60° (middle panels) and 60°–90° (lower panels).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



6-year global cloud  
climatology from  
AIRS

C. J. Stubenrauch et al.



**Fig. 14.** Average difference between the height of the tropopause and the cloud top as a function of real cloud geometrical thickness, separately for high opaque clouds (full circles), cirrus (open triangles) and thin cirrus (open circles) and for three different latitude bands: 30° N–30° S (upper panel), 30°–60° (middle panel) and 60°–90° (lower panel).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)