

Interactive
Comment

Interactive comment on “Formic acid above the

Jungfrauoch during 1985–2007: observed variability, seasonality, but no long-term background evolution” by R. Zander et al.

G.P. Stiller

gabriele.stiller@kit.edu

Received and published: 25 June 2010

Dear Rudy,

I am impressed on the high-quality long-term time series of HCOOH over Jungfrauoch. I would like to bring to your attention that HCOOH global distributions have been inferred from MIPAS-Envisat data for the years 2002 to 2008 and have been published recently (see: Grutter, M., N. Glatthor, G. P. Stiller, H. Fischer, U. Grabowski, M. Höpfner, S. Kellmann, A. Linden, and T. von Clar-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



mann (2010), Global distribution and variability of formic acid as observed by MIPAS-ENVISAT, *J. Geophys. Res.*, 115, D10303, doi:10.1029/2009JD012980; <http://www.agu.org/pubs/crossref/2010/2009JD012980.shtml>).

It might be interesting to you to see that MIPAS observes the seasonal cycle of HCOOH in the Northern hemisphere as well; maximum values are found at about June with ~ 100 pptv at 8 km altitude, minimum values at about December with ~ 50 pptv at 8 km, both decreasing rapidly with altitude (we have used the same spectroscopic data as you for the MIPAS retrievals).

I thought you might wish to refer to the MIPAS observations in the revised version of your paper. The MIPAS data are publicly available at <http://www.imk-asf.kit.edu/english/308.php> and are at your disposal for further comparisons.

Kind regards, Gabi Stiller

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 10, 14771, 2010.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

