

Interactive comment on “Sensitivity of a global model to the uptake of N₂O₅ by tropospheric aerosol” by H. L. Macintyre and M. J. Evans

Anonymous Referee #2

Received and published: 2 July 2010

This short paper presents a conceptual overview of the effect of revisions in the value for heterogeneous uptake coefficients for N₂O₅ on global models of NO_x, O₃ and OH. Several previous modeling papers have considered the effect of switching this chemical reaction on or off for fixed values of the heterogeneous uptake efficiency, and a few have tried to incorporate aerosol composition-dependent uptake coefficients in regional and global models. This paper is a unique contribution since it explores variation in the uptake coefficient over its entire plausible range, from very small to very high values. Although it clearly simplifies the picture considerably by fixing the value for each global simulation and neglecting any regional or seasonal variability, it does demonstrate the range over which global NO_x, O₃ and OH are sensitive to this reaction. The conclusion that the range of uptake coefficients determined from more recent laboratory and field studies is exactly the range over which these species are

C4785

most sensitive is compelling. For this reason I recommend publication in ACP.

A few comments that the authors should consider:

The paper considers competition between daytime photochemical conversion of NO_x to HNO₃ via OH and nighttime oxidation of NO_x in O₃ to produce N₂O₅. While this competition is certainly relevant – and likely an important metric in a global scale model – it neglects competition between nighttime heterogeneous reactions of N₂O₅ and those of the nitrate radical, NO₃. In fact, no details are actually given about how NO₃ chemistry is handled here. One obvious impact, for example, of changing N₂O₅ heterogeneous uptake would be the change in formation of other NO_x species, such as organic nitrates, from NO₃ oxidation reactions. Such chemistry would be less important in winter, when the largest effects due to N₂O₅ heterogeneous reaction are inferred. The authors should include some comment about variability in NO₃ chemistry and how this is handled in their model.

A related issue is the branching between HNO₃ and ClNO₂ formation in N₂O₅ hydrolysis. The latter is a relatively recent discovery – at least in terms of its potential importance in the troposphere. It is clear that this branching has not been considered in this paper. The authors should explicitly state this assumption.

Lastly, the effect of model resolution is not considered in this global-scale model study. Anthropogenic NO_x emissions tend to be highly localized, and the chemistry that leads to N₂O₅ formation is quadratic in NO₂ (i.e., two NO₂ required to form each N₂O₅). Therefore, inputs of average NO_x into a model grid cell may not represent how NO_x and N₂O₅ actually evolve overnight in the near source regions where this chemistry is likely to have the greatest impact. Such fine-scale interactions are clearly not the goal of this paper, which is intended to simulate global scale effects. However, some comment on the potential for these kinds of interactions to affect the larger scale conclusions would be appropriate.