

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Size-resolved CCN distributions

A. Bougiatioti et al.

Size-resolved CCN distributions and activation kinetics of aged continental and marine aerosol

A. Bougiatioti¹, A. Nenes^{2,3,4}, C. Fountoukis^{3,4}, N. Kalivitis¹, S. N. Pandis^{4,5}, and N. Mihalopoulos^{1,4}

¹Environmental Chemical Processes Laboratory, Department of Chemistry, University of Crete, Voutes, 71003, Heraklion, Greece

²Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

³Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

⁴Institute of Chemical Engineering and High Temperature Chemical Processes (ICE-HT), Foundation for Research and Technology Hellas (FORTH), Patras, 26504, Greece

⁵Department of Chemical Engineering, Carnegie Mellon Univ., Pittsburgh, PA 15213, USA

Received: 15 March 2011 – Accepted: 22 March 2011 – Published: 21 April 2011

Correspondence to: N. Mihalopoulos (mihalo@chemistry.uoc.gr) and A. Nenes (athanasios.nenes@gatech.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Abstract

We present size-segregated measurements of cloud condensation nucleus (CCN) activity of aged aerosol sampled at Finokalia, Crete, during the Finokalia Aerosol Measurement Experiment of summer 2007 (FAME07). From analysis of the data, hygroscopicity and activation kinetics distributions are derived. The CCN are found to be highly hygroscopic, (expressed by a size- and time-averaged hygroscopicity parameter $\kappa \sim 0.22$), with the majority of particles activating at ~ 0.5 – 0.6% supersaturation. Air masses originating from Central-Eastern Europe tend to be associated with higher CCN concentrations and slightly lower hygroscopicity ($\kappa \sim 0.18$) than for other air mass types. The particles were always well mixed, as reflected by the high activation ratios and narrow hygroscopicity distribution widths. Smaller particles (~ 30 nm) were found to be more hygroscopic (~ 0.1 κ units higher) than the larger ones (~ 100 nm). The particles with diameters less than 80 nm exhibited a diurnal hygroscopicity cycle (with κ peaking at $\sim 14:00$ h local time), consistent with photochemical aging and volatilization of less hygroscopic material from the aerosol. Use of bulk chemical composition and the aerosol number distribution results in excellent CCN closure when applying Köhler theory in its simplest form. Using asymptotic and threshold droplet growth analysis, the “aged” organics present in the aerosol were found not to suppress or delay the water uptake kinetics of particles in this environment.

1 Introduction

Predicting the global distribution of cloud condensation nuclei (CCN) is required for an improved understanding of aerosol-cloud-climate interactions (IPCC, 2007). Recent studies address the relative importance of aerosol size distribution, particle composition and mixing state on predictions of CCN concentrations (Cubison et al., 2008; Roberts et al., 2002; Ervens et al., 2005; Dusek et al., 2006; Quinn et al., 2008). These studies help evaluate theory and also quantify CCN prediction error associated with simplifications taken to calculate CCN in indirect effect studies; if propagated through

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a global model simulation of the aerosol indirect effect, this error can be expressed as an indirect forcing uncertainty (Sotiropoulou et al., 2006, 2007).

Many of the recent CCN studies have focused on the impact of organic compounds. These organics due to their chemical complexity and abundance in global ambient aerosol (Zhang et al., 2007) introduce uncertainties in predictions of CCN concentrations. Laboratory experiments suggest that the solubility of organic-rich particles in water is important for their CCN activity (Cruz and Pandis, 1997; Bilde and Svenningsson, 2004), although residual water, inorganic water uptake, mutual deliquescence and metastable states may allow sparingly-soluble organic material to be available for activation as if they were completely soluble in water (Chang et al., 2007; Raymond and Pandis, 2002; Broekhuizen et al., 2004; Padró et al., 2006; Marcolli et al., 2006; Chang et al., 2007; Raymond and Pandis, 2003; Bilde and Svenningsson, 2004). The existence of surfactants that depress surface tension has also been shown to enhance CCN activity (Kuwata et al., 2008; Moore et al., 2008; Asa-Awuku et al., 2008; Padró et al., 2010). Despite these effects, the converging view is that the compositional complexity of organic aerosol tends to yield material with little variability in hygroscopicity, so that bulk chemical metrics of the organic phase (such as oxidation state) may be a sufficient indicator of its CCN activity (Jimenez et al., 2009).

In lieu of the above, single-parameter approaches, such as the hygroscopicity parameter, κ , (Petters and Kreidenweis, 2007) have been effectively used to parameterize the cumulative effect of organics on the solute term of Köhler theory (the theoretical basis of all most approaches used to describe CCN activity). Typical values of this parameter are $\kappa = 0$ for insoluble wettable materials that do not significantly affect water activity via adsorption (Kumar et al., 2009, 2010), ~ 0.1 for oxidized secondary organic aerosols (Engelhart et al., 2008, 2011; Duplissy et al., 2008; Petters and Kreidenweis, 2007; Asa-Awuku 2008, 2009; King et al., 2010), ~ 0.6 for ammonium sulfate and nitrate, 0.95-1 for sea salt (Niedermeier et al., 2008) and ~ 1.25 for pure sodium chloride particles (Rose et al., 2010). The effective hygroscopicity parameter of multicomponent aerosols can be approximated by a volume-weighted average of the κ -values of the

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



individual chemical components (Kreidenweis et al., 2008; Gunthe et al., 2009). Continental and marine aerosols tend to cluster around $\kappa \sim 0.3$ and $\kappa \sim 0.7$, respectively (Andreae and Rosenfeld, 2008; Kreidenweis et al., 2008; Pöschl et al., 2009), with a fair amount of variability across space and time.

Size-resolved CCN activity measurements can quantify the role of composition on CCN activity. Applying this approach to Amazonian aerosol, Gunthe et al. (2009) found that the median value of the hygroscopicity parameter ($\kappa \approx 0.15$) was by a factor of two lower than the values observed for continental aerosols. Aitken mode particles were found to be less hygroscopic than accumulation mode particles, affected by the organic mass fraction which decreased with increasing particle diameter ($\sim 90\%$ in Aitken and $\sim 80\%$ in accumulation mode). During the Feldberg Aerosol Characterization Experiment (FACE-2005), Dusek et al. (2010) established that during new particle formation events, the organic fraction of particles in the Aitken mode augments and the CCN efficiency is reduced; this is consistent with secondary organic aerosol condensing upon the particles and reducing their hygroscopicity. During the CAREBeijing-2006 campaign, Wiedensohler et al. (2009) measured unusually high CCN concentrations during a new particle formation event. Size-resolved CCN measurements revealed that during periods of intense secondary aerosol formation, κ varied with particle size, from 0.1–0.3 to 0.35–0.5 in the 40–90 nm range, and ~ 0.45 for particles with ~ 190 nm diameter). During the Pearl River Delta (PRIDE-PR2006) experiment, Rose et al. (2010) characterized the aerosol near the Guangzhou megacity and found that the hygroscopicity of accumulation mode particles was generally higher than for nucleation and Aitken mode particles. During a local biomass burning event, κ dropped on average by 30% to 0.2, which can be considered characteristic for freshly emitted smoke from the burning of agricultural waste. Size-segregated CCN activity and subsaturated hygroscopic growth measurements were carried out by Cerully et al. (2011) at Hyytiälä, Finland; CCNc-derived κ values for 40–80 nm ranged between 0.10 and 0.40 with an average 0.20 ± 0.06 . Diurnal trends of κ show a minimum at sunrise and a maximum in the late afternoon; this trend was found to covary with inorganic volume fraction

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the m/z 44 organic mass fraction given by a Quadrupole Aerosol Mass Spectrometer. Size-segregated CCN activity and subsaturated hygroscopic growth measurements were carried out by Mochida et al. (2010) at Cape Hedo, Japan; the variation in the hygroscopicity of smaller range particles (Aitken-mode size) was found to be consistent with shifts in the relative abundances of ammoniated sulfate and carbonaceous material. They also found that enhancement in bulk hygroscopicity from surfactants was not important. Mochida et al. (2008) observed a rapid conversion of highly hygroscopic to intermediately hygroscopic particles, likely caused by the condensation of freshly formed secondary organics upon particles in Tokyo, Japan, significantly altering the ability of urban aerosols to form cloud droplets. Conversely, the mixing of non-hygroscopic primary organic aerosol (POA) and black carbon (BC) particles with photochemically produced hygroscopic species take place in a few hours during daytime, suggesting that away from primary sources, the number concentration of CCN may be predicted accurately enough for climate model applications assuming an internal mixture and using only the bulk chemical composition (Medina et al., 2007; Lance et al., 2009; Sotiropoulou et al., 2006, 2007; Wang et al., 2010).

Size-resolved CCN activity measurements can also be used to infer the origin of the observed hygroscopicity in ambient (and compositionally complex) aerosol samples, primarily through characterization of their water-soluble (WS) fraction. Padró et al. (2007) first introduced the method of Köhler Theory Analysis (KTA) to infer the molar volume and solubility of organic compounds from size-resolved CCN activity of aerosol generated from atomization of an aqueous solution and demonstrated its applicability to samples of known composition. Moore et al. (2008) extended KTA to include an inference of molar volume and surfactant characteristics for marine organic matter isolated from seawater collected near the Georgia coast. The analysis revealed that the solute contribution was negligible and the primary effect of the organic matter on CCN activity is through its impact on surface tension. When this analysis was applied to WS extracts of secondary organic aerosol (SOA) from biogenic and anthropogenic parent hydrocarbons (Engelhart et al., 2008, 2011; Asa-Awuku, 2009, 2010), a remarkably

constant κ emerged (0.15 ± 0.08 when surface tension effects are deconvoluted, ~ 0.3 when they are implicitly included in the κ calculation). Analysis of the WS fraction from biomass burning (Asa-Awuku et al., 2008) also revealed a similar hygroscopicity parameter ($\kappa \sim 0.3$). Padró et al. (2010) determined the hygroscopicity of the WS fraction of Mexico City aerosol collected during MILAGRO-2006; κ was 0.28 ± 0.06 and remained fairly constant regardless of location and organic fraction, reflecting a compensation between shifts in molar volume and surface tension depression. All together, these studies suggest that the hygroscopicity of oxidized organic aerosol may be predicted solely from knowledge of the water-soluble fraction of the material. Bougiatioti et al. (2009) tested this postulation for Eastern Mediterranean CCN and found that predictions of CCN notably improved when using the measured WS fraction and a prescribed $\kappa = 0.3$. Given that WS fraction of organic aerosol correlates with aerosol oxidation state (Weber et al., 2007), the constant value for WS κ would explain the linear correlation observed between the oxidation state and κ of organic aerosol (Jimenez et al., 2009).

Another uncertain aspect of cloud droplet formation is the impact of slowly-dissolving compounds, droplet surface films and aerosol amorphous states on the activation kinetics of CCN. If prevalent, these kinetic limitations could have an important impact on cloud droplet number and size distribution (e.g., Chuang et al., 1997; Nenes et al., 2001a, 2002; Lance et al., 2004). Observational studies on the activation kinetics of ambient or complex CCN (e.g., Moore et al., 2008; Ruehl et al., 2008, 2009; Sorooshian et al., 2008; Lance et al., 2009; Padró et al., 2010; Cerully et al., 2011) have tested this hypothesis by comparing the size of activated droplets (from the CCN in question) against those obtained from calibration salt aerosol (used as a standard of rapid activation). Differences in droplet sizes can be used to detect kinetic limitations and can be parameterized as changes in the water vapor uptake coefficient. Evidence to date suggests that dry organic-rich aerosol with low hygroscopicity may exhibit activation delays, while aerosol rich in inorganic salts or water-soluble organic carbon tends to exhibit rapid activation kinetics (Moore et al., 2008; Ruehl et al., 2008; Shantz

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2008; Asa-Awuku et al., 2009, 2011; Padró et al., 2010; Cerully et al., 2011). Size-resolved droplet growth analysis data provides information on the kinetic heterogeneity of CCN, allowing for the determination of kinetic parameter distributions (Ruehl et al., 2008, 2009) which would not necessarily be seen when applying “Threshold Droplet Growth Analysis” [TDGA] on polydisperse CCN measurements (Bougiatioti et al., 2009).

The Eastern Mediterranean is a climatically sensitive area of the globe. Aerosols play an important role, with a top of the atmosphere radiative cooling up to five times larger than warming from greenhouse gases (Vrekoussis et al., 2005). In the summertime, anthropogenic pollution from continental Europe (Lelieveld et al., 2002) interacts with strong mineral dust events from Northern Africa (Kalivitis et al., 2011). The combination of anthropogenic aerosol inflow, the sparsity of precipitation and persistently high RH values in the area can result in doubling (or more) of the total aerosol light scattering compared to winter (Kalivitis et al., 2007). The organic aerosol is always in a highly oxidized state, regardless of the aerosol source (Hildebrandt et al., 2010), with about 70% of the total organic mass being water-soluble (Bougiatioti et al., 2009). Earlier work has shown that most of the aerosol in this region activates at 0.6% supersaturation, characteristic of the aged nature of the aerosols; CCN calculations using the simplest form of Köhler theory (i.e., size-invariant composition, and insoluble organic fraction) agrees with measurements to within 10% (for supersaturations between 0.2 and 0.73%). Based on the aerosol “bulk” composition, an average κ of 0.24 ± 0.08 is found for the aerosol at Finokalia (Bougiatioti et al., 2009). The variation of CCN properties with size however is still unknown and was postulated to explain the supersaturation-dependent CCN closure bias seen in Bougiatioti et al. (2009). The present study addresses this issue and presents size-resolved CCN activity measurements and characteristics of Eastern Mediterranean atmospheric aerosols during the Finokalia Aerosol Measurement Experiment – 2007 (FAME07) campaign. The focus of the subsequent analysis is on the size-dependant hygroscopicity, mixing state and activation kinetics of the aerosol.

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2 Observational data set

2.1 Measurement site

Measurements took place at the Finokalia station (35°32' N, 25°67' E; <http://finokalia.chemistry.uoc.gr>) of the University of Crete and constituted part of the Finokalia Aerosol Measurement Experiment – 2007 (FAME07) campaign (Bougiatioti et al., 2009). A more detailed description of the site can be found in Mihalopoulos et al. (1997) and Sciare et al. (2003). This study focuses on the period during which size-resolved measurements of CCN were carried out (July to mid-August). HYSPLIT backtrajectory analysis (www.arl.noaa.gov/ready/hysplit4.html) showed that throughout the campaign a diversity of air masses was sampled, with influences from biomass burning events from Europe, the former Soviet Union and Asia Minor (Fig. 1a, b) to air masses with less anthropogenic influence originating from the marine boundary layer or the free troposphere (Fig. 1c, d).

2.2 Instrumentation

Prior to measurement, the ambient aerosol was dried down to ~5% relative humidity (RH) with Silica-gel diffusional dryers (Fig. 2); this ensures that particle sizing is not affected by residual aerosol water, which can bias the analysis of CCN activity data for very acidic particles (Murphy et al., 2008). The dried particles were subsequently charged by passing through a Kr-85 bipolar charger (TSI Model 3077) and classified by electrical mobility with a Differential Mobility Analyzer (DMA; TSI Model 3081). The DMA was operated at a 10:1 sheath-to-aerosol flow ratio (6 and 0.6 l min⁻¹ for the sheath and aerosol flows, respectively). The classified aerosol was then split into two streams, one sent to a Condensation Particle Counter (TSI Model 3010) to measure their concentration and the other to a Continuous Flow Streamwise Thermal Gradient CCN Chamber (CFSTGC; Roberts and Nenes, 2005; Lance et al., 2006) to measure the number of particles which act as CCN. The flows at the CPC and CFSTGC

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were monitored throughout the campaign. The activated droplets in the CFSTGC are counted and sized at the exit of its growth chamber with an Optical Particle Counter (OPC) that detects droplets and classifies them into 20 size bins (with diameter ranging from ~ 1.0 to $10\ \mu\text{m}$). During the campaign, two flow configurations were used (Fig. 2) which differed as to the point of entry of the filtered make-up air flow. In the first configuration (Fig. 2a) a make-up flow of $0.75\ \text{l min}^{-1}$ is supplied to the classified aerosol stream before splitting the stream for the CPC and the CCN instrument. The second configuration was used after 8 August (Fig. 2b) and involved introducing the make-up flow to the aerosol stream supplied to the CPC. This was done to reduce the CFSTGC measurement uncertainty due to low particle concentrations (Moore et al., 2010).

The instruments were operated in Scanning Mobility CCN Analysis mode (SMCA; Moore et al., 2010), where the DMA voltage is continuously cycled between a minimum and maximum value and the time series of CN and CCN concentrations are recorded. The particle size distribution is then obtained from the CN timeseries of a scan cycle using well-established SMPS inversion techniques (e.g., Wang and Flagan, 1989). The same inversion procedure is applied to the CCN timeseries to obtain the CCN size distribution and the size-resolved “activation fraction” (ratio of of CCN concentration to CN concentration) and the droplet size from activated CCN as a function of mobility diameter. Multiple charge corrections were applied in the CCN inversion, as detailed in Moore et al. (2010).

Throughout the campaign, the CFSTGC was operated at a total flow rate of $0.5\ \text{l min}^{-1}$, with a sheath-to-aerosol flow ratio of 10:1, and a top-bottom column difference, ΔT , between 4 and 15 K. CCN concentrations were measured at each supersaturation for 6 min, yielding size-resolved CCN activation curves that cycled through 5 different supersaturations every 30 minutes. Calibration of the instrument supersaturation was attained with the use of classified NaCl aerosol and following the procedure of Bougiatioti et al. (2009). The CCN instrument was calibrated numerous times throughout the campaign. For the lower supersaturations (0.2 and 0.38%), the relative variability in instrument supersaturation did not exceed 1% between calibrations, whereas

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

for the highest supersaturation (0.73%) the variability was under 3% (Bougiatioti et al., 2009).

2.3 Aerosol chemical composition and size distribution

The chemical composition of the aerosol was determined by analysis of polytetrafluoroethylene (PTFE) (PM₁₀, PM_{1,3} and PM_{1,3-10}) and quartz (PM₁ and PM₁₀) 4-h filter samples. PTFE and quartz filters were analyzed for water-soluble ions after extraction in ultrasonic bath with nanopure water. The solutions obtained were analyzed by ion chromatography (IC) for anions (Cl⁻, Br⁻, NO₃⁻, SO₄²⁻, C₂O₄²⁻) and cations (K⁺, Na⁺, NH₄⁺, Mg²⁺, Ca²⁺), using the procedure of Bardouki et al. (2003). Quartz filters were also analyzed for organic and elemental carbon using the OC/EC analyzer (SUNSET Laboratory Inc.) and for water-soluble organic carbon (TOC-V_{CSH}, Total Organic Carbon Analyzer, SHIMATZU). Auxiliary chemical measurements (O₃, NO_x, CO, BC, Rn-222) and meteorological parameters (wind speed and direction, pressure, temperature, relative humidity) were also continuously monitored.

A Scanning Mobility Particle Sizer (SMPS) was used to measure the particle size distribution for mobility diameters between 20 and 460 nm by placing them in 100 bins and using a sheath-to-aerosol ratio of 10:1. The SMPS consists of a scanning Differential Mobility Analyzer (DMA) running in series with a Condensation Particle Counter (CPC) for which a complete distribution is obtained every 3 min. The total aerosol concentration (CN) was measured by a Condensation Particle Counter (CPC; TSI model 3010) with a 50% particle detection efficiency at 10 nm diameter.

2.4 Analysis of SMCA data

Each constant supersaturation segment (~6-min long) was screened to minimize observational biases and examined for minimal fluctuations in the CFSTGC flow chamber temperature gradient and stability of the flows (data exhibiting maximum temperature deviation of more than 15% from the setpoint are discarded). The size-dependant

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



activation ratio function, $R_a(d_p)$, is then determined by computing the ratio of CCN over CN concentration as a function of dry particle diameter. $R_a(d_p)$ is then fit to a sigmoidal curve:

$$R_a(d_p) = \frac{\text{CCN}(d_p)}{\text{CN}(d_p)} = \frac{E}{1 + \left(\frac{d_{50}}{d_p}\right)^C} \quad (1)$$

5 where d_p is the particle dry diameter, E is the asymptote of $R_a(d_p)$ for large particle sizes, d_{50} is the characteristic dry diameter for which 50% of those particles act as CCN (and corresponds to the inflection point of the sigmoid) and C is an empirical coefficient. Eq. 1 describes the size-resolved CCN data well (e.g., Fig. 3a); the fitted sigmoids upon integration also reproduce the measured CCN concentration to within
 10 2.3% ($R^2 = 0.98$; not shown).

$R_a(d_p)$ can be used to study the size-dependent hygroscopicity and mixing state as follows. The function is evaluated at 20, 30, 40, 50, 60, 70, 80 and 100 nm for all sigmoids obtained during a supersaturation cycle. The supersaturation-dependant activation ratio function, $R_a(S)$, (the fraction of particles of size d_p that are CCN at S), is thus determined and fit to a sigmoidal function:
 15

$$R_a(S) = \frac{\text{CCN}(S, d_p = \text{const})}{\text{CN}(d_p = \text{const})} = \frac{E^*}{1 + (S/S_*)^{C^*}} \quad (2)$$

where E^* is the asymptote of the sigmoid for large supersaturations, S_* is the location of the inflection point and C^* is a fitting constant. E^* indicates the fraction of hygroscopic particles of size d_p , S_* indicates their “characteristic” (most probable) critical supersaturation, and, C is related to the extent of their chemical heterogeneity. Figure 3b presents typical examples of $R_a(S)$, for 60 and 100 nm particles, obtained at 3 August, with the error bars representing the observed daily variability.
 20

Equation (2) expresses the cumulative distribution function of critical supersaturations observed for particles of size d_p . Assuming that this distribution arises solely from

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chemical heterogeneity of the particles (for a sheath-to-aerosol ratio of 10:1 used in this study, Lance (2007) showed this to be a very good assumption for values of $C^* > -10$, as the width of the DMA transfer function introduces relatively small dispersion), one can obtain a probability distribution of hygroscopicities as follows. From the definition

5 of κ , one can express $S_* = \omega \kappa_*^{-1/2}$ and $S = \omega \kappa^{-1/2}$ where $\omega = \left(\frac{4 \left(\frac{4M_w \sigma_w}{RT \rho_w} \right)^3}{27 d_p^3} \right)^{1/2}$ and κ_* is the characteristic (most probable) hygroscopicity of the distribution. Substitution into Eq. (2) provides the hygroscopicity-dependant activation ratio function, $R_a(\kappa)$, (the fraction of particles of size d_p that are CCN with hygroscopicity κ),

$$R_a(\kappa) = \frac{\text{CCN}(\kappa, d_p = \text{const})}{\text{CN}(\kappa, d_p = \text{const})} = \frac{E^*}{1 + (\kappa_*/\kappa)^{C^*/2}} \quad (3)$$

10 Differentiation and normalization of Eq. (3) provides the probability distribution function of κ , $\rho(\kappa)$, for particles of diameter d_p (Lance et al., 2007; Cerully et al., 2011),

$$\rho(\kappa) = \frac{1}{E^*} \frac{dR_a(\kappa)}{d\kappa} = \frac{C^*}{2} \frac{\kappa_*^{C^*/2}}{\kappa^{(C^*/2+1)} \left[1 + \left(\frac{\kappa_*}{\kappa} \right)^{C^*/2} \right]^2} \quad (4)$$

from which one can compute the dispersion (standard deviation) of the hygroscopicity

parameter around κ^* as $\sigma_\kappa^2 = \frac{\int (\kappa - \kappa_*)^2 \rho(\kappa') d\kappa'}{\int \rho(\kappa') d\kappa'}$ (Lance et al., 2007; Cerully et al., 2011).

15 σ_κ describes the “chemical dispersion” or the degree of chemical heterogeneity (i.e., mixing state) for the CCN of diameter d_p . The upper limit of hygroscopicity in the integrals is set to unity (instead of ∞) to reflect the range of κ found in the atmosphere (Cerully et al., 2011). Both σ_κ and κ^* are determined using the method discussed above for each diameter and supersaturation cycle (~30 min).

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.5 Activation kinetics

CCN activation kinetics and changes thereof from the presence of organics can be determined from the droplet size of activated CCN measured at the OPC of the CFSTGC for all supersaturations considered. Most often, TDGA is applied to polydisperse CCN, which exhibit a range of critical supersaturations and droplet sizes at the exit of the CFSTGC. If activation kinetics is as fast as for the pure salt, the droplet sizes observed in the OPC will thus be equal to (or larger) than the standard for the range of supersaturations measured (Sorooshian et al., 2008; Moore et al., 2008). If the sizes of the droplets formed on ambient aerosol are statistically smaller than the sizes of those formed on pure NaCl aerosol, then the presence of organics likely delay CCN activation. In this study, shifts in activation kinetics are detected by comparing the size of the activated (on the timescale of exposure to supersaturation in the CFSTGC) ambient and $(\text{NH}_4)_2\text{SO}_4$ particles with (i) S_c equal to the instrument supersaturation, S_{in} , (i.e., for particles with diameter d_{p50}), and, (ii) particles of 100 nm dry diameter (which tend to yield droplet sizes insensitive to their critical supersaturation; Ruehl et al., 2009).

3 Results and discussion

3.1 CCN measurements

The complete time series of the CN and CCN measurements, obtained from the integration of the SMPS and CCN activity spectra can be seen in Fig. 4. The CCN generally correlate well with CN, with CCN concentrations increasing (as expected) with supersaturation. Throughout the measurement period, CCN concentrations at each supersaturation level varied by up to a factor of 7. The measured CCN concentrations at 0.51, 0.66 and 0.73% supersaturation were often very similar and close to CN concentrations (with average values of 2067 cm^{-3} , 2156 cm^{-3} and 2200 cm^{-3} , respectively). The high activation fractions at these supersaturations (0.8–1.0), and the

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

small change between 0.5% and 0.75% suggests that most particles are activated at $\sim 0.6\%$ supersaturation. This behavior is consistent with the findings of Bougiatioti et al. (2009) and reflects the characteristics of aged aerosol, where most particles are relatively large and contain significant amounts of soluble material.

CCN size distributions ($dN_{\text{CCN}}/d\log D_p$) were calculated by multiplying $R_a(d_p)$ (Eq. 1) with the corresponding aerosol particle number size distribution obtained from the SMPS (Moore et al., 2010). An example of CCN size distributions is shown in Fig. 5. As the supersaturation increases, the CCN concentrations also increase with the distributions converging towards the aerosol total number distribution. At the lowest supersaturation (0.2%), all particles with diameter above 90–120 nm activate, while all particles above 60 nm form droplets at 0.38% supersaturation.

Figure 6a shows campaign averages of $R_a(d_p)$ at different supersaturation levels. As expected, the diameter at which half of the particles activate (d_{p50}), increases with decreasing supersaturation, although $R_a(d_p)$ is remarkably consistent throughout the measurement period. The CCN activity (portrayed by the S_c versus d_{p50} relationship) for the whole measurement period is presented in Fig. 6b. The horizontal error bars represent the variability in d_{p50} throughout the study period; the variability reflects the shifts in aerosol mixing state and composition throughout the campaign. The vertical error bars denote the standard deviation of in the instrument supersaturation between calibrations. The aerosol sampled throughout the measurement period is highly CCN-active even at the lower supersaturations. Figure 6c and d show $R_a(S)$ for 40, 50, 60, 80 and 100 nm particles measured on 14 July, and on 24 July, respectively. The case of 24 July is chosen for being close to the campaign averages (within 15%) and representative of air masses from the northeast and southeast sector. For air masses coming from Central and Eastern Europe (Fig. 6c) the activation fractions are somewhat lower and do not exceed 80%, even at the higher supersaturations, even for the larger particle sizes. This is not the case for local air masses (Greece) and masses originating from the rest of the geographic sectors (Fig. 6d), where the majority of the particle sizes (≥ 50 nm) are activated and form CCN ($R_a \geq 0.8$) even at 0.5% supersaturation.

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Despite these variations, the shape of the activation curves does not change significantly which implies that the size dependence of the particle composition does not vary much over time. It is possible that the lower R_a values observed for air masses coming from Central and Eastern Europe are due to some external mixing with boundary layer aerosol (Fig. 1a). This is possible, as the particle size distributions (Fig. 7) during that day (14 July) exhibited a bimodal profile, with a pronounced maxima in the small particle range. This implies that two distinct particle types (from each mode) could be present in size-selected aerosol.

3.2 Time series and chemical composition

The complete time series of characteristic parameters derived from $R_a(S)$ (e.g., E^* , κ_* , σ_κ) and the chemical composition measured throughout the campaign are illustrated in Fig. 8. The average particulate matter concentration of the PM_1 fraction during the measurement period was of $13.75 \pm 4.8 \mu\text{g m}^{-3}$, accounting for $40 \pm 9.5\%$ of the total mass. Ammonium sulfate accounted for $89.9 \pm 4.2\%$ of the total inorganic PM_1 mass fraction, with the remaining inorganic mass consists mostly of K^+ , and on occasion, small amounts of NO_3^- and Cl^- . The sulfate volume fraction on average was 0.3 ± 0.06 (Fig. 8b) and the sulfate concentration had a campaign average of $6.6 \pm 2.8 \mu\text{g m}^{-3}$. As expected, air masses originating from Central-Eastern Europe presented a higher particle load, but with a smaller amount of sulfate ($5.3 \pm 1.6 \mu\text{g m}^{-3}$) and therefore this can explain the lower hygroscopicity values observed during that period. The particulate organic matter (OM) concentrations remained close to the mean value of $6 \pm 2.7 \mu\text{g m}^{-3}$. From the organic carbon analysis, the fine mode (PM_1) represented $72.3 \pm 17.7\%$ of the total organic carbon concentration, varying from 1.1 to $6.1 \mu\text{g m}^{-3}$. The WSOC analysis showed that $51.6 \pm 7.6\%$ of the total (bulk) organic carbon was water-soluble, with this ratio reaching $67.1 \pm 6.6\%$ in the fine fraction. Although high, this ratio is consistent with the aged nature of the aerosol (Jaffrezo et al., 2005 and references within) and is consistent with PM_1 WSOC reported for the area (Bougiatioti et al., 2009).

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effective hygroscopicity parameter κ (Fig. 8a) derived from the filter analysis was 0.22 ± 0.04 , which is similar to κ (0.24 ± 0.08) derived from PM_{10} measurements at the same location but over a different time period (Bougiatioti et al., 2009). From the SMCA analysis, we can derive a number-average κ ,

$$\bar{\kappa} = \frac{\int_0^{\infty} \kappa(d_p) n(d_p) dd_p}{\int_0^{\infty} n(d_p) dd_p} \quad (5)$$

where $n(d_p) = \left[\frac{dN}{dd_p} \right]$ is the aerosol number size distribution function (obtained from the SMPS observations). The average $\bar{\kappa}$ over the measurement period was 0.34 ± 0.04 , and is somewhat higher than the κ derived from the filter measurements, suggesting an on average higher hygroscopicity of small particles (which would affect $\bar{\kappa}$ more than the mass-weighted κ). Nevertheless, both hygroscopicity parameters are not considerably different and close to the average value of 0.3 proposed for continental locations (Andreae and Rosenfeld, 2008; Pöschl et al., 2008; Rose et al., 2008).

Figure 8c shows campaign-average values of E^* , κ^* and σ_{κ} deduced from $R_a(S)$; Fig. 8d, e present the same quantities for 14 and 24 July, respectively. Throughout the measurement period, all particles are highly hygroscopic. Larger particles exhibit somewhat larger E^* (reflective of their age, or complete activation at the maximum supersaturation measured) and lower κ^* . The latter is consistent with more sulfate condensation on the smaller particles, and confirms the postulation of Bougiatioti et al. (2009) on size-dependant hygroscopicity. The chemical dispersion, portrayed by σ_{κ} , represents the standard deviation of the fit around κ^* . It remains low and mostly constant throughout the whole measurement period for all particle sizes, consistent with the aged and well-mixed nature of the aerosol arriving at Finokalia. Comparing the characteristics of aerosol in different air masses, Central-Eastern European aerosol (Fig. 8d) exhibits 2.5% smaller hygroscopicity parameter than particles from Greece

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and eastern/southern sector (Fig. 8e). The main difference was a small decrease in the E^* values, to around 0.85. This decrease suggests that the particles exhibit a small (but detectable) chemical heterogeneity which leads to a small fraction of them not activating in the CFSTGC.

3.3 Diurnal behavior of CCN activity

The complete time series of the inferred characteristic hygroscopicity parameter κ_* , given as daily averages, for five of the dry particle sizes examined (40, 50, 60, 80 and 100 nm), is plotted in Fig. 9a. The vertical error bars represent the diurnal variability of the hygroscopicity values. We can see that the day-to-day variability of κ_* is less pronounced as we shift to larger dry particle sizes, with the smaller particles exhibiting greater hygroscopicity values (and variability) than the larger ones. After 20 July, the day-to-day variability within the dry particle sizes decreases, even at the smallest sizes. This is probably because air masses arriving at Finokalia after 20 July originated mainly from “non-polluted” areas, such as the Western Mediterranean, Eastern and Southern sector, as well as local Northern sector (Greece mainland, Southern France and Southern Italy), and, therefore, exhibited lower aerosol particle concentrations, especially at the smaller sizes.

Having in mind that the smallest activation fractions were found for air masses arriving from Central-Eastern Europe at the beginning of the measurement campaign, we examined the diurnal variability of the activated fraction as well as the hygroscopicity for the five dry particle sizes, for 14 to 15 July and for 30 to 31 July (which lies within the more “uniform” period). The results are presented in Fig. 9b, c, d and e. $R_a(S)$ substantially increases after 14:00 h local time, for all particle sizes considered. As for the hygroscopicity parameter, there is also an increase in the values after 14:00 h, but mostly only for particles <80 nm. Photochemical oxidation of gas phase chemical species and subsequent condensation of hygroscopic, low volatility products is a plausible explanation for the observed changes in E^* , κ_* . Furthermore, photochemical oxidation in the particle phase may add functional groups to the organic constituents already present,

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



making them more hygroscopic. This is consistent with the conclusions of Hildebrandt et al. (2010), who showed that the OA composition exhibits a statistically significant diurnal variation with more oxidized OA present in the afternoon (although there was no statistically significant diurnal variation observed in the bulk composition of the non-refractory submicron aerosol). The same trend is observed in the CCN activity (with an increase in κ_* values after 14:00 h) but is less pronounced (Fig. 7). Finally, the activation fraction on 30–31 July shows a slight decrease (10%) around 14:00 h local time, which may be caused by the mixing and dilution of the aerosol with air masses from aloft (Fig. 1d).

3.4 CCN closure

A closure study for the prediction of CCN concentrations is performed using different assumptions about the hygroscopic behavior of OM. The CCN concentration in each SMCA scan was calculated by integrating the CCN size distribution $R_a(d_p)n(d_p)$,

$$\text{CCN} = \int_0^{\infty} R_a(d_p)n(d_p)dd_p \quad (6)$$

The calculated CCN are then compared against predictions using κ -Köhler theory assuming a uniform composition with size and a hygroscopicity parameter $\kappa = \kappa_s \varepsilon_s + \kappa_o \varepsilon_o$, where κ_s , κ_o , is the hygroscopicity of the salt and organic fractions, respectively, and ε_s , ε_o their respective volume fraction in the dry aerosol. Three closure approaches are investigated: (a) assuming that the organic fraction is insoluble, i.e. $\kappa_o = 0$, (b) 70% of the organics are water-soluble (as suggested by a carbon mass balance of extracted filters; Bougiatioti et al., 2009) with a hygroscopicity of 0.3, i.e. $\kappa_o = 0.21 (= 0.7 \times 0.3)$, and, (c) using a prescribed CCN hygroscopicity parameter value of $\kappa = 0.34$, which is the average value for all investigated particle sizes over the whole measurement period.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1 presents the predicted against measured CCN for the whole measurement period, assuming that the organics are insoluble. Even with this simplified treatment, the CCN closure is very good (relative error $-2.8 \pm 14\%$, $R^2 = 0.93$) for the whole dataset. When the average organic water-soluble fraction is considered in the CCN calculations, the closure error is further reduced ($+1.8 \pm 12\%$) and the underprediction bias and scatter is diminished (best fit slope 0.97, $R^2 = 0.95$). The underprediction bias at the low supersaturations is also reduced at the higher supersaturations (Table 1). Using the time-dependant WSOC/OC ratio does not substantially affect the closure, as it changes CCN predictions by only 1% (not shown).

Table 1 also shows the predicted against measured CCN using a hygroscopicity parameter value of $\kappa = 0.34$. For this case also the closure is very good (relative error $2.6 \pm 22\%$, $R^2 = 0.92$). The scatter is almost twofold higher than the two cases examined above, reflective of the unresolved variability in κ , especially during the high-concentration periods (Fig. 9). Overall, the changes in the predicted κ between the 3 calculation scenarios do not affect closure considerably, largely because of the generally large CCN activation fraction and homogeneity of the aged aerosol population.

3.5 Droplet activation kinetics

Figure 10a shows the application of TDGA to the droplet growth data for the whole measurement period, for large aerosol (100 nm) while Fig. 10b shows the droplet sizes for CCN with S_c equal to the instrument supersaturation, S_{in} . The solid line represents the size of the activated calibration NaCl particles (with S_c equal to the instrument supersaturation), while the light-colored round points represent the average droplet size from the ambient aerosol activation data with the respective standard deviation. The droplet diameter from ambient large-aerosol measurements is on average above the lower limit established by the calibration aerosol. The percentile of droplets smaller than the lower limit of the calibration is $\sim 7.5\%$ at the lower supersaturations and does not exceed 12.5% at the higher supersaturations. The variability in the droplet size seen in the NaCl and ambient aerosol data is of the same order as the OPC bin resolution (0.5 μm).

The droplet sizes from d_{p50} particles exhibits larger scatter than for 100 nm aerosol (Fig. 10b). This can also be seen in Table 2, which summarizes the average values of droplet sizes for both cases. The difference ($\sim 67\%$) between the two average values (2.21 vs. 3.68 μm) at the lowest supersaturation diminishes as the latter increases ($\sim 14\%$ at the highest supersaturation). This is consistent with growth via water vapor condensation, which tends to narrow the droplet distribution as the supersaturation (and exposure time) increases. Activated CCN with $S_c \sim S_{in}$ occasionally yields droplets that are smaller than those from calibration NaCl particles. Given that $R_a(d_p)$ is steepest at d_{p50} , small fluctuations in the instrument conditions, the presence of some multiply-charged particles and the finite width of the DMA transfer function may broaden the distribution of ambient activated droplets beyond what was seen in the NaCl calibrations. Nevertheless, the percentile of droplets smaller than the ones formed from NaCl does not exceed 6.5% (of the total number) for the higher supersaturations and is $\sim 7.8\%$ for the lower supersaturations. For this, the scatter in the droplet values is most probably associated with experimental uncertainty rather than a kinetic limitation caused by the presence of the organics in the aerosol sampled.

Finally, in Fig. 10c we plot the average droplet diameter as a function of the CCN instrument supersaturation for 5 different dry particle sizes. The vertical error bars represent the variability in the average droplet diameter throughout the study period, calculated separately for each supersaturation and each dry particle diameter. The horizontal error bars represent the maximum supersaturation uncertainty (8%) quoted for the instrument (Rose et al., 2007). The droplet growth is slightly greater for larger dry particle diameters; this is because their critical supersaturation is low and generally allows more time for growth in the chamber. As the dry particle sizes augments however, their corresponding activated droplet diameter values converge, especially at the higher supersaturations. This is consistent with those droplets approaching the asymptotic limit of diffusional growth, $D_p \rightarrow 2G\tau S_{in}$, where τ is the exposure time of the CCN to supersaturation S_{in} , and, G is the water vapor mass transfer coefficient to the droplet (Nenes et al., 2001b). The convergence to a common limit for all ambient

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particles suggests that the mass transfer coefficient is the same for all particles, and implies that shifts in activation kinetics (which would manifest as changes in G) are not observed for the largest CCN.

Based on the above it can be concluded that for the range of supersaturations considered, the aged organics present in the Eastern Mediterranean aerosol do not significantly delay the CCN activation and growth process. This is consistent with the abundant amounts of sulfate and highly oxidized present in the aerosol, which tends to be associated with rapid activation kinetics (Padró et al., 2010; Engelhart et al., 2008, 2011; Asa-Awuku et al., 2009, 2010; Cerully et al., 2011). Although kinetic delays have been previously observed in aerosol with anthropogenic influence (Chang et al., 2007; Sorooshian et al., 2008; Murphy et al., 2009; Ruehl et al., 2009; Mochida et al., 2010; Asa-Awuku et al., 2011), this effect was not seen at Finokalia. This may be a consequence of the highly aged (oxidized) nature of the aerosol combined with its preconditioning (wetting) in the marine boundary layer prior to sampling.

4 Summary and conclusions

Size-resolved CCN measurements from 0.21 to 0.73% supersaturation, aerosol size distribution and chemical composition were carried out at the Finokalia measuring site of the University of Crete during the FAME07 campaign in summer 2007. The sampled aerosols arrived at the site after long-range transport with the organic fraction being mainly secondary, “aged” and highly oxidized. The majority of the particles seem to activate at a supersaturation of 0.6%. When air masses originated from Central-Eastern Europe aerosol particle concentrations were higher and the lowest hygroscopicity parameter value was observed ($\kappa = 0.18$ versus an average value of 0.22). The hygroscopicity of small particles (40 nm) is 0.1κ units higher than the one of larger particles (100 nm) with all particles being highly hygroscopic. Particles smaller than 80 nm exhibit higher hygroscopicity values after 14:00 h local time, which is consistent with the condensation of secondary aerosol and oxidation of organic constituents in the particle phase.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Application of Köhler theory using measurements of bulk composition and size distribution resulted in excellent CCN closure. Assuming that organics are insoluble, closure was attained to within $5 \pm 16\%$ for all supersaturations and with including the effects of the water-soluble organic fraction reduces the underprediction bias (to $2 \pm 7\%$) and the scatter (from $R^2 = 0.89$ to 0.91). This level of closure is consistent with the measurements carried out at Finokalia by Bougiatioti et al. (2009). Our results show a relatively lower error in closure than studies with aerosol with a variety of distinct local sources (Medina et al., 2007; Gunthe et al., 2009; Lance et al., 2009; Asa-Awuku et al., 2011), which tend to overestimate CCN concentrations (especially under polluted conditions). Finally, analysis of the droplet sizes using asymptotic and threshold droplet growth analysis suggests that the activation kinetics of ambient CCN does not vary, and is similar to NaCl calibration aerosol. This finding is consistent with evidence to date that highly oxidized aerosol grows as quickly as calibration salt aerosol. The present study results suggests that, in the case of hygroscopic aerosols, in areas not influenced by local sources, CCN activity and droplet activation kinetics can be successfully parameterized and predicted, even when size-resolved chemical composition measurements are not available.

Acknowledgements. This research project (PENED) is co-financed by E.U.-European Social Fund (75%) and the Greek Ministry of Development-GSRT (25%). A. Nenes acknowledges support from the US NSF (CAREER) and from NOAA-ACC. S. N. Pandis acknowledges support by NSF-ATM 0732598.

References

- Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth Sci. Rev.*, 89, 13–41, 2008.
- Asa-Awuku, A., Sullivan, A. P., Hennigan, C. J., Weber, R. J., and Nenes, A.: Investigation of molar volume and surfactant characteristics of water-soluble organic compounds in biomass burning aerosol, *Atmos. Chem. Phys.*, 8, 799–812, doi:10.5194/acp-8-799-2008, 2008.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Asa-Awuku, A., Engelhart, G. J., Lee, B. H., Pandis, S. N., and Nenes, A.: Relating CCN activity, volatility, and droplet growth kinetics of β -caryophyllene secondary organic aerosol, *Atmos. Chem. Phys.*, 9, 795–812, doi:10.5194/acp-9-795-2009, 2009.
- Asa-Awuku, A., Moore, R. H., Nenes, A., Bahreini, R., Holloway, J. S., Brock, C. A., Middlebrook, A. M., Ryerson, T., Jimenez, J., DeCarlo, P., Hecobian, A., Weber, R. Stickel, R., Tanner, D. J., and Huey, L. G: Airborne Cloud Condensation Nuclei Measurements during the 2006 Texas Air Quality Study, *J. Geophys. Res.*, in press, 2011.
- Bardouki, H., Liakakou, H., Economou, C., Sciare, J., Smolik, J., Zdimal, V., Eleftheriadis, K., Lazaridis, M., Dye, C., and Mihalopoulos, N.: Chemical composition of size-resolved atmospheric aerosols in the eastern Mediterranean during summer and winter, *Atmos. Environ.*, 37, 195–208, 2003.
- Bilde M. and Svenningsson, B.: CCN activation of slightly soluble organics: the importance of small amounts of inorganic salt and particle phase, *Tellus B*, 56, 128–134, 2004.
- Bougiatioti, A., Fountoukis, C., Kalivitis, N., Pandis, S. N., Nenes, A., and Mihalopoulos, N.: Cloud condensation nuclei measurements in the marine boundary layer of the Eastern Mediterranean: CCN closure and droplet growth kinetics, *Atmos. Chem. Phys.*, 9, 7053–7066, doi:10.5194/acp-9-7053-2009, 2009.
- Broekhuizen, K., Kumar P. P., and Abbatt, J. P. D.: Partially soluble organics as cloud condensation nuclei: role of trace soluble and surface active species, *Geophys. Res. Lett.*, 31, L01107, doi:10.1029/2003GL018203, 2004.
- Cerully, K.M., Raatikainen, T., Lance, S., Tkacik, D., Petäjä, T., Ehn, M., Kulmala, M., Worsnop, D. R., Laaksonen, A., Smith, J. N., and Nenes, A., Aerosol Hygroscopicity and CCN Growth Kinetics in a Boreal Forest Environment During the 2007 EUCAARI Campaign, *Atmos. Chem. Phys.*, in review, 2011.
- Chang, R. Y.-W., Liu, P. S. K., Leatch, W. R., and Abbatt, J. P. D.: Comparison between measured and predicted CCN concentrations at Egbert, Ontario: Focus on the organic aerosol fraction at the semi-rural site, *Atmos. Environ.*, 41, 8172–8182, 2007.
- Chuang, P., Charlson, R., and Seinfeld, J.: Kinetic limitations on droplet formation in clouds, *Nature*, 390 (6660), 594–596, 1997.
- Cruz, C. N. and Pandis, S. N.: A study of the ability of pure secondary organic aerosol to act as cloud condensation nuclei, *Atmos. Environ.*, 31, 2205–2214, 1997.
- Cubison, M. J., Ervens, B., Feingold, G., Docherty, K. S., Ulbrich, I. M., Shields, L., Prather, K., Hering, S., and Jimenez, J. L.: The influence of chemical composition and mixing state of

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Los Angeles urban aerosol on CCN number and cloud properties, *Atmos. Chem. Phys.*, 8, 5649–5667, doi:10.5194/acp-8-5649-2008, 2008.

Duplissy, J., Gysel, M., Alfarra, M. R., Dommen, J., Metzger, A., Prevot, A. S. H., Weingartner, E., Laaksonen, A., Raatikainen, T., Good, N., Turner, S. F., McFiggans, G., and Baltensperger, U.: Cloud forming potential of secondary organic aerosol under near atmospheric conditions, *Geophys. Res. Lett.*, 35, L03818, doi:10.1029/2007GL031075, 2008.

Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size matters more than chemistry for cloud-nucleating ability of aerosol particles, *Science*, 312, 1375–1378, 2006.

Dusek, U., Frank, G. P., Curtius, J., Drewnick, F., Schneider, J., Kürten, A., Rose, D., Andreae, M. O., Borrmann, S., and Pöschl, U.: Enhanced organic mass fraction and decreased hygroscopicity of cloud condensation nuclei (CCN) during new particle formation events, *Geophys. Res. Lett.*, 37, L03804, doi:10.1029/2009GL040930, 2010.

Engelhart, G. J., Asa-Awuku, A., Nenes, A., and Pandis, S. N.: CCN activity and droplet growth kinetics of fresh and aged monoterpene secondary organic aerosol, *Atmos. Chem. Phys.*, 8, 3937–3949, doi:10.5194/acp-8-3937-2008, 2008.

Engelhart, G. J., Moore, R. H., Nenes, A., and Pandis, S. N.: CCN Activity of Isoprene Secondary Organic Aerosol, *J. Geophys. Res.*, 116, D02207, doi:10.1029/2010JD014706, 2011.

Ervens, B., Feingold, G., and Kreidenweis, S.: Influence of water-soluble organic carbon on cloud drop number concentration, *J. Geophys. Res.*, 110, D18211, doi:10.1029/2004JD005634, 2005.

Gunthe, S. S., King, S. M., Rose, D., Chen, Q., Roldin, P., Farmer, D. K., Jimenez, J. L., Artaxo, P., Andreae, M. O., Martin, S. T., and Pöschl, U.: Cloud condensation nuclei in pristine tropical rainforest air of Amazonia: size-resolved measurements and modeling of atmospheric aerosol composition and CCN activity, *Atmos. Chem. Phys.*, 9, 7551–7575, doi:10.5194/acp-9-7551-2009, 2009.

Hennigan, C. J., Bergin, M. H., Russell, A. G., Nenes, A., and Weber, R. J.: Gas/particle partitioning of water-soluble organic aerosol in Atlanta, *Atmos. Chem. Phys.*, 9, 3613–3628, doi:10.5194/acp-9-3613-2009, 2009.

Hildebrandt, L., Engelhart, G. J., Mohr, C., Kostenidou, E., Lanz, V. A., Bougiatioti, A., DeCarlo, P. F., Prevot, A. S. H., Baltensperger, U., Mihalopoulos, N., Donahue, N. M., and Pandis, S.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N.: Aged organic aerosol in the Eastern Mediterranean: the Finokalia Aerosol Measurement Experiment 2008, *Atmos. Chem. Phys.*, 10, 4167–4186, doi:10.5194/acp-10-4167-2010, 2010.

Intergovernmental Panel on Climate Change, *Climate Change 2007: Synthesis Report*, 2007.

Jaffrezo, J.-L., Aymoz, G., Delaval, C., and Cozic, J.: Seasonal variations of the water soluble organic carbon mass fraction of aerosol in two valleys of the French Alps, *Atmos. Chem. Phys.*, 5, 2809–2821, doi:10.5194/acp-5-2809-2005, 2005.

Jimenez, J. L., Canagratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dulnea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.

Kalivitis, N., Gerasopoulos, E., Vrekoussis, M., Kouvarakis, G., Kubilay, N., Hatzianastassiou, N., Vardavas, I., and Mihalopoulos, N.: Dust transport over the eastern Mediterranean derived from Total Ozone Mapping Spectrometer, Aerosol Robotic Network, and surface measurements, *J. Geophys. Res.* 112, D03202, doi:10.1029/2006JD007510, 2007.

Kalivitis, N., Bougiatioti, A., Kouvarakis, G., and Mihalopoulos, N.: Long term measurements of atmospheric aerosol optical properties in the Eastern Mediterranean, *Atmos. Res.*, in review, 2011.

King, S. M., Rosenoern, T., Shilling, J. E., Chen, Q., Wang, Z., Biskos, G., McKinney, K. A., Pöschl, U., and Martin, S. T.: Cloud droplet activation of mixed organic-sulfate particles produced by the photooxidation of isoprene, *Atmos. Chem. Phys.*, 10, 3953–3964, doi:10.5194/acp-10-3953-2010, 2010.

Kreidenweis, S. M., Petters, M. D., and Chuang, P. Y.: Cloud particle precursors, *Clouds in the Perturbed Climate System: Their Relationship to Energy Balance, Atmospheric Dynamics, and Precipitation*. Strüngmann Forum Report, vol. 2, edited by: Heintzenberg, J. and Charlson, R. J., The MIT Press, Cambridge, MA, 2008.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kumar, P., Nenes, A., and Sokolik, I.: The Importance of Adsorption for CCN Activity and Hygroscopic Properties of Mineral Dust Aerosol, *Geophys. Res. Lett.*, 36, L24804, doi:10.1029/2009GL040827, 2009.

Kumar, P., Sokolik, I. N., and Nenes, A.: Measurements of cloud condensation nuclei activity and droplet activation kinetics of fresh unprocessed regional dust samples and minerals, *Atmos. Chem. Phys. Discuss.*, 10, 31039–31081, doi:10.5194/acpd-10-31039-2010, 2010.

Kuwata, M., Kondo, Y., Miyazaki, Y., Komazaki, Y., Kim, J. H., Yum, S. S., Tanimoto, H., and Matsueda, H.: Cloud condensation nuclei activity at Jeju Island, Korea in spring 2005, *Atmos. Chem. Phys.*, 8, 2933–2948, doi:10.5194/acp-8-2933-2008, 2008.

Lance, S., Nenes, A., and Rissman, T.: Chemical and dynamical effects on cloud droplet number: Implications for estimates of the aerosol indirect effect., *J. Geophys. Res.*, 109, D22208, doi:10.1029/2004JD004596, 2004.

Lance, S., Medina, J., Smith, J., and Nenes, A.: Mapping the operation of the DMT continuous flow CCN counter, *Aerosol Sci. Technol.*, 40, 242–254, 2006.

Lance, S.: Quantifying compositional impacts of ambient aerosol on cloud droplet formation, Doctoral Thesis, Georgia Institute of Technology, 2007.

Lance, S., Nenes, A., Mazzoleni, C., Dubey, M. K., Gates, H., Varutbangkul, V., Rissman, T. A., Murphy, S. M., Sorooshian, A., Flagan, T. A., Seinfeld, J. H., Feingold, G., and Jins-son, H.: Cloud condensation nuclei activity, closure, and droplet growth kinetics of Houston aerosol during the Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS), *J. Geophys. Res.*, 114, D00F15, doi:10.1029/2008JD011699, 2009.

Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrmann, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., De Reus, M., Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J., and Ziereis, H.: Global Air Pollution Crossroads over the Mediterranean, *Science*, 298, 794–799, 2002.

Marcollì, C. and Krieger, U. K.: Phase changes during hygroscopic cycles of mixed organic/inorganic model systems of tropospheric aerosols, *J. Phys. Chem.*, 110, 1881–1893, 2006.

Medina, J., Nenes, A., Sotiropoulou, R.-E. P., Cottrell, L. D., Ziemba, L. D., Beckman, P. J., and Griffin, R. J.: Cloud condensation nuclei closure during the International Consortium for Atmospheric Research on Transport and Transformation 2004 campaign: Effects of size-

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- resolved composition, *J. Geophys. Res.*, 112, D10S31, doi:10.1029/2006JD007588, 2007.
- Mihalopoulos, N., Stephanou, E., Kanakidou, M., Pilitsidis, S., and Bousquet, P.: Tropospheric aerosol ionic composition above the Eastern Mediterranean area, *Tellus*, 49B, 314–326, 1997.
- 5 Mochida, M., Miyakawa, T., Takegawa, N., Morino, Y., Kawamura, K., and Kondo, Y.: Significant alteration in the hygroscopic properties of urban aerosol particles by the secondary formation of organics, *Geophys. Res. Lett.*, 35, L02804, doi:10.1029/2007GL031310, 2008.
- Mochida, M., Nishita-Hara, C., Kitamori, Y., Aggarwal, S. G., Kawamura, K., Miura, K., and Takami, A.: Size-segregated measurements of cloud condensation nucleus activity and hygroscopic growth for aerosols at Cape Hedo, Japan, in spring 2008, *J. Geophys. Res.*, 115, D21207, doi:10.1029/2009JD013216, 2010.
- 10 Moore, R. H., Ingall, E. D., Sorooshian, A., and Nenes, A.: Molar mass surface tension, and droplet growth kinetics on marine organics from measurements of CCN activity, *Geophys. Res. Lett.*, 35, L07801, doi:10.1029/2008GL033350, 2008.
- 15 Moore, R., Nenes, A., and Medina, J.: Scanning Mobility CCN Analysis: A method for fast measurements of size-resolved CCN distributions and activation kinetics, *Aerosol Sci. Technol.*, 44, 861–871, 2010.
- Murphy, S. M., Agrawal, H., Sorooshian, A., Padró, L. T., Gates, H., Hersey, S., Welch, W. A., Jung, H., Miller, J. W., Cocker, D. R., Nenes, A., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Comprehensive Simultaneous Shipboard and Airborne Characterization of Exhaust from a Modern Container Ship at Sea, *Environ. Sci. Technol.*, 43, 4626–4640, 2009.
- 20 Nenes, A., Ghan, S., Abdul-Razzak, H., Chuang, P. Y., and Seinfeld, J. H.: Kinetic Limitations on Cloud Droplet Formation and Impact on Cloud Albedo, *Tellus*, 53B, 133–149, 2001a.
- Nenes, A., Chuang, P. Y., Flagan, R., and Seinfeld, J. H.: A Theoretical Analysis of Cloud Condensation Nucleus (CCN) Instruments, *J. Geophys. Res.*, 106, 3449–3474, 2001b.
- 25 Nenes, A., Charlson, R. J., Facchini, M. C., Kulmala, M., Laaksonen, A., and Seinfeld, J. H.: Can Chemical Effects on Cloud Droplet Number Rival the First Indirect Effect?, *Geophys. Res. Lett.*, 29(17), 1848, doi:10.1029/2002GL015295, 2002.
- Niedermeier, D., Wex, H., Voigtländer, J., Stratmann, F., Brüggemann, E., Kiselev, A., Henk, H., and Heintzenberg, J.: LACIS-measurements and parameterization of sea-salt particle hygroscopic growth and activation, *Atmos. Chem. Phys.*, 8, 579–590, doi:10.5194/acp-8-579-2008, 2008.
- 30 Padró, L. T. and Nenes, A.: Cloud droplet activation: solubility revisited, *Atmos. Chem. Phys.*

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Discuss., 7, 2325–2355, doi:10.5194/acpd-7-2325-2007, 2007.

Padró, L. T., Asa-Awuku, A., Morrison, R., and Nenes, A.: Inferring thermodynamic properties from CCN activation experiments: single-component and binary aerosols, *Atmos. Chem. Phys.*, 7, 5263–5274, doi:10.5194/acp-7-5263-2007, 2007.

5 Padró, L. T., Tkacik, D., Latham, T., Hennigan, C. J., Sullivan, A. P., Weber, R. J., Huey, L. G., and Nenes, A.: Investigation of cloud condensation nuclei properties and droplet growth kinetics of the water-soluble aerosol fraction in Mexico City, *J. Geophys. Res.*, 115, D09204, doi:10.1029/2009JD013195, 2010.

Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.

Pöschl, U., Rose, D., and Andreae, M. O.: Climatologies of Cloud-related Aerosols. Part 2. Particle Hygroscopicity and Cloud Condensation Nuclei Activity, *Clouds in the Perturbed Climate System: their relationship to Energy Balance, Atmospheric Dynamics and Precipitation*, Strüngmann Forum Report, vol. 2, edited by: Heizenberg, J. and Charlson, R. J., The MIT Press, Cambridge, MA, 2008.

Quinn, P. K., Bates, T. S., Coffman, D. J., and Covert, D. S.: Influence of particle size and chemistry on the cloud nucleating properties of aerosols, *Atmos. Chem. Phys.*, 8, 1029–1042, doi:10.5194/acp-8-1029-2008, 2008.

20 Raymond, T. M. and Pandis, S. N.: Cloud activation of single-component organic aerosol particles, *J. Geophys. Res.*, 107, 4787, doi:10.1029/2002JD002159, 2002.

Raymond, T. M. and Pandis, S. N.: Formation of cloud droplets by multicomponent organic particles, *J. Geophys. Res.*, 108, 4469, doi:10.1029/2003JD003503, 2003.

Roberts, G. and Nenes, A.: A continuous-flow streamwise thermal-gradient CCN chamber for atmospheric measurements, *Aerosol Sci. Technol.*, 39, 206–221, 2005.

25 Roberts, G. C., Artaxo, P., Zhou, J. C., Swietlicki, E., and Andreae, M. O.: Sensitivity of CCN spectra on chemical and physical properties of aerosol: A case study from the Amazon Basin, *J. Geophys. Res.*, 107, 8070, doi:10.1029/2001JD000583, 2002.

30 Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and Pöschl, U.: Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei counter (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol particles in theory and experiment, *Atmos. Chem. Phys.*, 8, 1153–1179, doi:10.5194/acp-8-1153-2008, 2008.

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rose, D., Nowak, A., Achtert, P., Wiedensohler, A., Hu, M., Shao, M., Zhang, Y., Andreae, M. O., and Pöschl, U.: Cloud condensation nuclei in polluted air and biomass burning smoke near the mega-city Guangzhou, China – Part 1: Size-resolved measurements and implications for the modeling of aerosol particle hygroscopicity and CCN activity, *Atmos. Chem. Phys.*, 10, 3365–3383, doi:10.5194/acp-10-3365-2010, 2010.
- Ruehl, C. R., Chuang, P. Y., and Nenes, A.: Distinct CCN activation kinetics above the marine boundary layer along the California coast, *Geophys. Res. Lett.*, 36(15), L15814, doi:10.1029/2009GL038839, 2009.
- Ruehl, C. R., Chuang, P. Y., and Nenes, A.: How quickly do cloud droplets form on atmospheric particles?, *Atmos. Chem. Phys.*, 8, 1043–1055, doi:10.5194/acp-8-1043-2008, 2008.
- Sciare, J., Bardouki, H., Moulin, C., and Mihalopoulos, N.: Aerosol sources and their contribution to the chemical composition of aerosols in the Eastern Mediterranean Sea during summertime, *Atmos. Chem. Phys.*, 3, 291–302, doi:10.5194/acp-3-291-2003, 2003.
- Shantz, N. C., Leatch, W. R., Phinney, L., Mozurkewich, M., and Toom-Sauntry, D.: The effect of organic compounds on the growth rate of cloud droplets in marine and forest settings, *Atmos. Chem. Phys.*, 8, 5869–5887, doi:10.5194/acp-8-5869-2008, 2008.
- Sotiropoulou, R. E. P., Medina, J., and Nenes A.: CCN predictions: is theory sufficient for assessments of the indirect effect?, *Geophys. Res. Lett.*, 33, L05816, doi:10.1029/2005GL025148, 2006.
- Sotiropoulou, R. E. P., Nenes, A., Adams, P. J., and Seinfeld, J. H.: Cloud condensation nuclei prediction error from application of Kohler theory: Importance for the aerosol indirect effect, *J. Geophys. Res.*, 112, D12202, doi:10.1029/2006JD007834, 2007.
- Sorooshian, A., Murphy, S. M., Hersey, S., Gates, H., Padro, L. T., Nenes, A., Brechtel, F. J., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Comprehensive airborne characterization of aerosol from a major bovine source, *Atmos. Chem. Phys.*, 8, 5489–5520, doi:10.5194/acp-8-5489-2008, 2008.
- Vrekoussis, M., Liakakou, E., Koçak, M., Kubilay, N., Oikonomou, K., Sciare, J., and Mihalopoulos, N.: Seasonal variability of optical properties of aerosols in the Eastern Mediterranean, *Atmos. Environ.*, 39(37), 7083–7094, 2005.
- Wang, S. and Flagan, R.: Scanning electrical mobility spectrometer, *J. Aerosol Sci.*, 20, 1485–1488, 1989.
- Wang, J., Cubison, M. J., Aiken, A. C., Jimenez, J. L., and Collins, D. R.: The importance of aerosol mixing state and size-resolved composition on CCN concentration and the variation

of the importance with atmospheric aging of aerosols, *Atmos. Chem. Phys.*, 10, 7267–7283, doi:10.5194/acp-10-7267-2010, 2010.

Weber, R. J., Sullivan, A. P., Peltier, R. E., Russell, A., Yan, B., Zheng, M., deGouw, J., Warneke, C., Brock, C., Holloway, J. S., Atlas, E. L., and Edgerton, E.: A study of secondary organic aerosol formation in the anthropogenic-influenced southeastern United States, *J. Geophys. Res.*, 112, D13302, doi:10.1029/2007JD008408, 2007.

Wiedensohler, A.: Technical note: An approximation of the bipolar charge distribution for particles in the submicron range, *J. Aerosol Sci.*, 19, 387–389, 1988.

Wiedensohler, A., Cheng, Y. F., Nowak, A., Wehner, B., Achtert, P., Berghof, M., Birmili, W., Wu, Z. J., Hu, M., Zhu, T., Takegawa, N., Kita, K., Kondo, Y., Lou, S. R., Hofzumahaus, A., Holland, F., Wahner, A., Gunthe, S. S., Rose, D., Su, H., and Pöschl, U.: Rapid aerosol particle growth and increase of cloud nucleus activity by secondary aerosol formation and condensation: A case study for regional air pollution in northeastern China, *J. Geophys. Res.*, 114, D00G08, doi:10.1029/2008JD010884, 2009.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007gl029979, 2007.

Size-resolved CCN distributions

A. Bougiatioti et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN distributions

A. Bougiatioti et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Table 1. Regression statistics for the CCN closure.

Supersaturation (%) (number of data points)	WSOC/OC = 0		WSOC/OC = 0.7		$\kappa = 0.34$	
	$\frac{CCN_{predicted}}{CCN_{observed}}$	R^2	$\frac{CCN_{predicted}}{CCN_{observed}}$	R^2	$\frac{CCN_{predicted}}{CCN_{observed}}$	R^2
0.21 (1012)	0.94	0.94	1.02	0.95	0.97	0.86
0.38 (865)	0.92	0.93	0.99	0.96	0.92	0.92
0.52 (996)	0.98	0.95	1.00	0.96	0.93	0.93
0.66 (845)	0.97	0.94	0.99	0.94	0.92	0.92
0.73 (761)	0.94	0.92	0.96	0.93	0.94	0.89

Size-resolved CCN distributions

A. Bougiatioti et al.

Table 2. Average droplet size (standard deviation) measured by the optical particle counter (μm) for different supersaturations. Results shown for Finokalia and NaCl calibration aerosol with dry particle diameter of size d_{p50} and 100 nm.

Supersaturation (%)	Dry diameter d_{p50}		Dry diameter 100 nm	
	Finokalia	NaCl	Finokalia	NaCl
0.21	2.2 (0.6)	3.0 (0.6)	3.7 (0.3)	4.0 (0.7)
0.38	4.9 (0.6)	5.4 (0.7)	6.4 (0.3)	6.4 (0.8)
0.52	6.8 (0.6)	6.7 (0.6)	7.9 (0.3)	8.1 (0.6)
0.66	7.9 (0.6)	7.5 (0.6)	8.7 (0.4)	9.0 (0.6)
0.73	8.1 (0.6)	8.0 (0.6)	9.3 (0.3)	9.9 (0.6)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 1. Maps indicating the location of the sampling site and three-day HYSPLIT back trajectories for characteristic types of air masses sampled during FAME07: (a) 14 July 2007, (b) 17 July 2007, (c) 24 July 2007, and (d) 1 August 2007.

Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 2. Schematic of the setup used for the size-resolved CCN measurements.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Fig. 3. (a) CCN activation fraction and distribution from 25 July at 11:49 a.m. (UTC+2) with sigmoidal fit at 0.3% supersaturation and (b) example of CCN activity spectrum from August 3rd (00:12-11:58 AM (UTC+2)) for 60 and 100 nm particles.

Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 4. Time series of the CN and CCN measurements throughout the measurement period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 5. Example of daily average CCN distributions from 2 August at different levels of supersaturation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 6. (a) Averaged size-resolved CCN efficiency spectra at different levels of supersaturation (0.2–0.73%): activated particle fraction plotted against particle dry diameter, (b) CCN activation spectrum (S_c vs. d_{p50}) for the whole measurement period, (c) $R_a(d_p)$ for 14 July, and, (d) $R_a(d_p)$ for 24 July.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 7. Daily average aerosol size distributions, characteristic of the air masses sampled.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 8. (a) Time series of the characteristic (SMCA, 100 nm diameter) and filter-derived hygroscopicity parameter κ for the totality of the measurement period, (b) sulfate volume fraction time series and (c) chemical and mixing state parameters averages for the whole measurement period and for 14 July (left) and 24th (right).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Size-resolved CCN
distributions

A. Bougiatioti et al.




Fig. 9. (a) Time series of the hygroscopicity parameter κ_* for the five dry particle sizes, (b) diurnal variability of the activated fraction (E_*) for 14–15 July, (c) inferred hygroscopicity parameter (κ_*) for 14–15 July, (d) E_* for 30–31 July, and (e) κ_* for 30–31 July, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Size-resolved CCN distributions

A. Bougiatioti et al.




Fig. 10. (a) Droplet sizes of CCN at the different supersaturations for ambient and calibration aerosol at the exit of the growth chamber, (b) droplet sizes of ambient CCN at d_{p50} and (c) average droplet diameter at different dry particle sizes.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

