

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Chlorine activation on stratospheric aerosols: uncertainties in parameterizations and surface area

T. Wegner<sup>1</sup>, J.-U. Grooß<sup>1</sup>, M. von Hobe<sup>1</sup>, F. Stroh<sup>1</sup>, O. Sumińska-Ebersoldt<sup>2</sup>,  
C. M. Volk<sup>3</sup>, E. Hösen<sup>3</sup>, V. Mitev<sup>4</sup>, G. Shur<sup>5</sup>, and R. Müller<sup>1</sup>

<sup>1</sup>Institute of Energy and Climate Research – Stratosphere (IEK-7), Forschungszentrum Jülich, Jülich, Germany

<sup>2</sup>Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

<sup>3</sup>Department of Physics, University of Wuppertal, Wuppertal, Germany

<sup>4</sup>Centre Suisse d'Electronique et de Microtechnique SA, Neuchâtel, Switzerland

<sup>5</sup>Central Aerological Observatory, Dolgoprudny, Russia

Received: 17 July 2012 – Accepted: 17 July 2012 – Published: 16 August 2012

Correspondence to: T. Wegner (t.wegner@fz-juelich.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Chlorine activation in the Arctic is evaluated by examining the different parameterizations for uptake coefficients on stratospheric aerosols, high-resolution in-situ measurements and vortex-wide satellite observations. The parameterizations for heterogeneous chemistry on liquid aerosols are most sensitive to temperature with the reaction rates doubling for each Kelvin increase in temperature. However, differences between the parameterizations are negligible. For Nitric Acid Trihydrate particles (NAT) the major factors of uncertainty are the number density of nucleated particles and different parameterization choices. These two factors induce an uncertainty that covers several orders of magnitude on the reaction rate. But as predicted reaction rates on liquid aerosols always exceed those on NAT the overall uncertainty is small. In-situ observations of  $\text{ClO}_x$  from Arctic winters in 2005 and 2010 are used to validate the heterogeneous chemistry parameterizations. The ambient conditions for these measurements proved to be very different between those two winters with HCl being the limiting reacting partner for the 2005 measurements and  $\text{ClONO}_2$  for the 2010 measurements. Modeled levels of chlorine activation are in very good agreement with the in-situ observations and the surface area provided by Polar Stratospheric Clouds (PSCs) has only a limited impact on modeled chlorine activation. This indicates that the parameterizations give a good representation of the processes in the atmosphere. Backtrajectories started on the location of the observations in 2005 indicate temperatures on the threshold for PSC formation, hence the surface area is mainly provided by the background aerosol. Still, the model shows additional chlorine activation during this time-frame, providing cautionary evidence for chlorine activation even in the absence of PSCs. Vortex-averaged satellite observations also show no definite connection between chlorine activation and PSC formation. The inter- and intra-annual variability of vortex-average HCl and  $\text{HNO}_3$  based on MLS observations is examined for the Arctic winters 2004/2005 to 2010/2011. These observations show that removal of HCl and  $\text{HNO}_3$  from the gas-phase are not correlated.  $\text{HNO}_3$  loss exhibits great inter-annual

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



variability depending on prevailing temperatures while HCl loss is continuous through December without considerable inter- or intra-annual variability. Only the recovery of HCl in late in winter depends on the level of denitrification. Hence, the occurrence of HNO<sub>3</sub> containing PSC particles does not seem to have a significant effect on the speed of initial chlorine activation on a vortex-wide scale.

## 1 Introduction

During polar night in the stratosphere the main chlorine reservoir species HCl and ClONO<sub>2</sub> are converted to photo-labile species through heterogeneous reactions on liquid and solid particles (Solomon et al., 1986), a process known as chlorine activation. Under the influence of sunlight, these photo-labile species are converted to ClO<sub>x</sub> (ClO + 2 · Cl<sub>2</sub>O<sub>2</sub>), which is referred to as “active chlorine”, as ClO<sub>x</sub> drives the catalytic ozone loss cycles which lead to severe depletion of ozone in polar spring (Molina and Molina, 1987; Solomon, 1999). The liquid and solid particles acting as reactions sites for heterogeneous chemistry are provided by Polar Stratospheric Clouds (PSCs) and the ubiquitous background sulfate aerosol layer. PSCs are composed of mixtures of Supercooled Ternary Solution (STS), Nitric Acid Trihydrate (NAT) and ice (e.g. Peter and Grooß, 2012). Clouds composed of STS and low number density NAT ( $3 \times 10^{-4} \text{ cm}^{-3}$  –  $10^{-3} \text{ cm}^{-3}$ ) are the most common type in cold Arctic winters and STS with high number density NAT ( $> 10^{-3} \text{ cm}^{-3}$ ) are more common for the Antarctic (Pitts et al., 2011). Ice formation is usually only observed over the Antarctic as the polar vortex over the Arctic tends to be warmer. An exception was the Arctic winter of 2009/2010, which was exceptionally cold leading to ice formation (Pitts et al., 2011; Khosrawi et al., 2011). An overview of our current understanding of PSC processes is given in Lowe and MacKenzie (2008) and Peter and Grooß (2012).

The greatest uncertainty in PSC formation is the nucleation mechanism for NAT which directly influences the NAT number density. However, based on CALIPSO observations (Pitts et al., 2011) and in-situ measurements (Fahey et al., 2001), NAT

### Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



number densities can be constrained to values between  $10^{-1}$  and  $10^{-4}$   $\text{cm}^{-3}$ . Background aerosol surface area density (SAD) depends on the COS and  $\text{SO}_2$  fluxes into the stratosphere and is significantly enhanced only after large volcanic eruptions (Robock, 2000). The influence of the sulfate aerosol on stratospheric ozone was particularly observable after the eruptions of El Chichón (Hofmann and Solomon, 1989) and Mt. Pinatubo (Portmann et al., 1996; Tilmes et al., 2008b). The importance of PSCs and the sulfate aerosol layer on heterogeneous chemistry is well established (e.g. Solomon, 1999), with the impact of an unperturbed aerosol layer on heterogeneous processing first discussed by Rodriguez et al. (1988). But recently Drdla and Müller (2012) suggested that even during volcanic quiescent times chlorine activation in polar night is dominated by reactions on cold binary sulfate aerosol. However, elevated levels of active chlorine usually coincide with the presence of PSCs which makes it difficult to attribute chlorine activation to heterogeneous processing on a particular aerosol or cloud type. Kawa et al. (1997) reported in-situ measurements of active chlorine from the Antarctic during the ASHOE/MAESA campaign in 1994 where backtrajectories indicated that during the ten days prior to the measurements, temperatures had been too high for PSC formation and therefore concluded that the observed chlorine activation must have occurred on the background aerosol. With the current high chlorine loading in the stratosphere and a deliberate enhancement of the stratospheric aerosol layer being discussed to counter a temperature increase at the surface (e.g. Crutzen, 2006; Rasch et al., 2008; Tilmes et al., 2008a), understanding the role of the background aerosol for chlorine activation is essential.

The following sections will discuss the existing parameterizations for heterogeneous chemistry on the various aerosol types and their uncertainty. These parameterizations are validated with in-situ measurements and the role of the background aerosol is constrained with model simulations and satellite observations.

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Chlorine activation

Following Portmann et al. (1996) the evolution of chlorine species in the polar regions can be divided into four phases. The “setup phase” preceding the polar winter when  $\text{Cl}_y$  is partitioned between the reservoir species ( $\text{HCl}$  and  $\text{ClONO}_2$ ), the “activation phase” during polar night when the reservoir species are partly converted into  $\text{ClO}_x$ , the “maintenance/further activation phase” when activation and deactivation are in competition and further activation occurs if temperatures are sufficiently low and finally the “termination phase” when active chlorine is converted back into the reservoir species.

The three heterogeneous reactions mainly responsible for chlorine activation are (Solomon, 1999; Peter and Grooß, 2012):



In the Arctic active chlorine is principally deactivated into  $\text{ClONO}_2$ , the speed of deactivation is thus limited by the availability of  $\text{NO}_x$  ( $\text{NO} + \text{NO}_2 + \text{NO}_3$ ) (Müller et al., 1994; Douglass et al., 1995)



The formation of  $\text{NO}_x$  is controlled by solar radiation intensity and thus, it is a function of the solar zenith angle and the availability of gas-phase  $\text{HNO}_3$ :



Large NAT particles can effectively remove  $\text{HNO}_3$  from the lower stratosphere (Fahey et al., 2001) which results in suppressed deactivation and prolongs the availability of  $\text{ClO}_x$  into late winter and spring (Harris et al., 2010). In the illuminated polar vortex in spring,  $\text{ClO}_x$  rapidly depletes ozone, causing the creation of the ozone hole over the

### Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Antarctic and severe depletion of ozone over the Arctic for very cold winters. Extensive denitrification of the lower polar stratosphere through sedimentation of NAT particles is more common in the Antarctic than in the Arctic. Temperatures in the Antarctic are lower and remain below the NAT equilibrium temperature ( $T_{\text{NAT}}$ , Hanson and Mauersberger, 1988) for longer periods than in the Arctic and thus, allow NAT particles to grow and sediment. In the Arctic temperatures are higher and more variable than in the Antarctic which results in less pronounced denitrification and faster deactivation of chlorine through Reaction (R4). In the Antarctic, the almost complete destruction of ozone in polar spring leads to an increase in Cl concentrations, which allows the deactivation reaction



to occur (Douglass et al., 1995). Recently, Grooß et al. (2011) reported that in the Antarctic, for very low ozone values ( $< 0.5$  ppmv) a balance is maintained for a certain period between rapid gas-phase production of HCl and HOCl and rapid heterogeneous reaction between these two compounds. This period ends by very rapid (on the order of one day) irreversible Cl deactivation into HCl and almost complete destruction of ozone.

The speed of the heterogeneous Reactions (R1) to (R3) is described by the rate constant  $k$  which depends on the uptake coefficient  $\gamma$ , aerosol surface area density SAD and mean gas velocity  $c_{\text{gas}}$ .

$$k = 0.25 \times \gamma \times c_{\text{gas}} \times \text{SAD} \quad (1)$$

The uptake coefficient  $\gamma$ , which describes the fraction of collisions of gas molecules with the particle surface which lead to a reaction, and the surface area density are the main factors controlling the heterogeneous reaction rate and both strongly depend on temperature.

In this work we use parameterizations of the uptake coefficient to model heterogeneous chemistry on liquid aerosols derived by Shi et al. (2001) and Hanson (1998), and for NAT particles the work of Carslaw and Peter (1997), which is based on laboratory

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measurements by Hanson and Ravishankara (1993) and Abbatt and Molina (1992). NAT reaction probabilities based on the scheme by Hanson and Ravishankara (1993) represent an upper limit while the scheme by Abbatt and Molina (1992) represents the lower limit. For liquid aerosols the parameterizations yield very similar reaction rates, with Shi et al. (2001) reporting an uncertainty of 40 % for Reaction (R1) and 32 % for Reaction (R2). As the parameterization by Shi et al. (2001) for liquid aerosols has only been derived for binary aerosols it has been extended to STS particles by assuming that there is no difference in the uptake coefficient for STS particles and binary aerosol particles that would exist in the absence of  $\text{HNO}_3$ , as shown by Elrod et al. (1995). Liquid aerosol surface area depends on the stratospheric  $\text{H}_2\text{SO}_4$  content and increases significantly only for temperatures below about 192 K with the uptake of  $\text{HNO}_3$  and the formation of STS (Carslaw et al., 1994). The surface area density for NAT particles depends primarily on the assumed particle density. The main reaction channel for chlorine activation is Reaction (R1) (Fig. 1a). Other reactions are less important for the initial chlorine activation phase as  $\text{ClONO}_2$  constitutes the second largest reservoir and is quantitatively removed (Müller et al., 1994; Douglass et al., 1995; Portmann et al., 1996). When  $\text{ClONO}_2$  starts to regenerate and  $\text{HCl}$  falls below  $\text{ClONO}_2$  mixing ratios, Reaction (R2) gains importance if temperatures are low enough.

Figure 1 shows the first order loss rates in these two species for typical stratospheric conditions for Reactions (R1) and (R2) on liquid aerosols and NAT particles. NAT surface area density is calculated assuming a uniform distribution of spherical particles and the liquid aerosol surface area assuming a log-normal distribution. The various heterogeneous chemistry parameterizations and possible number densities for NAT particles cover several orders of magnitude in first order loss rates, indicating a large uncertainty concerning their ability to act as reaction sites for chlorine activation. However, over most of the temperature range heterogeneous reaction rates on NAT are slower than on the background aerosol, even when a high NAT number density of  $10^{-1} \text{ cm}^{-3}$  (Fig. 1) is assumed. Only at temperatures around the thermodynamic equilibrium temperature for NAT, reaction rates on NAT are faster than on the background

aerosol. With the formation of STS or a NAT number density less than  $10^{-1} \text{ cm}^{-3}$ , reaction rates on the liquid aerosol always exceed those on NAT. While chlorine activation on liquid aerosols is not sensitive to the parameterizations, it is very sensitive to temperature. The reaction rate doubles for every Kelvin cooling and increases tenfold over a 2 K temperature range around 192 K with the uptake of  $\text{HNO}_3$  on the background aerosol (Carslaw et al., 1994; Peter, 1997). This causes high sensitivity of simulated chlorine activation to small variabilities in the temperature field. Therefore, for heterogeneous reactivity on liquid aerosols even a small bias in the temperature field has larger effects than the uncertainty of the uptake coefficients itself.

### 3 In situ observations

During a flight into the Arctic vortex by the high-altitude research aircraft Geophysica on 7 March 2005, high concentrations of active chlorine compounds ( $\text{ClO}_x$ ) were observed by the HALOX instrument (von Hobe et al., 2005), and back-trajectories from the flightpath indicate that chlorine in the observed air masses was reactivated just 24 h prior to the measurements. For the analysis of chlorine activation, chemistry and PSC microphysics are simulated along trajectories starting on 3rd March 2005 and ending on the flightpath. Trajectory and chemistry simulations are performed using the CLaMS Model (McKenna et al., 2002b,a) driven by ERA-INTERIM meteorological fields (Dee et al., 2011). To calculate the reaction rates JPL 2006 recommendations (Sander et al., 2006) are used. STS growth is simulated with the thermodynamic model developed by Carslaw et al. (1995), dynamic NAT growth with the model of Carslaw et al. (2002) and the NAT nucleation rate is taken from Voigt et al. (2005).

The initialization of these trajectories is based on in-situ measurements of  $\text{N}_2\text{O}$  and  $\text{CH}_4$  from HAGAR (Homan et al., 2010; Werner et al., 2010),  $\text{H}_2\text{O}$  from the FISH instrument (Zöger et al., 1999) and aerosol number density (size range: 10 nm–5  $\mu\text{m}$  in diameter) from the COPAS instrument (Weigel et al., 2009). Additional details of this flight can be found in von Hobe et al. (2006).  $\text{CH}_4$  observations by HAGAR indicate that

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a homogeneous air mass was sampled although the flight covered 65°–85° equivalent latitude. From these measurements we derive the total inorganic chlorine loading  $Cl_y$  using the  $CH_4$ - $Cl_y$  tracer correlation reported in Grooß et al. (2002) as a function of potential temperature ( $\Theta$ ) and equivalent latitude ( $\Phi$ ).  $ClONO_2$ , HCl, and  $HNO_3$  mixing ratios are taken from ACE-FTS v2.2update (Bernath et al., 2005) observations on 3rd and 4th March, interpolated on the trajectory positions on 3rd March 12 UTC as a function of potential temperature and equivalent latitude. The difference between the concentrations of  $Cl_y$  and the sum of  $ClONO_2 + HCl$  is initialized as  $ClO_x$ . For most of the flightpath  $HNO_3$  values, initialized from ACE-FTS, are between 6 and 7 ppbv. Calculated with the passive tracer subtraction method (Grooß et al., 2002), this yields a denitrification of  $\sim 50\%$  corroborating the conclusion by von Hobe et al. (2006) that the observed air masses were highly denitrified. All other chemical species are interpolated from the hemispheric CLaMS simulation for the Arctic winter 2004/2005 (Grooß and Müller, 2007). The background aerosol surface area is taken from the climatology compiled by David Considine (Eyring et al., 2010). The error of the initialization of  $HNO_3$  and the chlorine species is estimated by interpolating the satellite measurements on the potential temperature of the Geophysica observations and subsequently calculating the range of observations of the three nearest neighbors in the equivalent latitude space. Typically, this results in an error of less than 0.2 ppbv.

As heterogeneous chemistry is highly sensitive to temperature, accurate knowledge of it is necessary to model the activation of chlorine on stratospheric aerosols. For example, Brakebusch et al. (2012) have shown that temperatures in the specified dynamics version of the Whole Atmosphere Community Climate Model tend to be warm biased compared to MLS observations and that the agreement between model and satellite observations improves when a  $-1.5\text{ K}$  bias is applied to the heterogeneous chemistry calculations. Ambient temperature on board Geophysica is measured by the Thermodynamic Complex (TDC, Shur et al., 2006). These measurements similarly, show a  $\sim 1.5\text{ K}$  bias when compared to ERA-INTERIM temperatures (Fig. 2). Between 455–470 K ERA-INTERIM temperatures are consistently higher than the observations.

ERA-INTERIM also shows better agreement at higher temperatures and the warm bias appears to be limited to temperatures below 205 K. In Fig. 2b the vertical profile for the temperature bias is shown. The greatest warm bias is located at 460 K, on flight altitude, with 1.5 K. No in-situ temperature measurements exist over the course of the trajectories so we assume the warm bias exists for the whole trajectory length and adjust the temperatures below 205 K according to the kernel shown in Fig. 2.

Modeling PSCs along the trajectories indicates (Fig. 3) the presence of NAT and STS 24 h prior to the measurements. On the trajectories ending between 08:45 and 09:30 UTC typically less than 0.1 ppbv HNO<sub>3</sub> condenses on STS. This enhances the surface area density by about a factor of 1.5 compared to the background aerosol. The maximum calculated enhancement is about a factor of 2. For trajectories ending before 08:45 and after 09:30 all available HNO<sub>3</sub> is predicted to condense to STS, with a maximum enhancement of the surface area density by a factor of 5. However, these maximum enhancements only occur for a few hours along the trajectories. For most of the time when the trajectories experience temperatures below  $T_{\text{NAT}}$ , PSCs enhance the surface area by less than 10 % over the background aerosol. As temperatures are below  $T_{\text{NAT}}$  for only about 20 h modeled NAT particles cannot reach thermodynamic equilibrium. With temperatures below  $T_{\text{NAT}}$  for such a short time, NAT number density is on the order of 10<sup>-3</sup> cm<sup>-3</sup>, which leads to a negligible increase in surface area density by NAT. Therefore, any modeled increase in surface area density is caused by STS and not NAT particles.

To assess the ability of the model to reproduce the measured extent of chlorine activation two simulations are performed. The first calculates the heterogeneous reaction rates with a surface area density that includes all PSCs (Full PSC) and the second uses only the surface area density of the background aerosol without any enhancement due to STS or NAT (Binary only). Temperatures are well above the frost point so ice PSCs do not form in either of these simulations. Figure 4 shows that the difference between initialized and measured ClO<sub>x</sub> on the flight path is on average about 1 ppbv. Maximum activation is simulated for trajectories ending before 08:45 and after

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

09:30 UTC. These trajectories also exhibit the maximum uptake of  $\text{HNO}_3$  on PSCs as minimum temperatures are about 1–2 K lower than for the other trajectories. For trajectories ending between 08:45 and 09:30 about 0.7 ppbv additional chlorine is activated within 92 h. The model cannot reproduce all of the fine structures of the measurements which is likely due to the coarse initialization and resolution of the ERA-INTERIM meteorological fields. However, overall both model simulations show good agreement with measured  $\text{ClO}_x$ , within the uncertainty of initialization and measurements. For trajectories ending before 08:45 modeled chlorine activation tends to be at the upper limit of uncertainty and afterwards at the lower limit. The difference between both simulations is minimal with the “Full PSC” simulation activating slightly more chlorine than the “Binary only” simulation. However, this difference is on the order of 10 % indicating that the heterogeneous reactivity provided by the binary aerosol in the model is sufficient to produce an activation that corresponds to most of the observed activation of chlorine within the considered time-frame. Even though modeled PSCs increase the surface area density by up to 500 % the effect on chlorine activation is limited, as this maximum increase only lasted for a short time-frame. In the model, neither HCl nor  $\text{ClONO}_2$  are completely depleted thus, chlorine activation along the trajectories is limited by the heterogeneous reaction rates and not by the availability of either reservoir species. As the model results overlap with the observations, within the uncertainties of this simulation and measurements, the temperature dependence of the current parameterizations for heterogeneous chlorine activation is in agreement with the processes in the real atmosphere.

In addition to the Geophysica flight from 7 March 2005 we examined three Geophysica flights in January 2010 from the RECONCILE campaign in the Arctic (Fig. 5). The temperature histories along 7 day back-trajectories ending on the flightpath of these flights indicate temperatures low enough for efficient heterogeneous chemistry. Temperatures are below  $T_{\text{NAT}}$  at the beginning of the trajectories and during the flights. In between temperatures exceed 200 K so all PSCs that formed initially very likely evaporated. Prior to the flights temperatures for most trajectories are below  $T_{\text{NAT}}$  for about

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

30 h consecutively with some trajectories during flight 2 residing below  $T_{\text{NAT}}$  over 50 h. Trajectories for flight 3 are the longest time below  $T_{\text{NAT}}$ . Here temperatures are below  $T_{\text{NAT}}$  for 40 h at the beginning of the trajectories and again for 30 h prior to the flight. In contrast to the 2005 flight, minimum temperatures for all three flights are below 190 K, low enough for extensive PSC formation and thus low enough for extensive  $\text{HNO}_3$  uptake from the gas-phase. Measurements during the three considered flights in January 2010 from the up- and downward facing LIDAR MAL (Matthey et al., 2003) on-board Geophysica show that backscatter ratios are elevated by a factor of 3–4 and a depolarization of less than 4 %. The high backscatter combined with the low depolarization is an indicator that PSCs mostly consisted of STS droplets and that no or only very few NAT particles were present. Chlorine activation is simulated by running box-model calculations along these 7 day trajectories. The initialization is taken directly from a hemispheric 3-D-CLaMS simulation. Total  $\text{Cl}_y$  is estimated from measured  $\text{CH}_4$  by HAGAR via the  $\text{CH}_4$ - $\text{Cl}_y$  correlation (Grooß et al., 2002). Model  $\text{HCl}$  and  $\text{ClONO}_2$  are then scaled accordingly so that model  $\text{Cl}_y$  agrees with observations. Generally, model  $\text{Cl}_y$  has to be scaled up by about 15 % to agree with observations. For all three flights the model activates significant amounts of chlorine over the 7 day period considered. For flights 1 and 2 additional chlorine is activated along the majority of trajectories during the 7 days of simulation, generally around 0.4 ppbv with some trajectories showing up to 0.8 ppbv additional  $\text{ClO}_x$ . Only a minor fraction of the trajectories shows no additional chlorine activation. The trajectories for flight 3 all show additional chlorine activation, generally around 0.5 ppbv with maximum additional  $\text{ClO}_x$  of 1.2 ppbv. The limiting factor for chlorine activation for these three flights is the availability of  $\text{ClONO}_2$ . The model simulates that  $\text{ClONO}_2$  is completely depleted at the time of the flights, hence, the additionally available surface area density provided by PSCs cannot have an effect on  $\text{ClO}_x$  levels. Consequently, the simulation only using the background aerosol surface area density to calculate heterogeneous reaction rates produces the same results as the “Full PSC” simulation. In contrast to the flight on 7 March 2005, no reliable  $\text{Cl}_2\text{O}_2$  measurements exist for the flights in January 2010 so we are limited to comparing modeled

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and measured ClO. ClO possesses a distinct diurnal cycle and the parameters describing the equilibrium between ClO and Cl<sub>2</sub>O<sub>2</sub> are still subject to some uncertainty (von Hobe et al., 2007). To calculate ClO from simulated ClO<sub>x</sub> we use the results from Sumińska-Ebersoldt et al. (2012) who constrained the ClO/Cl<sub>2</sub>O<sub>2</sub> equilibrium parameters to in-situ observations. The thermal equilibrium constant is taken from Plenge et al. (2005) and the Cl<sub>2</sub>O<sub>2</sub> absorption cross-section from von Hobe et al. (2009) scaled to the absolute measurements of Lien et al. (2009). Figure 5 shows modeled and measured ClO for the three flights in January 2010. The first two flights have decreasing solar zenith angles leading to increasing ClO values and the third flight was carried out in darkness. Overall, the model shows excellent agreement with the measurements with flights 1 and 2 showing the dependence of ClO on the solar zenith angle and flight 3 showing nighttime, thermal equilibrium measurements.

#### 4 Vortex average

To further examine, which effect the surface area enhancement caused by the HNO<sub>3</sub> uptake has on chlorine activation vortex averaged satellite observations from MLS for the Arctic winters 2004/2005 to 2010/2011 are analyzed. We focus on the vortex core, described by equivalent latitudes poleward of 75° N. A high inter-annual variability, common for the Arctic, is evident in the observations of HCl and HNO<sub>3</sub> (Fig. 6). The winters 2004/2005, 2007/2008 and 2010/2011 had very cold and stable vortices leading to strong denitrification and depressed values of gas-phase HCl extending into spring. The observations for the winters 2005/2006 and 2008/2009, on the other hand, show hardly any denitrification and HCl starts increasing in the middle of January. The onset of chlorine activation at the beginning of winter is controlled by the quantitative titration of the available ClONO<sub>2</sub> through the heterogeneous Reaction (R1). Thus, a decrease in gas-phase HCl can serve as an indicator for chlorine activation.

$$\frac{d[\text{ClO}_x]}{dt} \approx -\frac{d[\text{HCl}]}{dt} - \frac{d[\text{ClONO}_2]}{dt} \approx -2\frac{d[\text{HCl}]}{dt} \quad (2)$$

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Although HCl shows high inter-annual variability for the period of January to April, observations show only little variability in December when chlorine begins to become activated. At the end of December HCl has decreased by about 1 ppbv, 50 % of the initially available HCl. Significant differences in HCl depletion between the different winters emerge not until the middle of January. In winters 2005/2006 and 2008/2009 HCl starts recovering in January, while in winters 2004/2005 and 2007/08 the final recovery does not begin until February. This corresponds to observations of  $\text{HNO}_3$ . A decrease in gas-phase  $\text{HNO}_3$  is an indicator for PSC formation and irreversible denitrification. Pitts et al. (2009, 2011) have shown that the first PSCs forming in Arctic winter are mainly composed of STS and low number density NAT ( $< 10^{-3} \text{ cm}^{-3}$ ). Therefore, the observed decrease in gas-phase  $\text{HNO}_3$  in December can mostly be attributed to condensation of  $\text{HNO}_3$  on STS and a lesser extent to NAT and, possibly irreversible denitrification.

In early winter,  $\text{HNO}_3$  shows, in contrast to HCl, high inter-annual variability indicating that the additional surface area provided by PSCs has no detectable direct influence on the rate of chlorine activation on a vortex wide scale. For the winters 2004/2005 and 2009/2010 75 % of gas-phase  $\text{HNO}_3$  are removed throughout the winter. In 2004, gas-phase  $\text{HNO}_3$  shows a steep decline mid of December and at the end of December in 2009. For 2004 we can assume that the surface area density was significantly enhanced through  $\text{HNO}_3$  uptake throughout December while for 2009 surface area density was for most of the time at background levels. In winters 2005/2006 and 2008/2009 observations show only a gradual decrease of gas-phase  $\text{HNO}_3$ . This decrease of gas-phase  $\text{HNO}_3$  indicates that PSCs formed only to a minor extent compared to the winters 2004/2005 and 2009/2010. Pitts et al. (2011) show that maximum PSC coverage in the winter 2008/2009 was only one third of the maximum coverage in 2009/2010. Still, the decrease of HCl in December is very similar for all winters despite the high variability in gas-phase  $\text{HNO}_3$ .

Figure 7 shows the probability density distribution of the maximum enhancement of the surface area by STS in the absence of NAT formation for the winters 2004, 2008 and 2009. These three winters cover the whole spectrum in HCl and  $\text{HNO}_3$  variability. STS

formation is calculated according to Carslaw et al. (1995) along 7 day back-trajectories from MLS observations in December 2005, 2008 and 2009 poleward of 75° N equivalent latitude, assuming that no irreversible denitrification or dehydration has occurred yet. We show the fraction of trajectories that fall into 6 different bins. The first bin contains the fraction of trajectories with a maximum enhancement of the surface area density of less than 1.1, i.e. showing no significant enhancement over background aerosol levels. For the next two bins (maximum enhancement factor 1.1–1.5 and 1.5–2) only a small fraction of gas-phase HNO<sub>3</sub> has condensed on STS but has already significantly enhanced the surface area density. In the bins with an enhancement factor of 2–5 and 5–10 most available gas-phase HNO<sub>3</sub> is condensed on STS and in the last bin virtually all HNO<sub>3</sub> is in STS leading to an increase of surface area density over the background aerosol of more than a factor of 10. Figure 7 shows this distribution on 3 different days in December and for 5 altitudes. The trajectories ending 15 December show no or only a minor increase of surface area density but already some differences emerge between the different winters. In 2009 none of the trajectories show any increase in SAD and for 2008 less than 20 % show a small enhancement. In 2004, however, only less than 30 % between 475 and 550 K show no increase in surface area density.

Differences between the winters become more apparent for 22 December. For 2008 and 2009 the distribution hardly changes compared to 15 December, most trajectories still show little enhancement of surface area density. In 2004, however, most trajectories show some enhancement of surface area density and between 475 and 550 K all do. At 500 K more than 50 % of the trajectories already show an enhanced surface area density by more than a factor of 2. The difference between these three winters becomes most obvious for the trajectories started on 29 December. In 2004 the distribution has completely shifted from trajectories only showing minor enhancement of surface area density on 15 December to the majority showing an enhancement of a factor greater than 10. On the 450 and 475 K isentropes more than 80 % of the backward trajectories exhibit temperatures which lead to a more than tenfold increase in surface area density.

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This is also visible in the sharp drop in MLS gas-phase  $\text{HNO}_3$  at the end of December 2004 (Fig. 6). Nonetheless, MLS HCl measurements show no drastic change in the loss rate suggesting that the heterogeneous loss of HCl is not connected to the available surface area density. The change in probability density for 2008 and 2009 towards the end of December is less pronounced. Most trajectories show only a minor increase in surface area density but still their distribution is shifted towards higher surface area densities. This is more apparent in 2008 than in 2009, on 29 December 2009 the maximum increase in surface area density for almost all trajectories was less than a factor 1.5.

These three winters show the huge variability in PSC occurrence and  $\text{HNO}_3$  uptake for December in the Arctic. Nevertheless, HCl loss rates observed by MLS are fairly constant throughout December with little inter- and intra-annual variability. Only the vertical extent of HCl loss shows inter-annual variability which is correlated with the vertical extent of the low temperatures necessary for efficient heterogeneous chemistry.

## 5 Conclusions

We have examined the importance of the stratospheric background aerosol for chlorine activation. The evaluation of heterogeneous chemistry parameterizations has shown that uncertainties in NAT microphysics contribute most to the overall uncertainty in modeling heterogeneous chemistry. The uncertainties for NAT reactivity cover several orders of magnitude. However, using the most commonly observed NAT number densities, heterogeneous processing on these particles is significantly slower than on the background aerosol. For modeling chemistry on the binary background aerosol the greatest uncertainty results from temperature uncertainties as the reaction rate doubles for every Kelvin and increases tenfold with the formation of STS over a 2 K temperature range. To study heterogeneous chemistry on a synoptic timescale we analyzed backtrajectories from the flightpath of the Geophysica flight on 7 March 2005. These trajectories show chlorine reactivation about 30 h before the flight. With the initialization

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 derived from satellite data, simulating chlorine activation along trajectories ending on the flightpath allows the influence of the various aerosols on modeled chlorine activation to be assessed. Even though NAT and STS could form under the prevailing conditions the additional surface area provided by PSCs does not significantly enhance chlorine activation. 90 % of additionally activated chlorine during this time-frame originate from heterogeneous chemistry on a surface area provided by the background aerosol. This shows that for modeling heterogeneous chemistry the increase in reaction rate with decreasing temperature is more important than the increase of surface area density and that to correctly model heterogeneous chemistry on synoptic timescales accurate knowledge about the prevailing temperatures is essential. This flight also showed that a considerable amount of chlorine can be activated on a timescale of hours when both HCl and ClONO<sub>2</sub> are available. The three flights from January 2010 corroborate these results albeit under very different ambient conditions. For the 2005 flight neither HCl nor ClONO<sub>2</sub> are completely depleted in the model and therefore, the temperature is the decisive factor determining the level of chlorine activation. In 2010, however, ClONO<sub>2</sub> is the limiting factor as it is completely depleted while temperatures remain low enough for efficient heterogeneous chemistry. Thus, heterogeneous chemistry on the background aerosol surface area yields identical results as calculations with full PSC surface area and both simulations show excellent agreement with observations.

20 Not only can heterogeneous processing on the background aerosol surface area explain the bulk of chlorine activation on synoptic timescales along individual trajectories, but based on satellite observations we could demonstrate that the vortex-average chlorine activation rate for the last seven Arctic winters is not correlated with the occurrence of PSCs and the associated uptake of HNO<sub>3</sub> from the gas-phase. The observed HCl loss rate in December is similar for all considered Arctic winters despite their high variability in PSC occurrence. Even intra-annual variability of gas-phase HNO<sub>3</sub> is not reflected in observed HCl. During the December 2004 gas-phase HNO<sub>3</sub> dropped sharply while the HCl loss rate remained constant. Thus, the vortex-average

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

observations confirm the findings from the in-situ observations that the surface area provided by PSCs does not significantly impact chlorine activation rates.

*Acknowledgements.* This work and the 2010 flights were funded under the EU-FP7 project RECONCILE (RECONCILE-226365-FP7-ENV-2008-1). The 2005 flight was funded by the EU (APE-INFRA) as a test flight for the ESA MARSCHALS instrument. We are grateful to MDB for their support of the Geophysica flights and supply of avionic data, Ralf Weigel for COPAS, Nicole Spelten and Cornelius Schiller for FISH data, to ECMWF for meteorological analyses and to the ACE-FTS and EOS MLS teams for their high-quality data products. The Atmospheric Chemistry Experiment (ACE), also known as SCISAT, is a Canadian-led mission mainly supported by the Canadian Space Agency.

The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

## References

- Abbatt, J. P. D. and Molina, M. J.: Heterogeneous interactions of ClONO<sub>2</sub> and HCl on nitric acid trihydrate at 202 K, *J. Phys. Chem.*, 96, 7674–7679, doi:10.1021/j100198a036, 1992. 20567
- Bernath, P. F., McElroy, C., Abrams, M., Boone, C., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P., Colin, R., DeCola, P., DeMazière, M., Drummond, J., Dufour, D., Evans, W., Fast, H., Fussen, D., Gilbert, K., Jennings, D., Llewellyn, E., Lowe, R., Mahieu, E., McConnell, J., McHugh, M., McLeod, S., Michaud, R., Midwinter, C., Nassar, R., Nichitiu, F., Nowlan, C., Rinsland, C., Rochon, Y., Rowlands, N., Semeniuk, K., Simon, P., Skelto, R., Sloan, J., Soucy, M.-A., Strong, K., Tremblay, P., Turnbull, D., Walker, K., Walkty, I., Wardle, D., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry Experiment (ACE): mission overview, *Geophys. Res. Lett.*, 32, L15S01, doi:10.1029/2005GL022386, 2005. 20569
- Brakebusch, M., Randall, C. E., Kinnison, D. E., Tilmes, S., Santee, M. L., and Manney, G. L.: Evaluation of whole atmosphere community climate model simulations of ozone during Arctic winter 2004–2005, *J. Geophys. Res.*, in preparation, 2012. 20569
- Carlsaw, K. S. and Peter, T.: Uncertainties in reactive uptake coefficients for solid stratospheric particles – 1. surface chemistry, *Geophys. Res. Lett.*, 24, 1743–1746, 1997. 20566

20578

ACPD

12, 20561–20591, 2012

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carslaw, K. S., Luo, B. P., Clegg, S. L., Peter, T., Brimblecombe, P., and Crutzen, P. J.: Stratospheric aerosol growth and  $\text{HNO}_3$  gas phase depletion from coupled  $\text{HNO}_3$  and water uptake by liquid particles, *Geophys. Res. Lett.*, 21, 2479–2482, 1994. 20567, 20568

5 Carslaw, K. S., Luo, B., and Peter, T.: An analytic expression for the composition of aqueous  $\text{HNO}_3$ - $\text{H}_2\text{SO}_4$  stratospheric aerosols including gas phase removal of  $\text{HNO}_3$ , *Geophys. Res. Lett.*, 22, 1887–1880, 1995. 20568, 20575

Carslaw, K. S., Kettleborough, J. A., Northway, M. J., Davies, S., Gao, R., Fahey, D. W., Baumgardner, D. G., Chipperfield, M. P., and Kleinböhl, A.: A vortex-scale simulation of the growth and sedimentation of large nitric acid hydrate particles, *J. Geophys. Res.*, 107, 8300–8418, 2002. 20568

10 Crutzen, P.: Albedo enhancement by stratospheric sulfur injections: a contribution to resolve a policy dilemma?, *Climatic Change*, 77, 211–220, doi:10.1007/s10584-006-9101-y, 2006. 20564

Dameris, M., Peter, T., Schmidt, U., and Zellner, R.: Das Ozonloch und seine Ursachen, *Chem. unserer Zeit*, 41, 152–168, doi:10.1002/ciuz.200700423, 2007. 20585

15 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011. 20568

20 Douglass, A. R., Schoeberl, M. R., Stolarski, R. S., Waters, J. W., Russell III, J. M., Roche, A. E., and Massie, S. T.: Interhemispheric differences in springtime production of  $\text{HCl}$  and  $\text{ClONO}_2$  in the polar vortices, *J. Geophys. Res.*, 100, 13967–13978, doi:10.1029/95JD00698, 1995. 20565, 20566, 20567

25 Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Ann. Geophys.*, 30, 1055–1073, doi:10.5194/angeo-30-1055-2012, 2012. 20564

30 Elrod, M. J., Koch, R. E., Kim, J. E., and Molina, M. J.:  $\text{HCl}$  vapour pressures and reaction probabilities for  $\text{ClONO}_2 + \text{HCl}$  on liquid  $\text{H}_2\text{SO}_4$ - $\text{HNO}_3$ - $\text{HCl}$ - $\text{H}_2\text{O}$  solutions, *Faraday Discuss.*, 100, 269–278, 1995. 20567

Eyring, V., Sheperd, T. G., and Waugh, D. W.: SPARC report on the evaluation of chemistry-climate models, in: SPARC Rep. No. 5, WRCP-132, WMO-TD No. 1526, World Meteorol. Organ., Geneva, 434 pp., 2010. 20569

5 Fahey, D. W., Gao, R. S., Carslaw, K. S., Kettleborough, J., Popp, P. J., Northway, M. J., Holecck, J. C., Ciciora, S. C., McLaughlin, R. J., Thompson, T. L., Winkler, R. H., Baumgardner, D. G., Gandrud, B., Wennberg, R. O., Dhaniyala, S., McKinney, K., Peter, T., Salawitch, R. J., Bui, T. P., Elkins, J. W., Webster, C. R., Atlas, E. L., Jost, H., Wilson, J. C., Herman, R. L., Kleinböhl, A., and von König, M.: The detection of large HNO<sub>3</sub>-containing particles in the winter Arctic stratosphere, *Science*, 291, 1026–1031, doi:10.1126/science.1057265, 2001. 20563, 20565

10 Grooß, J.-U. and Müller, R.: Simulation of ozone loss in Arctic winter 2004/2005, *Geophys. Res. Lett.*, 34, L05804, doi:10.1029/2006GL028901, 2007. 20569

15 Grooß, J.-U., Günther, G., Konopka, P., Müller, R., McKenna, D. S., Stroh, F., Vogel, B., Engel, A., Müller, M., Hoppel, K., Bevilacqua, R., Richard, E., Webster, C. R., Elkins, J. W., Hurst, D. F., Romashkin, P. A., and Baumgardner, D. G.: Simulation of ozone depletion in spring 2000 with the Chemical Lagrangian Model of the Stratosphere (CLaMS), *J. Geophys. Res.*, 107, 8295, doi:10.1029/2001JD000456, 2002. 20569, 20572

20 Grooß, J.-U., Brauttsch, K., Pommrich, R., Solomon, S., and Müller, R.: Stratospheric ozone chemistry in the Antarctic: what determines the lowest ozone values reached and their recovery?, *Atmos. Chem. Phys.*, 11, 12217–12226, doi:10.5194/acp-11-12217-2011, 2011. 20566

Hanson, D. R.: Reaction of ClONO<sub>2</sub> with H<sub>2</sub>O and HCl in sulfuric acid and HNO<sub>3</sub>/H<sub>2</sub>SO<sub>4</sub>/H<sub>2</sub>O mixtures, *J. Phys. Chem. A*, 102, 4794–4807, doi:10.1021/jp972767s, 1998. 20566

Hanson, D. R. and Mauersberger, K.: Laboratory studies of the nitric acid trihydrate: implications for the south polar stratosphere, *Geophys. Res. Lett.*, 15, 855–858, 1988. 20566

25 Hanson, D. R. and Ravishankara, A. R.: Reaction of ClONO<sub>2</sub> with HCl on NAT, NAD and frozen sulfuric acid and hydrolysis of N<sub>2</sub>O<sub>5</sub> and ClONO<sub>2</sub> on frozen sulfuric acid, *J. Phys. Chem.*, 97, 12309–12319, doi:10.1021/j100149a035, 1993. 20567

Harris, N. R. P., Lehmann, R., Rex, M., and von der Gathen, P.: A closer look at Arctic ozone loss and polar stratospheric clouds, *Atmos. Chem. Phys.*, 10, 8499–8510, doi:10.5194/acp-10-8499-2010, 2010. 20565

30 Hofmann, D. J. and Solomon, S.: Ozone destruction through heterogenous chemistry following the eruption of El Chichón, *J. Geophys. Res.*, 94, 5029–5041, 1989. 20564

---

## Chlorine activation on stratospheric aerosols

T. Wegner et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Homan, C. D., Volk, C. M., Kuhn, A. C., Werner, A., Baehr, J., Viciani, S., Ulanovski, A., and Ravegnani, F.: Tracer measurements in the tropical tropopause layer during the AMMA/SCOUT-O3 aircraft campaign, *Atmos. Chem. Phys.*, 10, 3615–3627, doi:10.5194/acp-10-3615-2010, 2010. 20568

5 Kawa, S. R., Newman, P. A., Lait, L. R., Schoeberl, M. R., Stimpfle, R. M., Kohn, D. W., Webster, C. R., May, R. D., Baumgardner, D., Dye, J. E., Wilson, J. C., Chan, K. R., and Loewenstein, M.: Activation of chlorine in sulfate aerosol as inferred from aircraft observations, *J. Geophys. Res.*, 102, 3921–3933, 1997. 20564

10 Khosrawi, F., Urban, J., Pitts, M. C., Voelger, P., Achtert, P., Kaphlanov, M., Santee, M. L., Manney, G. L., Murtagh, D., and Fricke, K.-H.: Denitrification and polar stratospheric cloud formation during the Arctic winter 2009/2010, *Atmos. Chem. Phys.*, 11, 8471–8487, doi:10.5194/acp-11-8471-2011, 2011. 20563

Lien, C.-Y., Lin, W.-Y., Chen, H.-Y., Huang, W.-T., Jin, B., Chen, I.-C., and Lin, J. J.: Photodissociation cross sections of ClOCl at 248.4 and 266 nm, *J. Chem. Phys.*, 131, 174301, doi:10.1063/1.3257682, 2009. 20573

15 Lowe, D. and MacKenzie, A. R.: Polar stratospheric cloud microphysics and chemistry, *J. Atmos. Sol.-Terr. Phys.*, 70, 13–40, 2008. 20563

20 Matthey, R., Cacciani, M., Fiocco, G., Martinez, A. A., Martucci, G., Mitev, V., Pace, G., and Stefanutti, L.: Observations of aerosol and clouds with the ABLE and MAL lidars during the mid-latitude and Arctic ENVISAT validation campaigns, in: *European Rocket and Balloon Programmes and Related Research*, vol. 530, edited by: Warmbein, B., ESA Special Publication, 579–584, 2003. 20572

25 McKenna, D. S., Groöf, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 2. formulation of chemistry scheme and initialization, *J. Geophys. Res.*, 107, 4256, doi:10.1029/2000JD000113, 2002a. 20568

30 McKenna, D. S., Konopka, P., Groöf, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 1. formulation of advection and mixing, *J. Geophys. Res.*, 107, 4309, doi:10.1029/2000JD000114, 2002b. 20568

Molina, L. T. and Molina, M. J.: Production of Cl<sub>2</sub>O<sub>2</sub> from the self-reaction of the ClO radical, *J. Phys. Chem.*, 91, 433–436, doi:10.1021/j100286a035, 1987. 20563

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Müller, R., Peter, T., Crutzen, P. J., Oelhaf, H., Adrian, G. P., von Clarmann, T., Wegner, A., Schmidt, U., and Lary, D.: Chlorine chemistry and the potential for ozone depletion in the arctic stratosphere in the winter of 1991/1992, *Geophys. Res. Lett.*, 21, 1427–1430, 1994. 20565, 20567
- 5 Peter, T.: Microphysics and heterogeneous chemistry of polar stratospheric clouds, *Annu. Rev. Phys. Chem.*, 49, 785–822, 1997. 20568
- Peter, T. and Grooß, J. U.: *Polar Stratospheric Clouds and Sulfate Aerosol Particles: Microphysics, Denitrification and Heterogeneous Chemistry*, chap. 4, RSC Publishing, Cambridge, 2012. 20563, 20565
- 10 Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, doi:10.5194/acp-9-7577-2009, 2009. 20574
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, doi:10.5194/acp-11-2161-2011, 2011. 20563, 20574
- 15 Plenge, J., Köhl, S., Vogel, B., Müller, R., von Hobe, M., Flesch, R., and Rühl, R.: Bond strength of chlorine peroxide, *J. Phys. Chem. A*, 109, 6730–6734, doi:10.1021/jp044142h, 2005. 20573
- Portmann, R. W., Solomon, S., Garcia, R. R., Thomason, L. W., Poole, L. R., and McCormick, M. P.: Role of aerosol variations in anthropogenic ozone depletion in the polar regions, *J. Geophys. Res.*, 101, 22991–23006, 1996. 20564, 20565, 20567
- 20 Rasch, P. J., Crutzen, P. J., and Coleman, D. B.: Exploring the geoengineering of climate using stratospheric sulfate aerosols: the role of particle size, *Geophys. Res. Lett.*, 35, L02809, doi:10.1029/2007GL032179, 2008. 20564
- 25 Robock, A.: Volcanic eruptions and climate, *Rev. Geophys.*, 38, 191–219, doi:10.1029/1998RG000054, 2000. 20564
- Rodriguez, J. M., Ko, M. K. W., and Sze, N. D.: Antarctic chlorine chemistry: possible global implications, *Geophys. Res. Lett.*, 15, 257–260, 1988. 20564
- 30 Sander, S., Friedl, R. R., Golden, D. M., Kurylo, M. J., Moortgat, G. K., Wine, P. H., Ravishankara, A. R., Kolb, C. E., Molina, M. J., Finlyason-Pitts, B. J., Huie, R. E., and Orkin, V. L.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, evaluation number 17, JPL Publication, 10-6, 2006. 20568

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Shi, Q., Jayne, T., Kolb, C. E., Worsnop, D. R., and Davidovits, P.: Kinetic model for reaction of  $\text{ClONO}_2$  with  $\text{H}_2\text{O}$  and  $\text{HCl}$  and  $\text{HOCl}$  with  $\text{HCl}$  in sulfuric acid solutions, *J. Geophys. Res.*, 106, 24259–24274, 2001. 20566, 20567
- Shur, G. N., Yushkov, V. A., Drynkov, A. V., Fadeeva, G. V., and Potertikova, G. A.: Study of thermodynamics of the stratosphere at high latitudes of the Northern Hemisphere on the M-55 geofizika flying laboratory, *Russ. Meteorol. Hydrol.*, 8, 43–53, 2006. 20569
- Solomon, S.: Stratospheric ozone depletion, a review of concepts and history, *Rev. Geophys.*, 37, 275–316, 1999. 20563, 20564, 20565
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, 1986. 20563
- Sumińska-Ebersoldt, O., Lehmann, R., Wegner, T., Groß, J.-U., Hösen, E., Weigel, R., Frey, W., Griessbach, S., Mitev, V., Emde, C., Volk, C. M., Borrmann, S., Rex, M., Stroh, F., and von Hobe, M.:  $\text{ClOOCl}$  photolysis at high solar zenith angles: analysis of the RECONCILE self-match flight, *Atmos. Chem. Phys.*, 12, 1353–1365, doi:10.5194/acp-12-1353-2012, 2012. 20573
- Tilmes, S., Müller, R., and Salawitch, R.: The sensitivity of polar ozone depletion to proposed geoengineering schemes, *Science*, 32, 1201–1204, doi:10.1126/science.1153966, 2008a. 20564
- Tilmes, S., Müller, R., Salawitch, R. J., Schmidt, U., Webster, C. R., Oelhaf, H., Camy-Peyret, C. C., and Russell III, J. M.: Chemical ozone loss in the Arctic winter 1991–1992, *Atmos. Chem. Phys.*, 8, 1897–1910, doi:10.5194/acp-8-1897-2008, 2008b. 20564
- Voigt, C., Schlager, H., Luo, B. P., Dörnbrack, A., Roiger, A., Stock, P., Curtius, J., Vössing, H., Borrmann, S., Davies, S., Konopka, P., Schiller, C., Shur, G., and Peter, T.: Nitric Acid Trihydrate (NAT) formation at low NAT supersaturation in Polar Stratospheric Clouds (PSCs), *Atmos. Chem. Phys.*, 5, 1371–1380, doi:10.5194/acp-5-1371-2005, 2005. 20568
- von Hobe, M., Groß, J.-U., Müller, R., Hrechanyy, S., Winkler, U., and Stroh, F.: A re-evaluation of the  $\text{ClO}/\text{Cl}_2\text{O}_2$  equilibrium constant based on stratospheric in-situ observations, *Atmos. Chem. Phys.*, 5, 693–702, doi:10.5194/acp-5-693-2005, 2005. 20568
- von Hobe, M., Ulanovsky, A., Volk, C. M., Groß, J.-U., Tilmes, S., Konopka, P., Günther, G., Werner, A., Spelten, N., Shur, G., Yushkov, V., Ravegnani, F., Schiller, C., Müller, R., and Stroh, F.: Severe ozone depletion in the cold Arctic winter 2004–2005, *Geophys. Res. Lett.*, 33, L17815, doi:10.1029/2006GL026945, 2006. 20568, 20569

**Chlorine activation  
on stratospheric  
aerosols**


T. Wegner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- von Hobe, M., Salawitch, R. J., Canty, T., Keller-Rudek, H., Moortgat, G. K., Grooß, J.-U., Müller, R., and Stroh, F.: Understanding the kinetics of the ClO dimer cycle, *Atmos. Chem. Phys.*, 7, 3055–3069, doi:10.5194/acp-7-3055-2007, 2007. 20573
- 5 von Hobe, M., Stroh, F., Beckers, H., Benter, T., and Willner, H.: The UV/Vis absorption spectrum of matrix-isolated dichlorine peroxide, ClOOCl, *Phys. Chem. Chem. Phys.*, 11, 1571–1580, 2009. 20573
- 10 Weigel, R., Hermann, M., Curtius, J., Voigt, C., Walter, S., Böttger, T., Lepukhov, B., Belyaev, G., and Borrmann, S.: Experimental characterization of the COndensation PArticle counting System for high altitude aircraft-borne application, *Atmos. Meas. Tech.*, 2, 243–258, doi:10.5194/amt-2-243-2009, 2009. 20568
- 15 Werner, A., Volk, C. M., Ivanova, E. V., Wetter, T., Schiller, C., Schlager, H., and Konopka, P.: Quantifying transport into the Arctic lowermost stratosphere, *Atmos. Chem. Phys.*, 10, 11623–11639, doi:10.5194/acp-10-11623-2010, 2010. 20568
- Zöger, M., Afchine, A., Eicke, N., Gerhards, M.-T., Klein, E., McKenna, D., Mörschel, U., Schmidt, U., Tan, V., Tuitjer, F., Woyke, T., and Schiller, C.: Fast in situ stratospheric hydrometers: a new family of balloon-borne and airborne Lyman alpha photofragment fluorescence hydrometers, *J. Geophys. Res.*, 104, 1807–1816, 1999. 20568

## Chlorine activation on stratospheric aerosols

T. Wegner et al.



**Fig. 1.** First order loss rates for  $\text{ClONO}_2 + \text{HCl}$  (a) and  $\text{ClONO}_2 + \text{H}_2\text{O}$  (b) for different parameterizations and aerosol types for typical stratospheric conditions (50 hPa, 5 ppmv  $\text{H}_2\text{O}$ , 1 ppbv HCl, 0.5 ppbv  $\text{ClONO}_2$ , 10 ppbv  $\text{HNO}_3$ , 0.15 ppbv  $\text{H}_2\text{SO}_4$  and 10 background aerosol particles  $\text{cm}^{-3}$ ). Solid red and blue lines depict ternary aerosol, dashed lines binary aerosol. Solid green and orange lines represent NAT particles with density  $10^{-1} \text{ cm}^{-3}$ , dashed lines  $10^{-4} \text{ cm}^{-3}$ . Adapted from Dameris et al. (2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.



**Fig. 2.** Top: temperatures interpolated from ERA-INTERIM data on the flightpath versus TDC temperature measurements on board the Geophysica as function of potential temperature. Bottom: vertical profile of temperature differences averaged into 3 K bins.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 3.** Temperature history (a), simulated condensation of HNO<sub>3</sub> on STS (b), NAT (c) and corresponding enhancement of the surface area density over the background aerosol (d) along 92 h back-trajectories ending on the Geophysica flightpath. Areas under  $T_{\text{NAT}}$  are enclosed by the black line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.



**Fig. 4.** Comparison of HALOX in-situ data and CLaMS simulations. The solid black line marks the HALOX ClO measurement with its accuracy shaded grey and blue the simulations with the error bars showing their uncertainty.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Chlorine activation on stratospheric aerosols

T. Wegner et al.




**Fig. 5.** Top: temperature history along 7 day back-trajectories from HALOX measurements for 3 flights in January 2010. Middle:  $\text{ClO}_x$  initialization (black) and modeled  $\text{ClO}_x$  on the flightpath with full PSC surface area density (red) and background aerosol only (blue dashed). Bottom: measured (black) and modeled values of  $\text{ClO}$  (red).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.




**Fig. 6.** HCl and HNO<sub>3</sub> observations by MLS for the Arctic winters 2004/2005 to 2010/2011 on 500 K potential temperature in the vortex core (equivalent latitude > 75° N).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Chlorine activation  
on stratospheric  
aerosols**

T. Wegner et al.



**Fig. 7.** Probability density distribution of maximum surface area density enhancement by STS along 7 day back-trajectories starting on MLS observations poleward of  $75^{\circ}$  N equivalent latitude.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

