

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

The effect of regional changes in anthropogenic aerosols on rainfall of the East Asian Summer Monsoon

L. Guo¹, E. J. Highwood¹, L. C. Shaffrey², and A. G. Turner²

¹Department of Meteorology, University of Reading, Reading, UK

²NCAS-Climate, University of Reading, Reading, UK

Received: 31 July 2012 – Accepted: 24 August 2012 – Published: 5 September 2012

Correspondence to: L. Guo (l.guo@reading.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The response of East Asian Summer Monsoon (EASM) precipitation to long term changes in regional anthropogenic aerosols (sulphate and black carbon) is explored in an atmospheric general circulation model, the atmospheric component of the UK High-Resolution Global Environment Model v1.2 (HiGAM). Separately, sulphur dioxide (SO₂) and black carbon (BC) emissions in 1950 and 2000 over East Asia are used to drive model simulations, while emissions are kept constant at year 2000 level outside this region. The response of the EASM is examined by comparing simulations driven by aerosol emissions representative of 1950 and 2000. The aerosol radiative effects are also determined using an off-line radiative transfer model. During June, July and August, the EASM was not significantly changed as either SO₂ or BC emissions increased from 1950 to 2000 levels. However, in September, precipitation is significantly decreased when emissions are at the 2000 level. The cooler land surface temperature over China in September due to increased aerosols reduces the surface thermal contrast that supports the EASM circulation. However, mechanisms causing the surface temperature decrease in September are different between sulphate and BC experiments. In the sulphate experiment, the sulphate direct and the 1st indirect radiative effects contribute to the surface cooling. In the BC experiment, the BC direct effect is the main driver of the surface cooling, however, a decrease in low cloud cover due to the increased heating by BC absorption partially counteracts the direct effect. This results in a weaker land surface temperature response to BC changes than to sulphate changes. The resulting precipitation response is also weaker, and the responses of the monsoon circulation are different for sulphate and black carbon experiments. This study demonstrates a mechanism that links regional aerosol emission changes to the precipitation changes of the EASM, and it could be applied to help understand the future changes in EASM precipitation in CMIP5 simulations.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Monsoons play a key role in global mass and heat transport (Trenberth et al., 2000). The Asian monsoon (including both the Indian monsoon and East Asian monsoons) is the strongest monsoon system due to the combination of thermal contrast between the largest continent and the Pacific and Indian Oceans and the presence of the highest terrain (Wang and Ding, 2006). It affects the lives of more than a third of the world's population in South and East Asia.

Over the past several decades, rapid industrialisation over Asia has resulted in a dramatic increase in emissions of aerosols and aerosol precursor gases (Smith et al., 2004; Nozawa et al., 2007). Aerosols have the potential to affect precipitation by modulating radiation in the atmosphere and at the surface (Haywood and Boucher, 2000) and by changing cloud microphysics (Twomey, 1977; Albrecht, 1989).

In recognition of the need to understand the variability of the monsoon and the industrialisation mentioned above, there have been a number of studies concerning aerosol and monsoon interaction in the past decade. Early studies considered only scattering aerosol, like sulphate (e.g. Boucher et al., 1998; Iwasaki and Kitagawa, 1998). An increase in scattering aerosols results in a negative direct radiative forcing at the top of the atmosphere (TOA), which reduces the short-wave radiation reaching the surface (also called “solar dimming”) and acts to cool the global surface temperature. During summer, the land-sea surface temperature gradient over India is weakened in the presence of sulphate aerosol. Since the monsoon is driven by thermal contrast between continent and ocean, any weakening of this would weaken the monsoon circulation and decrease precipitation. Even though different models and methods are used in different studies, the result that increased sulphate aerosol suppresses monsoon circulation and decreases precipitation over South and East Asia is a consistent conclusion (Boucher et al., 1998; Iwasaki and Kitagawa, 1998; Huang et al., 2007; Randles and Ramaswamy, 2008). In those transient climate change simulations which consider sulphate, the cooling effects of sulphate are found to partially counteract the effects of

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



increased carbon dioxide on monsoon precipitation (e.g. Roeckner et al., 1999; Ashrit et al., 2003; Johns et al., 2003).

The role of some aerosols in absorbing short-wave radiation over South Asia and the north Indian Ocean (the Asian Blown Cloud, ABC) was identified by Ramanathan et al. (2001). Subsequently, many researchers began to consider the mechanisms by which absorbing aerosol affects the monsoons over South and East Asia (Menon et al., 2002; Chung and Zhang, 2004; Chung and Ramanathan, 2006; Ramanathan et al., 2005; Lau et al., 2006; Wang, 2007; Meehl et al., 2008; Randles and Ramaswamy, 2008; Ming et al., 2010). Most of these studies are based on GCM simulations which compare the results of experiments with and without aerosol effects. However, unlike the consistent results for sulphate aerosol, the results of black carbon (BC) studies are diverse. In particular, the results appear to differ depending on whether an atmosphere-only model or a coupled ocean-atmosphere model is used. In coupled models, the Indian monsoon is weakened as the meridional sea surface temperature (SST) gradient over the north Indian Ocean is reduced by surface cooling caused by the increased BC (Ramanathan et al., 2005; Meehl et al., 2008). However, in atmosphere-only models, BC increases the precipitation over Indian subcontinent because the ascent is intensified by atmospheric heating due to the increased BC concentration (Menon et al., 2002; Randles and Ramaswamy, 2008). Chung and Ramanathan (2006) compared the relative importance of the weakening meridional SST gradient over the Northern Indian Ocean and the atmospheric heating due to BC. They concluded that the precipitation patterns were opposite for the two cases and that the weakening meridional SST gradient over the northern Indian Ocean dominated the precipitation pattern.

The above mentioned studies of impacts have focused on the Indian monsoon; so far there are far fewer studies focusing on East Asia. Those that do exist (e.g. Wang, 2007; Kim et al., 2007; Lau et al., 2006) have drawn different conclusions as to impacts of aerosol (both sulphate and BC) on the East Asian monsoon based on modelling studies. For example, Wang (2007) suggested that the BC heating in a few specific areas such as the east and the west Pacific regions can have a remote impact on

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the East Asian Summer Monsoon (EASM) by causing an “ENSO-like” effect on the atmospheric circulation. Kim et al. (2007) suggested that the direct effect globally increased sulphate emissions induces changes in the north Pacific storm track during boreal spring, which in turn changes circulation and reduces rainfall over Central East Asia. Lau et al. (2006) suggested that changes of the East Asia rain belt are associated with the increased loading of absorbing aerosol over northern India during late spring. These proposed mechanisms are large-scale and not necessarily related to local aerosol change in the East Asian monsoon region, which is in contrast to the studies of the aerosol impact on the Indian monsoon. In this study, we will focus on whether local changes in aerosol can affect the EASM.

It is important to note that GCM simulations studies (e.g. Boer and Yu, 2003) suggest that the regional pattern of temperature response to a particular forcing cannot be mapped directly from the regional pattern of forcing. Changes in temperature and other fields seen in East Asia may therefore be influenced by changes to radiation and clouds in remote locations. However, in this study, both the global aerosol impacts and global responses of surface temperature have been examined and seem to be small outside the region in which aerosol emissions are changed (East Asia), therefore, discussion will focus on the regional changes over East Asia.

Experiments for sulphate and BC aerosols are carried out separately. The model descriptions and experimental design are given in Sect. 2. The changes in aerosol concentration and cloud properties are given in Sect. 3. The responses of the EASM in surface temperature, precipitation and circulation to sulphate and BC are given in Sect. 4. Conclusions and discussions are given in Sect. 5.

2 Models and experiments

2.1 Models

2.1.1 HiGAM

The general circulation model used in this study is the atmospheric component of the UK High-resolution Global Environment Model (HiGAM), version 1.2 (Shaffrey et al., 2009). The horizontal resolution of HiGAM is 0.83° latitude \times 1.25° longitude; vertically HiGAM has 38 levels and the top of the atmosphere is 39 km. HiGAM uses a non-hydrostatic, fully compressible dynamical core, with semi-Lagrangian advection. It is discretized horizontally on the Arakawa C-grid and vertically on the Charney-Phillips staggered grid (Davies et al., 2005). The physical parametrisations in HiGAM are similar to those in HadGAM1 (Hadley Centre Global Environmental Model, with horizontal resolution of 1.25° latitude \times 1.875° longitude) which has been summarised by Martin et al. (2006). Performance of the coupled version of HiGAM has been evaluated by Shaffrey et al. (2009), in which they noted that the higher resolution decreases SST errors, improves small scale coupling and the simulation of ENSO.

There are four types of aerosol species included in HiGAM: sulphate, black carbon, biomass burning material and sea salt. Sulphur dioxide (SO_2 , a precursor gas for sulphate) and BC are two major anthropogenic aerosol sources over East Asia. Figure 1a shows the annual mean emissions of SO_2 in 2000 (data from Smith et al., 2004). East Asia (defined here as $5\text{--}60^\circ$ N, $90\text{--}150^\circ$ E) is one of the three strongest emission regions in the world, the annual mean emission intensity being about $39.5 \text{ Tg}(\text{SO}_2) \text{ yr}^{-1}$; the other two important regions are Western Europe and the east coast of USA. Compared to the other two regions, East Asia has been a rapidly industrialising region since the 1950s, the increase of emissions has been dramatic. According to the historical emission data from Smith et al. (2004), in 1950 over East Asia the annual mean emission intensity of SO_2 was $3.5 \text{ Tg}(\text{SO}_2) \text{ yr}^{-1}$; since then, the emissions of SO_2 have been increased by ten-fold in the second half of the twentieth century. However, over

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the other two intensive emission regions, the peaks of emissions appeared during the 1970s, and were twice the magnitude of emissions in 1950. BC emissions have a similar geographical pattern as SO_2 , as both of them are primarily emitted from fossil fuel burning due to anthropogenic activities. According to historical emission data from Nozawa et al. (2007), the annual emission intensity of BC over East Asia in 1950 was $0.3 \text{ Tg(C) yr}^{-1}$, which increased to $2.4 \text{ Tg(C) yr}^{-1}$ by 2000.

The sulphate scheme is described by Jones et al. (2001) and Woodage et al. (2003), in which the precursor gases, SO_2 and dimethyl sulphate ($(\text{CH}_3)_2\text{SO}_4$), undergo chemical reactions with oxidants in the atmosphere to form sulphate aerosol in both gaseous and aqueous phases. These are then finally removed from the atmosphere by dry and wet deposition. Sulphate aerosol is treated as hydrophilic in HiGAM, and the optical properties of sulphate vary with the relative humidity. Sulphate aerosol can act as a cloud condensation nuclei (CCN) and change cloud reflectivity and cloud lifetime by changing cloud droplet number concentration (CDNC). The BC scheme (Roberts and Jones, 2004) is different in that BC is assumed to be emitted as primary particles. The BC is treated as hydrophobic and therefore insufficiently soluble to act as CCN in the model. Thus, BC cannot affect clouds micro-physically in HiGAM, but it can still interact with clouds by absorbing solar radiation and changing the heating rate in the atmosphere, which is defined as the semi-direct effect by previous studies (Hansen et al., 1997; Ackerman et al., 2000).

2.1.2 Edwards-Slingo radiative transfer model

An off-line radiative transfer model is also used in this work to estimate aerosol radiative effects. This is the same radiation scheme used in HiGAM. This model was first introduced by Edwards and Slingo (1996) (E-S code hereafter) and it calculates radiative fluxes by summing the results of a number of quasi-monochromatic fluxes, each carried out using a two-stream approximation.

To estimate the aerosol radiative effect in HiGAM, the single column version of E-S code has been expanded into a 3-dimensional version. The atmospheric state vari-

ables (temperature, pressure and relative humidity), various radiative agents (greenhouse gases, aerosols and clouds) on each grid point are taken from the monthly mean output of HiGAM. Myhre et al. (2002) pointed out that calculation of the radiative effect is sensitive to spatial and temporal resolution. The coarser the resolution, the more underestimated the radiative effect will be. The coarser resolution averages and smooths out the small scale relative humidity fluctuations, as sulphate optical properties increase non-linearly with relative humidity, the sulphate radiative effect estimated using smoothed relative humidity is smaller than the estimates with high value small scale relative humidity fluctuations. The aerosol radiative effects shown here use the monthly output of HiGAM. Therefore, the magnitude is likely to be underestimated, but it can still be used qualitatively to explain the different impacts from different aerosol species.

The definition of the aerosol radiative effect in this study is similar to the definition of aerosol radiative forcing in other studies, including the IPCC report (e.g. Forster et al., 2007; Haywood and Shine, 1997). However, there are two differences between the effect used in this study and the forcing in others. Temporally, the radiative effect refers to the change in monthly (or even shorter time interval) net radiation rather than annual mean. Spatially, the radiative effect is a regional estimate over East Asia rather than global mean.

2.2 Experiment design

The experiments in this study are designed to explore impacts of aerosol on the EASM. Three experiments are carried out (see Table 1). In the Control experiment, global emissions of both SO₂ and BC are used at the 2000 level. Sea surface temperature and sea ice data are from AMIP-II (<http://www-pcmdi.llnl.gov/projects/amip/>), greenhouse gases except ozone are given as fixed values, while ozone uses the time series of the Stratospheric Processes and Their Role in Climate (SPARC) program. More details of model configuration can be found in Martin et al. (2006); Shaffrey et al. (2009). In the SO₂-1950 experiment, SO₂ emissions at the 1950 level are used over East Asia (5–

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



60° N, 90–150° E), but SO₂ emissions are kept at 2000 level over the rest of the world (as shown in Fig. 1b). In the BC_1950 experiment, BC emissions at the 1950 level are used over East Asia, while those over the rest of world are at the 2000 level. The Control experiment is an eighteen-year AMIP-type run (Gates et al., 1999), driven by observed monthly mean SST from 1983 to 2000. The SO₂-1950 and BC_1950 simulations consist of an 18 member ensembles of the period April to September, beginning in each April of the control run. The validity of this experimental design is based on the fact that the lifetimes of both sulphate and BC aerosols are short. In HiGAM, the global mean lifetime of sulphate is four days and it is six days for BC. Aerosol concentration in the atmosphere adjusts quickly to the change in emissions and long continuous runs are therefore not necessary.

2.3 Evaluation of East Asian Summer Monsoon in HiGAM

Figure 2a and b compare the global JJA precipitation in HiGAM with GPCP data (Adler et al., 2003). HiGAM generally captures the global pattern of precipitation, although there are specific biases. Over East Asia, the Mei-yu/Baiyu/Changma front in HiGAM is weaker than observed. This is due to the weakness of the subtropical high over the western North Pacific Ocean. The Indian monsoon precipitation has not been simulated well in HiGAM (Shaffrey et al., 2009) but a similar problem are also shown in other GCMs (Annamalai et al., 2007). Figure 2c and d show the 850 hPa geopotential height and wind over East Asia. Comparing to ERA-40 data, the Somali jet is too strong over the Indochina Peninsula and the South China Sea, and reaches further east. This is associated with the southeasterlies along the western flank of the Northwestern Pacific Subtropical Anticyclone. However, HiGAM captures the most prominent features of the EASM, the Somali jet over the northern Indian Ocean and the subtropical anticyclone over the western North Pacific Ocean.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



droplet size: smaller cloud drops make cloud more reflective. The radiation budgets at TOA and the surface are therefore sensitive to the changes in r_e , as we will discuss in more detail in Sect. 4.

However, the column burdens of black carbon in both Control and BC_1950 experiments do not show any clear seasonal variation (Fig. 3a). This is because BC emissions no have seasonal variation in HiGAM, also that BC is emitted as primary particles in HiGAM and its concentration depends more on physical conditions of the atmosphere rather than photochemistry. As BC cannot act as CCN in HiGAM, changes of BC column burden hardly effect r_e and cloud drops remain at a similar size to those in the Control experiment (Fig. 3b).

Since sources of anthropogenic emission are mainly confined to the surface, and the lifetimes of both sulphate and black carbon are short, the vertical distributions of sulphate and black carbon are uneven with larger concentrations confined within 2 km above the surface. This vertical distribution of aerosols makes the aerosol indirect effect on low level cloud more important than other types of cloud, thus the strong aerosol-cloud interactions in HiGAM occur at the bottom of the atmosphere.

Figure 3c shows eighteen-year monthly and area-mean low cloud fraction over East Asia in the three experiments. A similar seasonal cycle can be observed in Fig. 3c for all the experiments. Low cloud fraction decreases from April, stays small during JJA, and increases again in September when the EASM withdraws from the region under consideration. Yu et al. (2004) suggested that the larger low cloud fraction (mainly stratus) in pre- and post-monsoon season are generated and maintained by the frictional and blocking effect of the Tibetan Plateau. During the summer monsoon season, as more active convection is occurring over East Asia, mid and high-level clouds are dominant.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Climate responses of the East Asian Summer Monsoon

4.1 Sulphate experiment

Figure 4 shows the monthly change of surface temperature and precipitation averaged over East Asia; results are subjected to Student's t-test and changes exceeding 95 % significance are hatched. As sulphate has increased (Fig. 4a), both surface temperature and precipitation decrease in each month (Control minus SO₂-1950). Significant decreases of surface temperature occur in May and September only, while significant decreases of precipitation occur in June and September only. The surface temperature decrease in May is due to the increased low cloud cover with smaller cloud droplet size which significantly increases reflectivity in the Control experiment. However, there is no corresponding significant decrease in precipitation in May, because the EASM has not yet advanced over this region. The small change in precipitation is mainly due to changes in local evaporation. Precipitation in June has decreased significantly by nearly 13 %, however there is no corresponding significant decrease in surface temperature. The decrease of precipitation is distributed rather locally in June (not shown), and is mainly confined to Southern China where the monsoon front is located. This decrease of precipitation in June indicates a delayed onset of the EASM due to cooler land surface temperatures in May and a weaker monsoon onset. However, as monsoon precipitation is dominated by convection, which could occur above the aerosol layer, and as monsoon precipitation has strong inter-annual variations, the impacts of aerosol in these experiments are not significant on the EASM during JJA.

Both surface temperature and precipitation show their biggest decrease in September. These changes in surface temperature and precipitation are related to aerosol radiative effects. Two different aerosol radiative effects have been calculated in this study using the E-S code. The “direct radiative effect”, is calculated as the difference of net radiation at both TOA and surface due to the change in sulphate concentration, but with the atmospheric state variables (temperature, pressure and relative humidity) and other radiative agents unchanged. The “combined radiative effect”, is calculated from

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the difference of the net radiation at both TOA and surface by considering changes in both sulphate concentration and cloud droplet size predicted by the model. Note that, since the changes in cloud cover and cloud liquid water content are not significant, only changes in cloud droplet size are considered in the off-line model.

5 The black lines in Fig. 5a and b show the monthly sulphate direct radiative effect at TOA and the surface averaged over East Asia (20–45° N, 100–122° E). Sulphate exerts a negative radiative effect over East Asia as SO₂ emissions increase from 1950 to 2000. The magnitude of the sulphate direct radiative effects at TOA and the surface are comparable, as sulphate is a scattering aerosol and little radiative energy is absorbed in the atmosphere. As shown in Fig. 5a and b, the negative radiative effect of sulphate is stronger during the summer monsoon season (JJA). This is consistent with the difference of sulphate column burden between the Control and SO₂_1950 experiment (shown as the solid blue line and the dashed blue lines of Fig. 3a), and indicates that more sulphate in the atmosphere during JJA reflects more solar radiation back to space, and reduces the incident solar radiation at the surface. However, the sulphate direct radiative effect is small in September, so it would appear that this does not explain the stronger decreases in surface temperature and precipitation at that time.

15 The blue lines in Fig. 5a and b show the monthly sulphate combined radiative effect at TOA and the surface averaged over East Asia. When the decrease of cloud droplet size is taken into account, the negative radiative effect of sulphate is significantly intensified in April, May and September. The monthly variation of the sulphate combined radiative effect is similar to the variation of surface temperature and precipitation, especially in September, the stronger negative sulphate radiative effects being co-located with stronger cooling at the surface and the stronger reduction in precipitation. In general, the sulphate combined effect is larger than the sulphate direct effect. This is because with larger low cloud fraction in April and September over East Asia, the sulphate-cloud interaction becomes important, especially in September.

25 Figure 7a shows the geographical distribution of the eighteen-year sulphate combined effect at the surface in September. It is negative over East Asia with the centre

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

over the lee side of the Tibetan Plateau where low cloud fraction is large. Figure 7b shows the eighteen-year mean changes in surface temperature in September due to sulphate increase (Control minus SO_2_{1950}). Surface temperature has a negative change which is consistent with the sulphate combined effect at the surface. Surface temperature decreases over South-Eastern China, Central China, the southern slopes of the Tibetan Plateau and parts of the Indochina Peninsula. Figure 7c shows the eighteen-year mean changes in precipitation in September (Control minus SO_2_{1950}). The precipitation significantly decreases over Central China, South-Western China and parts of the Indochina Peninsula.

There are two mechanisms that can be used to explain the decrease of precipitation over East Asia. One is that the local evaporation will decrease as the surface temperature decreases. However, the decrease of local evaporation in September ($-0.11 \text{ mm day}^{-1}$ over East Asia) is too small to explain the precipitation decrease ($-0.82 \text{ mm day}^{-1}$). Another explanation is that the change in surface temperature will induce changes in monsoon circulation. As the land surface temperature is decreased, the thermal contrast between land and ocean is also weakened. Therefore, the moisture transport is also weakened. For the EASM, the transportation of moisture mainly comes from two passages: from the Pacific Ocean in the east and Indian Ocean in the west.

To examine the changes in EASM circulation, the eighteen-year mean change of vertically integrated (from surface to 700 hPa) moisture flux and moisture flux divergence are examined for sulphate change, and is shown in Fig. 7d. Both the geographical patterns and magnitudes of changes in moisture flux divergence are consistent with changes in precipitation, suggesting that changes in the moisture transported from the adjacent oceans is the major factor contributing to precipitation decrease over East Asia. The changes of moisture flux as sulphate increases show a weakening of the EASM low level circulation. Both of the moisture transportation passages are weakened.

4.2 Black carbon experiment

Figure 4b shows monthly changes of surface temperature and precipitation due to the increase of BC averaged over East Asia. Unlike the responses in the sulphate experiment in which both surface temperature and precipitation consistently decrease, the sign of responses of surface temperature and precipitation in the black carbon experiment are varied. However, similar to the sulphate experiment, both surface temperature and precipitation decrease significantly in September.

The black lines in Fig. 5c and d show the monthly BC direct radiative effects at TOA and the surface averaged over East Asia (20–45° N, 100–122° E). Unlike the sulphate direct radiative effect which has comparable value between TOA and the surface, the BC direct radiative effect has opposite signs at the TOA and surface: positive at TOA and negative at the surface. An increase in the BC concentration led to more solar radiation being absorbed by black carbon. As more solar radiation is absorbed, less reaches the surface, thus a negative radiative effect is shown at surface. However, the surface-atmosphere as a whole gains energy, therefore, a positive radiative effect at TOA.

The blue lines in Fig. 5c and d show the monthly BC combined effects at TOA and the surface averaged over East Asia. Since BC does not act as CCN in the model, the change of cloud droplet size is not considered in the combined effect (as shown in Fig. 3b). Even though the low cloud fraction is not significantly changed in BC_1950 experiments (red line in Fig. 3c), the increases of low level cloud in September are bigger than that in Control experiment. Figure 6 shows the change of low cloud fraction due to BC changes (Control minus BC_1950) in September. The significant decrease of low level cloud is over the lee-side of the Tibetan Plateau, where the low cloud cover is building up during September. This local decrease in low cloud fraction cannot be seen when averaged over East Asia as in Fig. 3c, but it will change the BC radiative effects regionally.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 8a shows the BC combined radiative effect at the surface in September. The BC direct radiative effect is compensated by a decrease in low cloud cover over the lee side of the Tibetan Plateau. This is because the increased BC concentration exerts a negative radiative effect at the surface, but as low cloud cover is reduced due to BC heating, more solar radiation reaches the surface which partially compensates for the BC direct radiative effect. Comparing the sulphate combined radiative effect to the BC combined radiative effect at the surface, the patterns are different, and the responses of the EASM are also different.

Figure 8b shows the eighteen-year mean changes of surface temperature in September in the BC experiment. Compared to the sulphate experiment, the extent of the surface temperature decrease is smaller; it is only significant over South-Eastern China and parts of Central China; the magnitude is also smaller. Figure 8c shows the eighteen-year mean changes of precipitation in September in the BC experiment. The decrease of precipitation in the BC experiment is weaker, and is mainly over Central China. Figure 8d shows the eighteen-year mean change of vertically integrated (from surface to 700 hPa) moisture flux and moisture flux divergence in September in the BC experiment. In comparison with the sulphate experiment, only the eastern passage of moisture transport is weakened. Thus the weakening of the EASM circulation is less in the BC experiment, and the reduction in precipitation is less when compared to the sulphate experiment. As the low cloud cover decrease compensates the BC direct radiative effect, the temperature is changed less over the lee side of the Tibetan Plateau and the Indochina Peninsula than in sulphate experiment. The change in land-sea surface temperature contrast with the Indian Ocean is not as strong as the change further east with the Pacific Ocean, thus only the eastern moisture transport route is affected.

5 Conclusions and discussions

In this study, experiments using HiGAM with an interactive aerosol scheme show that the EASM in the model is altered when local emissions of both sulphate and BC

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



aerosols separately are increased from 1950 to 2000 levels. These impacts of aerosols are more significant during the withdrawal phase (September) of the EASM. For different aerosol species (sulphate and BC), the mechanisms that cause the change of precipitation in September differ.

5 For sulphates:

- The direct radiative effect cools the surface but not enough to alter the EASM.
- The indirect radiative effect on cloud droplet size is large in September due to the vertical coincidence of aerosol and low level cloud.
- The cloud albedo is increased and the surface temperature cools over a wide
10 region of East Asia.
- The decrease in land-sea thermal contrast weakens the circulation, transporting less moisture from both ocean basins.

For black carbon:

- The direct radiative effect is the major factor contributing to changes in EASM in
15 September.
- Low level cloud cover decreases as the absorption of solar radiation by BC heats the atmosphere.
- The direct radiative effect is slightly compensated by low level cloud decrease as more solar radiation reaches the surface.
- A small decrease in the land-sea thermal contrast in the BC experiment compared
20 to the sulphate experiment results in a weakening of moisture transport from the Pacific Ocean only.

How robust is this study? The atmospheric GCM means that the aerosol impact on SST and feedbacks of SST changes in aerosols have not been considered. However,

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Ackerman, A., Toon, O., Stevens, D., Heymsfield, A., Ramanathan, V., and Welton, E.: Reduction of tropical cloudiness by soot, *Science*, 288, 1042, 2000. 23013
- Adler, R., Huffman, G., Chang, A., Ferraro, R., Xie, P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P., and Nelkin, E.: The Version-2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979–present), *J. Hydrometeorol.*, 4, 1147–1167, 2003. 23015
- Albrecht, B.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227, 1989. 23009
- Annamalai, H., Hamilton, K., and Sperber, K. R.: The South Asian summer monsoon and its relationship with ENSO in the IPCC AR4 simulations, *J. Climate*, 20, 1071–1092, doi:10.1175/JCLI4035.1, 2007. 23015
- Ashrit, R., Douville, H., and Kumar, K.: Response of the Indian monsoon and ENSO-monsoon teleconnection to enhanced greenhouse effect in the CNRM coupled model, *J. Meteor. Soc. Japan*, 81, 779–803, 2003. 23010
- Boer, G. and Yu, B.: Climate sensitivity and response, *Clim. Dyn.*, 20, 415–429, 2003. 23011
- Boucher, O., Pham, M., and Sadourny, R.: General circulation model simulations of the Indian summer monsoon with increasing levels of sulphate aerosols, *ANGEO*, 16, 346–352, 1998. 23009
- Chung, C. and Ramanathan, V.: Weakening of North Indian SST gradients and the monsoon rainfall in India and the Sahel, *J. Climate*, 19, 2036–2045, 2006. 23010
- Chung, C. and Zhang, G.: Impact of absorbing aerosol on precipitation: Dynamic aspects in association with convective available potential energy and convective parameterization closure and dependence on aerosol heating profile, *J. Geophys. Res.*, 109, D22103, 2004. 23010
- Davies, T., Cullen, M., Malcolm, A., Mawson, M., Staniforth, A., White, A., and Wood, N.: A new dynamical core for the Met Office’s global and regional modelling of the atmosphere, *Quart. J. Roy. Meteor. Soc.*, 131, 1759–1782, 2005. 23012
- Edwards, J. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model, *Quart. J. Roy. Meteor. Soc.*, 122, 689–719, 1996. 23013
- Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D., Haywood, J., Lean, J., Lowe, D., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I*

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, chap. Changes in Atmospheric Constituents and in Radiative Forcing, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 23014

Fujino, J., Nair, R., Kainuma, M., Masui, T., and Matsuoka, Y.: Multi-gas mitigation analysis on stabilization scenarios using AIM global model, Multigas Mitigation Climate Policy, Special Issue, 3, p. 343, 2006. 23024

Gates, W., Boyle, J., Covey, C., Dease, C., Doutriaux, C., Drach, R., Fiorino, M., Gleckler, P., Hnilo, J., Marlais, S., Phillips, T., Potter, G., Santer, B., Sperber, K., Taylor, K., and Williams, D., An overview of the results of the Atmospheric Model Intercomparison Project (AMIP I), B. Amer. Meteor. Soc., 80, 29–56, 1999. 23015

Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, J. Geophys. Res.-Atmos., 102, 6831–6864, 1997. 23013

Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: a review, Rev. Geophys., 38, 513–543, 2000. 23009

Haywood, J. and Shine, K.: Multi-spectral calculations of the direct radiative forcing of tropospheric sulphate and soot aerosols using a column model, Quart. J. Roy. Meteor. Soc., 123, 1907–1930, 1997. 23014

Huang, Y., Chameides, W., and Dickinson, R.: Direct and indirect effects of anthropogenic aerosols on regional precipitation over East Asia, J. Geophys. Res., 112, D03212, 2007. 23009

Iwasaki, T. and Kitagawa, H.: A possible link of aerosol and cloud radiations to Asian summer monsoon and its implication in long-range numerical weather prediction, J. Meteor. Soc. Japan, 76, 965–982, 1998. 23009

Johns, T., Gregory, J., Ingram, W., Johnson, C., Jones, A., Lowe, J., Mitchell, J., Roberts, D., Sexton, D., Stevenson, D., Tett, S., and Woodage, M., Anthropogenic climate change for 1860 to 2100 simulated with the HadCM3 model under updated emissions scenarios, Clim. Dyn., 20, 583–612, 2003. 23010

Jones, A., Roberts, D., Woodage, M., and Johnson, C.: Indirect sulphate aerosol forcing in a climate model with an interactive sulphur cycle, J. Geophys. Res., 106, 20293–20310, 2001. 23013

Kim, M., Lau, W., Kim, K., and Lee, W.: A GCM study of effects of radiative forcing of sulfate aerosol on large scale circulation and rainfall in East Asia during boreal spring, Geophys. Res. Lett., 34, L24701, 2007. 23010, 23011

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Lau, K., Kim, M., and Kim, K.: Asian summer monsoon anomalies induced by aerosol direct forcing: the role of the Tibetan Plateau, *Clim. Dyn.*, 26, 855–864, doi:10.1007/s00382-006-0114-z, 2006. 23010, 23011
- Lee, E. and Sohn, B.: Recent increasing trend in dust frequency over Mongolia and Inner Mongolia regions and its association with climate and surface condition change, *Atmos. Environ.*, 45, 4611–4616, 2011. 23024
- Lesins, G., Chylek, P., and Lohmann, U.: A study of internal and external mixing scenarios and its effect on aerosol optical properties and direct radiative forcing, *J. Geophys. Res.*, 107, 4094, 2002. 23024
- Martin, G., Ringer, M., Pope, V., Jones, A., Dearden, C., and Hinton, T.: The physical properties of the atmosphere in the new Hadley Centre Global Environmental Model (HadGEM1). Part I: Model description and global climatology, *J. Climate*, 19, 1274–1301, 2006. 23012, 23014
- Meehl, G., Arblaster, J., and Collins, W.: Effects of black carbon aerosols on the Indian monsoon, *J. Climate*, 21, 2869–2882, 2008. 23010
- Menon, S., Hansen, J., Nazarenko, L., and Luo, Y.: Climate effects of black carbon aerosols in China and India, *Science*, 297, 2250, 2002. 23010
- Ming, Y., Ramaswamy, V., and Persad, G.: Two opposing effects of absorbing aerosols on global-mean precipitation, *Geophys. Res. Lett.*, 37, L13701, 2010. 23010
- Myhre, G., Jonson, J., Bartnicki, J., Stordal, F., and Shine, K.: Role of spatial and temporal variations in the computation of radiative forcing due to sulphate aerosols: a regional study, *Quart. J. Roy. Meteor. Soc.*, 128, 973–989, 2002. 23014
- Nozawa, T., Nagashima, T., Ogura, T., Yokohata, T., Okada, N., and Shiogama, H.: Climate change simulations with a coupled ocean-atmosphere GCM called the model for interdisciplinary research on climate: MIROC, Tech. Rep. Vol. 12, Center for Global Environmental Research, National Institute for Environmental Studies, Japan, 2007. 23009, 23013
- Ramanathan, V., Crutzen, P., Lelieveld, J., Mitra, A., Althausen, D., Anderson, J., Andreae, M., Cantrell, W., Cass, G., Chung, C., Clarke, A., Coakley, J. and Collins, W. and Conant, W., Dulac, F., Heintzenberg, J., Haymsfield, A., Holben, B., Howell, S., Hudson, J., Jayaraman, A., Kiehl, J., Krishnamuri, T., Lubin, D., McFarquhar, G., Novakov, T., Ogren, J., Podgorny, J., Prather, K., Priestley, K., Prospero, J., Quinn, P., Rajeev, K., Rasch, P., Rupert, S., Sadourny, R., Satheesh, S., Shaw, G., Sheridan, P., and Vaterno, F.: Indian Ocean Experiment: an integrated analysis of the climate forcing and effects of the great Indo-Asian haze, *J. Geophys. Res.*, 106, 28371–28398, 2001. 23010

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J., Washington, W., Fu, Q., Sikka, D., and Wild, M.: Atmospheric brown clouds: Impacts on South Asian climate and hydrological cycle, *Proc. Nat. Acad. Sci. USA*, 102, 5326, 2005. 23010

Randles, C. and Ramaswamy, V.: Absorbing aerosols over Asia: a Geophysical Fluid Dynamics Laboratory general circulation model sensitivity study of model response to aerosol optical depth and aerosol absorption, *J. Geophys. Res.*, 113, D21203, 2008. 23009, 23010

Roberts, D. and Jones, A.: Climate sensitivity to black carbon aerosol from fossil fuel combustion, *J. Geophys. Res.*, 109, D16202, 2004. 23013

Roeckner, E., Bengtsson, L., Feichter, J., Lelieveld, J., and Rodhe, H.: Transient climate change simulations with a coupled atmosphere-ocean GCM including the tropospheric sulfur cycle, *J. Climate*, 12, 3004–3032, 1999. 23010

Shaffrey, L., Stevens, I., Norton, W., Roberts, M., Vidale, P., Harle, J., Jrrar, A., Stevens, D., Woodage, M., Demory, M., Donners, J., Clark, D., Clayton, A., Cole, J., Wilson, S., Connolleg, W., Davies, T., Iwi, A., Johns, T., King, J., New, A., Slingo, J., Slingo, A., Steenman-Clark, L., and Martin, G., UK HiGEM: the new UK high-resolution global environment model-model description and basic evaluation, *J. Climate*, 22, 1861–1896, 2009. 23012, 23014, 23015

Smith, S., Andes, R., Conception, E., and Lurz, J.: Historical sulfur dioxide emissions, 1850–2000: methods and results, Tech. Rep. 14537, Pacific Northwest National Laboratory, Joint Global Change Research Institute 8400 Baltimore Avenue College Park, Maryland 20740, 2004. 23009, 23012, 23031

Trenberth, K., Stepaniak, D., and Caron, J.: Global monsoon as seen through the divergent atmospheric circulation, *J. Climate*, 13, 3969–3993, 2000. 23009

Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, 1977. 23009

van Vuuren, D. P., Den Elzen, M. G. J., Lucas, P. L., Eickhout, B., Strengers, B. J., van Ruijven, B., Wonink, S., and van Houdt, R.: Stabilizing greenhouse gas concentrations at low levels: an assessment of reduction strategies and costs, *Climatic Change*, 81, 119–159, 2007. 23024

Wang, B. and Ding, Q.: Changes in global monsoon precipitation over the past 56 years, *Geophys. Res. Lett.*, 33, L06711, 2006. 23009

Wang, C.: Impact of direct radiative forcing of black carbon aerosols on tropical convective precipitation, *Geophys. Res. Lett.*, 34, L05709, 2007. 23010

Woodage, M., Davison, P., and Roberts, D.: Aerosol processes, Tech. Rep. 20, Met Office, Exeter, UK 2003. 23013

Yu, R., Wang, B., and Zhou, T.: Climate effects of the deep continental stratus clouds generated by the Tibetan Plateau, J. Climate, 17, 2702–2713, 2004. 23017

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Summary of aerosol emissions used in experiments. East Asia is defined as a region of 5–60° N, 90–150° E. In SO₂-1950 experiment, BC emissions are same as Control experiment. In BC-1950 experiment, SO₂ emissions are same as Control experiment.

Experiment	Aerosol emissions over East Asia	Aerosol emissions over rest of the world
Control	SO ₂ in 2000 BC in 2000	SO ₂ in 2000 BC in 2000
SO ₂ -1950	SO ₂ in 1950	SO ₂ in 2000
BC-1950	BC in 1950	BC in 2000

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 1. Annual mean SO₂ emissions. Units: Tg(S) yr⁻¹. **(a)** Emissions in 2000. **(b)** Same as **(a)**, except that, within the inner box (5–60° N, 90–150° E), the emissions are in the level of 1950. A linear transition is applied between inner and outer boxes. Emission data is from Smith et al. (2004).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 2. (a) JJA precipitation in HiGAM (1983–2000, units: mm day^{-1}). (b) JJA precipitation from GPCP 2.5×2.5 monthly data (1979–2010, units: mm day^{-1}) (c) JJA 850 hPa geopotential height (contour, units: gpm) and wind (vector, units: m s^{-1}) in HiGMA (1983–2000) (d) Same as (c) but from ERA-40 (1958–2002).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 3. (a) 18-yr (1983–2000) monthly and area-mean aerosol column burden over East Asia. Units: mg m^{-2} . (b) 18-yr (1983–2000) monthly area-mean vertical mean cloud drop effective radius over East Asia. Units: μm . (c) 18-yr (1983–2000) monthly and area-mean low cloud fraction over East Asia. Units: 1. Shading colours along each line are standard deviations calculated using eighteen-yr monthly data.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 4. Changes in monthly area mean surface temperature (red, Units: K) and precipitation (green, Units: mm day⁻¹) over East Asia continent (20–45° N, 100–122° E). **(a)** Change in sulphate experiment (Control minus SO₂-1950). **(b)** Change in black carbon experiment (Control minus BC-1950). The percentage over each precipitation bar is percentage changes of precipitation against Control experiment. Significant changes excess 95% confident level are hatched.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 5. (a) 18-yr monthly area-mean sulphate radiative effect (Control minus $\text{SO}_2\text{-1950}$) at TOA over East Asia ($20\text{--}45^\circ\text{N}$, $100\text{--}122^\circ\text{E}$). (b) Same as (a), but for surface. (c) 18-yr monthly area-mean black carbon radiative effect (Control minus $\text{SO}_2\text{-1950}$) at TOA over East Asia. (d) Same as (c), but for surface. Units: W m^{-2} . Black line is direct-only effect, grey shading is standard deviation. Blue line is combined effect, blue shading is standard deviation.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 6. Change of low cloud fraction due to black carbon changes (Control minus BC_1950) in September. Units: 1. Significant changes excess 95% confidence level are hatched.

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 7. (a) 18-yr mean net radiative flux (downward is positive) in September due to sulphate combine effect (Control minus SO₂_1950). Units: W m⁻². The blue box indicates East Asia (20–45° N, 100–122° E), the area mean value is shown above the to-right corner of the blue box. (b) 18-yr mean change in surface temperature in September for sulphate experiment. Units: K. (c) 18-yr mean change in precipitation in September for sulphate experiment. Units: mm day⁻¹. (d) 18-yr mean change in vertical integrated (surface to 700 hPa) moisture flux (vector, Units: kg m⁻¹ s⁻¹) and moisture flux divergence (shaded, Units: Units: kg m⁻² s⁻¹) for sulphate experiment. Significant changes excess 95 % confident level are hatched.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

The effect of regional changes in anthropogenic aerosols on rainfall

L. Guo et al.




Fig. 8. (a) 18-yr mean net radiative flux (downward is positive) in September due to BC combine effect (Control minus BC_1950). Units: $W m^{-2}$. The blue box indicates East Asia ($20\text{--}45^\circ N$, $100\text{--}122^\circ E$), the area mean value is shown above the to-right conner of the blue box. (b) 18-yr mean change in surface temperature in September for BC experiment. Units: K. (c) 18-yr mean change in precipitation in September for BC experiment. Units: $mm day^{-1}$. (d) 18-yr mean change in vertical integrated (surface to 700 hPa) moisture flux (vector, Units: $kg m^{-1} s^{-1}$) and moisture flux divergence (shaded, Units: $kg m^{-2} s^{-1}$) for BC experiment. Significant changes excess 95 % confident level are hatched.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	