

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**Tropical tropopause
ice clouds**

P. Spichtinger and
M. Krämer

Tropical tropopause ice clouds: a dynamic approach to the mystery of low crystal numbers

P. Spichtinger¹ and M. Krämer²

¹Institut für Physik der Atmosphäre, Johannes Gutenberg-Universität, Mainz, Germany

²Institut für Energie- und Klimaforschung (IEK-7), Forschungszentrum Jülich, Jülich, Germany

Received: 24 July 2012 – Accepted: 15 October 2012 – Published: 25 October 2012

Correspondence to: P. Spichtinger (spichtin@uni-mainz.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The occurrence of high, persistent ice supersaturation inside and outside cold cirrus in the tropical tropopause layer (TTL) remains an enigma that is intensely debated as the “ice supersaturation puzzle”. However, it was recently confirmed that observed supersaturations are consistent with very low ice crystal concentrations, which is incompatible with the idea that homogeneous freezing is the major method of ice formation in the TTL. Thus, the tropical tropopause “ice supersaturation puzzle” has become an “ice nucleation puzzle”. To explain the low ice crystal concentrations, a number of mainly heterogeneous freezing methods have been proposed. Here, we reproduce in situ measurements of frequencies of occurrence of ice crystal concentrations by extensive model simulations, driven by the special dynamic conditions in the TTL, namely the superposition of slow large-scale updraughts with high-frequency short waves. From the simulations, it follows that the full range of observed ice crystal concentrations can be explained when the model results of the scenarios are mixed for both heterogeneous/homogeneous and pure homogeneous ice formation occurring in very slow ($< 1 \text{ cm s}^{-1}$) and faster ($> 1 \text{ cm s}^{-1}$) large-scale updraughts. This statistical analysis shows that about 80 % of TTL cirrus can be explained by “classical” homogeneous ice nucleation, while the remaining 20 % stem from heterogeneous and homogeneous freezing occurring within the same environment. The mechanism limiting ice crystal production via homogeneous freezing in an environment full of gravity waves is the shortness of the gravity waves, which stalls freezing events before a higher ice crystal concentration can be formed.

1 Introduction

Water vapour is the most important natural greenhouse gas. However, in the stratosphere, an increase in water vapour would possibly result in cooling. Most trace substances enter the stratosphere through the tropical tropopause layer (TTL), localized

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between the main level of convective outflow, 150 hPa, and about 70 hPa (Fueglistaler et al., 2009a). The TTL water vapour budget, and thus the exchange with the stratosphere, depends crucially on the occurrence and properties of ice clouds in this cold region ($T < 200$ K). It is believed that homogeneous freezing of liquid solution particles (Koop et al., 2000), which predominate the particle population, is the preferred pathway of ice formation in the low-temperature regime ($T < 235$ K). High water vapour supersaturation with respect to ice is required to initiate homogeneous ice nucleation. The number of emerging ice crystals depends on temperature and the ambient relative humidity over ice (RH_i). A strong increase in RH_i due to rising vertical velocity will produce large amounts of ice crystals (see Sect. 3.2).

In the TTL, very slow large-scale updraughts prevail (≤ 0.01 ms⁻¹, e.g. Virts et al., 2010), which lead to low ice crystal concentrations (≤ 0.1 cm⁻³). However, tropical deep convection is a source of intrinsic gravity waves in this region (e.g. Fritts and Alexander, 2003), which consequently initiate much higher vertical velocities and therefore higher ice crystal number concentrations (mostly > 0.3 cm⁻³, e.g. Jensen et al., 2010) are expected. Since the many ice crystals grow rapidly by water vapour diffusion (Korolev and Mazin, 2003), it is expected that the initially high ice supersaturation quickly reduces to thermodynamical equilibrium (ice saturation) after ice formation.

In contrast, over the last few years high and persistent ice supersaturations were observed inside and outside ice clouds in the cold TTL in several airborne field campaigns, creating a discussion called the “supersaturation puzzle” (Peter et al., 2006). A step forward in this discussion was made recently: Krämer et al. (2009) observed ice crystal concentrations much lower than expected (most often < 0.1 cm⁻³), but consistent with the high supersaturations measured also. These observations turned the “supersaturation” into a “nucleation puzzle”, which is supported by earlier measurements of low ice crystal numbers (McFarquhar et al., 2000; Thomas et al., 2002; Lawson et al., 2008). The “nucleation puzzle” is currently being intensely discussed and other nucleation pathways suppressing, modifying or replacing homogeneous freezing have been proposed (Kärcher and Koop, 2005; Zobrist et al., 2008; Murray, 2008; Gensch

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2008; Spichtinger and Gierens, 2009c; Krämer et al., 2009; Murray et al., 2010). Most of the former approaches explaining the TTL ice nucleation are of a chemical or microphysical nature. Here, we present intense model studies of ice cloud formation under dynamical conditions typical for the TTL. By directly comparing model simulations and observations, we claim that the special TTL dynamics – namely a superposition of very slow large-scale updraughts with high-frequency short waves – can produce about 80 % of the observed low numbers of ice crystals by the “classical” homogeneous freezing of solution droplets (Koop et al., 2000), while about 20 % is formed via homogeneous and heterogeneous freezing occurring within the same environment.

The study is structured as follows. In the next section, we present an introduction to ice formation processes in homogeneous freezing events and the impact of local dynamics. Section 3 presents idealized ice cloud simulations in order to show the impact of the superposition of motions on different scales. In Sect. 4, the special dynamic conditions in the TTL are briefly summarized (large-scale motion vs. gravity waves) before realistic simulations of ice formation in the TTL are presented and discussed in comparison to observations in Sect. 5. In Sect. 6, the results and their interpretation are discussed in combination with other approaches. We then end with some conclusions.

2 Basics of ice formation

2.1 The ice cloud model

The simulations in this study are carried out using a box model with a state-of-the-art bulk microphysics scheme developed for cold ice clouds ($T < 235$ K). The model is described in detail in Spichtinger and Gierens (2009a) and Spichtinger and Cziczo (2010), thus here only some key properties are briefly described. The double-moment scheme is consistent, based on general moments of the underlying mass distribution(s) of the ice crystals, which are assumed to be lognormal with a fixed geometric standard deviation of mass $\sigma_m = 2.85$, corresponding to a geometric standard deviation

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of length $\sigma_L \sim 1.4$ – 1.6 . The dominant processes in the low-temperature regime are parameterized, as nucleation of ice crystals, diffusional growth, and evaporation and sedimentation of ice crystals. The model can arbitrarily treat many classes of ice, each of them containing prognostic variables for ice as well as background aerosol mass and number concentrations. Thus, for the ice particles, there are complementary aerosol particles which control the ice formation process. In contrast to the usual treatment in bulk moment schemes used in other models (e.g. COSMO, see Seifert and Beheng, 2006), the different classes of ice are differentiated by their formation mechanism (e.g. heterogeneously vs. homogeneously formed ice). The aggregation of ice crystals is not included here, because this process is assumed to be of minor importance at low temperatures (see e.g. Kajikawa and Heymsfield, 1989). For homogeneous freezing of aqueous solution droplets (in short: homogeneous ice nucleation), sulphuric acid is assumed to be the background aerosol. For details of the freezing process, see Sect. 2.2. The H_2SO_4 background aerosol droplets are distributed according to a lognormal size distribution with a fixed geometric standard deviation; the modal radius of the distribution is initialized at the beginning but can change due to nucleation processes, since for frozen aerosol droplets, it is assumed that the aerosol is trapped in the ice crystals thus removed from the background aerosol. The parameterization of the diffusional growth process was extended to the temperature and pressure range as relevant for the TTL (see also Krämer et al., 2009), i.e. $T \geq 178$ K, $p \geq 50$ hPa.

2.2 Homogeneous freezing of solution droplets: RH_i variations and ice crystal numbers

First, we will briefly recapitulate the mechanism of homogeneous freezing of aqueous solution droplets. At low temperatures, pre-existing soluble aerosol particles (e.g. H_2SO_4) can take up some water via the Koehler theory, thus growing to form aqueous solution droplets, i.e. the weight fraction of the solute is quite small. These droplets can be supercooled by a few degrees until they freeze spontaneously. The nucleation rate for homogeneous freezing of solution droplets can be described using water activity

a_w (see e.g. Koop et al., 2000). Thus, the nucleation rate for a given droplet of a certain diameter depends only on temperature and on the environmental relative humidity. Starting with a “dry” size distribution of soluble aerosol particles $f(r)$, a size distribution of aqueous solution droplets $f_d(r)$ results from Koehler theory. Note that the distribution $f_d(r)$ depends on environmental temperature and humidity. The distributions are normalized by the total number of aerosol particles N_a . Using the volume nucleation rate $J = J(a_w, T) = J(\text{RH}_i, T)$, we can describe the number of nucleated ice crystals per time interval Δt using

$$\Delta N = N_a \int_0^{\infty} f_d(r) \left(1 - \exp\left(-J \frac{4}{3} \pi r^3 \Delta t\right) \right) dr \quad (1)$$

Assuming the existence of the third general moment of the distribution $f_d(r)$, we can describe the ice crystal production rate using the following expression as derived from Eq. (1):

$$\frac{dN}{dt} = N_a \frac{4}{3} \pi \mu_3[r] \cdot J \quad (2)$$

where $\mu_k[r] := \int_0^{\infty} f_d(r) r^k dr$. Note that here an only marginal change of temperature and

humidity causes J to vary over several orders of magnitude (see Koop et al., 2000). The nucleation process, involving the nucleation rate based on the environmental conditions, is one important factor for determining the number of ice crystals formed in a typical cooling event, but other processes might be important as well. For a detailed analysis, we investigated a typical nucleation event. We assumed an isolated air parcel (i.e. no exchange with the environment is allowed) rising with a prescribed vertical velocity. Here we also assumed a constant vertical velocity. To evaluate the time evolution of the relative humidity, we considered the total derivative of the relative humidity with

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



respect to ice, i.e.

$$\text{RH}_i = 100\% \frac{p \cdot q}{e \cdot p_{\text{ice}}(T)} \quad (3)$$

$$\frac{d\text{RH}_i}{dt} = \underbrace{\frac{\partial \text{RH}_i}{\partial T} \frac{dT}{dt} + \frac{\partial \text{RH}_i}{\partial p} \frac{dp}{dt}}_{\approx \text{adiabatic expansion}} + \underbrace{\frac{\partial \text{RH}_i}{\partial q} \frac{dq}{dt}}_{\text{growth}} \quad (4)$$

- 5 Then, we inspected the terms in Eq. (4) in order to examine a homogeneous nucleation event. First, we investigated the change in temperature and pressure, assuming just external changes driven by the vertical updraught. The temperature rate dT/dt can be described using the vertical updraught w , reading

$$\frac{dT}{dt} = \frac{dT}{dz} \frac{dz}{dt} = -\frac{g}{c_p} w = -\Gamma w \quad (5)$$

- 10 where g and c_p denotes the gravity acceleration and the specific heat capacity for constant pressure, respectively. Γ is the dry adiabatic lapse rate. Using the Clausius-Clapeyron equation and the isentropic coefficient $\kappa = R/c_p = 2/7$ for air and assuming pure adiabatic temperature changes, we can reformulate the first two terms on the right-hand side of Eq. (4) in a more suggestive way:

$$15 \frac{\partial \text{RH}_i}{\partial T} \frac{dT}{dt} + \frac{\partial \text{RH}_i}{\partial p} \frac{dp}{dt} = -\frac{\text{RH}_i}{T} \left(\frac{L}{R_v T} - \frac{1}{\kappa} \right) \frac{dT}{dt} \quad (6)$$

Please note that the first term in brackets is usually about one order of magnitude larger than the absolute value of the second term for the temperature range relevant for the existence of ice crystals ($180 \leq T \leq 273$ K). From Eqs. (5) and (6), we can clearly see that adiabatic expansion (i.e. vertical updraughts with positive w -values) lead to an
 20 increase in relative humidity.

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The last term on the right-hand side of Eq. (4) describes changes in specific humidity, which only occur if ice particles with a bulk mass mixing ratio q_c exist inside the air parcel (because exchange with the environment is not permitted), i.e. $dq/dt = -dq_c/dt$, constituting a sink for relative humidities in the range $RHi > 100\%$. Thus, in the case of ice particles inside the box and for relative humidities larger than saturation, there are two competing terms; a source and a sink. We then investigated their relative role for a homogeneous nucleation event.

For this purpose, we simulated a situation of a constant updraught, i.e. a standard cooling event, indicated by red lines in all three panels of Fig. 1. The box model is used including processes of (a) homogeneous freezing of aqueous solution droplets and (b) diffusional growth/evaporation only; sedimentation is ignored in order to reduce the complexity of the experiment. We specified one class of homogeneously frozen ice crystals. For the soluble background aerosol, which forms the solution droplets, that trigger homogeneous nucleation later on, we prescribed a dry H_2SO_4 aerosol with log-normal distribution, a geometric standard deviation of $\sigma_r = 1.5$, and initial modal radius $r_m = 25$ nm. The background aerosol number concentration was set to $N_a = 300$ cm^{-3} , which is more representative for extra-tropical tropospheric conditions. For the standard cooling event with initial conditions of $T_{init} = 220$ K and $p_{init} = 300$ hPa, and a prescribed persistent updraught of $w = 0.1$ ms^{-1} , the relative humidity increased (red line in upper panel of Fig. 1). At some point, the threshold RHi for homogeneous nucleation of droplets of a certain size (black line: threshold for ~ 250 nm droplets) was reached and thus ice crystals were formed via homogeneous freezing. However, these ice crystals were quite small. Their surface was small so that their ability to deplete water vapour was quite limited; therefore, the relative humidity continued to increase i.e. the source of supersaturation due to adiabatic cooling dominated over the sink of diffusional growth. The increasing relative humidity led to further ice crystal nucleation. After some time (around $t \sim 350$ s), the size of the ice crystals increased, allowing the crystals to deplete the water vapour efficiently, balancing the source via adiabatic cooling. After $t \sim 355$ s, the sink was stronger than the source and the relative humidity decreased (i.e. net

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sink), although there was still a source of supersaturation by adiabatic cooling. At later times, the relative humidity crossed the homogeneous nucleation threshold, approaching from higher values, thus ice crystal formation was shut down. Until this threshold was reached, there was still ice crystal nucleation. This scenario is used quite often in many idealized studies in order to investigate ice formation at constant updraughts under certain conditions (see, e.g. Kärcher and Lohmann, 2002; Hoyle et al., 2005; Barahona and Nenes, 2008; Spichtinger and Gierens, 2009a; Spichtinger and Cziczo, 2010), leading to a one-to-one relationship between vertical updraughts and ice crystal number concentrations produced in such cooling events. The main issue here is that the vertical updraught was kept constant during the whole cooling event, which lasted about $\tau_{\text{nuc}} \approx 140$ s. In the middle panel of Fig. 1, the ice crystal nucleation rate is shown, including the nucleation time τ_{nuc} , whereas in the right panel of the figure, the evolution of the ice crystal number concentration is shown. Since we show a scenario at quite high temperatures (initially $T = 220$ K), the ice crystal number concentration formed in the cooling event is quite low ($N \approx 105 \text{ l}^{-1}$). Again it should be noted that this type of cooling event (constant vertical updraught/cooling rate) has been used in many previous studies to investigate ice nucleation (see, e.g. Kärcher and Lohmann, 2002; Hoyle et al., 2005; Barahona and Nenes, 2008; Spichtinger and Cziczo, 2010).

Next, we will look the case of a change in vertical updraught during the nucleation event (i.e. within the time interval τ_{nuc}). To clearly demonstrate this issue, we investigated the following idealized scenario. We started with a constant updraught of $w = 0.1 \text{ ms}^{-1}$, but at $t = 320$ s we switched to the value of $w = -0.1 \text{ ms}^{-1}$, i.e. the updraught was turned into a downdraught of equal strength (Fig. 1, blue lines). Until the change in updraught, the simulations were identical. However, when the cooling process was turned into a warming process, the relative humidity decreased with approximately the same rate as it increased before. The homogeneous nucleation threshold was reached soon after this (at $t \sim 355$ s, incidentally approximately the time where the maximum RHi values were reached in the reference simulation), thus the nucleation event was stopped much earlier, as can be clearly seen in the ice crystal nucleation

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rate in the middle panel. This led to a significantly lower ice crystal number concentration, compared to the reference simulation. When vertical velocities were changed, the ice crystal number concentration was about $N \approx 20 \text{ l}^{-1}$.

The key finding of these investigations is that a changing vertical updraught could massively influence the number of ice crystals produced during a nucleation event. This finding was used to simulate the small ice crystal number concentrations found in the TTL. The updraught was then changed by superimposing a wave structure to a very slow vertical motion. As we will show later (Sect. 4), a changing updraught in the TTL is triggered by high-frequency gravity waves.

3 Idealized TTL ice cloud simulations

In this section, the role of different superimposed vertical velocity components in the formation of ice crystals will be investigated by means of idealized model simulations relevant to the situation in the TTL.

3.1 Model setup

To perform the idealized TTL ice cloud simulations, we used the box model described in Sect. 2.1 with the following model setup: as in the simulations described in the previous section, only one class of ice formed by homogeneous nucleation was assumed; the same background aerosol conditions were prescribed as in Sect. 2, i.e. we assume pure homogeneous freezing of aqueous solution droplets containing sulphuric acid. In the first step, we established a reliable relationship between vertical updraught and ice crystal number concentrations for the temperature range $185 \leq T \leq 200 \text{ K}$ and the vertical velocity range of $0.01 \leq w \leq 10 \text{ ms}^{-1}$. In our model simulations, the aerosol particles were removed from the background as soon as ice crystals were formed. Hence, the background aerosol was reduced during nucleation events and acted as

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a limiting factor for the ice crystal number concentrations formed. As the initial pressure of all simulations we used $p_{\text{init}} = 100 \text{ hPa}$.

In a second step, we investigated the influence of changing updraughts in a more relevant setup. For this purpose, we specified two different types of vertical velocity; a large-scale component w_{LS} and a wave-like signature, given by a pure sine wave with a time period of $\tau = 500 \text{ s}$, i.e. $w_{\text{wave}}(t) = A \cdot \sin(2\pi t/\tau)$, where the amplitude had a value of $A = 1.25 \text{ ms}^{-1}$. For the large scale component we distinguished between two different regimes, i.e. a low regime ($w_{\text{LS}} = 0.008 \text{ ms}^{-1}$) and a high regime ($w_{\text{LS}} = 0.03 \text{ ms}^{-1}$). In Sect. 5, we will show that the choice of these values was not arbitrary but motivated by the realistic dynamic conditions in the TTL (see Sect. 4).

3.2 Results

In Fig. 2, the relationship between constant and persistent vertical updraughts and ice crystal number concentrations are shown, representative of conditions in the TTL. For low vertical velocities, we can see a kind of power law (note the double-logarithmic plot). However, for larger values of w the increase slows down as a result of the limitation of the nucleation process by the pre-existing background aerosol (see also Spichtinger and Gierens, 2009a). Under the usual assumption that the TTL is full of gravity waves with vertical velocities $w \geq 0.2 \text{ ms}^{-1}$, it could be argued that only ice crystal number concentrations in the range $N \geq 1 \text{ cm}^{-3}$ should occur. However, as stated in the introduction, this is the essence of the nucleation puzzle, because much smaller values have been obtained in observations (see Fig. 3). To clarify this central point in our study, we then superimposed a constant large-scale vertical motion w_{LS} and a high-frequency wave $w_{\text{wave}}(t)$, i.e. $w(t) = w_{\text{LS}} + w_{\text{wave}}(t)$.

The results of ice crystal formation in the second scenario are shown in Fig. 4. We started with the faster motion regime ($w_{\text{LS}} = 0.03 \text{ ms}^{-1}$, top right panel), which constitutes the well-known case described by Lin et al. (1998): The relative humidity increased because of the slow vertical updraught and the high-frequency wave (see red

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



line in top right panel of Fig. 4). At about $t \sim 1400$ sec and about $t \sim 1950$ s, the relative humidity peaked as the wave was pushed over the thresholds for homogeneous ice nucleation by the large-scale updraught. The combination of the short wave and the large-scale updraught resulted in a strong change in RH_i, causing quite a vigorous increase in the ice nucleation rate. High ice crystal number concentrations were formed in two overshoots, which can be seen in the middle right panel of Fig. 4 (red line). Additionally, the two nucleation events can be seen clearly in the ice crystal production rate dN_i/dt (Fig. 4, bottom right panel) The high ice crystal number concentrations led to an effective diffusional growth, the gas phase water was depleted quite fast causing the ice supersaturation to relax to values around saturation. The final ice crystal number concentration was much higher than that in the scenario with a constant updraught only $w_{LS} = 0.03 \text{ ms}^{-1}$ (blue line in the middle right panel of Fig. 4). The maximum ice crystal number concentrations for these two cases were $N_i \sim 0.33 \text{ cm}^{-3}$ (constant updraught, blue line) and $N_i \sim 11.1 \text{ cm}^{-3}$ (induced by the short wave, red line). Therefore, there was an enhancement of about two orders of magnitude.

In the case of a much slower updraught ($w_{LS} = 0.008 \text{ ms}^{-1}$) with a superimposed short wave (left panel of Fig. 4, red lines), the picture was different than expected. The very slow lifting of the air parcel pushed the RH_i peaks of the wave very slowly towards the homogeneous nucleation thresholds (red line in left panel of Fig. 4). The first nucleation events were very weak (see production rate dN_i/dt (Fig. 4), bottom left panel), because the overshoots were very small and the change in RH_i was smaller than for a faster large-scale updraught. However, in these first nucleation events a few ice crystals were formed, which began depleting the available water vapour. In subsequent nucleation events, more ice crystal were formed, but the rates remained quite low because of the gentle overshoots. The available ice crystals also reduced the ice supersaturation quite slowly, resulting in a slow decrease in RH_i and thus in a gentle shutdown of the nucleation. This process finally resulted in a small ice crystal number concentration, which is comparable to concentrations produced by the weak constant updraught alone (blue line). Consequently, the RH_i decreased only slowly in the further lifetime of

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the ice cloud, causing persistent high ice supersaturation. The values for maximum ice crystal number concentrations for these two cases were by $N_i \sim 0.025 \text{ cm}^{-3}$ (constant updraught alone, blue line) and $N_i \sim 0.048 \text{ cm}^{-3}$ (constant updraught plus wave, red line). Thus, there was a slight enhancement by a factor of about two.

5 In summary, both the slow and fast motion scenarios showed enhanced ice crystal number concentrations when a short wave was superimposed on the constant updraught. However, slowing down the constant updraught significantly reduced the enhancement of ice crystal number concentrations induced by the short wave. This is the central finding, which causes the main effect in our investigations. We will discuss the
10 relevance of such an idealized case for the situation in the TTL in the next Sect. (4).

Last but not least, it should be noted that for low-frequency waves, there was no significant difference in simulations with superimposed waves for different large-scales updraughts. In all cases, the nucleation event was not stopped by dynamics but by
15 diffusional growth leading to much higher ice crystal number concentrations. The difference in the produced ice crystal number concentrations was due to the randomly occurring phase of the wave, as already mentioned in previous studies (see e.g. Lin et al., 1998).

4 TTL dynamics

20 In this section, we will look at the different types of vertical motion in the TTL, as driven by very different processes. In the first part, we will explore the large-scale components mainly driven by Kelvin waves, whereas in the second part we will evaluate the conditions in the TTL favouring a special kind of gravity wave.

4.1 Large-scale dynamics

25 First, we will the question of whether low vertical updraughts are common in the TTL, especially in the vertical velocity range $|\omega| \leq 0.01 \text{ ms}^{-1}$. There are two different sources

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for large-scale updraughts in the TTL. The first involves ascents to compensate diabatic heating (e.g. Fueglistaler et al., 2009b; Salby and Callaghan, 2004); this compensating effect led to very small updraughts with vertical velocity values $w \sim 0.001 \text{ ms}^{-1}$. The second source involves large-scale equatorial waves, e.g. Kelvin waves (Webster, 1972; Gill, 1980; Wheeler et al., 2000; Ryu and Lee, 2010; Thompson et al., 2011), which are usually triggered by convection and organized convective systems. Equatorial Kelvin waves with time periods in the order of a few days lead to temperature anomalies of up to 8 K with typical values around 2–3 K (Immler et al., 2008), which translate into updraughts in the order of $0.002 \leq w_{\text{LS}} \leq 0.01 \text{ ms}^{-1}$. There is now evidence from several studies (Boehm and Verlinde, 2000; Immler et al., 2008; Fujiwara et al., 2009; Virts et al., 2010) that there is a correlation between cold anomalies in the TTL and cirrus clouds. From recent studies on cirrus clouds in the TTL using the LIDAR in space CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation), the vertical motions in the regions of cirrus cloud occurrence were estimated (Virts et al., 2010). Here, values of the large-scale motion at $p = 100 \text{ hPa}$ were estimated to be in the range $-0.0175 \text{ Pa s}^{-1} \leq \omega \leq 0$ (see Fig. 8.8 in Virts, 2009), which translates into vertical velocities in the range $0 \leq w \leq 0.01 \text{ ms}^{-1}$. Thus, from observational and theoretical considerations, the occurrence of low large-scale motions in the range $0 \leq w \leq 0.01 \text{ ms}^{-1}$ can be confirmed and the occurrence of cirrus clouds in such updraught regions is quite common. It should be mentioned here that in case of large-scale updraughts in the range $0 \leq w \leq 0.01 \text{ ms}^{-1}$ without any superposition of motions on smaller scales, only small ice crystal number concentrations ($N_i \leq 0.1 \text{ cm}^{-3}$) would occur, as can be derived from Fig. 2.

4.2 Gravity waves

In contrast to the (tropical) troposphere, the TTL is much more stable and its dynamics are less vigorous. Convection only very rarely reaches the TTL, exciting gravity waves at its lower boundary by triggering internal gravity waves in the stable stratified layers above the convective cells (Fritts and Alexander, 2003). Additionally, gravity

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



waves are excited at lateral boundaries of the convective updraught regions in order to balance subsidence (Bretherton and Smolarkiewicz, 1989). The stratification of the TTL is very high ($0.01 \leq N \leq 0.025 \text{ s}^{-1}$). Figure 5 shows a climatological vertical profile of the potential temperature (left) and the Brunt-Vaisala frequency (right), as derived from SHADOZ measurements (Thompson et al., 2003). The TTL should be full of gravity waves (see, e.g. Fritts and Alexander, 2003; Jensen and Pfister, 2004), although measurements are complicated and usually cover only larger wave lengths, i.e. low-frequency waves. The frequency of gravity waves depends on the thermal stratification as well as on their horizontal and vertical propagation. Recent high-resolution measurements at latitudes $\sim 26^\circ \text{ N}$ (Wang et al., 2006) support the occurrence of such short high-frequency, mostly horizontally propagating waves. For almost horizontally propagating waves, the horizontal wave lengths can be estimated using $\lambda_x = 2\pi U/N$, where U is horizontal wind. Since the mean horizontal wind is quite low (i.e. $U \leq 12 \text{ ms}^{-1}$) in the equatorial region (Fueglistaler et al., 2009a), this leads to very small horizontal wave lengths in the order of $3 \text{ km} \leq \lambda_x \leq 7.6 \text{ km}$ or roughly $\lambda_x < 10 \text{ km}$, which are difficult to see both in situ and from satellite instruments. Thus, we can assume that some of these waves have very high internal frequencies. Typical vertical amplitudes of these waves have been estimated to be in the range $\Delta z \sim 100 \text{ m}$ or even much smaller (Gary, 2006; Gierens et al., 2007), which translates roughly into temperature changes of $\Delta T \leq 1 \text{ K}$. These waves represent the prototype for TTL gravity waves in our investigations. In addition to the boundary conditions from thermal stratification and winds in the TTL, we added more information about a possible restriction of the frequency spectrum in gravity waves occurring. From recent theoretical investigations (Ruprecht et al., 2010; Ruprecht and Klein, 2011) on the propagation of gravity waves in a convectively dominated environment, we know that the full spectrum of frequencies, i.e. $0 \leq \omega \leq N$ does not occur. The convective towers act as high-pass filters, depending on the fractional coverage of the convective cells compared to the cloud-free environment, allowing only part of the frequency to occur. The frequency spectrum is then given by $f \cdot N \leq \omega \leq N$, where f depends on the fractional coverage σ , i.e. $f \approx \sqrt{\sigma}$. For typical

values of $\sigma \sim 0.1$, we get a frequency cut-off factor of $f \approx 0.32$. Since there is a lot of convective activity in the tropical regions, we can assume higher fractional coverage of convective cells. Thus, $\sigma = 0.1$ is a lower limit. In later applications of this “frequency window”, we will use a lower limit of $f = 0.4$, i.e. f will be in the range $0.4 \leq f \leq 1$, corresponding to cloud cover values of $\sim 0.16 \leq \sigma \leq 1$.

5 Realistic TTL ice cloud simulations

After observations and theoretical investigations have revisited that the conditions as specified in Sect. 3 may occur in the TTL, we will now define more realistic scenarios for a large number of box model simulations. The results of these simulations will be presented in terms of ice crystal number concentrations. Additionally, we will discuss the plausibility of heterogeneous ice nucleation for the situation in the TTL.

5.1 Model setup

For the realistic model simulations, we ran the ice cloud box model along idealized air mass trajectories. The setup for the microphysics as well as for the idealized dynamics, i.e. trajectories, is described in the next two sections.

5.1.1 Model setup for microphysics

For a realistic description of the background aerosol conditions in the TTL, we used the output of a 2-D aerosol model (Weisenstein et al., 2007) to calculate size distributions of sulphate aerosols at different pressure levels typical for the TTL ($p \sim 70/100/120/150$ hPa). The size distributions were fitted with three modes using log-normal distributions. The fits are shown in Fig. 6, representing the initial size distributions of the sulphuric acid background aerosol. The three size modes of aerosols led to three classes of ice, formed by homogeneous nucleation. The total background aerosol number concentration was of the order $N_a \sim 50\text{--}80 \text{ cm}^{-3}$, which agrees well

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with measurements. However, these values are smaller than the typical aerosol concentrations in the extra-tropical tropopause region ($N_a \sim 300 \text{ cm}^{-3}$). For the environmental conditions in the TTL, we derived the initial temperature/pressure from the climatological values (see Fig. 5) and the spread in the measurements (see, e.g. Thompson et al., 2003; Fueglistaler et al., 2009a; Selkirk et al., 2010). The values are given in Table 1. Note that the initial temperatures were set about 10 K lower than the mean climatological values shown in Fig. 5. We shifted the temperature profile in order to be consistent with the in situ measurements (Krämer et al., 2009) as otherwise the temperatures in our simulations would have been too high.

We wanted to investigate ice nucleation at specific environmental conditions typical of the TTL (temperature/pressure) as given by values in table 1. Thus, the model initialization was specified to ensure that ice nucleation starts nearby the given values at around half the simulation time (i.e. after ~ 5 h) in the case of a pure large-scale updraught. For example, when $T_{\text{nuc}} = 196 \text{ K}$ and $p_{\text{nuc}} = 150 \text{ hPa}$ with a constant updraught of $w_{\text{LS}} = 0.01 \text{ ms}^{-1}$, we set $T_{\text{init}} = 197.8 \text{ K}$ and $p_{\text{init}} = 154.78 \text{ hPa}$. Using this procedure, we ensured that when high-frequency waves with high amplitudes were superimposed to the large-scale updraught (see Sect. 5.1.2) nucleation would occur close to the typical temperature/pressure values.

The box model also included a module for the sedimentation of ice crystals (Spichtinger and Cziczo, 2010). To activate this module, additional information on the vertical extension of the box Δz as well as on the sedimentation factor f_{sed} was required. Here, we specified $\Delta z = 50 \text{ m}$, $f_{\text{sed}} = 0.5$ in the upper part of the cloud (i.e. the main nucleation zone) and $f_{\text{sed}} = 0.9$ in the middle part of the cloud (see discussion in Spichtinger and Cziczo, 2010).

5.1.2 Model setup for dynamics

We saw in Sect. 4 that the TTL dynamics were dominated by motions on two different scales, namely (a) very low large-scale updraughts in the order of $w_{\text{LS}} \leq 1 \text{ cm s}^{-1}$ and (b) high-frequency gravity waves triggered by convective activity. The large-scale

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



updraughts were triggered by large-scale features such as Kelvin waves, occupying large horizontal regions in the tropical atmosphere. Thus, gravity waves are most likely lifted by the large-scale motions. In terms of an ascending air parcel along a trajectory, we thus assumed a slowly lifted parcel, which also experiences a superimposed (high-frequency) oscillation driven by gravity waves. As described above, the source for the gravity waves is most likely convection. Either waves are triggered by convective plumes bouncing against the TTL from below or waves are radiated laterally from the convective tower. In both cases, we can assume intrinsic gravity waves with frequencies near the stratification. Thus, in our simulations we concentrated on monochromatic waves with a distinct frequency as determined below. For our box model calculations we specified – similar to the idealized simulations in Sect. 3 – a vertical velocity for lifting the (closed) air parcel adiabatically as a superposition of the two motions mentioned above: $w(t) = w_{\text{LS}} + w_{\text{wave}}(t)$. For the large-scale component w_{LS} we then specified the following ranges, split into a very slow and a faster velocity range

$$w_{\text{LS}} = \underbrace{0.003/0.005/0.008/0.01}_{\text{very slow}} / \underbrace{0.02/0.03/0.05}_{\text{faster}} \text{ ms}^{-1} \quad (7)$$

To determine the monochromatic wave component $w_{\text{wave}}(t)$, we introduced temperature changes instead of changes in the vertical velocity due to observational constraints on the amplitude of temperature variations in the TTL. We assumed an adiabatic process, i.e. that the vertical velocity was connected to temperature changes via Eq. (5). Thus, the temperature driven by this motion is given by $T(t) = T(t=0) + A_T \cos(\omega t)$ via integration of Eq. (5), with A_T : amplitude of the temperature variation. From observations, we know that this amplitude can reach values up to $A_T = 1 \text{ K}$, thus we specified the range $A_T = 0.1/0.25/0.5/0.75/1.0 \text{ K}$, respectively. As stated for the large-scale component, the value of A_T was fixed for one simulation. From Ruprecht et al. (2010) (see Sect. 4), we know that the frequency spectrum is bounded by the Brunt-Vaisala frequency N and a lower value $f \cdot N$, where f depends on the area covered by convective cells, i.e. on the fraction σ . By assuming a typical lower limit of $\sigma \leq 0.1$, we set the

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



range for the real frequency of the waves, $\omega = f \cdot N$, $f = 0.4/0.5/0.6/0.7/0.8/0.9/1$; as stated for the other parameters, f and N were fixed for a whole single simulation. To specify the typical Brunt-Vaisala frequency, we used the climatological mean derived from Fig. 5. In order to investigate a realistic range of frequency spectra, we set the maximum value N_{\max} (as given in table 1) so that the mean climatological value N_{mean} was approximately given by $N_{\text{mean}} \approx 0.9 \cdot N_{\max}$. Consequently, the lower limit of the frequency window is given by $N_{\min} = 0.4 \cdot N_{\max}$. The range of frequencies for the different pressure levels is shown in Fig. 7. In summary, we ran single box model simulations with a superposition of a constant large-scale component and one monochromatic wave so that the temperature evolution of a single run was

$$T(t) = T_{\text{init}} \underbrace{-\Gamma \cdot w_{\text{LS}} \cdot t}_{\text{large scale}} + \underbrace{A_T \cos(f \cdot N_{\max} \cdot t)}_{\text{wave}} \quad (8)$$

In our simulations, we used all possible combinations as stated above. The total simulation time for each single case was $t_{\text{sim}} = 36\,000$ s (10 h) with a time step of $\Delta t = 0.1$ s in order to resolve nucleation events as best as possible (see discussion in Spichtinger and Gierens, 2009a).

For a statistical investigation, we collected values from all simulations, filtered by a certain criterion (e.g. slow vs. fast large-scale component or pure homogeneous nucleation vs. homogeneous and heterogeneous nucleation).

5.2 TTL ice crystal numbers from homogeneous nucleation

Figure 8 shows the frequencies of occurrence of ice crystal number concentrations derived from all simulations with homogeneous ice nucleation (HOM) described in the previous section together with the in situ measurements from Krämer et al. (2009). The results are split into simulations with superimposed very slow large-scale motion ($w_{\text{LS}} \leq 1 \text{ cm s}^{-1}$, green) and simulations with superimposed faster large-scale motion ($w_{\text{LS}} > 1 \text{ cm s}^{-1}$, blue). The observations were surprisingly well represented by the simulations

with very slow w_{LS} . Ice crystal number concentrations from the simulations were in the range $0.001 \leq N_i \leq 0.3 \text{ cm}^{-3}$ (note that the detection limit of the measurements was about 0.005 cm^{-3} ; lower concentrations from the simulations were clipped for better comparison). However, the peak in the measurements around $N_i \sim 0.005\text{--}0.007 \text{ cm}^{-3}$ was missing in the simulations and the higher observation values were not reached. Nevertheless, the shape of the frequency distribution in the range $0.01 \leq N_i \leq 0.3 \text{ cm}^{-3}$ was almost perfectly matched.

In contrast to the simulations for very slow large-scale velocities, the measurements were not reproduced by faster large-scale vertical motions. In these simulations, the probability density function (pdf) of ice crystal number concentration had a similar shape as the very slow w_{LS} pdf. However, the distribution shifted to very high values of N_i . The broad peak of the pdf was somewhere in the range $0.5 \leq N_i \leq 5 \text{ cm}^{-3}$ with quite frequent values up to $N_i \sim 20 \text{ cm}^{-3}$. The pdf compares to that shown by Jensen et al. (2010, their Fig. 6), though in the study of Jensen et al. (2010) the peak and maximum ice concentrations were smaller, probably because a large-scale motion of 0.3 cm s^{-1} (superimposed with a broad spectrum of gravity waves) was used, which ranges here up to 0.5 cm s^{-1} . From the comparison of pdfs from very slow and faster large-scale updraughts, we clearly see that the transition from low to high number concentrations in the case of high-frequency waves is determined by the large-scale component of the vertical velocity. There is a clear separation between these different regimes, and the critical values seem to be around $w \sim 1 \text{ cm s}^{-1}$.

Additionally, it should be noted that in both sets of simulations very high vertical velocities up to values of $w \sim 3 \text{ m s}^{-1}$ occurred. The statistics for vertical velocities in the different regimes (very slow vs. faster w_{LS}) are shown in Fig. 9. It can be seen from the figure that the vertical velocity distributions are almost identical except of for “shift” of about $\Delta w \sim 0.03 \text{ m s}^{-1}$, given by the large-scale component. However, the ice crystal number concentrations were decoupled from the vertical velocities, since the nucleation events in high frequency waves were stopped by downdraughts (see Sects. 2 and 3) and not by diffusional growth. Thus, it makes no sense anymore to evaluate

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the vertical velocity to estimate ice crystal number concentrations as the simple 1–1 relationship shown, e.g. in Fig. 2.

5.3 TTL ice crystal numbers from heterogeneous and homogeneous nucleation

Heterogeneous nucleation was brought into the discussion to explain the low ice crystal number concentrations in the TTL (Khvorostyanov et al., 2006; Gensch et al., 2008; Jensen et al., 2008; Krämer et al., 2009; Jensen et al., 2010; Murray et al., 2010). Thus, we extended our simulations by including heterogeneous nucleation in the following way: we added another class of ice representing heterogeneous nucleation on solid particles, although we did not specify the nature of the ice nuclei (IN) but merely the effect of freezing ice crystals on them. One could think of these IN as a mixture of glassy particles (Murray et al., 2010), crystallized ammonium sulphate particles (Jensen et al., 2010) and mineral dust (Froyd et al., 2009), which are all possible candidates causing heterogeneous freezing in the TTL. In Fig. 6, the additional heterogeneous IN mode is indicated by the grey curve. Here, we again use a lognormally distributed aerosol with modal diameter $L_m = 0.5 \mu\text{m}$, a geometrical standard deviation of $\sigma_L = 1.4$ and a total number concentration of $N_{\text{IN}} = 7 \text{ l}^{-1}$ (see e.g. DeMott et al., 2003). The nucleation threshold is size-dependent, i.e. $\text{RH}_{\text{het}} = (a/L)^b + \text{RH}_0$ (Spichtinger and Cziczo, 2010), with $a = 1 \mu\text{m}$, $b = 0.65$, $\text{RH}_0 = 143.5\%$; thus, the thresholds are quite high (in the order of $\text{RH}_i \sim 145\%$), mimicking quite bad IN. We chose this option because good IN should be washed out by freezing and sedimentation at lower altitudes, while only bad IN will survive the transport to higher altitudes in the TTL. We then ran the same set of simulations as described in Sect. 5.1 with additional heterogeneous freezing. Note that when the homogeneous freezing threshold was reached after heterogeneous ice nucleation, a second (homogeneous) freezing event occurred in the simulations. We did not specify the type of heterogeneous INs. They could stem (classically) from advected mineral dust or even from glassy particles, as proposed by Murray et al. (2010).

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In Fig. 10, the results for superimposed very slow large-scale motions are shown for pure homogeneous nucleation (hom, same green line as in Fig. 8) and for competing heterogeneous-homogeneous nucleation (het/hom, blue curve) as described above. For a better comparison, Fig. 10 also shows the sum of all simulations (all: hom + het/hom, black curve) and the in situ measurements (red curve). The most remarkable feature of competing heterogeneous nucleation can be seen in the blue pdf. Heterogeneous freezing formed ice crystals before homogeneous nucleation was triggered. Thus, few ice crystals (in order of $N_i \sim 7l^{-1}$, see peak in Fig. 10) grew in a highly supersaturated environment depleting water vapour in a considerable way. Additional homogeneous nucleation events were rare and produced the ice crystal numbers between $N_i \sim 0.01\text{--}0.1\text{ cm}^{-3}$, or nucleation was quenched (see discussion in Spichtinger and Gierens, 2009a,b; Spichtinger and Cziczo, 2010). This led to a general reduction of the produced ice crystal number concentrations, which can be seen clearly in the pdf (Fig. 10, blue curve). For simulations with waves superimposed on faster large-scale components there was almost no effect compared to pure homogeneous nucleation (not shown). It is also remarkable that the peak in the measurements at $N_i \sim 0.005\text{--}0.008\text{ cm}^{-3}$ (red curve) was reproduced by heterogeneous ice nucleation. To sum up all simulations (black curve), there is an agreement with the observations, although part of the ice events in the concentration range $N_i \sim 0.01\text{--}0.3\text{ cm}^{-3}$ are missing and the events at $N_i > 0.3\text{ cm}^{-3}$ are not represented by the model simulations.

Seemingly, pure homogeneous freezing events and heterogeneous/homogeneous ice formation in very slow large-scale motions superimposed with short, high-frequency gravity waves both contribute significantly to the TTL ice clouds, but cannot explain the full observed ice crystal concentration spectrum.

Instead, a mixture of different scenarios might lead to better agreement with the measurements. For this purpose, we used normalized pdfs of the different scenarios, i.e. (1) pure homogeneous nucleation for slow vertical updraughts $f_1 := f_{\text{hom,slow}}$, (2) pure homogeneous nucleation for faster vertical updraughts $f_2 := f_{\text{hom,fast}}$, (3) homogeneous/heterogeneous nucleation for slow vertical updraughts $f_3 := f_{\text{het/hom,slow}}$, and

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(4) pure homogeneous and homogeneous/heterogeneous nucleation for fast vertical updraughts $f_4 := f_{\text{hom+het/hom,fast}}$, respectively. The selection of these pdfs is motivated by the analysis later on, i.e. these are the pdfs we will use for a good agreement with the measurements. For brevity, we numbered the different scenarios as described in table 2. We combined these pdfs to form one pdf using weighting coefficients via

$$f_{\text{comb}} := \sum_{i=1}^4 a_i f_i / \sum_{i=1}^4 a_i \quad (9)$$

such that the mixture of the scenarios best matched the observations.

Some combinations of the full set of model simulations are shown in Fig. 11. Here, the blue curve represents a mixture of all simulations in the very slow and faster large-scale motion regimes with pure homogeneous freezing, using coefficients $a_1 = 0.99$, $a_2 = 0.01$, $a_3 = a_4 = 0$. In this case, it can be seen that the observations were reproduced well for ice concentrations $N_i > 0.01 \text{ cm}^{-3}$. Only the peak at $N_i \sim 0.005\text{--}0.008 \text{ cm}^{-3}$ was still missing. However, adding the simulations with heterogeneous/homogeneous ice formation to the mixture (black curve) ensured agreement between model simulations and observations.

The combination of pdfs of the simulated ice concentration frequency spectra best matching the observations showed that the lowest ice crystal numbers ($< 0.01 \text{ cm}^{-3}$) stemmed most likely from heterogeneous as well as homogeneous freezing in the very slow large-scale updraughts (about $\sim 20\%$, i.e. $a_3 = 0.2$), that the middle ice crystal numbers ($0.01\text{--}0.3 \text{ cm}^{-3}$) were solely from homogeneous freezing in the same updraught regime (about $\sim 79\%$, i.e. $a_1 = 0.79$), and that the highest TTL ice crystal concentrations ($> 0.3 \text{ cm}^{-3}$) also evolved from homogeneous freezing or even heterogeneous and homogeneous freezing occurring within the same environment, but in the faster large-scale updraughts (about $\sim 1\%$, i.e. $a_4 = 0.01$). Consequently, we also set $a_2 = 0$. Hence, about 80% of the TTL cirrus can be explained by “classical” homogeneous ice nucleation (Koop et al., 2000) and the remaining 20% by a combination of

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



heterogeneous and homogeneous freezing within the same environment. The mechanism limiting the ice crystal production from homogeneous freezing in an environment full of gravity waves is the shortness of the waves, causing the freezing events to stall before a higher number of ice crystals can be formed. As stated above, the superposition of two different dynamic motion regimes is crucial for this kind of mechanism explaining low ice crystal number concentrations.

6 Discussion

The model study presented here shows that the measured low TTL ice crystal number concentrations can be represented using a realistic setup for the special dynamics in that region and “classical” homogeneous ice nucleation (e.g. Koop et al., 2000). Additionally, we were able to represent the shape of the ice crystal number frequency distribution in a very reasonable way.

It was found that the ice crystal number densities are bound to the velocity regimes in the TTL. Superimposing the TTL high-frequency short waves with slow large-scale vertical velocities ($w_{LS} \leq 0.01 \text{ m s}^{-1}$) yielded the observed low ice crystal number concentrations, since the ice nucleation events were stalled early on by downdraughts. On the other hand, lower frequencies of the short gravity waves or faster large-scale motions ($w_{LS} > 0.01 \text{ m s}^{-1}$) triggered higher ice crystal number concentrations. These cases reflect what we would expect from theory (Lin et al., 1998; Hoyle et al., 2005) and what is usually discussed in studies about the impact of homogeneous nucleation on the TTL ice crystal numbers, concluding that this formation mechanism is unable to produce low ice concentrations (Jensen and Pfister, 2004). The major new part of our study is the special combination of high frequency waves *and* very slow large-scale updraughts.

Furthermore, we found that the shape of the measured ice crystal frequency distribution can be represented by mixing different possible scenarios: in many cases there were no heterogeneous IN available, and thus pure homogeneous nucleation was the

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dominant pathway (~ 79 %). In some more rare scenarios, heterogeneous nucleation took place in addition to homogeneous nucleation (~ 20 %), and finally, sometimes the dynamic conditions also differed, i.e. the more unlikely case of faster large-scale up-draughts appeared (~ 1 %). Generally, we can state that heterogeneous nucleation is not the main pathway for the formation of very low ice crystal number concentrations, but that the major mechanism is homogeneous ice nucleation driven by the special TTL dynamics; heterogeneous nucleation simply modifies the nucleation events but does not qualitatively change the results.

In many recent studies, the contamination of the sulphuric acid droplets by organics and thus the inhibition of homogeneous nucleation is considered to be the main reason for very low ice crystal number concentrations in the TTL. However, homogeneous nucleation would only be suppressed, if almost all background aerosol droplets were contaminated by these substances. From recent measurements (see e.g. Froyd et al., 2009), we know that there is a significant contribution of organics in the TTL. However, the relative mass fraction of organics in the TTL is never higher than 80 % (case Pre-AVE¹, where exceptionally high values were measured), and in many measurements, the relative fraction of organics in the TTL is in the range of 0.1–0.5 (e.g. CR-AVE² cases). Thus, it is very unlikely – at least from real measurements – that organics can suppress homogeneous nucleation completely, since the non-contaminated particles will nucleate to ice crystals.

In former studies on the impact of dynamics, a spectrum of gravity waves was usually used to drive the box models (see e.g. Jensen and Pfister, 2004). Although these spectra were determined from measurements, it is unclear if they are really representative of the TTL conditions and high-frequency waves were not included in these studies. The main reason for neglecting these waves was the lack of measurement techniques capable of detecting these very short wavelengths. Thus, the effects discussed in this study were not included in former studies. As we pointed out in our discussion, the

¹Pre-Aura Validation Experiment, 2004.

²Costa Rica Aura Validation Experiment, 2006.

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



short wavelengths are extremely important for very short freezing events. For longer wavelengths, as used in former studies, the impact of dynamics increases the ice crystal number concentrations tremendously.

In our study, we concentrated on monochromatic waves, which is a reasonable restriction. However, if we were to assume a spectrum of superimposed waves within the narrow spectral window as indicated by the theoretical considerations (Ruprecht et al., 2010), this might lead to a spread in the number concentrations towards larger values. In some sensitivity studies (not shown), the quality of the results driven by monochromatic waves did not change.

In contrast to former studies (see e.g. Jensen and Pfister, 2004; Jensen et al., 2010; Barahona and Nenes, 2011) the amplitude of the waves was not restricted to very small amplitudes to obtain low ice crystal number concentrations. For our approach this restriction was unnecessary, because the nucleation event was determined by the short wavelengths and not by the amplitude of the waves. Thus, our simulations were more realistic in this sense, because the whole spread of amplitudes in the range up to $\Delta T \sim 1$ K can be observed from measurements (Gary, 2006).

We already mentioned the contamination of the background aerosol solution droplets by organic substances. Although the contamination and thus suppression of homogeneous nucleation works well in laboratory experiments (Zobrist et al., 2008; Murray et al., 2010), at the moment there is no indication from in situ measurements in the TTL (a) that the respective substances (e.g. citric acid) are available in the TTL and (b) that there is contamination of almost all aerosol particles in order to guarantee total suppression of homogeneous nucleation (see e.g. discussion in Kärcher and Koop, 2005). There are just indications that the process of total suppression of homogeneous nucleation might work, the corroboration by measurements is still lacking. On the other hand, contamination of some aerosol particles could lead to additional heterogeneous IN. Since we did not specify the type of heterogeneous INs used in this study, there could be room for such glassy particles.

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, in our opinion, the total suppression of homogeneous nucleation by contamination with the suitable material is a very complex and rigid theory. Our proposal allows classical homogeneous and heterogeneous ice nucleation to happen in a very simple dynamic setup, given by the special environment of the TTL. Applying Ockham's razor (principle of parsimony) we tend to prefer the simpler dynamic approach, particularly because our simulations show very good agreement with in situ measurements.

7 Conclusions

In summary, we showed that the superposition of high-frequency internal gravity waves with very slow large-scale motions (e.g. driven by Kelvin waves) could trigger homogeneous freezing events in such a way that the low ice crystal number concentrations measured in the TTL are reasonably represented: $\sim 79\%$ of the ice crystals were formed by "classical" homogeneous freezing in very-slow large-scale motions, $\sim 20\%$ stemmed from combined heterogeneous/homogeneous ice nucleation in very slow large-scale motions, and only $\sim 1\%$ came from homogeneous freezing with or without simultaneously occurring heterogeneous nucleation in faster large-scale motions.

There is therefore an interplay of the different complementary mechanisms (e.g. TTL dynamics, homogeneous and heterogeneous ice nucleation, as well as the suppression of nucleation by organic substances or crystallization of ammonium sulphate) may lead to the phenomenon of low ice crystal number concentrations. In order to verify the relative contributions of these different effects, more measurements in the TTL are necessary. To investigate these effects in a more general treatment, additional two- or even three-dimensional model simulations including a realistic treatment of gravity waves are important.

Acknowledgements. We would like to thank Stephan Borrmann for providing the ice crystal data and Klaus Gierens and Ben Murray for fruitful discussions. We also greatly acknowledge Mrs. Hazel Bulet from the Language Service of the Research Center Jülich for improving the

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



English language of the manuscript. The output from the aerosol model was kindly provided by Debra Weisenstein and Ana Cirisan. This study contributes to the COST action Atmospheric Water Vapour in the Climate System (WaVaCS).



This publication is supported by COST – www.cost.eu

References

- Barahona, D. and Nenes, A.: Parameterization of cirrus cloud formation in large-scale models: homogeneous nucleation, *J. Geophys. Res.*, 113, D11211, doi:10.1029/2007JD009355, 2008. 28117
- Barahona, D. and Nenes, A.: Dynamical states of low temperature cirrus, *Atmos. Chem. Phys.*, 11, 3757–3771, doi:10.5194/acp-11-3757-2011, 2011. 28134
- Boehm, M. T. and Verlinde, J.: Stratospheric influence on upper tropospheric tropical cirrus, *Geophys. Res. Lett.*, 27, 3209–3212, 2000. 28122
- Bretherton, C. S. and Smolarkiewicz, P. K.: Gravity waves, compensating subsidence and detrainment around cumulus clouds, *J. Atmos. Sci.*, 46, 740–759, 1989. 28123
- DeMott, P., Cziczo, D., Prenni, A., Murphy, D., Kreidenweis, S., Thomson, D., Borys, R., and Rogers, D.: Measurements of the concentration and composition of nuclei for cirrus formation, *Proc. Nat. Acad. Sciences*, 100, 14655–14660, 2003. 28129
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009a. 28111, 28123, 28125
- Fueglistaler, S., Legras, B., Beljaars, A., Morcrette, J.-J., Simmons, A., Tompkins, A. M., and Uppala, S.: The diabatic heat budget of the upper troposphere and lower/mid stratosphere in ECMWF reanalyses, *Q. J. R. Meteorol. Soc.* 135, 21–37, 2009b. 28122
- Froyd, K. D., Murphy, D. M., Sanford, T. J., Thomson, D. S., Wilson, J. C., Pfister, L., and Lait, L.: Aerosol composition of the tropical upper troposphere, *Atmos. Chem. Phys.*, 9, 4363–4385, doi:10.5194/acp-9-4363-2009, 2009. 28129, 28133
- Fritts, D. C. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, *Rev. Geophys.*, 41, 1003, doi:10.1029/2001RG000106, 2003. 28111, 28122, 28123

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fujiwara, M., Iwasaki, S., Shimizu, A., Inai, Y., Shiotani, M., Hasebe, F., Matsui, I., Sugimoto, N., Okamoto, H., Nishi, N., Hamada, A., Sakazaki, T., and Yoneyama, K.: Cirrus observations in the tropical tropopause layer over the Western Pacific, *J. Geophys. Res.*, 114, D09304, doi:10.1029/2008JD011040, 2009. 28122
- 5 Gary, B. L.: Mesoscale temperature fluctuations in the stratosphere, *Atmos. Chem. Phys.*, 6, 4577–4589, doi:10.5194/acp-6-4577-2006, 2006. 28123, 28134
- Gensch, I., Bunz, H., Baumgardner, D., Christensen, L., Fahey, D., Hermann, R., Lawson, P., Popp, P., Smith, J., Webster, C., Weinstock, E., Wilson, J., Peter, T., and Krämer, M.: Supersaturations, microphysics and nitric acid partitioning in a cold cirrus observed during CR-AVE 2006: an observation-modeling intercomparison study, *Environ. Res. Lett.*, 3, 035003, doi:10.1088/1748-9326/3/3/035003, 2008. 28111, 28129
- 10 Gierens, K., Kohlhepp, R., Dotzek, N., Smit, H. G.: Instantaneous fluctuations of temperature and moisture in the upper troposphere and tropopause region. Part 1: probability densities and their variability, *Meteorol. Z.*, 16, 221–231, 2007. 28123
- 15 Gill, A. E.: Some simple solutions for heat-induced tropical circulation, *Q. J. R. Meteorol. Soc.*, 106, 447–462, 1980. 28122
- Hoyle, C., Luo, B., and Peter, T.: The origin of high ice crystal number densities in cirrus clouds, *J. Atmos. Sci.*, 62, 2568–2579, 2005. 28117, 28132
- 20 Immler, F., Krüger, K., Fujiwara, M., Verver, G., Rex, M., and Schrems, O.: Correlation between equatorial Kelvin waves and the occurrence of extremely thin ice clouds at the tropical tropopause, *Atmos. Chem. Phys.*, 8, 4019–4026, doi:10.5194/acp-8-4019-2008, 2008. 28122
- Jensen, E. and Pfister, L.: Transport and freeze-drying in the tropical tropopause layer, *J. Geophys. Res.*, 109, D02207, doi:10.1029/2003JD004022, 2004. 28123, 28132, 28133, 28134
- 25 Jensen, E. J., Pfister, L., Bui, T. V., Lawson, P., Baker, B., Mo, Q., Baumgardner, D., Weinstock, E. M., Smith, J. B., Moyer, E. J., Hanisco, T. F., Sayres, D. S., St. Clair, J. M., Alexander, M. J., Toon, O. B., and Smith, J. A.: Formation of large ($\sim 100\ \mu\text{m}$) ice crystals near the tropical tropopause, *Atmos. Chem. Phys.*, 8, 1621–1633, doi:10.5194/acp-8-1621-2008, 2008. 28129
- 30 Jensen, E. J., Pfister, L., Bui, T.-P., Lawson, P., and Baumgardner, D.: Ice nucleation and cloud microphysical properties in tropical tropopause layer cirrus, *Atmos. Chem. Phys.*, 10, 1369–1384, doi:10.5194/acp-10-1369-2010, 2010. 28111, 28128, 28129, 28134

- Kajikawa, M. and Heymsfield, A.: Aggregation of ice crystals, *J. Atmos. Sci.*, 46, 3108–3121, 1989. 28113
- Kärcher, B. and Koop, T.: The role of organic aerosols in homogeneous ice formation, *Atmos. Chem. Phys.*, 5, 703–714, doi:10.5194/acp-5-703-2005, 2005. 28111, 28134
- 5 Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *J. Geophys. Res.*, 107(D2), 4010, doi:10.1029/2001JD000470, 2002. 28117
- Khvorostyanov, V., Morrison, H., Curry, J., Baumgardner, D., and Lawson, P.: High supersaturation and modes of ice nucleation in thin tropopause cirrus: simulation of the 13 July 2002 cirrus regional study of tropical anvils and cirrus layers case, *J. Geophys. Res.*, 111, D02201, doi:10.1029/2004JD005235, 2006. 28129
- 10 Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature* 406, 611–614, 2000. 28111, 28112, 28114, 28131, 28132
- 15 Korolev A. and Mazin, I.: Supersaturation of water vapor in clouds, *J. Atmos. Sci.*, 60, 2957–2974, 2003. 28111
- Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505–3522, doi:10.5194/acp-9-3505-2009, 2009. 28111, 28112, 28113, 28125, 28127, 28129, 28144, 28145, 28150, 28152, 28153
- 20 Lawson, R. P., Pilson, B., Baker, B., Mo, Q., Jensen, E., Pfister, L., and Bui, P.: Aircraft measurements of microphysical properties of subvisible cirrus in the tropical tropopause layer, *Atmos. Chem. Phys.*, 8, 1609–1620, doi:10.5194/acp-8-1609-2008, 2008. 28111
- 25 Lin, H., Noone, K. J., Ström, J., and Heymsfield, A. J.: Dynamical influences on cirrus cloud formation process, *J. Atmos. Sci.*, 55, 1940–1949, 1998. 28119, 28121, 28132
- McFarquhar, G. M., Heymsfield, A. J., Spinhirne, J., and Hart, B., : Thin and subvisual tropopause tropical cirrus: observations and radiative impacts, *J. Atmos. Sci.*, 57, 1841–1853, 2000. 28111
- 30 Murray, B. J.: Inhibition of ice crystallisation in highly viscous aqueous organic acid droplets, *Atmos. Chem. Phys.*, 8, 5423–5433, doi:10.5194/acp-8-5423-2008, 2008. 28111
- Murray, B., Wilson, T. W., Dobbie, S., Cui, Z., Al-Jumur, S. M. R. K., Möhler, O., Schnaiter, M., Wagner, R., Benz, S., Niemand, M., Saathoff, H., Ebert, V., Wagner, S., and Kärcher, B.:

**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Heterogeneous nucleation of ice particles on glassy aerosols under cirrus conditions, *Nature Geosci.*, 3, 233–237, doi:10.1038/NGEO817, 2010. 28112, 28129, 28134
- Peter, T., Marcolli, C., Spichtinger, P., Corti, T., Baker M. B., and Koop, T.: When dry air is too humid, *Science*, 314, 1399–1402, 2006. 28111
- 5 Ruprecht, D. and Klein, R.: A model for nonlinear interactions of internal gravity waves with saturated regions, *Meteorol. Z.*, 20, 243–252, 2011. 28123
- Ruprecht, D., Klein, R., and Majda, A. J.: Modulation of internal gravity waves in a multiscale model for deep convection on mesoscales, *J. Atmos. Sci.*, 67, 2504–2519, 2010. 28123, 28126, 28134
- 10 Ryu, J.-H. and Lee, S.: Effect of tropical waves on the tropical tropopause transition layer upwelling, *J. Atmos. Sci.*, 67, 3130–3148, 2010. 28122
- Salby, M. and Callaghan, P.: Control of the tropical tropopause and vertical transport across it, *J. Climate*, 17, 965–985, 2004. 28122
- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds, Pt. 1: model description, *Meteorol. Atmos. Phys.*, 92, 45–66, 2006. 28113
- 15 Selkirk, H. B., Vömel, H., Valverde Canossa, J. M., Pfister, L., Diaz, J. A., Fernández, W., Amador, J., Stolz, W., and Peng, G. S.: Detailed structure of the tropical upper troposphere and lower stratosphere as revealed by balloon sonde observations of water vapor, ozone, temperature, and winds during the NASA TCSP and TC4 campaigns, *J. Geophys. Res.*, 115, D00J19, doi:10.1029/2009JD013209, 2010. 28125
- 20 Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmos. Chem. Phys.*, 9, 685–706, doi:10.5194/acp-9-685-2009, 2009a. 28112, 28117, 28119, 28127, 28130
- Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1b: Structuring cirrus clouds by dynamics, *Atmos. Chem. Phys.*, 9, 707–719, doi:10.5194/acp-9-707-2009, 2009b. 28130
- 25 Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 2: Competition of different nucleation mechanisms, *Atmos. Chem. Phys.*, 9, 2319–2334, doi:10.5194/acp-9-2319-2009, 2009c. 28112
- Spichtinger, P. and Cziczo, D. J.: Impact of heterogeneous ice nuclei on homogeneous freezing events in cirrus clouds, *J. Geophys. Res.*, 15, D14208, doi:10.1029/2009JD012168, 2010. 28112, 28117, 28125, 28129, 28130
- 30 Thomas, A., Borrmann, S., Kiemle, C., Cairo, F., Volk, M., Beuermann, J., Lepuchov, B., Santacesaria, V., Matthey, R., Rudakov, V., Yushkov, V., MacKenzie, A. R., and Stefanutti, L.: In

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

situ measurements of background aerosol and subvisible cirrus in the tropical tropopause region, *J. Geophys. Res.*, 107, 4763, doi:10.1029/2001JD001385, 2002. 28111

Thompson, A. M., Witte, J. C., McPeters, R. D., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W. J. H. Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S. Ogawa, T., Johnson, B. J., Vömel H., and Labow, G.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. comparison with Total Ozone Mapping Spectrometer (TOMS) and ground-based measurements, *J. Geophys. Res.*, 108, 8238, doi:10.1029/2001JD000967, 2003. 28123, 28125, 28147, 28149

Thompson, A. M., Allen, A. L., Lee, S., Miller, S. K., and Witte, J. C.: Gravity and Rossby wave signatures in the tropical troposphere and lower stratosphere based on Southern Hemisphere Additional Ozonesondes (SHADOZ), 1998–2007, *J. Geophys. Res.*, 116, D05302, doi:10.1029/2009JD013429, 2011. 28122

Virts, K.: Cirrus in the tropical tropopause transition layer: formation mechanisms and influence of the local and planetary-scale environment, MSc thesis, University of Washington, 108 pp., available at: <http://www.atmos.washington.edu/~kvirts/docs/Thesis.pdf> (last access: 23 October 2012), 2009. 28122

Virts, K., Wallace, J., Fu, Q., and Ackermann, T.: Tropical tropopause transition layer cirrus as represented by CALIPSO lidar observations, *J. Atmos. Sci.*, 67, 3113–3129, 2010. 28111, 28122

Wang, L., Alexander, M. J., Bui, T. B., and Mahoney, M. J.: Small-scale gravity waves in ER-2 MMS/MTP wind and temperature measurements during CRYSTAL-FACE, *Atmos. Chem. Phys.*, 6, 1091–1104, doi:10.5194/acp-6-1091-2006, 2006. 28123

Webster, P. J.: Response of the tropical atmosphere to local, steady forcing, *Mon. Wea. Rev.*, 100, 518–541, 1972. 28122

Weisenstein, D. K., Penner, J. E., Herzog, M., and Liu, X.: Global 2-D intercomparison of sectional and modal aerosol modules, *Atmos. Chem. Phys.*, 7, 2339–2355, doi:10.5194/acp-7-2339-2007, 2007. 28124, 28148

Wheeler, M., Kiladis, G. N., and Webster, P. J.: Large-scale dynamical fields associated with convectively coupled equatorial waves, *J. Atmos. Sci.*, 57, 613–640, 2000. 28122

Zobrist, B., Marcolli, C., Pedernera, D. A., and Koop, T.: Do atmospheric aerosols form glasses?, *Atmos. Chem. Phys.*, 8, 5221–5244, doi:10.5194/acp-8-5221-2008, 2008. 28111, 28134

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Table 1. Settings for realistic simulations to be fulfilled at ice nucleation.

p (hPa)	70	100	120	150
T (K)	190	184	188	196
N_{\max} (s^{-1})	0.029	0.022	0.0172	0.0128
$N_{\text{mean}} = N_{\max} \cdot 0.9$ (s^{-1})	0.0261	0.0198	0.0155	0.0115
$N_{\min} = N \cdot 0.4$ (s^{-1})	0.0116	0.0088	0.0069	0.0051

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Numbering for the different pdfs.

index	1	2	3	4
scenario	hom,slow	hom,fast	het/hom,slow	hom+het/hom,fast




Fig. 1. Example of the impact of changing updraughts on the formation of ice crystals via homogeneous freezing of solution droplets. The three panels show the time evolution of **(a)** relative humidity wrt ice, **(b)** ice crystal nucleation rate, and **(c)** ice crystal number concentration. Red curves indicate a constant persistent updraught of $w = 0.1 \text{ ms}^{-1}$, whereas blue curves indicate a scenario with a constant updraught of $w = 0.1 \text{ ms}^{-1}$ until $t = 320 \text{ s}$, followed by a constant downdraught of $w = -0.1 \text{ ms}^{-1}$. The black line in panel **(a)** indicates the homogeneous nucleation threshold for solution droplets of size $r = 0.25 \mu\text{m}$; this threshold corresponds to a production rate of $dN_i/dt = 0.04 \text{ l}^{-1} \text{ s}^{-1}$.

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer




Fig. 2. Ice crystal number concentration as formed in idealized simulations with a prescribed constant updraught under conditions typical for the TTL ($T = 185/190/195/200$ K, $p = 100$ hPa). Note that due to a finite background aerosol concentration, the ice crystal number concentration is limited by the aerosol concentration. This leads to the asymptotic behaviour of curves for $w > 0.5$ ms^{-1} . In addition, the measured ice crystal number concentrations are indicated (Krämer et al., 2009); the orange bar indicates minimal and maximal concentrations and the mean value, whereas the black bar indicates the 25 % and 75 % percentile as well as the median value.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Fig. 3. Measured ice crystal number concentration distribution in the temperature range $T < 200$ K, from Krämer et al. (2009).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Exemplary cases of updraught scenarios. Left column: large-scale updraught $w_{LS} = 0.008 \text{ m s}^{-1}$ (slow regime), right column: large-scale updraught $w_{LS} = 0.03 \text{ m s}^{-1}$ (fast regime). Blue lines indicate constant updraught, red lines denote superposition of large-scale updraught and a short wave. Upper panel: time evolution of the relative humidity with respect to (wrt) ice (in %); saturation is indicated by a black line, the grey line indicates relative humidity values where the ice crystal concentration rate dN/dt exceeds the value $1 \text{ m}^{-3} \text{ s}^{-1}$. Middle panel: time evolution of ice crystal number concentration. Lower panel: time evolution of ice crystal production rate dN/dt ; the value $1 \text{ m}^{-3} \text{ s}^{-1}$ is indicated by a black line. Here, the stepwise character of the nucleation for the superimposed short wave (red line) can be seen clearly in contrast to the continuous nucleation during a constant large-scale updraught (blue line).


**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Fig. 5. Mean climatological vertical profiles in the tropical atmosphere as derived from measurements (Thompson et al., 2003); left: potential temperature, right: Brunt-Vaisala frequency.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Fig. 6. Initial size distributions of sulphuric acid droplets in the TTL for the pressure levels 70/100/120/150 hPa (coloured lines), as derived from a 2-D aerosol model (Weisenstein et al., 2007); additionally, the size distribution of the heterogeneous ice nuclei as used for heterogeneous nucleation is indicated (grey line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer




Fig. 7. Vertical profile of static stability (mean Brunt-Vaisala frequency) as derived from measurements (Thompson et al., 2003). Additionally, the range for the initialization of the simulations is indicated by black bars, where we set $N_{\text{mean}} \approx 0.9 \cdot N_{\text{max}}$; thus, the bars show the interval $0.4 \cdot N_{\text{max}} \leq N \leq N_{\text{max}}$ as described in detail in Sect. 5.1; the values are reported in Table 1.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



Tropical tropopause
ice cloudsP. Spichtinger and
M. Krämer

Fig. 8. Frequencies of occurrence of ice crystal number concentrations from simulations with pure homogeneous freezing in very slow and faster large-scale motions ($w_{LS} \leq 1 \text{ cm s}^{-1}$, green; $w_{LS} > 1 \text{ cm s}^{-1}$, blue); the red line represents the measurements of Krämer et al. (2009). Note that the detection limit of the measurements is $N_i \sim 0.005 \text{ cm}^{-3}$; thus, the lowest concentration bins of the model simulations are clipped for better comparison with the measurements.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Fig. 9. Vertical velocity from simulations with pure homogeneous freezing in very slow ($w_{LS} \leq 1 \text{ cm s}^{-1}$) and faster ($w_{LS} > 1 \text{ cm s}^{-1}$) large-scale motions. The absolute values and the distributions are nearly equal, except for the shift of about $\Delta w \sim 0.03 \text{ m s}^{-1}$, given by the setup.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropopause ice clouds

P. Spichtinger and
M. Krämer




Fig. 10. Frequencies of occurrence of ice crystal number concentrations from simulations in very slow large-scale motions ($w_{LS} \leq 1 \text{ cm s}^{-1}$) with homogeneous freezing (hom, green), heterogeneous/homogeneous freezing (het/hom, blue) and a mixture of both (all); the red line represents the measurements of Krämer et al. (2009). Note that the lowest concentration bins of the model simulations are clipped for better comparison with the measurements.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Tropical tropopause
ice clouds**P. Spichtinger and
M. Krämer

Fig. 11. Frequencies of occurrence of ice crystal number concentrations: mixture of simulations in very slow ($w_{\text{LS}} \leq 1 \text{ cm s}^{-1}$) and faster ($w_{\text{LS}} > 1 \text{ cm s}^{-1}$) large-scale motions with homogeneous freezing (hom, blue) and with heterogeneous/homogeneous freezing (het/hom, black); the red line represents the measurements of Krämer et al. (2009); the lowest concentration bins of the model simulations are clipped off for a better comparison with the measurements.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

