

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Exploring the atmospheric chemistry of O₂SO₃⁻ and assessing the maximum turnover number of ion catalysed H₂SO₄ formation

N. Bork^{1,2}, T. Kurtén^{1,3}, and H. Vehkamäki¹

¹Division of Atmospheric Sciences and Geophysics, Department of Physics, P.O. Box 64, 00014 University of Helsinki, Finland

²Department of Chemistry, H. C. Ørsted Institute, University of Copenhagen, Denmark

³Laboratory of Physical Chemistry, Department of Chemistry, University of Helsinki, P.O. Box 55, 00014, Finland

Received: 24 August 2012 – Accepted: 2 November 2012 – Published: 22 November 2012

Correspondence to: N. Bork (nicolai.bork@helsinki.fi)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

It has recently been demonstrated that the O_2SO_3^- ion forms in the atmosphere as a natural consequence of ionizing radiation. Here, we present a density functional theory-based study of the reactions of O_2SO_3^- with O_3 . The most important reactions are (a) oxidation of O_2SO_3^- to O_3SO_3^- and (b) cluster decomposition into SO_3 , O_2 and O_3^- . The former reaction is highly exothermic and the nascent O_3SO_3^- will rapidly decompose into SO_4^- and O_2 . If the origin of O_2SO_3^- is SO_2 oxidation by O_3^- , the latter reaction closes a catalytic cycle wherein SO_2 is oxidized to SO_3 . The relative rates between the two major sinks for O_2SO_3^- is assessed, thereby providing a measure of the maximum turnover number of ion catalysed SO_2 oxidation, i.e. how many SO_2 can be oxidized per free electron. The rate ratio between reactions (a) and (b) is significantly altered by the presence or absence of a single water molecule, but reaction (b) is in general much more probable. Although we are unable to assess the overall importance of this cycle in the real atmosphere due to the unknown influence of CO_2 and NO_x , we roughly estimate that ion induced catalysis may contribute with several percent of H_2SO_4 levels in typical CO_2 free and low NO_x reaction chambers, e.g. the CLOUD chamber at CERN.

1 Introduction

One of the most significant uncertainties in weather and climate forecasts are related to the processes leading to cloud formation (Simpson and Wiggert, 2009; Rosenfeld, 2006; Spracklen et al., 2008; Carslaw et al., 2002). It is well known that each cloud droplet forms around a cloud condensation nucleus (CCN). CCN's may originate from a solid particle e.g. soot, salt, or pollen, but recently it has become increasingly clear that a significant part originate from clustering of purely gaseous molecules (Kulmala et al., 2004; Wiedensohler et al., 2009; Yu and Luo, 2009; Kazil et al., 2010).

Although molecular clustering and aerosol particle formation has been intensively studied, fundamental uncertainties prevail. One of the most significant uncertainties

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concerns the chemical composition of the very smallest molecular clusters. Sulfuric acid has repeatedly been found in atmospheric clusters and the concentration of sulfuric acid is known to correlate well with nanoparticle formation rates (Sipilä et al., 2010; Nieminen et al., 2009). Traditionally, binary nucleation of H₂SO₄ and H₂O has been the most studied mechanism, but during the last couple of decades it has become evident that a third species e.g. ammonia or an amine is needed as well (Kirkby et al., 2011; Ortega et al., 2012). However, the critical importance of sulfuric acid remains undisputed.

The main source of atmospheric sulfuric acid is UV induced oxidation of SO₂ which is emitted from volcanoes and through fossil fuel combustion. The oxidation mechanism involves several elementary reactions, but is relatively well understood (Bondybey and Beyer, 2002). The rate limiting step is the initial S(IV) to S(V) oxidation (Li and McKee, 1997)



where *M* is a stabilizing molecule. HSO₃ is then further oxidized to SO₃ by O₂ before hydration to H₂SO₄. The final hydration is known to be catalyzed by one or more excess water molecules and is the main sink of atmospheric SO₃ (Morokuma and Muguruma, 1994). In addition to this, a mechanism where SO₂ is oxidized by Criegee biradicals has been proposed (Welz et al., 2012; Mauldin III et al., 2012).

During the last decades, the influence of ions in aerosol formation has been targeted by several chamber studies (Kirkby et al., 2011; Enghoff and Svensmark, 2008). Here, it has been firmly established that ions enhance formation of nanometer-sized aerosol particles in SO₂- and O₃- containing atmospheres under UV exposure. One possible explanation, based on classical nucleation theory, is that the energy barrier of nucleation is lowered by the charge hereby increasing the total nucleation rate (Tohmfor and Volmer, 1938; Lovejoy et al., 2004). The lower barrier might result from reduced evaporation rates from ionic clusters due to charge delocalization or from increased collision rates between ionic particles and dipolar molecules or between oppositely charged particles (Su and Bowers, 1973; Kupiainen et al., 2011).

O₂SO₃⁻ chemistry and catalytic SO₂ oxidation

N. Bork et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A fundamentally different approach to explain ion induced nucleation is through ion induced chemistry altering the composition of the metastable gas phase. This idea is supported by several experimental studies where mainly collision limited ionic gas-phase chemical reactions were found, see e.g. Fehsenfeld and Ferguson (1974) and references therein. Further, a successful chamber study lead Svensmark et al. (2007) to suggest an ion induced SO₂ oxidation mechanism. Despite of this, little research has been dedicated to enhanced nucleation from ion induced chemistry, see however Sorokin and Arnold (2007, 2009) and Enghoff et al. (2012).

Atmospheric ions are mainly products of radioactivity and galactic cosmic rays. Although both cations and free electrons are produced, most attention have gathered around the influence of the electrons and subsequent anions (Nadykto et al., 2006; Kurtén et al., 2009). It is well known that some of the first stable products are O₂⁻(H₂O)_m and O₃⁻(H₂O)_n clusters (Luts and Parts, 2002). The first 5 water molecules are strongly bound to the ion, but several more may attach under cold and/or humid conditions (Bork et al., 2011b).

Further, it has been shown that O₃⁻(H₂O)_n may oxidize SO₂ to SO₃⁻ via the reaction



This reaction is probably the dominant sink of O₃⁻ in most reaction chambers since CO₂ most often is omitted. Initially, the product cluster might partially decompose due to the large release of internal energy in the oxidation. However, the high concentrations of both O₂ and H₂O ensures that the equilibrium quickly settles. Hereafter, the predominant products are O₂SO₃⁻ and O₂SO₃⁻(H₂O) (Bork et al., 2011a; Möhler et al., 1992). This is supported by a recent field study by Ehn et al. (2010), finding that after HSO₄⁻, O₂SO₃⁻ is the second most common sulfur based anion in a boreal forest.

The further chemical fate of O₂SO₃⁻ and the extra electron is largely unknown, but is potentially important. We have investigated the most probable reactions of de- and monohydrated O₂SO₃⁻ under standard conditions focusing mainly on the reaction with O₃. The two main reactions are,



where *W* is shorthand for water. The latter reaction is of particular interest since a catalytic cycle of SO_2 oxidation thereby is closed, see Fig. 1.

Using ab initio calculations we have determined the structures, energies and thermodynamics of reactants, transition states and products. We proceed to evaluate the rates of the relevant reactions and hereby evaluate the total turnover number, i.e. how many SO_2 oxidations a single electron on average will induce. Finally, we estimate the maximum fraction of H_2SO_4 originating from ionic catalysis under typical conditions in CO_2 free reaction chambers.

2 Computational details

As always when treating weakly bound systems, anionic systems and radical reactions, special care must be taken to ensure reliable results. Of special interest of this study is the relative adiabatic electron affinity of SO_3 and O_3 since this energy difference will be the thermodynamic driving force behind Reaction (R3b). The experimental electron affinity of O_3 has been determined to $202.9 \text{ kJ mol}^{-1}$ with high certainty (Novick et al., 1979; Arnold et al., 1994). On the contrary, despite several studies, the electron affinity of SO_3 remains highly uncertain. The most recent studies find values of $183 \pm 10 \text{ kJ mol}^{-1}$ (Gleason, 1987) and $190 \pm 10 \text{ kJ mol}^{-1}$ (Dobrin et al., 2000). However, the accumulated experimental evidence suggest that most likely the electron affinity of O_3 exceeds that of SO_3 by ca. 15 kJ mol^{-1} .

While scanning a variety of methods, we found that most ab initio methods actually predict a larger electron affinity of SO_3 than of O_3 , despite large basis sets. However, the CAM-B3LYP functional is an exception to this (Yanai et al., 2004). At basis sets of at least triple zeta quality, both the order and magnitude of the electron affinity difference is in accordance with the experimental value. Therefore, we have used

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the CAM-B3LYP functional with aug-cc-pVTZ basis set throughout (Dunning, 1989), predicting the difference in electron affinity between O₃ and SO₃ to be 18 kJ mol⁻¹. Further, it has been shown that CAM-B3LYP is superior to B3LYP with respect to determining activation energies for a range of typical chemical reactions (Peach et al., 2006; Yanai et al., 2004). All DFT calculations were performed using the Gaussian09 package (<http://gaussian.com/>).

The present reactions involve both neutral and charged ozone as well as highly oxidized sulfur species with unknown electronic properties. Therefore, we have performed CCSD(T)-F12 calculations with the VDZ-F12 basis set to test the DFT results (Adler et al., 2007; Peterson et al., 2008). These calculations have been performed using the MOLPRO package (<http://molpro.net/>) and are described in detail elsewhere (Bork et al., 2011b).

Energy barriers were determined by scans of the potential energy surface and confirmed by frequency analyses showing exactly one imaginary frequency. Further, by following the reaction coordinate in both directions it was ensured that transition states indeed connected the desired reactants and products.

The temperature dependent reaction rate constants, $k(T)$, were determined using harmonic transition state theory (Hänggi et al., 1990; Billing and Mikkelsen, 1996),

$$k(T) = A \times \exp\left(\frac{-E_A}{RT}\right), \text{ where} \quad (1)$$

$$A = \frac{\prod v_{\text{reac}}}{\prod v_{TS}^\dagger}. \quad (2)$$

E_A is energy barrier separating reactants and products, R is the molar gas constant and v are the harmonic frequencies for the reactant and transition state. The † indicates that the imaginary frequency should be omitted.

We finally note that in all cases molecular oxygen is treated as an electronic triplet and all reactions are spin allowed.

3 Results and discussion

3.1 Self dissociation

The simplest reaction of O_2SO_3^- based clusters is spontaneous dissociation into O_2 and SO_3^- , i.e.



This reaction is potentially important since the reaction



is distinctly exothermic, thus providing a direct path to complete the catalytic cycle in Fig. 1. Further, both SO_3^- and O_2SO_3^- have recently been measured by Ehn et al. (2010) in a Finnish boreal forest, where typical ratios of 1:10 was found, suggesting that this reaction may be significant.

After optimizing all species we found Gibbs free O_2 binding energies of 50.4 and 41.3 kJ mol^{-1} for the de- and monohydrated clusters, respectively. Little experimental data is available but Möhler et al. (1992), based on mass spectroscopical data, concluded that O_2 is bound stronger to SO_3^- than H_2O is to CO_3^- , HCO_3^- , and NO_3^- . This implies that $\Delta G_{(\text{R3})} \leq -25 \text{ kJ mol}^{-1}$ at room temperature (Keese et al., 1979; Payzant et al., 1971).

Since these binding energies suggest much lower $[\text{SO}_3^-]:[\text{O}_2\text{SO}_3^-]$ ratios than the observed ratio of 1:10, the field data may be explained by SO_3^- being stabilized by more than one water or that the sources and sinks of SO_3^- and O_2SO_3^- induces a dynamic rather than a thermal equilibrium. Although interesting, these considerations are beyond the scope of this study.

We conclude that although some SO_3^- exists, the population of O_2SO_3^- is significantly higher and we will thus focus on this species in the remaining of this article. Finally, we note that, contrary to all previous studies, we generally avoid referring to O_2SO_3^-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as SO_5^- since the $\text{O}_2\text{-SO}_3^-$ bond resembles that of a molecular cluster more than a molecular covalent bond, concerning both distance and strength.

3.2 O_3 collision and formation of $\text{O}_2\text{SO}_3^- \text{W}_{0-1} \text{O}_3$

First, the $\text{O}_2\text{SO}_3^- \text{O}_3$ and $\text{O}_2\text{SO}_3^- \text{WO}_3$ molecular complexes were optimized, see Fig. 2. In the dehydrated system, only a few different stable configurations were found, while at least 10 stable configurations were found in the monohydrated cluster. Most configurations were within 10 kJ mol^{-1} and are thus predicted to co-exist at typical atmospheric temperatures. However, for the remaining of this article mainly the single most stable configurations with respect to ΔG will be considered.

Structurally, only small changes were apparent upon ozonation since the binding preference of the O_3 and H_2O are very different. In accordance with this, the binding energy of O_3 to O_2SO_3^- is only minutely affected by hydration. Considering the equilibrium,



and regardless of the presence or absence of water, we found $\Delta H_{(\text{R5})}$ values of ca. -12 kJ mol^{-1} and $\Delta G_{(\text{R5})}$ values of ca. $+12 \text{ kJ mol}^{-1}$ implying a short lifetime towards de-ozonation.

Not only does the water molecule not change the thermodynamics of ozonation, but neither does the ozone change the thermodynamics of hydration. Considering the condensation-evaporation equilibrium, i.e.



we find $\Delta H_{(\text{R6})}$ and $\Delta G_{(\text{R6})}$ of -42 and -7 kJ mol^{-1} , respectively. Also this reaction was investigated by Möhler et al. (1992) whom determined $\Delta G_{(\text{R6})}$ to $-17.2 \text{ kJ mol}^{-1}$. To clarify this discrepancy, we optimized all relevant species with CCSD(T)-F12 and found an electronic binding energy of 51.3 kJ mol^{-1} . Assuming $\Delta S_{(\text{R6})} = -120 \text{ J mol}^{-1}$

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



K^{-1} , which is typical of water condensation reactions we find $\Delta G_{(R6)} = -15 \text{ kJ mol}^{-1}$, in good accordance with Möhler et al. (1992).

Using the law of mass action

$$\frac{[\text{O}_2\text{SO}_3^- \text{W}]}{[\text{O}_2\text{SO}_3^-]} = [\text{W}] \times \exp\left(\frac{-\Delta G_{R7}}{RT}\right) \quad (3)$$

we find that roughly 90% of O_2SO_3^- are hydrated at $T = 298 \text{ K}$ and 50% relative humidity and the fate of the hydrated cluster is thus of primary importance. However, in all but extremely dry conditions both hydrated and dehydrated clusters co-exist and should both be considered when evaluating Reaction (??).

Having established the structures and relative populations of the relevant clusters, we proceed to evaluate the possible chemical reactions.

3.3 Cluster decomposition


The most important reaction considered here is decomposition of the reactant cluster as



As mentioned, the binding energy of the O_2SO_3^- cluster is at least 25 kJ mol^{-1} , which is only partially compensated by the transferring of the electron. Hence, the dehydrated Reaction (R4) is ca. 15 kJ mol^{-1} exothermic.

However, the water molecule is bound significantly stronger to the O_3^- ion than to the SO_3^- ion. Considering also this effect we find that the monohydrated cluster decomposition reaction is ca. 10 kJ mol^{-1} exothermic at $T = 298 \text{ K}$. The Gibbs free potential energy surface is shown in Fig. 3 for $T = 298 \text{ K}$ and tabulated in the supplement.

Considering first the dehydrated reaction, we identified three normal modes of the $\text{O}_2\text{SO}_3^- \text{O}_3$ clusters clearly leading towards dissociation via



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



These modes are shown in Fig. 2. Sometimes the modes will interfere constructively and in other cases destructively. Assuming that the modes are all in the ground state, we manually determined an effective frequency of 540 cm^{-1} . We use this as prefactor, A , in Eq. (1).


By analysing the Mulliken charges (Mulliken, 1955) during these vibrations it was confirmed that electronic density was smoothly being transferred to O_3 from SO_3^- during a vibration and that no electronic barrier separated the charge transfer Reaction (R4). Hence, the minimum energy required to dissociate the dehydrated cluster is exactly the energy of the net reaction, here 31.9 kJ mol^{-1} above the reactants. This implies an overall rate constant of $1.6 \times 10^{-12}\text{ cm}^3\text{ molecule}^{-1}\text{ s}^{-1}$ at $T = 298\text{ K}$, see also Fig. 4.

Contrary to the dehydrated cluster, in the monohydrated cluster the dissociation into SO_3 , O_2 , and $\text{O}_3^- \text{W}$ is almost thermo neutral. In this case the rate limiting step is formation of the $\text{O}_3 \text{O}_2\text{SO}_3^- \text{W}$ cluster itself and the effective barrier is hence just 12.3 kJ mol^{-1} at 298 K . The mechanism leading to cluster decomposition is however identical to the dehydrated cluster and the modes leading to dissociation are almost identical. These are given in parenthesis in Fig. 2. Due to the small energy barrier we find that this reaction essentially is collision limited, see also Fig. 4.

Comparing the decomposition rates of the dehydrated and monohydrated systems, we see that the water molecule is significantly facilitating cluster decomposition. The main reason is the more favourable thermodynamics since H_2O is bound significantly stronger to O_3^- than to O_2SO_3^- .

3.3.1 Formation of O_3SO_3^-

The other major reaction sink of O_2SO_3^- considered here is oxidation by O_3 . The sulfur is already tetrahedrally coordinated by oxygen, so the only possible oxidation is from further O-O binding. Most likely is the formation of O_3SO_3^- since the reaction



O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is ca. 175 kJ mol^{-1} exothermic. The de- and monohydrated O_3SO_3^- structures are shown in the supplement.

The two transition states of Reaction (R9) were then determined and the monohydrated structure is shown in Fig. 5a. Besides the absent water molecule, the dehydrated structure differs only slightly from the monohydrated and is shown in the supplement. However, considering the barrier heights, the effect of water is very significant. At standard conditions the de- and monohydrated energy barriers are 27.1 and 39.6 kJ mol^{-1} respectively, implying that the water is significantly hindering the oxygen transfer due to destabilisation of the transition state, see Fig. 3. Finally, the pre-factors (Eq. 1) were determined to $3.12 \times 10^{10} \text{ s}^{-1}$ and $1.90 \times 10^9 \text{ s}^{-1}$ respectively.

The reaction rate constant of Reaction (R9) was readily determined from Eq. (1) using the parameters described above, see Fig. 4. It is thus clear that the water molecule, while favouring charge transfer, effectively is hindering the oxidation of O_2SO_3^- . The main reason is the increased energy barrier. The effect is largest at high temperatures but at all temperatures investigated here the hydrated Reaction (R9) is at least 2.5 orders of magnitude slower than the dehydrated.

To our knowledge O_3SO_3^- has not previously been described in the literature, nor has it been observed in experiments or in field studies. This suggests that O_3SO_3^- is reactive and has too short lifetime to allow for observation. We therefore proceeded to investigate the possible decomposition of O_3SO_3^- via



The monohydrated transition state is shown in Fig. 5(b) and the dehydrated transition state is shown in the supplement. We hereby determine the energy barrier to ca. 22 kJ mol^{-1} at standard conditions, practically independent of hydration, see Fig. 3. Since the energy release of Reaction (R9) is so much larger than the energy barrier, it is evident that Reaction (R10) is almost instantaneous.

 **O_2SO_3^- chemistry
and catalytic SO_2
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Overall catalytic turnover

The presence of an ion catalysed SO₂ oxidation cycle is of interest since an extra source of atmospheric H₂SO₄ significantly may alter the conditions for aerosol particle formation and growth. Although this assessment is the overall goal of this line of studies, several unresolved factors remain.

Considering the real atmosphere the most noticeable shortcoming is the influence of CO₂. This is known to be the dominant sink of O₃⁻ via



and cannot be neglected in any realistic model (Dotan et al., 1977; Luts and Parts, 2002). The chemical fate of CO₃⁻ is uncertain although the reaction



is known to proceed at collision rates, similar to Reaction (R2) (Fehsenfeld and Ferguson, 1974). However, extending the catalytic cycle in Fig. 1 to include these reactions is beyond the scope of this study.

Presently, we are able to assess the overall catalytic turnover in CO₂ free atmospheres, including most of the until now conducted chamber studies of ion induced nucleation. Taking all investigated clusters, their relative concentrations, and all reactions into account we find that the probability for cluster decomposition through Reaction (R3b) is 10⁵-10⁶ times more likely than other reactions between O₃SO₃⁻W₀₋₁ and O₃. Details are presented in the supplement. This thus suggests that each free electron may induce up to 10⁵-10⁶ SO₂ oxidations in atmospheres consisting of N₂, O₂, O₃, H₂O and SO₂ only.

This is consistent with recent experimental results from Enghoff et al. (2012). In the absence of UV light it was found that each electron yielded a production of 10⁶ sulfates in atmospheres heavily enriched in SO₂ and O₃. Further it was found that the isotopic fingerprint of this ion produced sulfate was very different of sulfate from UV induced SO₂ oxidation, clearly pointing at two different chemical mechanisms.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



More commonly chamber atmospheres are made to resemble typical atmospheric conditions and hence various pollutants become important. In those cases, the catalytic cycle may be terminated by mechanisms different from the ones presented so far, most likely involving species different from SO_2 and O_3 . Considering e.g. a collision between $\text{O}_3^-(\text{H}_2\text{O})_n$ and a NO_2 radical, this will rapidly be oxidized to NO_3^- with large energy gain,



Due to the stability of NO_3^- , the catalytic cycle for that particular electron is most likely terminated. Other potential scavengers include any species in which an electron may be more stabilized than on O_3^- or O_2SO_3^- , either through immediate charge transfer or after some chemical reaction. These include most radicals, NO_x , cations and pre-existing aerosol particles or molecular clusters. Typically, NO_x will be the dominant electron scavenger.

We here outline two methods of determining the rate of ion induced H_2SO_4 formation. Most simply, if the actual concentrations of the involved ions are known, the rate of ion induced H_2SO_4 formation may be determined based on the mechanism outlined in Fig. 1. Assuming $[\text{O}_3] \gg [\text{SO}_2]$ we obtain

$$r_{\text{ion cat.}} = Z \times [\text{O}_3^-(\text{H}_2\text{O})_n] \times [\text{SO}_2] \quad (4)$$

where Z denote the collision rate.

Alternatively, the ion induced H_2SO_4 formation rate may be determined by realising that the incoming flux of electrons, J_{ion} , is the primary limiting factor. As shown in Fig. 1, the electron will remain catalytically active until scavenged, which occurs when the ionic cluster is hit by a scavenger. Again assuming $[\text{O}_3] \gg [\text{SO}_2]$ and that NO_x is the dominant

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scavenger, the number of catalytic cycles per electron is thus determined as

$$\text{Max.cyc.} = \frac{Z(\text{SO}_2)}{Z(\text{electron scavenger})} \quad (5)$$

$$\sim \frac{[\text{SO}_2]}{[\text{NO}_x]} \quad (6)$$

where Z denotes the collision rate. The overall catalytic efficiency and is then given as

$$r_{\text{ion cat.}} = J_{\text{ion}} \times \text{Max.cyc.} \quad (7)$$

We now proceed to evaluate the relevance of ion catalyzed SO_2 oxidation in a typical CO_2 free chamber. Since we have no data for the ionic concentrations we will use the latter approach. The well known UV induced H_2SO_4 production rate via Reaction (R1) is

$$r_{\text{UV}} = k_{\text{UV}} \times [\text{M}] \times [\text{OH}] \times [\text{SO}_2], \quad (8)$$

where $k_{\text{UV}} = 4.5 \times 10^{-31} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ (Atkinson et al., 2004).

We here assume the following conditions: $[\text{OH}] = 5 \times 10^5 \text{ cm}^{-3}$ (day and night average), $[\text{NO}_x] = 10 \text{ ppt} = 2.5 \times 10^8 \text{ cm}^{-3}$, $[\text{SO}_2] = 5 \text{ ppb} = 1.25 \times 10^{11} \text{ cm}^{-3}$, $[\text{O}_3] \gg [\text{SO}_2]$, and $J_{\text{ion}} = 20 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ corresponding to mid-troposphere. We thus obtain

$$\frac{[\text{H}_2\text{SO}_4]_{\text{ion cat.}}}{[\text{H}_2\text{SO}_4]_{\text{total}}} = \frac{r_{\text{ion cat.}}}{r_{\text{ion cat.}} + r_{\text{UV}}} = 1.5\%. \quad (9)$$

Obviously, this is merely a rough estimate intended mainly as an illustrative example. However, variations in $[\text{H}_2\text{SO}_4]$ on the order of a few percent are important and the lack of this contribution in models of gaseous sulfur chemistry may lead to significant inaccuracies in predicting aerosol formation and growth rates.

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Conclusions

It has previously been determined that de- or monohydrated clusters of O_2SO_3^- are natural products of free atmospheric electrons. We have investigated the chemical fate of O_2SO_3^- after various reactions with O_3 .

Using a combination of DFT and coupled cluster levels of theory we have optimized several possible reactants and products. We have linked the most important of these either through transition states or shown that the reactions are barrierless with respect to ΔE . We have determined pre-factors and free energy barrier heights and, using transition state theory, also the reaction rates.

We find two major reaction mechanisms leading to fundamentally different products, shown in Fig. 3. The first mechanism is a conventional oxidation eventually leading to SO_4^- with large energy gain. However, significant energy barriers must be overcome and we find that this reaction is of minor importance.

Of primary importance is the transferring of the electron from O_2SO_3^- to O_3 . In the dehydrated system this is distinctly endothermic but still requires a lower energy than crossing the energy barrier of oxidation. In the hydrated systems, the water molecule can transfer with the charge from $\text{O}_2\text{SO}_3^-(\text{H}_2\text{O})$ to $\text{O}_3^-(\text{H}_2\text{O})$ with significant energy gain. We find that this path is 10^5 to 10^6 times more probable than $\text{O}_2\text{SO}_3^-(\text{H}_2\text{O})$ oxidation. This reaction closes a catalytic cycle of SO_2 oxidation shown in Fig. 1. Both the chemical mechanism and the catalytic turnover number is in accordance with a recent chamber study by Enghoff et al. (2012).

Considering also the concentrations of other typical reactants we conclude that the cycle is more probable to terminate due to reactions with an electron scavenger, most likely some NO_x , than due to its own inherent side reactions. We are unable to assess the influence of this cycle in the real atmosphere, most noticeable due to CO_2 . In chamber studies under UV exposure and in realistic gas mixtures we roughly estimate the contribution of ion catalysed H_2SO_4 to be on the order of several percent.

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Besides being a source of atmospheric H₂SO₄, we note that the previously unknown origin of both O₂SO₃⁻ and SO₄⁻, seen in several experiments and field studies, readily is explained through this mechanism. Finally, this mechanism is also in accordance with the absence of SO₆⁻ from any observations since it explains why SO₆⁻ immediately decomposes to SO₄⁻ and O₂.

Acknowledgements. The authors thank the Academy of Finland for funding and the CSC IT Centre for Science in Espoo, Finland for computer time. We acknowledge the Villum foundation, University of Helsinki Chancellors travel grant, the Academy of Finland (CoE Project No. 1118615, LASTU Project No. 135054), ERC Project No. 257360- MOCAPAF for funding. We finally thank Martin B. Enghoff and Henrik Svensmark for valuable comments.

References

- Adler, T. B., Knizia, G., and Werner, H. J.: A simple and efficient CCSD (T)-F12 approximation, *J. Chem. Phys.*, 127, 221–106, 2007. 30182
- Arnold, D., Xu, C., Kim, E., and Neumark, D.: Study of low-lying electronic states of ozone by anion photoelectron spectroscopy of O, *J. Chem. Phys.*, 101, 912–922, 1994. 30181
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004. 30190
- Billing, G. D. and Mikkelsen, K. V.: Introduction to molecular dynamics and chemical kinetics, Wiley, New York, USA, 1996. 30182
- Bondybey, V. and Beyer, M.: How many molecules make a solution?, *Int. Rev. Phys. Chem.*, 21, 277–306, 2002. 30179
- Bork, N., Kurtén, T., Enghoff, M. B., Pedersen, J. O. P., Mikkelsen, K. V., and Svensmark, H.: Structures and reaction rates of the gaseous oxidation of SO₂ by an O₃-(H₂O)₀–5 cluster – a density functional theory investigation, *Atmos. Chem. Phys.*, 12, 3639–3652, doi:10.5194/acp-12-3639-2012, 2012. 30180



**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bork, N., Kurtén, T., Enghoff, M. B., Pedersen, J. O. P., Mikkelsen, K. V., and Svensmark, H.: Ab initio studies of O₂⁻(H₂O)_n and O₃⁻(H₂O)_n anionic molecular clusters, n ≥ 12, Atmos. Chem. Phys., 11, 7133–7142, doi:10.5194/acp-11-7133-2011, 2011. 30180, 30182
- Carlsaw, K., Harrison, R. G., and Kirkby, J.: Cosmic rays, clouds, and climate, Science, 298, 1732–1737, 2002. 30178
- Dobrin, S., Boo, B., Alconcel, L., and Continetti, R.: Photoelectron Spectroscopy of SO₃⁻ at 355 and 266 nm, J. Phys. Chem. A, 104, 10695–10700, 2000. 30181
- Dotan, I., Davidson, J. A., Streit, G.E. Albritton, D., and Fehsenfeld, F.: A study of the reaction O₃⁻ + CO₂ ↔ CO₃⁻ + O₂ and its implication on the thermochemistry of CO₃ and O₃ and their negative ions, J. Phys. Chem., 67, 2874–2879, 1977. 30188
- Dunning, T. H. J.: Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen, J. Chem. Phys., 90, 1007–1023, 1989. 30182
- Ehn, M., Junninen, H., Petäjä, T., Kurtén, T., Kerminen, V.-M., Schobesberger, S., Manninen, H. E., Ortega, I. K., Vehkamäki, H., Kulmala, M., and Worsnop, D. R.: Composition and temporal behavior of ambient ions in the boreal forest, Atmos. Chem. Phys., 10, 8513–8530, doi:10.5194/acp-10-8513-2010, 2010. 30180, 30183
- Enghoff, M. B., Bork, N., Hattori, S., Meusinger, C., Nakagawa, M., Pedersen, J. O. P., Danielache, S., Ueno, Y., Johnson, M. S., Yoshida, N., and Svensmark, H.: An isotopic analysis of ionising radiation as a source of sulphuric acid, Atmos. Chem. Phys., 12, 5319–5327, doi:10.5194/acp-12-5319-2012, 2012. 30179
- Enghoff, M. B., Bork, N., Hattori, S., Meusinger, C., Nakagawa, M., Pedersen, J. O. P., Danielache, S., Ueno, Y., Johnson, M. S., Yoshida, N., and Svensmark, H.: An isotopic analysis of ionising radiation as a source of sulphuric acid, Atmos. Chem. Phys., 12, 5319–5327, doi:10.5194/acp-12-5319-2012, 2012. 30180, 30188, 30191
- Fehsenfeld, F. C. and Ferguson, E. E.: Laboratory studies of negative ion reactions with atmospheric trace constituents, J. Chem. Phys., 61, 3181–3193, 1974. 30180, 30188
- Gleason, J.: Ph.D. dissertation, Univ. of Colorado, U.S.A, 1987. 30181
- Hänggi, P., Talkner, P., and Borkovec, M.: Reaction-rate theory: fifty years after Kramers, Rev. Mod. Phys., 62, 251–342, 1990. 30182
- Kazil, J., Stier, P., Zhang, K., Quaas, J., Kinne, S., O'Donnell, D., Rast, S., Esch, M., Ferrachat, S., Lohmann, U., and Feichter, J.: Aerosol nucleation and its role for clouds and Earth's radiative forcing in the aerosol-climate model ECHAM5-HAM, Atmos. Chem. Phys., 10, 10733–10752, doi:10.5194/acp-10-10733-2010, 2010. 30178

Keesee, R., Lee, N., and Castleman Jr, A.: Properties of clusters in the gas phase. 3. Hydration complexes of carbonate (1-) and bicarbonate (1-) ions, *J. Am. Chem. Soc.*, 101, 2599–2604, 1979. 30183

5 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A.,
10 Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–435, 2011. 30179

15 Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V., Birmili, W., and McMurry, P.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35, 143–176, 2004. 30178

Kupiainen, O., Ortega, I. K., Kurtén, T., and Vehkamäki, H.: Amine substitution into sulfuric acid – ammonia clusters, *Atmos. Chem. Phys. Discuss.*, 11, 30853–30875, doi:10.5194/acpd-11-30853-2011, 2011. 30179

20 Kurtén, T., Ortega, I. K., and Vehkamäki, H.: The sign preference in sulfuric acid nucleation, *J. Mol. Struct.-Theochem.*, 901, 169–173, 2009. 30180

Li, W. and McKee, M.: Theoretical Study of OH and H₂O Addition to SO₂, *J. Phys. Chem. A*, 101, 9778–9782, 1997. 30179

25 Lovejoy, E., Curtius, J., and Froyd, K.: Atmospheric ion-induced nucleation of sulfuric acid and water, *J. Geophys. Res.*, 109, D08 204, doi:10.1029/2003JD004460, 2004. 30179

Luts, A. and Parts, T.: Evolution of negative small air ions at two different temperatures, *J. Atmos. Sol.-Terr. Phys.*, 64, 763–774, 2002. 30180, 30188

Mauldin III, R., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F., Kerminen, V., and Kulmala, M.: A new atmospherically relevant oxidant of sulphur dioxide,
30 *Nature*, 488, 193–196, 2012. 30179

Möhler, O., Reiner, T., and Arnold, F.: The formation of SO₅⁻ by gas phase ion-molecule reactions, *J. Chem. Phys.*, 97, 8233–8239, 1992. 30180, 30183, 30184, 30185

**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Morokuma, K. and Muguruma, C.: Ab initio molecular orbital study of the mechanism of the gas phase reaction SO₃+ H₂O: importance of the second water molecule, *J. Am. Chem. Soc.*, 116, 10316–10317, 1994. 30179

Mulliken, R.: Electronic Population Analysis on LCAO-MO Molecular Wave Functions, *J. Phys. Chem.*, 23, 1831–1840, 1955. 30186

Nadykto, A. B., Al Natsheh, A., Yu, F., Mikkelsen, K. V., and Ruuskanen, J.: Quantum nature of the sign preference in ion-induced nucleation, *Phys. Rev. Lett.* 96, 125701, doi:10.1103/PhysRevLett.96.125701, 2006. 30180

Nieminen, T., Manninen, H., Sihto, S., Yli-Juuti, T., Mauldin, III, R., Petaja, T., Riipinen, I., Kerminen, V., and Kulmala, M.: Connection of sulfuric acid to atmospheric nucleation in boreal forest, *Environ. Sci. Technol.*, 43, 4715–4721, 2009. 30179

Novick, S., Engelking, P., Jones, P., Futrell, J., and Lineberger, W.: Laser photoelectron, photodetachment, and photodestruction spectra of O, *J. Phys. Chem.*, 70, 2652, 1979. 30181

Ortega, I. K., Kupiainen, O., Kurtén, T., Olenius, T., Wilkman, O., McGrath, M. J., Loukonen, V., and Vehkamäki, H.: From quantum chemical formation free energies to evaporation rates, *Atmos. Chem. Phys.*, 12, 225–235, doi:10.5194/acp-12-225-2012, 2012. 30179

Payzant, J., Yamdagni, R., and Kebarle, P.: Hydration of CN⁻, NO₂⁻, NO₃⁻, and OH⁻ in the Gas Phase, *Can. J. Chemistry*, 49, 3308–3314, 1971. 30183

Peach, M. J. G., Helgaker, T., Sałek, P., Keal, T. W., Lutnæs, O. B., Tozer, D. J., and Handy, N. C.: Assessment of a Coulomb-attenuated exchange–correlation energy functional, *Phys. Chem. Chem. Phys.*, 8, 558–562, 2006. 30182

Peterson, K. A., Adler, T. B., and Werner, H. J.: Systematically convergent basis sets for explicitly correlated wavefunctions: The atoms H, He, B–Ne, and Al–Ar, *J. Chem. Phys.* 128, 084102, doi:10.1063/1.2831537, 2008. 30182

Rosenfeld, D.: Aerosols, clouds, and climate, *Science*, 312, 1323–1324, 2006. 30178

Simpson, J. and Wiggert, V.: Models of precipitating cumulus towers, *Mon. Weather Rev.*, 97, 471–489, 2009. 30178

Sipilä, M., Berndt, T., Petäjä, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin III, R., Hyvärinen, A., Lihavainen, H., Kulmala, M.: The role of sulfuric acid in atmospheric nucleation, *Science*, 327, 1243–1246, 2010. 30179

Sorokin, A. and Arnold, F.: Laboratory study of cluster ions formation in H₂SO₄-H₂O system: Implications for threshold concentration of gaseous H₂SO₄ and ion-induced nucleation kinetics, *Atmos. Environ.*, 41, 3740–3747, 2007. 30180

**O₂SO₃⁻ chemistry
and catalytic SO₂
oxidation**

N. Bork et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sorokin, A. and Arnold, F.: Analysis of experiments on ion-induced nucleation and aerosol formation in the presence of UV light and ionizing radiation, *Atmos. Environ.*, 43, 3799–3807, 2009. 30180

Spracklen, D. V., Bonn, B., and Carslaw, K. S.: Boreal forests, aerosols and the impacts on clouds and climate, *Philos. T. R. Soc. A*, 366, 4613–4626, 2008. 30178

Su, T. and Bowers, M.: Theory of ion-polar molecule collisions. Comparison with experimental charge transfer reactions of rare gas ions to geometric isomers of difluorobenzene and dichloroethylene, *J. Chem. Phys.*, 58, 3027, doi:10.1063/1.1679615, 1973. 30179

Svensmark, H., Pedersen, J. O. P., Marsh, N. D., Enghoff, M. B., and Uggerhøj, U. I.: Experimental evidence for the role of ions in particle nucleation under atmospheric conditions, *Proc. R. Soc. A*, 463, 385–396, doi:10.1098/rspa.2006.1773, 2007. 30180

Tohmfor, G. and Volmer, M.: *Annalen der Physik*, doi:10.1002/andp.19384250204, 425, 109–131, 1938. 30179

Welz, O., Savee, J., Osborn, D., Vasu, S., Percival, C., Shallcross, D., and Taatjes, C.: Direct Kinetic Measurements of Criegee Intermediate (CH₂OO) Formed by Reaction of CH₂I with O₂, *Science*, 335, 204–207, 2012. 30179

Wiedensohler, A., Cheng, Y., Nowak, A., Wehner, B., Achtert, P., Berghof, M., Birmili, W., Wu, Z., Hu, M., Zhu, T., et al.: Rapid aerosol particle growth and increase of cloud condensation nucleus activity by secondary aerosol formation and condensation: A case study for regional air pollution in northeastern China, *J. Geophys. Res.*, 114, D00G08, doi:10.1029/2008JD010884, 2009. 30178

Yanai, T., Tew, D. P., and Handy, N. C.: A new hybrid exchange-correlation functional using the Coulomb-attenuating method (CAM-B3LYP), *Chem. Phys. Lett*, 393, 51–57, 2004. 30181, 30182

Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, doi:10.5194/acp-9-7691-2009, 2009. 30178

O_2SO_3^- chemistry and catalytic SO_2 oxidation

N. Bork et al.




Fig. 1. Schematics of a simple catalytic SO_2 oxidation cycle. The net reaction is $\text{O}_3 + \text{SO}_2 \rightarrow \text{O}_2 + \text{SO}_3$ and the catalyst is the electron. All reaction rates are mainly collision limited.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


O₂SO₃⁻ chemistry and catalytic SO₂ oxidation

N. Bork et al.




Fig. 2. Top: the molecular complex between O₂SO₃⁻, H₂O and O₃ including some descriptive bond lengths. Removing either the H₂O or O₃ causes little structural rearrangement. Bottom: Directions and frequencies of the three normal vibrational modes along the SO₃⁻-O₂ bond in the dehydrated (monohydrated) cluster. Sulfur is yellow, oxygen is red, and hydrogen is white.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

O₂SO₃⁻ chemistry and catalytic SO₂ oxidation

N. Bork et al.




Fig. 3. Schematic overview of the Gibbs free energy of the relevant species at standard conditions. The barrier height for O₂SO₃⁻ oxidation is increased by ca. 12 kJ mol⁻¹ by the presence of a water molecule while the thermodynamics of charge transfer and cluster decomposition is changed from distinctly endothermic to almost thermo neutral. At standard conditions and 50 % relative humidity ca. 90 % of O₂SO₃⁻ is monohydrated.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



O₂SO₃⁻ chemistry and catalytic SO₂ oxidation

N. Bork et al.




Fig. 4. Bimolecular rate constants of cluster decomposition mechanism and O₂SO₃⁻ oxidation in cm³ molecule⁻¹ s⁻¹ as function of temperature. For both the dehydrated and the monohydrated systems, cluster breakup is favoured although much more so in the hydrated systems. At standard conditions and 50 % relative humidity, ca. 90 % of O₂SO₃⁻ is monohydrated. The collision rate with H₂O is indicated (50 % relative humidity).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 5. Transition states for **(a)** the oxidation of $\text{O}_2\text{SO}_3^- \text{W}$ to $\text{O}_3\text{SO}_3^- \text{W}$ and **(b)** the following dissociation to $\text{SO}_4^- \text{W}$. The active bonds are shown as broken grey and some descriptive bond lengths are indicated. Sulfur is yellow, oxygen is red and hydrogen is white.