

**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

Overview of the Mount Tai Experiment (MTX2006) in Central East China in June 2006: studies of significant regional air pollution

Y. Kanaya¹, H. Akimoto^{1,*}, Z.-F. Wang², P. Pochanart^{1,**}, K. Kawamura³, Y. Liu^{1,2}, J. Li^{1,2}, Y. Komazaki¹, H. Irie^{1,***}, X.-L. Pan¹, F. Taketani¹, K. Yamaji¹, H. Tanimoto⁴, S. Inomata⁴, S. Kato⁵, J. Suthawaree⁵, K. Okuzawa³, G. Wang^{3,****}, S. G. Aggarwal^{3,*****}, P. Q. Fu^{3,2}, T. Wang⁶, J. Gao⁶, Y. Wang², and G. Zhuang⁷

¹Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, 3173-25 Showa-machi, Kanazawa-ku, Yokohama 2360001, Japan

²LAPC, Institute for Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

³Institute of Low Temperature Science, Hokkaido University, Sapporo 0600819, Japan

⁴National Institute for Environmental Studies, Tsukuba 3058506, Japan

⁵Tokyo Metropolitan University, Hachioji 1920397, Japan

⁶The Hong Kong Polytechnic University, Hong Kong, China

⁷Fudan University, Shanghai 200433, China

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



* now at: Asia Center for Air Pollution Research, Niigata 9502144, Japan

** now at: National Institute of Development Administration, Bangkok 10240, Thailand

*** now at: Chiba University, Chiba 2638522, Japan

**** now at: Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710075, China

***** now at: National Physical Laboratory, New Delhi 110012, India

Received: 25 December 2012 – Accepted: 4 January 2013 – Published: 15 January 2013

Correspondence to: Y. Kanaya (yugo@jamstec.go.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We conducted an intensive field campaign at the summit of Mt. Tai (1534 m a.s.l.), Shandong Province, located at the center of Central East China, during the period 28 May to 30 June 2006, to study seasonal maxima of regional air pollution with respect to ozone (O_3) and aerosols. The specific objectives, campaign design, and major findings are summarized. High concentrations of O_3 and its precursors, and aerosols, were detected and studied in the context of annual variations. Most importantly, we identified that emissions from regional-scale open crop residue burning after the harvesting of winter wheat, together with photochemical aging, strongly increased the concentrations of O_3 , aerosols, and primary species relevant to air quality in this month of the year. Studies of in-situ photochemical activity, regional source attribution of O_3 , O_3 -aerosol interactions, validation of satellite observations of tropospheric NO_2 , behaviors of volatile organic compounds, organic/inorganic aerosol species, loss rates of black carbon (BC), and instrument inter-comparisons are also summarized. The observed BC levels must have a strong impact on the regional climate.

1 Introduction

Asian regional air pollution is a major element of global air pollution (Akimoto, 2003) and is of great interest from the viewpoints of urban and rural air quality, trans-boundary air pollution, hemispherical air pollution, and climate impacts. Central East China (CEC), covering the North China Plain (NCP) and the Yangtze River delta region, stands out as the most highly polluted area in this region, as evidenced by satellite observations of NO_2 , which is regarded as representative of air pollutants (Beirle et al., 2004; Martin et al., 2006), and aerosols (Chin et al., 2004). It has also been observed that the tropospheric column density of NO_2 has been increasing rapidly since 2000 (Richter et al., 2005). Measurements of the aerosol optical depth (AOD) at 550 nm, performed by the satellite sensors MODIS/Terra and MODIS/Aqua from 2000, suggest that the

ACPD

13, 1527–1573, 2013

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NCP has one of the highest AOD values. The observed AOD has seasonal variations, with a maximum (~ 1.0) in June/July and a minimum (~ 0.5) in November–January. In contrast to satellite remote-sensing data, however, ground-based in-situ observational data (except in urban or near-urban areas) are very scarce, and information on the actual circumstances of regional air pollution, particularly for ozone (O_3), aerosols, and their precursors, has not been available for this region.

Among the regional air pollutants in CEC, O_3 and black carbon (BC) are of particular importance in the context of global climate change, since they are short-lived climate forcers with positive radiative forcing (RF), and are thought to contribute as much as long-lived greenhouse gases do to global warming (IPCC, 2007). High concentrations of O_3 are very toxic to humans (WHO, 2003) as well as to natural ecosystem and agricultural crops (Fuhrer et al., 1997; USEPA, 2006). In this context, to obtain in-situ observational data of air pollutants such as O_3 , BC, other aerosol components, and their precursors, the Frontier Research Center for Global Change (FRCGC) of the Japan Agency for Marine-Earth Science and Technology (JAMSTEC), and the Institute of Atmospheric Physics (IAP), Chinese Academy of Sciences (CAS), organized a field campaign named MTX2006 (Mount Tai Experiment 2006) at the summit of Mt. Tai (36.26° N, 117.11° E, 1534 m above sea level (a.s.l.)) in Shandong Province, China in June 2006. Mt. Tai is located at the center of the NCP, and is expected to provide regionally representative data for CEC. The obtained data and their analysis are presented in this special issue.

Prior to the MTX2006 campaign, we identified seasonal variations in surface O_3 at three mountain sites in CEC: at Mt. Tai, Mt. Hua (34.48° N, 110.08° E, 2065 m a.s.l., Shaanxi Province), and Mt. Huang (30.16° N, 118.26° E, 1840 m a.s.l., Anhui Province), as shown in Fig. 1. From the results, we found that the O_3 concentration was highest at Mt. Tai, with a monthly peak in June, followed by Mt. Hua and Mt. Huang, for which the peaks were smaller and appeared earlier, i.e. in late spring (Li et al., 2007). The highest concentration of O_3 at Mt. Tai was in accord with the spatial distribution of emissions of O_3 precursors, NO_x , and volatile organic compounds (VOCs), obtained in the emission

inventory database (Ohara et al., 2007). The data show that Mt. Tai is in the middle of the highest emission area, Mt. Hua is at the western edge, and Mt. Huang is just outside the heavy emission area. We therefore selected Mt. Tai as our campaign site and chose June as the study period, to represent the most serious regional air pollution in CEC.

Wang et al. (2006) simulated a regional high O₃ episode observed at two mountain sites (Mt. Tai and Mt. Huang) in May 2004, using the nested air quality prediction modeling system (NAQPMS) and found that it was the transport of O₃ and its precursors from the Yangtze delta that caused the high O₃ episode at the two sites, with contributions of 20–50 % during the episode. However, the lack of measurements of O₃ precursors prevented detailed studies.

Gao et al. (2005) reported O₃ and CO observations at the summit of Mt. Tai during the period June–November 2003, showing average concentrations of 58 ± 16 ppbv for O₃ and 393 ± 223 ppbv for CO during the study period. Another paper reported precipitation chemistry at Mt. Tai (Wang et al., 2008b). Observations at Shangdianzi (a rural site in northern CEC) showed that the average maximum hourly O₃ concentration in June exceeded 120 ppbv (Lin et al., 2008). Xu et al. (2008) summarized surface O₃ data from the Lin'an Regional Background Station, Zhejiang Province in the Yangtze delta region, during six periods between August 1991 and July 2006 (Luo et al., 2000; Wang et al., 2001, 2004). The seasonal cycle shows two peaks at around 40–50 ppbv, with the primary peak in May and the secondary peak in October. A maximum hourly mean O₃ concentration of 156 ppbv occurred in June 2006. Intensive field campaigns including measurements of O₃, CO, NO_y, SO₂, and VOCs were performed at Lin'an, in 1999–2000 and in spring 2001 (Wang et al., 2002, 2004) and key emission factors (e.g. CO/NO_x) were studied.

To our knowledge, there have been no other papers reporting air quality with respect to O₃ and its precursors in 2006 or earlier at regionally representative sites in NCP, which can be used for comparisons with results from regional or global chemical transport models. MTX2006 is therefore the first intensive campaign to provide

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a comprehensive set of near-surface air pollutant concentrations at a regionally representative site in NCP.

The MTX2006 study is a component of the O₃ and BC project sponsored by the Global Environmental Research Fund of the Ministry of the Environment, Japan. This paper presents an overview of the campaign and the obtained results, and a synthesis of the individual papers presented in this special issue on MTX2006, encompassing from field observations to modeling, remote sensing, and laboratory studies. Also, we newly explore comparisons of $\Delta\text{CO}/\Delta\text{NO}_y$ and $\Delta\text{O}_3/\Delta\text{NO}_z$ ratios between observations and model simulations, correlations between VOCs and levoglucosan, dependence of OC mass concentrations on air mass age, and fractions of OC which were molecularly identified.

2 Specific objectives of MTX2006

The general objective of MTX2006 was to quantify air quality in the middle of the NCP. To attain this objective, the specific aims were to answer the following questions.

1. What are the concentrations of surface O₃, aerosols (including BC), their precursors, and related species at the top of Mt. Tai, located in the middle of the regional pollution area of CEC, in the highest pollution season of June?
2. Why does the surface O₃ concentration peak sharply in June, as opposed to showing a broad plateau in the summer, as observed in Europe, another continental source region?
3. How much of the O₃ is transported to Mt. Tai from other regions, and how much is produced locally?
4. How fast is the in-situ photochemical production of O₃ at Mt. Tai? Is the O₃ production regime NO_x-limited or VOC-limited?

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5. What are the characteristics of the VOC composition compared with those of other urban/rural areas?
6. How well do the BC concentrations measured using different techniques agree with each other?
- 5 7. What is the major constituent of the aerosol particles? Are secondary organic aerosols and oxygenated VOCs (OVOCs) such as HCHO present abundantly, as inferred from the high O₃ levels?
8. How reliable are the satellite observations of NO₂ in CEC? Do they agree with MAX-DOAS (multi-axis-differential optical absorption spectroscopy) observations?
- 10 9. What are the effects of aerosols on spectral actinic fluxes and *J* values (photolysis frequencies) of several atmospheric molecules relevant to photochemistry?
10. Can we identify sources of OC (organic carbon) aerosol particles using levoglucosan as a biomass and biofuel tracer, lignin products as tracers for pine trees, and hopanoid hydrocarbons as automobile tracers? What are their temporal variations? What is the fraction of OC that we can identify molecularly?
- 15

During the first half of the campaign, we easily recognized that the air quality at the field site was strongly influenced by large-scale post-harvest open crop residue burning (OCRB, winter wheat). As a result, a large number of the papers in this issue are devoted to studies of the impact of open biomass burning on O₃, its precursors, and aerosol compositions, although this was not specifically targeted in the planning stage.

20

3 Geography of Mt. Tai

The campaign site is the meteorological observatory at the summit of Mt. Tai (36.26° N, 117.11° E, 1534 m a.s.l.), which is located in Shandong Province in the NCP in CEC. Figure 1 shows the location of Mt. Tai, superimposed on the emission inventory map of NO₂ (EDGAR ver. 4.2, source: EC-JRC/PBL EDGAR version 4.2, <http://edgar.jrc.ec.europa.eu/>, 2011) for the year 2006, showing that the campaign site is located in the middle of a high NO₂ emission area of CEC. Mt. Tai is an isolated peak in a UNESCO (United Nations Educational, Scientific, and Cultural Organization) World Heritage site, free from local emission sources. Although there are several small restaurants, shops, and hotels for tourists ~ 500 m away, emissions from these sources are thought to be too small to affect the observational data. Mt. Tai is about 6 km north of Tai'an city (population in its central region: ca. 1.5 million) and is about 60 km southeast of the provincial capital city, Jinan (population: ca. 4.3 million). So, although the air quality of Mt. Tai would be affected by regional emissions when the boundary layer builds up, covering the top of the mountain, in the afternoon, as will be described later, the site is believed to be free from local anthropogenic pollution, and is expected to provide regionally representative data.

4 Measured species investigators and campaign period

Table 1 lists the measured species and parameters along with methods, responsible investigator(s) and affiliation. Detailed descriptions of each instrument are included in companion papers of this special issue. They included O₃, NO_x/NO_y, NMHC (non-methane hydrocarbons), VOCs, CO and spectral actinic flux, and composition of aerosols including BC/EC (elemental carbon) and OC mass concentration, molecular composition of organic aerosols, ionic composition, and AOD. CO₂ was included to assist discussion on sources of aerosols and gases. Conventional meteorological parameters were also obtained. In addition to in situ measurements, remote sensing

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measurements of NO₂ vertical column density (VCD) and AOD were made by MAX-DOAS instruments located both at the top and foothill of the mountain. The foothill site is located at 36.16° N, 117.15° E, 126 m a.s.l.

In addition to the main organizing institutes, namely FRCGC/JAMSTEC and IAP/CAS, scientists from several other universities and research institutes participated in the campaign. As seen from Table 1, these included the National Institute for Environmental Studies (NIES), Tokyo Metropolitan University, and Hokkaido University in Japan, and Shandong University, Fudan University, and Nanjing University in China.

The MTX2006 campaign was conducted during the period from 28 May to 30 June 2006. The actual measurement period depends on the instrument. The meteorological conditions are summarized in Fig. 2. On 9, 15, and 29 June, low-pressure systems passed over the region. During the periods of passage, the wind direction was north. During the period 22–25 June, the main wind direction was east, as a result of a high-pressure system located in the east, near Japan. The wind direction in the other periods was normally south or southwest. Precipitation was recorded on 7, 13, 14, 21, 22, 26, 28, and 29 June. The average daytime maxima of $J(\text{NO}_2)$ and $J(\text{O}^1\text{D})$ (2π sr) were 7.4×10^{-3} and $2.8 \times 10^{-5} \text{ s}^{-1}$, respectively. The temperature and relative humidity (RH) ranges were 9–25 °C and 20–100 %, with averages of 17 °C and 67 %, respectively.

5 Modeling activities

Modeling activities were included as part of MTX2006 as post-mission analysis. Table 2 lists the modeling systems used, targeted subjects, and principal investigator and affiliation. A one-box model was used to study photochemical production and loss of O₃ by fixing the key species and parameters to observed values (Kanaya et al., 2009). Two different three-dimensional chemical transport models (3-D CTM) were used; one is NAQPMS, developed by Z. Wang et al. (2006), and the other is Community Multi-scale Air Quality Modeling System (CMAQ), developed by Byun and Ching (1999). The NAQPMS was used for studying source attribution and outflow of O₃ at Mt. Tai (Li et al.,

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2008a). The CMAQ was used to evaluate the impact of crop residue burning on O₃, CO, BC and OC over CEC (Yamaji et al., 2010), and to study the impact of the Asian monsoon on O₃ and precipitation acidity in Eastern and Central China (He et al., 2008; Ge et al., 2011).

5 6 Overview of MTX2006 findings

6.1 High concentrations of O₃ and aerosols at Mt. Tai and related features

Figure 3 shows the overall temporal variations in O₃, CO, benzene, CO₂, NO_x, NO_y, BC, OC, sulfate, and nitrate concentrations during the campaign period. Table 3 lists their average, maximum, and minimum concentrations. In Fig. 3, two high-concentration episodes are clearly seen for most gases and aerosols during the periods 6–7 and 12–13 June. These episodes are caused by the strong impact of agricultural waste burning plumes, as will be discussed later. A regular diurnal pattern with a daytime maximum and nighttime minimum can be seen particularly for NO_y, BC, and OC in the latter half of the campaign period. This pattern is the result of a polluted air mass being transported to the mountain top during the daytime; this is associated with the buildup of the planetary boundary layer. During the night, on the other hand, the observed air mass is isolated from pollution sources.

Very high O₃ concentrations, with an hourly maximum of 160 ppbv and a monthly average of 82 ppbv, were observed at the top of Mt. Tai in June 2006. There are not many reports in the literature of similarly high hourly concentrations of O₃ detected by continuous observations in China, even if sites influenced by urban plumes are included. T. Wang et al. (2006) presented data for a mountainous area (Changping) 50 km north of Beijing, for June/July 2005 with a monthly average of about 60 ppbv and an hourly maximum record of 285 ppbv. Wang et al. (2008a) reported an hourly maximum of about 170 ppbv with monthly averages of about 67 ppbv at Miyun, a rural site ~ 80 km northeast of Beijing, for June, July, and August 2006. The monthly maximum

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations occurred in June, the same as at Mt. Tai. The monthly average O₃ mixing ratio at Mt. Tai is therefore even higher than in the outskirts of Beijing, and the maximum concentration is almost the same as that at Miyun.

The monthly mean mixing ratio of CO at Mt. Tai, 560 ppbv, is about the same as the 600 ppbv and 600–700 ppbv levels reported at Miyun and Changping, respectively, but the maximum level of 1950 ppbv at Mt. Tai is much higher than that in the outskirts of Beijing (~ 1500 ppbv). In contrast, the mixing ratios of NO_y at Mt. Tai, with average and maximum values of 6.7 and 21 ppbv, were much lower than those at Changping, i.e. 15 and 50 ppbv, respectively (T. Wang et al., 2006). The ΔCO/ΔNO_y ratio at Mt. Tai, around 50 ppbv/ppbv, shown as the slope of Fig. 4a, is 1.4 times that at Lin'an (30–40, Wang et al., 2002, 2004), about twice that at Changping (20–30, T. Wang et al., 2006) and in a plume from Mexico City (20.6, Kleinman et al., 2008), and about five times that in US vehicular emissions (~10, Parrish et al., 2002). The large ΔCO/ΔNO_y ratio at Mt. Tai could be attributed to open biomass burning. However, the ratio was similar for the latter half of the campaign, where the influence from biomass burning was minimal. Figure 4d shows that the high ΔCO/ΔNO_y ratio is well reproduced by the CMAQ model simulation (Yamaji et al., 2010). Figure 4a and d also indicate that high O₃ concentrations are associated with relatively high CO and NO_y concentrations.

The median NO_x/NO_y ratio of 0.15, and the 15 and 85 percentile values of 0.07 and 0.27 at Mt. Tai, are much lower than the values at Miyun, predicted by a model to be 0.4–0.7 (Wang et al., 2008a). Our range is even lower than that of the NO_x/NO_y ratio reported by Kleinman et al. (2008) over the Mexico City plateau of between 0.11 and 0.83. From these findings, it was concluded that air at Mt. Tai is affected less by emissions of NO_x but more by emissions of CO, and is photochemically more aged, than is the case for air in studied regions surrounding the megacities of Beijing and Mexico City. It should be noted that CEC is not simply a source region; photochemical aging has great importance there.

The mean concentrations of BC (PM₁), EC (PM₁, NIOSH temperature protocol, 2–20 June 2006), and OC (the same as EC) at Mt. Tai were 3.4, 2.6, and 9.2 μgCm⁻³,

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



respectively. At Shangdianzi, the average BC mass concentration from April 2003 to January 2005 was $2.1 \mu\text{gCm}^{-3}$ (Yan et al., 2008). The mass concentrations of EC and $\text{PM}_{2.1}$ OC were found to be 1.3 and $5.1 \mu\text{gCm}^{-3}$ in summer 2004 (Yan et al., 2012). At Xianghe, located southeast of Beijing, Yang et al. (2009) reported that EC and OC mass concentrations were ~ 5 and $\sim 15 \mu\text{gCm}^{-3}$ on average in March 2005. Lin et al. (2009) and Pathak et al. (2011) reported that the average concentrations of EC and OC were 4.5 and $10 \mu\text{gCm}^{-3}$ in Beijing in summer (June to September) 2006 and 4.9 and $8.2 \mu\text{gCm}^{-3}$ at Changping in June/July 2005, which are comparable to those at Mt. Tai. However, it should be noted that time series plots of BC and OC at Mt. Tai show episodically very high-concentration events, as seen in Fig. 3. Two prominent events during the periods 6–7 and 12–13 June gave peak concentrations of 29 (22) and $77 \mu\text{gCm}^{-3}$ for BC (EC) and OC, respectively; these can be ascribed to the plumes of agricultural waste burning, as will be described in Sect. 6.2.

The mean concentrations of $\text{PM}_{2.5}$ and total suspended particles (TSP) during the campaign period were 123 and $135 \mu\text{g m}^{-3}$, respectively. These values are even higher than the average values of 78 and $115 \mu\text{g m}^{-3}$ observed in downtown Beijing in summer 2007 (Deng et al., 2011). The mass concentrations of sulfate and nitrate (both for $\text{PM}_{2.5}$) during the campaign period were 21.0 and $8.5 \mu\text{g m}^{-3}$ (Deng et al., 2011), which were similar to 22.5 and $9.7 \mu\text{g m}^{-3}$ at Changping in June/July 2005 (Pathak et al., 2011). In conclusion, the air quality at the top of Mt. Tai is comparable to that in the outflow region or even the central urban area of Beijing, and such high concentrations would prevail over the central area of CEC.

The high pollutant concentrations recorded during this campaign are evident in the year-round observational records of O_3 and BC at Mt. Tai (Fig. 5). The O_3 concentration peaks in June at Mt. Tai in 2004, 2005, and 2006 are all steep. The peak timings are generally later than those at Mt. Huang and Mt. Hua, mentioned earlier (see locations in Fig. 1), and that at Mondy in East Siberia, Russia (51.66°N , 101.00°E , 2000 m a.s.l.), an EANET (acid deposition monitoring network in East Asia) background remote site. The BC concentrations at Mt. Tai and Mt. Huang were high in winter, spring, and

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



autumn, and low in summer (Fig. 5b). The BC concentration levels were higher at Mt. Tai than at Mt. Huang throughout the year. The BC level at Mt. Tai in June 2006 was one of the highest monthly averages.

Figure 6 shows the average diurnal variations of selected species. Ozone showed a broad peak in the late afternoon (14:00–19:00 LT, LT = UTC + 8 h) as high as 92–94 ppbv. NO_y and CO had earlier peaks in the afternoon, coinciding with the transport of pollutants from lower altitudes with respect to the buildup of the boundary layer. BC showed high concentrations in the early morning (04:00–06:00 LT) as well as a peak in the afternoon. OC showed relatively flat diurnal variations in the early half-period (until 20 June, when the NIOSH temperature protocol was used with PM_{10} sampling), but showed regular diurnal variations with early afternoon maxima in the latter half-period, when an IMPROVE-like protocol was used with $\text{PM}_{2.5}$ sampling. CO_2 showed a regular daytime decrease, mainly in response to the source/sink provided by vegetation. Figure 7 shows three photographs looking westward from the observatory at the summit of Mt. Tai. In the upper panel, the outlines of distant mountains are clearly visible, when aerosols are not abundantly present. Once regional pollution occurs, as shown in the middle panel, the top boundary of the polluted layer is clearly visible below the mountain altitude level in the morning. The pollution layer develops in the later hours, as shown in the bottom panel, and then the mountain top is incorporated in the polluted layer, at around noontime.

The high concentrations of BC must have important implications for the regional climate in CEC. Here the RF is roughly estimated. First we simply calculate its column concentration (5.2 mg m^{-2}) by multiplying the monthly mean concentration ($3.4 \mu\text{g C m}^{-3}$) observed at the mountain top by the altitude (i.e. 1534 m), assuming a box-shaped vertical profile. Then by assuming that 85% of BC is anthropogenic (Bond et al., 2011) and multiplying by a sensitivity factor (1.25 W mg^{-1} , IPCC, 2007), we obtain an RF as high as 55 W m^{-2} . This value far exceeds the RF from CO_2 estimated to be $\sim 1.6 \text{ W m}^{-2}$. The regional-scale BC air pollution therefore greatly affects the

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



radiative budget. More detailed studies using radiative transfer models and chemistry–climate coupled models are needed

6.2 Importance of agricultural waste burning and regional transport

Figure 8 shows backward trajectories starting from 1500 m and 750 m at Mt. Tai for six selected times: 12:00 LT on 6, 9, 12, 14, 19, and 27 June 2006. The kinematic backward trajectories were calculated using the CGER-METEX program (<http://db.cger.nies.go.jp/metex/>), using the National Centers for Environmental Prediction meteorological reanalysis field as input. They are shown together with geographical distributions of hotspots detected by sensors on board NOAA and FY-1D satellites during the preceding five days. The color-coded map shows the winter-wheat-harvesting period. These figures are based on those produced by SEPA (State Environmental Protection Administration, currently Ministry of the Environment) of the Chinese government. As seen in Fig. 8, the winter-wheat-harvesting region shifts from the southern part in April/May to northern and western areas in June/July. June is the most active burning season south of Mt. Tai, after the harvesting of winter wheat. Figure 3, particularly for BC and OC, clearly shows more polluted days in the first half and cleaner days in the latter half of the campaign period. Two prominent pollution episodes are seen during the periods 6–7 June and 12–13 June, when the backward trajectory passed over the burnt areas. These characteristics strongly suggest the impact of agricultural waste burning on air quality at Mt. Tai. Suthawaree et al. (2010), Yamaji et al. (2010), and Fu et al. (2012), on this issue, similarly report the impact of burning using backward trajectories that passed over hotspots. On 8–10 June, the wind was from the north and thus the influence was small, although hotspots were continuously found south of Mt. Tai. Pan et al. (2012) combined the geographical distribution of the hotspots with backward Weather Research and Forecasting (WRF)-FLEXPART model simulations and found that the influential source area was 33–34° N and 116–118° E, about 300 km south of the observatory. They also suggested that the transport time of the air mass from the source region ranged from 10 to 50 h. The observed evidence of burning, in addition

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to high CO and BC concentrations was elevated concentrations of levoglucosan and other anhydrosugars (Fu et al., 2008), K^+ , and oxalic acid in $PM_{2.5}$ (Xu et al., 2009; Deng et al., 2011), CH_3Cl (Suthawaree et al., 2010), CH_3CN (Inomata et al., 2010), and lowered stable carbon isotope ratios ($\delta^{13}C$) of organic aerosol particles (Fu et al., 2012).

The impact of OCRB on O_3 , CO, BC, and OC concentrations over CEC during MTX2006 was evaluated using a regional chemical transport model, CMAQ, employing estimated daily emissions from OCRB, based on hotspot data obtained by MODIS (Yamaji et al., 2010). The impact of OCRB on regional emissions during the period 6–9 June amounted to 31–44% (NO_x), 47–61% (CO), 50–63% (EC) and 56–69% (primary organic aerosols) of the total sources in CEC. After 17 June, the impact of OCRB was almost zero. Figure 6 in Yamaji et al. (2010) demonstrates that the daily emissions from OCRB conveyed by southerly winds were an essential factor in reproducing atmospheric concentrations of pollutants during MTX2006. These emissions have a large impact not only on primary pollutants but also on secondary pollutants, such as O_3 , in the first half of June. The run with OCRB emissions (O6DS) clearly captured the observed variations in daily average O_3 concentrations, with a correlation coefficient R of 0.61 between the model and the observations, which is much better than that obtained without burning (NOCRB; $R = 0.34$). The monthly O_3 concentration simulated using O6DS (80.8 ppbv) was much closer than that obtained using NOCRB (73.9 ppbv) to the observed monthly O_3 concentration. The average impacts of OCRB emissions contributed 6% of O_3 , 20% of CO, 43% of EC and 53% of OC concentrations over CEC for the whole month of June; for the episodic period 6–9 June, the impacts increased to 12% for O_3 , 35% for CO, 56% for BC daily emissions, and 80% for OC over CEC. The real cause of such elevated O_3 concentrations in June at Mt. Tai, which had been a mystery, was therefore identified as biomass burning, by conducting a comprehensive field campaign covering the precursors of O_3 and related species.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

6.3 Attribution of nearground O₃ over Mt. Tai to the transport from other source regions

A regional chemical transport model, NAQPMS, with an online tracer tagging module, was used to study the source areas of near-ground O₃ at Mt. Tai during the MTX2006 period (Li et al., 2008a). During this time period, CEC was covered with air masses with high O₃ concentrations (65–85 ppbv) in the model simulation Mt. Tai was at the center of a high O₃ region with the prevailing wind from the south. The source regions were divided into CEC and outer regions (see Fig. 1 of Li et al., 2008a). The former consists of Local, South, East and North Shandong (SSD, ESD and NSD), Hebei/Beijing/Tianjin (HBT), Shanxi (SX), Henan (HN), Anhui (AH), Jiangsu (JS) and East China Sea (CHNSEA). The latter consists of Siberia (SIB), Northwest China (NWCHN), Southwest China (SWCHN), South China (SCHN), East China (ECHN), Northeast China (NECHN), the western Pacific rim (PAC), Korea (KOR), Japan (JPN), and Southeast Asia (SEASA). Source attributions from the regional scale down to sub-province levels were therefore made.

In general, regional photochemical production of O₃ within CEC was the most important mechanism, showing a contribution of 51.4 ppbv (60.2%) to a monthly mean O₃ mixing ratio of 85.4 ppbv. Specifically, the largest fractions of O₃ were formed in SSD, AH, and JS (32.4 ppbv or 37.9%), influenced by high emission rates (including those from biomass burning) and favorable meteorological conditions in these areas. They concluded that local photochemical production (9.8 ppbv or 11.5%) is a minor contribution, suggesting that emissions from local cities (e.g. Tai'an) could be of minor importance. The calculated in-situ O₃ production rate was in good agreement with the results of Kanaya et al. (2009), which will be described in Sect 6.4. Li et al. (2008b) showed that the same NAQPMS model was able to reproduce the observed diurnal variations in O₃ at Mt. Tai, and studied the roles of chemical production and transport. They suggested that the photochemistry in the surrounding region is more dominant than transport, giving diurnal variations in O₃ mixing ratios, with a maximum in

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the afternoon. Li et al. (2011), using a similar model framework, studied the impact of aerosols on O₃ photochemistry through the reduction of ultraviolet (UV) radiation over this period, with a strong influence from biomass burning. They estimated that the $J(\text{O}^1\text{D})$ values and OH concentrations are reduced by 37% and 40%, respectively, in the middle troposphere over CEC, leading to lower O₃ concentrations (by a maximum of 9% in highly polluted regions).

6.4 Photochemical O₃ production processes over CEC in June 2006

An observation-based boxmodel approach was undertaken to estimate concentrations of OH, HO₂, and RO₂ radicals and the net photochemical production rate of O₃, i.e. $F - D(\text{O}_3)$, the difference between the rates of gross formation [$F(\text{O}_3)$] and loss [$D(\text{O}_3)$] (Kanaya et al., 2009). The model calculation was constrained by measurements of O₃, H₂O, CO, NO, NO₂, hydrocarbons, HCHO, and CH₃CHO concentrations, and temperature and J values. Figure 3 of Kanaya et al. (2009) shows the calculated $F(\text{O}_3)$, $D(\text{O}_3)$ and $F - D(\text{O}_3)$ separately for high O₃-days with hourly values exceeding 100 ppbv and low O₃-days, and they argued that the difference was not large, suggesting that O₃ transportation is more important than in-situ photochemistry for high O₃ concentrations. The net production rate of O₃ was estimated to be 6.4 ppbv h⁻¹ as a 6-h average (09:00–15:00 LT), in rough agreement with the insitu production rate (~ 5 ppbv h⁻¹) estimated in a three-dimensional regionalscale modeling study (Li et al., 2008b). These values fall within the range of the rates estimated for the boundary layer over major cities in the United States, 3.5–11.3 ppbv h⁻¹ (Kleinman et al., 2002).

The estimated 6-h averages of $F - D(\text{O}_3)$ suggest that 58 ± 37 ppbv of O₃ are produced in 1 d. A sensitivity model run was made incorporating heterogeneous loss of HO₂ on aerosol particle surfaces, which could impede photochemical O₃ production. Pure water extracts of aerosol samples collected on quartz filters were used to regenerate aerosol particles in an aerosol flow tube in the laboratory to study the loss kinetics of HO₂ radicals as a result of heterogeneous reactions on the aerosol particles

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Taketani et al., 2012). The uptake coefficients (γ values), determined for real atmospheric aerosol particle samples (> 10 samples) for the first time had an average value of 0.25, larger than the values determined for particles with major single components (e.g. ammonium sulfate), suggesting the possibility that minor components of aerosols play important roles. When this heterogeneous loss was included, the daily integral of $F - D(O_3)$ fell from 58 to 39 ppbv. This quantity was still larger than the observed daytime increases in the O_3 concentration observed at the mountain top, 23 ppbv on average (Fig. 6) suggesting that the daytime buildup could be explained by in-situ photochemistry. Another sensitivity model runs with altered NO_x and hydrocarbon concentrations suggested that O_3 production occurred normally under NO_x -limited conditions. Figure 4b shows that NO_x is almost depleted by photochemical aging (in comparison with Fig. 4a, using NO_y as the x-axis), and thus the abundance of NMHCs, correlated with CO and used for the y-axis in Fig. 4b, becomes relatively high, resulting in NO_x -limited conditions. This is consistent with the box-model analysis in this study. An exception was 10 June, when fresh pollution from the north affected the site and the O_3 production was VOC-limited. This analysis suggests that the O_3 pollution over the CEC region will become even worse when NO_x emissions increase in the future. Figure 4e shows the modeled dependence of O_3 on NO_x and CO. Although the modeled dependence is similar to the observations (Fig. 4b), the model tends to underestimate CO and therefore the CO/ NO_x ratio. Because the O_3 concentration is positively dependent on the CO concentration (see color codes in Fig. 4e, b), an increase in CO in the model to reproduce the observed levels will increase the O_3 concentration.

Figure 4c shows that hourly O_3 concentrations increase with NO_2 ($[NO_2] = [NO_y] - [NO] - [NO_2]$) up to about 10 ppbv, but saturation occurs with further increases in NO_2 . This feature is consistent with the above-mentioned analysis using a box model which suggested general NO_x -limited conditions. The slope $\Delta[O_3]/\Delta[NO_2]$ was 5.8 ± 0.5 ppbv ppbv $^{-1}$ when the NO_2 mixing ratio was below 10 ppbv, implying efficient O_3 production from NO_x . This is quite similar to the value of 3–6 ppbv ppbv $^{-1}$ suggested for selected episodes studied at Changping near Beijing

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(T. Wang et al., 2006), and to that of $6.2 \text{ ppbv ppbv}^{-1}$ found for the Pico de Tres Padres (PTP) site near Mexico City (Wood et al., 2009). The observed slope is lower than the modeled slope (Fig. 4f, slope = $8.7 \pm 0.5 \text{ ppbv ppbv}^{-1}$). This suggests that the O_3 production efficiency per unit NO_x molecule oxidation is slightly lower than that predicted by the model. Although not shown, high O_3 concentrations are sometimes associated with aged air masses ($\ln([\text{NO}_y]/[\text{NO}_x]) > 2$ or $[\text{NO}_x]/[\text{NO}_y] < 0.14$).

6.5 NO_2 measurements by MAX-DOAS: comparisons with satellite and in-situ observations

During the campaign period, MAX-DOAS measurements of vertical profiles of NO_2 were performed from the top of Mt. Tai as well as from the foothills in the city of Tai'an (Irie et al., 2008). The mean difference between the NO_2 volume mixing ratios measured by MAX-DOAS instruments at Tai'an and Mt. Tai and from insitu data based on chemiluminescence detection coupled with a photolytic converter were as small as $-0.01 \pm 0.60 \text{ ppbv}$ and $-0.29 \pm 0.65 \text{ ppbv}$, respectively (see Fig. 5 of Irie et al., 2008). The mean NO_2 volume mixing ratio for the 0–1 km altitude layer above the surface derived from the MAX-DOAS measurements at Tai'an was about 4 ppbv.

The MAX-DOAS measurements showed complete diurnal variations in the tropospheric NO_2 VCD in the daytime. Typically, the value was highest at $\sim 15 \times 10^{15} \text{ molecules cm}^{-2}$ in the early morning (6:00–9:00 LT) and dropped to a minimum of $\sim 7 \times 10^{15} \text{ molecules cm}^{-2}$ at 13:00–15:00 LT. This indicates that the OMI measurements (at about 13:40 LT) were made when the tropospheric NO_2 VCD was lowest in its typical diurnal cycle, and that the systematic difference between tropospheric NO_2 VCDs derived from OMI and those observed by SCIAMACHY and GOME2 in the morning could be explained by this strong diurnal variation. Critical comparisons between tropospheric NO_2 VCDs derived from MAX-DOAS and OMI were made for the first time in CEC, with different spatial coincidence criteria. It was revealed that

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the use of a stricter coincident criterion of 0.1° resulted in better agreement within $(+1.6 \pm 0.6) \times 10^{15}$ molecules cm^{-2} ($+20 \pm 8\%$) (OMI minus MAX-DOAS).

6.6 Volatile organic compound and oxygenated volatile organic compound (VOC/OVOC) compositions and concentrations

5 The relative concentration ratios of individual NMHCs (10 alkanes, six aromatic compounds, seven alkenes, and acetylene), normalized by ethane, at Mt. Tai, derived from daily canister sampling followed by gas chromatography-flame ionization detector analysis, were compared with those observed in Beijing city center on 20 November 2005 (see Fig. 7 of Suthawaree et al., 2010). Here, the average concentration of ethane at
10 Mt. Tai was 2.4 ppbv, comparable to those in other areas of China (Barletta et al., 2005) reflecting its long lifetime and the absence of local sources. As expected, we found that the ratios of major alkanes and alkenes to ethane are much lower for observations at Mt. Tai than they are in Beijing, reflecting the fact that the air mass at Mt. Tai is isolated from sources and is more aged. The measured benzene/toluene ratio at Mt. Tai was
15 3.2 ± 1.0 , much higher than those (0.6) for other Chinese cities (Barletta et al., 2005) and other parts of the world where vehicular emissions make significant contributions. A high ratio to ethane was also found for acetylene probably reflecting the contribution of biomass/biofuel combustion in the surrounding areas. The analysis of halo-
20 genated compounds (C_2Cl_4 , CH_3Cl , CH_3Br) by gas chromatography-mass spectrometry showed low concentrations of C_2Cl_4 , 8 pptv on average, suggesting insignificant impacts of industrial emissions, and relatively high concentrations of CH_3Cl , 990 pptv on average, suggesting the influence of biomass burning. The isoprene concentrations were 209 pptv on average ($n = 25$) for daytime samples (collected at $\sim 14:40$ LT) sug-
25 gesting that biogenic activity was moderate and less active than in other mountainous areas in China (e.g. 480 pptv at Mt. Jianfeng; Tang et al., 2007).

Real-time measurements of NMHCs and OVOCs were made using a PTR-MS (proton-transfer reaction mass spectrometry) instrument during the period 12–30 June 2006, the latter half of the campaign period (Inomata et al., 2010). One of the major

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



findings is that the observed mixing ratios of acetaldehyde were approximately 1.6 times higher than those of formaldehyde during the whole observation period. This is in contrast to the finding that the mixing ratios of formaldehyde are generally higher than those of acetaldehyde in Beijing (Shao et al., 2009) and other urban, suburban and rural sites in the world, suggesting the possibility of primary emission of acetaldehyde. The concentrations of methanol were high, ~ 10 ppbv on average, compared with aldehydes and acetone. This result is similar to that observed in Beijing (Liu et al., 2009) and near biomass burning (Karl et al., 2007). The HCHO concentrations derived from PTR-MS were in good agreement with those from MAX-DOAS observations (Inomata et al., 2008). Figure 9 shows that the PTR-MS signal at m/z 97 showed a strong positive correlation ($R = 0.95$) with levoglucosan concentrations in the TSP samples. The correlation coefficient was even larger than that for CH_3CN and levoglucosan ($R = 0.89$). Karl et al. (2007) studied tropical biomass burning and assigned the peaks at m/z 97 to substituted furans and furfurals. This may be valid for our study. However, our (m/z 97)/(m/z 42) ratio of $1.6 \text{ ppbv ppbv}^{-1}$ was slightly lower than the value of $2.4\text{--}3.2 \text{ ppbv ppbv}^{-1}$ reported by Karl et al. (2007); this suggests that either the furans were removed by fast OH reactions during travel for 10–50 h and the original emission ratio was much higher, or that the species we detected at m/z 97 was different and had more chemical stability, resulting in the high correlation coefficient. Additionally, m/z 85 and m/z 87 also showed strong correlations ($R > 0.90$) with levoglucosan; the (m/z 85)/(m/z 42) and (m/z 87)/(m/z 42) ratios were 1.2 and $1.6 \text{ ppbv ppbv}^{-1}$, respectively, assuming a typical ion–molecule reaction rate constant of $2 \times 10^{-9} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. These analyses were made possible by the comprehensive coverage of gas and aerosol species during this MTX2006 field campaign. More studies are needed in the vicinity of biomass burning sources for better characterization.

6.7 BC measurement comparisons and $\Delta\text{BC}/\Delta\text{CO}$ ratio analysis

In addition to the year-round observations of BC using a Multi-Angle Absorption Photometer (MAAP, model 5012, Thermo Scientific, Waltham, MA, USA) instrument at the

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



summit of Mt. Tai, we installed several more instruments, namely a particle soot absorption photometer (PSAP; Radiance Research, Shoreline, WA, USA) and an ECOC semi-continuous analyzer (Sunset Laboratory, Tigard, OR, USA), operated either with IMPROVE-like or NIOSH temperature programs, for the purpose of instrument inter-comparison during the campaign period (Kanaya et al., 2008). These instruments were operated either with PM₁ or PM_{2.5} cyclones, depending on the time period. An aethalometer (AE-21, Magee Scientific, Berkeley, CA, USA) was also operated for several days during the first part of the campaign.

The regression analysis for each pair showed that the correlations were strong and the slopes ranged from 1.03 to 1.54 for the first period (before 20 June) with PM₁ sampling, and from 1.07 to 1.46 for the second period (after 20 June), using the IMPROVE-like temperature program for EC analysis with PM_{2.5} sampling (see Fig. 8 of Kanaya et al., 2008). Considering that disagreement up to a factor of 4 (e.g. Jeong et al., 2004) was reported in the United States among similar instruments, the spread at Mt. Tai was thought to be relatively narrow. We therefore obtained BC concentration data suitable for constraining BC emission rates from China, whose uncertainty was thought to be large (e.g. factors of 4–5; Bond et al., 2004; Streets et al., 2003).

We also found that enhanced MAAP_BC/EC ratios (e.g. higher than 2) occurred only when low NO_x/NO_y ratios were recorded, implying that aging is related to large MAAP_BC/EC ratios. Given that the tendency to underestimate EC is not straightforwardly explained in terms of aging, it is reasonable to conclude that the BC particles are coated by transparent materials after aging and that the MAAP results based on absorbance measurements gave overestimates possibly as a result of a lens effect induced by the coating.

Pan et al. (2012) studied $\Delta\text{BC}/\Delta\text{CO}$ ratios in detail, using CO as a conservative tracer, during the biomass-burning events and found that the ratio had a clear tendency to decrease with increasing transport time of the air mass from the biomass-burning source area. Here, the transport time was estimated using the WRF-FLEXPART model.

The estimated lifetime of BC was ca. 4.1 d, near the lower limit of the range suggested by past studies.

6.8 PM and inorganic ion/element concentrations

The daily mass concentrations of TSP and PM_{2.5} and their ionic and element compositions, were measured during MTX2006 and compared with the data obtained in spring (from March to early May) 2006 and 2007 at the same site (Deng et al., 2011). The concentrations of TSP and PM_{2.5}, determined using the filter-weighing method, ranged from 49 to 277 $\mu\text{g m}^{-3}$ and from 48 to 236 $\mu\text{g m}^{-3}$ respectively during MTX2006. The average PM_{2.5}/TSP ratio was 0.37 in spring and 0.91 in summer in 2006, indicating that fine particles dominated in summer whereas coarse particles dominated in spring. This ratio in summer is much higher than those in other cities in China (0.27–0.68), indicating that the air mass at Mt. Tai is chemically very aged consistent with other evidence discussed earlier.

Water-soluble ions contributed 10.8% of TSP and 24.0% of PM_{2.5} by mass in spring, compared with 40.9% and 41.1% during MTX2006. Sulfate, nitrate, and ammonium ions were the major water-soluble species in PM_{2.5}, accounting for 61.5% and 72.7% of the total measured ions in spring and during MTX2006, respectively. Also, higher concentrations of K⁺ a tracer of biomassburning sources, were observed during MTX2006, accounting for 8.3% of the total ions in TSP with high correlations with BC ($R = 0.90$) and oxalic acid ($R = 0.87$).

Xu et al. (2009) reported size-segregated water-soluble ions and metals in aerosol particles. Sulfate, ammonium, and K ions, and Pb, Zn, and Ti peaked in the accumulation mode (0.43–1.1 μm) Ca, Mg, Fe, Al, Ba, and Mn had peaks in the coarse mode (4.7–5.8 μm). In contrast, nitrate, Na, and Cl ions, and Co, Ni, Mo, and Cu showed bimodal distributions (0.43–0.65 and 4.7–5.8 μm).

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6.9 Organic aerosols: photochemical formation and molecular composition

Figure 10 shows that the $\Delta[\text{OC}]/\Delta[\text{CO}]$ ratio during the latter period of the campaign (from 16 June 2006) with negligible influence from biomass burning increased with increasing air-mass age, indicated by the $\ln([\text{NO}_y]/[\text{NO}_x])$ value. Here CO is considered to be a relatively inert chemical species and its background concentration ($[\text{CO}]_{\text{bg}}$) was assumed to be 90 ppbv. The increase in the $\Delta[\text{OC}]/\Delta[\text{CO}]$ ratio with photochemical aging suggests that carbon mass flux from gas-to-particle conversion is present. Robinson et al. (2007) assumed that SVOC and IVOC (semi-volatile and intermediate volatility organic compound) vapors are oxidized via a gas-phase OH reaction with a rate constant of $4 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ to yield next generation compounds that are 10 times less volatile, and calculated larger partitioning from gas-to-particle phases. The assumed reaction rate is about four times faster than the oxidation of NO_x to NO_y . To cover eight generations, corresponding to the eight orders of magnitude range in volatility considered in their model study, a $\ln([\text{NO}_y]/[\text{NO}_x])$ range between 0 and 2 needs to be studied. The study at Mt. Tai actually covered this range and the data are suitable for the quantitative investigation of possible increases in OC with photochemical aging.

Fu et al. (2008) summarized the organic molecular compositions (except dicarboxylic acids) of TSP samples collected on quartz filters using a high-volume air sampler and analyzed their temporal variations. They found that *n*-alkanes, fatty acids, fatty alcohols, sugars, glycerol polyacids, and phthalate esters are the major species, and that lignin and resin products, sterols, aromatic acids, hopanes, and polycyclic aromatic hydrocarbons (PAHs) are minor species. They showed the importance of plant emissions of waxes and soil resuspension as well as biomass burning, as source processes. From the same TSP samples, Kawamura et al. (2012) reported that water-soluble OC (WSOC)/total carbon (TC) ratio was 0.41 ± 0.09 in average, which is larger than 0.35 or less observed for $\text{PM}_{2.5}$ samples at Changping or other Chinese cities, Shanghai, Lanzhou, and Guangzhou (Pathak et al., 2011). Kawamura et al. (2012) also

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

reported that very high concentrations of saturated and unsaturated diacids (C_2 – C_{11} up to $61 \mu\text{g m}^{-3}$) were detected among WSOC (ranged from 1 to $37 \mu\text{g m}^{-3}$), with a predominance of oxalic (C_2) acid followed by malonic (C_3) and succinic (C_4) acids as well as C_2 – C_9 ω -oxocarboxylic acids, pyruvic acid and α -dicarbonyls (glyoxal and methylglyoxal). The highest concentrations of total diacids ($> 6 \mu\text{g m}^{-3}$) found at Mt. Tai were several times higher than those reported at ground levels in Chinese megacities. The temporal variations in diacids were interpreted to be caused by a combination of direct emissions from field burning of agricultural wastes and secondary photochemical production via oxidation of volatile and semi-volatile organic precursors emitted from field burning of wheat straw during transport to the summit of the mountain. Fu et al. (2010) quantified the oxidation products from isoprene (e.g. 2-methylerythritol), α/β -pinene (e.g. 3-hydroxyglutaric acid), and β -caryophyllene (e.g. β -caryophyllinic acid) in the filter samples and suggested the importance of isoprene-derived secondary organic carbons. Fu et al. (2012) studied the diurnal variations in 12 organic compound classes consisting of more than 130 species and their stable isotopic compositions. During the biomass-burning period in early June, the diurnal trends of most of the primary and secondary organic aerosol tracers were characterized by concentration peaks observed at midnight or in the early morning, whereas in late June (without the influence of intense biomass burning) most of the organic species peaked in the late afternoon. A strong anti-correlation was found between levoglucosan and $\delta^{13}\text{C}$ values, again suggesting the importance of crop residue burning. Altogether, 165 molecularly identified organic compounds including diacids, oxo/ketocarboxylic acids, dicarbonyls, n -alkanes, fatty acids, fatty alcohols, sugars, phthalates, polyols polyacids, aromatic acids, lignin and resin products, sterols, hopanes, PAHs, and biogenic secondary organic aerosol tracers (Fu et al., 2012; Kawamura et al., 2012) accounted for $14 \pm 6\%$ of the total OC mass in the TSP. The OC mass concentrations in the TSP were 2.6 ± 0.8 times larger than those in PM_1 (before 20 June, using the NIOSH protocol, $n = 46$) and 2.4 ± 0.7 times larger than those in $\text{PM}_{2.5}$ (after 20 June, IMPROVE-like protocol, $n = 22$) analyzed using a semi-continuous ECOC analyzer (Fig. 11). This suggests the presence of coarse

organic aerosol particles, and the presence of molecularly unidentified organics, even in PM₁.

Wang et al. (2009) investigated the size distributions of *n*-alkanes, PAHs, and hopanes collected using an Andersen eight-stage air sampler during the period 22–29 June 2006. The size distributions were compared with those sampled in winter at the summit of Mt. Tai and those sampled in the marine boundary layer and urban atmosphere. During MTX2006, bimodal distributions with peak sizes of 0.7–1.1 μm and 4.7–5.8 μm were recorded for both plant-wax and anthropogenic *n*-alkanes. All the PAHs had maximums in the range 0.7–1.1 μm. Hopanes showed bimodal distributions with peaks in the ranges 0.7–1.1 μm and > 3.3 μm.

7 Summary

MTX2006 was the first field campaign performed at a site geographically representing regional air pollution over the NCP. Comprehensive measurements and model simulations of O₃ and its precursors, and aerosol chemical species were performed. The period of the study (June) was chosen to target high O₃ concentrations (monthly average 82 ppbv). Together with high O₃ levels, we found very high concentrations of organic aerosols, probably as a result of extensive photochemistry. We found that OCRB after the harvesting of winter wheat was one of the most important factors leading to these high concentrations, and this was supported by coincident observations of peak concentrations of biomassburning tracers such as levoglucosan, K⁺, CH₃Cl, and CH₃CN, and by low δ¹³C values of organic aerosols. A large fraction of high O₃ was photochemically produced in CEC, generally under NO_x-limited conditions, with some slowing effects as a result of O₃-aerosol interactions through *J* value reduction and heterogeneous chemistry. The validation of satellite-based observations of tropospheric NO₂ VCDs was performed using MAX-DOAS observations. Instrument intercomparisons were made for BC measurements, which provided better constraints for BC emission rates from this region. Strong positive RF by BC was implied. The BC

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lifetime in the atmosphere was estimated using the transport time from biomassburning source areas. The inorganic and organic compositions of aerosol particles were studied in detail. Molecularly identified organic compounds accounted for 14 % of total OC in the TSP.

- 5 *Acknowledgements.* Support from local staff at the observatory is gratefully acknowledged. This work was supported by the Global Environment Research Fund and by the Environmental Research and Technology Development Fund (S-7, C-081, B-051) of the Ministry of the Environment, Japan.

References

- 10 Akimoto, H.: Global air quality and pollution, *Science*, 302, 1716–1719, 2003.
Barletta, B., Meinardi, S., Rowland, F. S., Chan, C. Y., Wang, X., Zou, S., Chan, L. Y., and Blake, D. R.: Volatile organic compounds in 43 Chinese cities, *Atmos. Environ.*, 32, 5979–5990, 2005.
- 15 Beirle, S., Platt, U., Wenig, M., and Wagner, T.: Highly resolved global distribution of tropospheric NO₂ using GOME narrow swath mode data, *Atmos. Chem. Phys.*, 4, 1913–1924, doi:10.5194/acp-4-1913-2004, 2004.
- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109, D14203, doi:10.1029/2003JD003697, 2004.
- 20 Bond, T. C., Zarzycki, C., Flanner, M. G., and Koch, D. M.: Quantifying immediate radiative forcing by black carbon and organic matter with the Specific Forcing Pulse, *Atmos. Chem. Phys.*, 11, 1505–1525, doi:10.5194/acp-11-1505-2011, 2011.
- Byun, D. W. and Ching, J. K. S. (eds.): Science algorithms of the EPA Models-3 community multiscale air quality (CMAQ) modeling system, EPA/600/R-99/030, USEPA NERL, Research Triangle Park, USA, 1999.
- 25 Chin, M., Chu, A., Levy, R., Remer, L., Kaufman, Y., Holben, B., Eck, T., Ginoux, P., and Gao, Q. X.: Aerosol distribution in the Northern Hemisphere during ACE-Asia: results from global model, satellite observations, and sun photometer measurements, *J. Geophys. Res.*, 109, D23S90, doi:10.1029/2004JD004829, 2004.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Deng, C., Zhuang, G., Huang, K., Li, J., Zhang, R., Wang, Q., Liu, T., Sun, Y., Guo, Z., Fu, J. S., and Wang, Z.: Chemical characterization of aerosols at the summit of Mountain Tai in Central East China, *Atmos. Chem. Phys.*, 11, 7319–7332, doi:10.5194/acp-11-7319-2011, 2011.

Fu, P. Q., Kawamura, K., Okuzawa, K., Aggarwal, S. G., Wang, G. H., Kanaya, Y., and Wang, Z. F.: Organic molecular compositions and temporal variations of summertime mountain aerosols over Mt. Tai, North China Plain, *J. Geophys. Res.-Atmos.*, 113, D19107, doi:10.1029/2008JD009900, 2008.

Fu, P. Q., Kawamura, K., Kanaya, Y., and Wang, Z. F.: Contributions of biogenic volatile organic compounds to the formation of secondary organic aerosols over Mt. Tai, Central East China, *Atmos. Environ.*, 44, 4817–4826, 2010.

Fu, P. Q., Kawamura, K., Chen, J., Li, J., Sun, Y. L., Liu, Y., Tachibana, E., Aggarwal, S. G., Okuzawa, K., Tanimoto, H., Kanaya, Y., and Wang, Z. F.: Diurnal variations of organic molecular tracers and stable carbon isotopic composition in atmospheric aerosols over Mt. Tai in the North China Plain: an influence of biomass burning, *Atmos. Chem. Phys.*, 12, 8359–8375, doi:10.5194/acp-12-8359-2012, 2012.

Fuhrer, J., Skarby, L., and Ashmore, M.: Critical levels of ozone effect in Europe, *Environ. Pollut.* 97, 91–106, 1997.

Gao, J., Wang, T., Ding, A. J., and Liu, C. B.: Observational study of ozone and carbon monoxide at the summit of mount Tai (1534 m a.s.l.) in central-eastern China, *Atmos. Environ.*, 39, 4779–4791, 2005.

Ge, B. Z., Wang, Z. F., Xu, X. B., Tang, J., He, Y. J., Uno, I., and Ohara, T.: Impact of the East Asian summer monsoon on long-term variations in the acidity of summer precipitation in Central China, *Atmos. Chem. Phys.*, 11, 1671–1684, doi:10.5194/acp-11-1671-2011, 2011.

He, Y. J., Uno, I., Wang, Z. F., Pochanart, P., Li, J., and Akimoto, H.: Significant impact of the East Asia monsoon on ozone seasonal behavior in the boundary layer of Eastern China and the west Pacific region, *Atmos. Chem. Phys.*, 8, 7543–7555, doi:10.5194/acp-8-7543-2008, 2008.

Inomata, S., Tanimoto, H., Kameyama, S., Tsunogai, U., Irie, H., Kanaya, Y., and Wang, Z.: Technical Note: Determination of formaldehyde mixing ratios in air with PTR-MS: laboratory experiments and field measurements, *Atmos. Chem. Phys.*, 8, 273–284, doi:10.5194/acp-8-273-2008, 2008.

Inomata, S., Tanimoto, H., Kato, S., Suthawaree, J., Kanaya, Y., Pochanart, P., Liu, Y., and Wang, Z.: PTR-MS measurements of non-methane volatile organic compounds during an

**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- intensive field campaign at the summit of Mount Tai, China, in June 2006, *Atmos. Chem. Phys.*, 10, 7085–7099, doi:10.5194/acp-10-7085-2010, 2010.
- IPCC: Working Group I, *Climate Change 2007, The Physical Science Basis*, Cambridge University Press, 2007.
- 5 Irie, H., Kanaya, Y., Akimoto, H., Tanimoto, H., Wang, Z., Gleason, J. F., and Bucsel, E. J.: Validation of OMI tropospheric NO₂ column data using MAX-DOAS measurements deep inside the North China Plain in June 2006: Mount Tai Experiment 2006, *Atmos. Chem. Phys.*, 8, 6577–6586, doi:10.5194/acp-8-6577-2008, 2008.
- Jeong, C.-H., Hopke, P. K., Kim, E., and Lee, D.-W.: The comparison between thermal-optical transmittance elemental carbon and Aethalometer black carbon measured at multiple monitoring sites, *Atmos. Environ.*, 38, 5193–5204, 2004.
- 10 Kanaya, Y., Komazaki, Y., Pochanart, P., Liu, Y., Akimoto, H., Gao, J., Wang, T., and Wang, Z.: Mass concentrations of black carbon measured by four instruments in the middle of Central East China in June 2006, *Atmos. Chem. Phys.*, 8, 7637–7649, doi:10.5194/acp-8-7637-2008, 2008.
- 15 Kanaya, Y., Pochanart, P., Liu, Y., Li, J., Tanimoto, H., Kato, S., Suthawaree, J., Inomata, S., Taketani, F., Okuzawa, K., Kawamura, K., Akimoto, H., and Wang, Z. F.: Rates and regimes of photochemical ozone production over Central East China in June 2006: a box model analysis using comprehensive measurements of ozone precursors, *Atmos. Chem. Phys.*, 9, 7711–7723, doi:10.5194/acp-9-7711-2009, 2009.
- 20 Karl, T. G., Christian, T. J., Yokelson, R. J., Artaxo, P., Hao, W. M., and Guenther, A.: The Tropical Forest and Fire Emissions Experiment: method evaluation of volatile organic compound emissions measured by PTR-MS, FTIR, and GC from tropical biomass burning, *Atmos. Chem. Phys.*, 7, 5883–5897, doi:10.5194/acp-7-5883-2007, 2007.
- 25 Kawamura, K., Tachibana, E., Okuzawa, K., Aggarwal, S. G., Kanaya, Y., and Wang, Z. F.: High abundance of water-soluble dicarboxylic acids, ketoacids and α -dicarbonyls in the mountain aerosols over the North China Plain during wheat-burning season, *Atmos. Chem. Phys. Discuss.*, submitted, 2012.
- 30 Kleinman, L. I., Daum, P. H., Imre, D., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J., and Rudolph, J.: Ozone production rate and hydrocarbon reactivity in 5 urban areas: a cause of high ozone concentration in Houston, *Geophys. Res. Lett.*, 29, 1467, doi:10.1029/2001GL014569, 2002.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Kleinman, L. I., Springston, S. R., Daum, P. H., Lee, Y.-N., Nunnermacker, L. J., Senum, G. I., Wang, J., Weinstein-Lloyd, J., Alexander, M. L., Hubbe, J., Ortega, J., Canagaratna, M. R., and Jayne, J.: The time evolution of aerosol composition over the Mexico City plateau, *Atmos. Chem. Phys.*, 8, 1559–1575, doi:10.5194/acp-8-1559-2008, 2008.
- 5 Li, J., Wang, Z. F., Akimoto, H., Gao, C., Pochanart, P., and Wang, X.: Modeling study of the seasonal cycle of ozone in the boundary layer over East Asia, *J. Geophys. Res.*, 112, D22S25, doi:10.1029/2006JD008209, 2007.
- Li, J., Wang, Z., Akimoto, H., Yamaji, K., Takigawa, M., Pochanart, P., Liu, Y., Tanimoto, H., and Kanaya, Y.: Near-ground ozone source attributions and outflow in central eastern China during MTX2006, *Atmos. Chem. Phys.*, 8, 7335–7351, doi:10.5194/acp-8-7335-2008, 2008a.
- 10 Li, J., Pochanart, P., Wang, Z., Liu, Y., Yamaji, K., Takigawa, M., Kanaya, Y., and Akimoto, H.: Impact of chemical production and transport on summertime diurnal ozone behavior at a mountainous site in North China Plain, *SOLA*, 4, 121–124, 2008b.
- Li, J., Wang, Z., Yamaji, K., Takigawa, M., Kanaya, Y., Pochanart, P., Liu, Y., Irie, H., Hu, B., Tanimoto, H., and Akimoto, H.: Impact of aerosols on summertime tropospheric photolysis frequencies and photochemistry over Central Eastern China, *Atmos. Environ.*, 45, 1817–1829, 2011.
- 15 Lin, P., Hu, M., Deng, Z., Slanina, J., Han, S., Kondo, Y., Takegawa, N., Miyazaki, Y., Zhao, Y., and Sugimoto, N.: Seasonal and diurnal variations of organic carbon in PM_{2.5} in Beijing and the estimation of secondary organic carbon, *J. Geophys. Res.*, 114, D00G11, doi:10.1029/2008JD010902, 2009.
- 20 Lin, W., Xu, X., Zhang, X., and Tang, J.: Contributions of pollutants from North China Plain to surface ozone at the Shangdianzi GAW Station, *Atmos. Chem. Phys.*, 8, 5889–5898, doi:10.5194/acp-8-5889-2008, 2008.
- 25 Liu, Y., Shao, M., Kuster, W. C., Goldan, P. D., Li, X., Lu, S., and de Gouw, J. A.: Source identification of reactive hydrocarbons and oxygenated VOCs in the summertime in Beijing, *Environ. Sci. Technol.*, 43, 75–81, 2009.
- Luo, C., St. John, J. C., Xiuji, Z., Lam, K. S., Wang, T., and Chameides, W. L.: A nonurban ozone air pollution episode over eastern China: observations and model simulations, *J. Geophys. Res.*, 105, 1889–1908, doi:10.1029/1999JD900970, 2000.
- 30 Martin, R. V., Sioris, C. E., Chance, K., Ryerson, T. B., Timothy Bertram, H., Wooldridge, P. J., Cohen, R. C., Neuman, J. A., Swanson, A. and Flocke, F. M.: Evaluation of space-based constraints on global nitrogen oxide emissions with regional aircraft measurements

**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

over and downwind of eastern North America, *J. Geophys. Res.*, 111, D15308, doi:10.1029/2005JD006680, 2006.

Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X., and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, doi:10.5194/acp-7-4419-2007, 2007.

Pan, X. L., Kanaya, Y., Wang, Z. F., Komazaki, Y., Taketani, F., Akimoto, H., Pochanart, P., and Liu, Y.: Estimation of lifetime of carbonaceous aerosol from open crop residue burning during Mount Tai Experiment 2006 (MTX2006), *Atmos. Chem. Phys. Discuss.*, 12, 14363–14392, doi:10.5194/acpd-12-14363-2012, 2012.

Parrish, D. D., Trainer, M., Hereid, D., Williams, E. J., Olszyna, K. J., Harley, R. A., Meagher, J. F., and Fehsenfeld, F. C.: Decadal change in carbon monoxide to nitrogen oxide ratio in U. S. vehicular emissions, *J. Geophys. Res.*, 107, ACH 5-1–ACH 5-9, doi:10.1029/2001JD000720, 2002.

Pathak, R. K., Wang, T., Ho, K. F., and Lee, S. C.: Characteristics of summertime PM_{2.5} organic and elemental carbon in four major Chinese cities: implications of high acidity for water-soluble organic carbon (WSOC), *Atmos. Environ.*, 45, 318–325, 2011.

Richter, A., Burrows, J. P., Nuss, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129–132, doi:10.1038/nature04092, 2005.

Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pandis, S. N., and Pierce, J. R.: Rethinking organic aerosols: semivolatile emissions and photochemical aging, *Science*, 315, 1259–1262, 2007.

Shao, M., Lu, S., Liu, Y., Xie, X., Chang, C., Huang, S., and Chen, Z.: Volatile organic compounds measured in summer in Beijing and their role in ground level ozone formation, *J. Geophys. Res.*, 114, D00G06, doi:10.1029/2008JD010863, 2009.

Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Elson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J. H., and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *J. Geophys. Res.*, 108, D218809, doi:10.1029/2002JD003093, 2003.

Suthawaree, J., Kato, S., Okuzawa, K., Kanaya, Y., Pochanart, P., Akimoto, H., Wang, Z., and Kajii, Y.: Measurements of volatile organic compounds in the middle of Central East China during Mount Tai Experiment 2006 (MTX2006): observation of regional background and

impact of biomass burning, *Atmos. Chem. Phys.*, 10, 1269–1285, doi:10.5194/acp-10-1269-2010, 2010.

Taketani, F., Kanaya, Y., Pochanart, P., Liu, Y., Li, J., Okuzawa, K., Kawamura, K., Wang, Z., and Akimoto, H.: Measurement of overall uptake coefficients for HO₂ radicals by aerosol particles sampled from ambient air at Mts. Tai and Mang (China), *Atmos. Chem. Phys.*, 12, 11907–11916, doi:10.5194/acp-12-11907-2012, 2012.

Tang, J. H., Chan, L. Y., Chan, C. Y., Li, Y. S., Chang, C. C., Liu, S. C., Wu, D., and Li, Y. D.: Characteristics and diurnal variations of NMHCs at urban, suburban, and rural sites in the Pearl River Delta and a remote site in South China, *Atmos. Environ.*, 41, 8620–8632, 2007.

USEPA: Air Quality Criteria for Ozone and Related Photochemical Oxidants (final), EPA/600/R-05/004aF-cF, US Environmental Protection Agency, Washington DC, 2006.

Wang, G., Kawamura, K., Xie, M., Hu, S., Gao, S., Cao, J., An, Z., and Wang, Z.: Size-distributions of *n*-alkanes, PAHs and hopanes and their sources in the urban, mountain and marine atmospheres over East Asia, *Atmos. Chem. Phys.*, 9, 8869–8882, doi:10.5194/acp-9-8869-2009, 2009.

Wang, T., Cheung, V. T. F., Anson, M., and Li, Y. S.: Ozone and related gaseous pollutants in the boundary layer of eastern China: overview of the recent measurements at a rural site, *Geophys. Res. Lett.*, 28, 2373–2376, doi:10.1029/2000GL012378, 2001.

Wang, T., Cheung, T. F., Li, Y. S., Yu, X. M., and Blake, D. R.: Emission characteristics of CO, NO_x, SO₂ and indications of biomass burning observed at a rural site in eastern China, *J. Geophys. Res.*, 107, D124157, doi:10.1029/2001JD000724, 2002.

Wang, T., Wong, C. H., Cheung, T. F., Blake, D. R., Arimoto, R., Baumann, K., Tang, J., Ding, G. A., Yu, X. M., Li, Y. S., Streets, D. G., and Simpson, I. J.: Relationships of trace gases and aerosols and the emission characteristics at Lin'an, a rural site in eastern China, during spring 2001, *J. Geophys. Res.*, 109, D19S05, doi:10.1029/2003JD004119, 2004.

Wang, T., Ding, A., Gao, J., and Wu, W. S.: Strong ozone production in urban plumes from Beijing, China, *Geophys. Res. Lett.*, 33, L21806, doi:10.1029/2006GL027689, 2006.

Wang, Y., McElroy, M. B., Munger, J. W., Hao, J., Ma, H., Nielsen, C. P., and Chen, Y.: Variations of O₃ and CO in summertime at a rural site near Beijing, *Atmos. Chem. Phys.*, 8, 6355–6363, doi:10.5194/acp-8-6355-2008, 2008a.

Wang, Y., Wai, K. M., Gao, J., Liu, X. H., Wang, T., and Wang, W. X.: The impacts of anthropogenic emissions on the precipitation chemistry at an elevated site in North-eastern China, *Atmos. Environ.*, 42, 2959–2970, 2008b.

ACPD

13, 1527–1573, 2013

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**Y. Kanaya et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Wang, Z. F., Li, J., Wang, X. Q., Pochanart, P., and Akimoto, H.: Modeling of regional high ozone episode observed at two mountain sites (Mt. Tai and Huang) in East China, *J. Atmos. Chem.*, 55, 253–272, 2006.

WHO: Health Aspects of Air Pollution with Particulate Matter, Ozone and Nitrogen Dioxide, World Health Organization, Bonn, Germany, 98 pp., 2003.

Wood, E. C., Herndon, S. C., Onasch, T. B., Kroll, J. H., Canagaratna, M. R., Kolb, C. E., Worsnop, D. R., Neuman, J. A., Seila, R., Zavala, M., and Knighton, W. B.: A case study of ozone production, nitrogen oxides, and the radical budget in Mexico City, *Atmos. Chem. Phys.*, 9, 2499–2516, doi:10.5194/acp-9-2499-2009, 2009.

Xu, H., Wang, Y., Wen, T., Yang, Y., and Zhao, Y.: Characteristics and source apportionment of atmospheric aerosols at the summit of Mount Tai during summertime, *Atmos. Chem. Phys. Discuss.*, 9, 16361–16379, doi:10.5194/acpd-9-16361-2009, 2009.

Xu, X., Lin, W., Wang, T., Yan, P., Tang, J., Meng, Z., and Wang, Y.: Long-term trend of surface ozone at a regional background station in eastern China 1991–2006: enhanced variability, *Atmos. Chem. Phys.*, 8, 2595–2607, doi:10.5194/acp-8-2595-2008, 2008.

Yamaji, K., Li, J., Uno, I., Kanaya, Y., Irie, H., Takigawa, M., Komazaki, Y., Pochanart, P., Liu, Y., Tanimoto, H., Ohara, T., Yan, X., Wang, Z., and Akimoto, H.: Impact of open crop residual burning on air quality over Central Eastern China during the Mount Tai Experiment 2006 (MTX2006), *Atmos. Chem. Phys.*, 10, 7353–7368, doi:10.5194/acp-10-7353-2010, 2010.

Yan, P., Tang, J., Huang, J., Mao, J. T., Zhou, X.J., Liu, Q., Wang, Z. F., and Zhou, H. G.: The measurement of aerosol optical properties at a rural site in Northern China, *Atmos. Chem. Phys.*, 8, 2229–2242, doi:10.5194/acp-8-2229-2008, 2008.

Yan, P., Zhang, R., Huan, N., Zhou, X., Zhang, Y., Zhou, H., and Zhang, L.: Characteristics of aerosols and mass closure study at two WMO GAW regional background stations in eastern China, *Atmos. Environ.*, 60, 121–131, 2012.

Yang, M., Howell, S. G., Zhuang, J., and Huebert, B. J.: Attribution of aerosol light absorption to black carbon, brown carbon, and dust in China – interpretations of atmospheric measurements during EAST-AIRE, *Atmos. Chem. Phys.*, 9, 2035–2050, doi:10.5194/acp-9-2035-2009, 2009.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Observed species and methods, and investigators responsible for observations.

Observed species	Method	Investigator, Affiliation
O ₃ , CO	UV absorption, NDIR	P. Pochanart, JAMSTEC
O ₃	UV absorption	T. Wang, J. Gao, Shandong Univ.
NO/NO _x /NO _y	LED/Molybdenum converter/ Chemiluminescence	H. Tanimoto, NIES
NMHCs	Canister/GC-FID/GC-MS	S. Kato, Tokyo Metropolitan Univ.
NMHCs	Canister/GC-MS	Y. Wang, IAP
NMHCs, VOCs	PTR-MS	H. Tanimoto, S. Inomata, NIES
CO ₂	NDIR	Y. Komazaki, JAMSTEC
NO ₂ VCD & AOD from mountain	MAX-DOAS	H. Irie, Y. Kanaya, JAMSTEC
NO ₂ VCD & AOD from foothill	MAX-DOAS	H. Irie, Y. Kanaya, JAMSTEC
Actinic flux & <i>J</i> values	Spectral radiometry	Y. Kanaya, JAMSTEC
<i>T</i> , RH, <i>P</i> , wind	Conventional instruments	Meteorological observatories
Black Carbon (PM ₁ , PM _{2.5})	Multi-Angle Absorption Photometry	P. Pochanart, JAMSTEC
EC/OC	Thermal-Optical-Transmittance	Y. Komazaki, JAMSTEC
Black Carbon	Particle Soot Absorption Photometry	Y. Komazaki, JAMSTEC
Black Carbon	Aethalometer	T. Wang, J. Gao, Shandong Univ.
Ions and metals of size segregated aerosol particles	Cascade impactor/ post analysis	Y. Wang, IAP
Ions and metals of TSP and PM _{2.5}	Sampling/ post analysis	G. Zhuang, Fudan Univ.
Organic gases/aerosols	High volume air sampler, filterpack collections/post analysis	K. Kawamura, Hokkaido Univ.
PAHs/size segregated aerosols	Sampling/ post analysis	G. Wang, Nanjing/Hokkaido Univ.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. MTX2006 modeling activities.

Model Type	Model Name	Horizontal Resolution	Research Subject	PI, Affiliation
0-D Box Model	RACM	NA	Photochemical O ₃ production	Y. Kanaya, JAMSTEC
3-D CTM	NAQPMS/MM5	27 × 27 km nested into 81 km grids	Source attribution of O ₃	J. Li, JAMSTEC/IAP
3-D CTM	CMAQ v.4.4/RAMS v.4.4	80 × 80 km	Impact of crop residual burning on air quality	K. Yamaji, JAMSTEC
3-D CTM	CMAQ v.4.4/	20 × 20 km nested into 80 km grids	Impact of Asian monsoon on O ₃	Y. J. He, JAMSTEC
3-D CTM	CMAQ v. 4.4/RAMS v.4.3	80 × 80 km	Impact of Asian monsoon on acidity of precipitation	B. Z. Ge, IAP

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Average, maximum, and minimum concentrations of O₃, CO, NO_x, NO_y, PM_{2.5}, TSP, BC and OC, sulfate, and nitrate at the summit of Mt. Tai in June 2006.

Species	Average	Maximum	Minimum
O ₃	82 ppbv	160 ppbv	12 ppbv
CO	560 ppbv	1769 ppbv	90 ppbv
NO _x	1.1 ppbv	8.8 ppbv	< DL
NO _y	6.7 ppbv	21.6 ppbv	< DL
PM _{2.5}	123.1 μg m ⁻³	235.7 μg m ⁻³	48.0 μg m ⁻³
TSP	135.0 μg m ⁻³	276.9 μg m ⁻³	49.2 μg m ⁻³
BC (PM ₁)	3.4 μg C m ⁻³	29.5 μg C m ⁻³	< DL
OC (PM ₁ , NIOSH)	9.2 μg C m ⁻³	76.7 μg C m ⁻³	1.4 μg C m ⁻³
Sulfate (PM _{2.5})	21.0 μg m ⁻³	45.1 μg m ⁻³	4.0 μg m ⁻³
Nitrate (PM _{2.5})	8.5 μg m ⁻³	21.6 μg m ⁻³	1.8 μg m ⁻³

< DL indicates below detection limit.
PM_{2.5} and TSP: 2–30 June.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 1. Geographical locations of Mt. Tai, Mt. Hua and Mt. Huang superimposed on EDGAR ver. 4.2 emission inventory map of NO_x (2006).




Fig. 2. Time series of atmospheric pressure, temperature, RH, precipitation, wind direction and speed, $J(\text{NO}_2)$ and $J(\text{O}^1\text{D})$ values.

**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.




Fig. 3. Overall time series of O_3 , CO, benzene, CO_2 , NO, NO_2 , NO_x , NO_y , NO_z , BC, OC, sulfate and nitrate. Hatched areas indicate two episodes with strong influences from OCRB.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Observed correlations between **(a)** NO_y and CO concentrations, **(b)** NO_x and CO concentrations, and **(c)** NO_z and O₃ concentrations. Symbols are colored by O₃ concentrations for **(a)** and **(b)**, and by CO concentrations for **(d)–(f)** same as **(a)–(c)** but using modeled concentrations.

Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.




Fig. 5. Year-round observations of **(a)** O₃ and **(b)** BC at Mt. Tai and other mountain sites. Data for Mt. Tai in June are surrounded by squares.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Overview of the Mount Tai Experiment in China, June 2006

Y. Kanaya et al.




Fig. 6. Average diurnal variations in O₃, CO, CO₂, NO, NO₂, NO_x, NO_y, NO_z, BC, and OC concentrations during the MTX2006 campaign.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Fig. 7. Photographs taken from the observatory looking westward. **(a)** Day with good visibility. **(b)** Pollution layer seen below the mountain in the morning. **(c)** Polluted boundary layer builds up in the morning and covers the mountain top before noon.

Overview of the
Mount Tai Experiment
in China, June 2006

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 8. Backward trajectories with starting altitudes of 1500 m (black) and 750 m (gray) at six different times (YYMMDDHH in LT, shown above individual panels). Plus marks indicate the locations of air mass every 12 h. Hotspots detected by satellite sensors during the 5 preceding days are shown in red (numbers indicate the detection day in June). Harvest areas of winter wheat in different months are shown by colored areas.

**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.




Fig. 9. Correlations between levoglucosan concentrations from TSP filter samples (plus marks) and m/z 97 and m/z 42 (CH_3CN) signals (colored solid lines) derived from continuous PTR-MS measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Overview of the
Mount Tai Experiment
in China, June 2006

Y. Kanaya et al.




Fig. 10. $\Delta[\text{OC}]/\Delta[\text{CO}]$ ratio during the latter period of the campaign (from 16 June 2006) without influence from biomass burning and increasing air mass age, indicated by $\ln([\text{NO}_y]/[\text{NO}_x])$. The CO background concentration, $[\text{CO}]_{\text{bg}}$ was assumed to be 90 ppbv.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Overview of the
Mount Tai Experiment
in China, June 2006**

Y. Kanaya et al.




Fig. 11. Time series of OC concentrations measured for TSP filter samples (green) and size-segregated OC concentrations (PM_{10} and $\text{PM}_{2.5}$) measured using a semi-continuous ECOC analyzer (red). The sum of molecularly identified OCs (165 species) for TSP samples is shown in blue. The time intervals of the TSP samplings are used in all cases

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)