

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Investigation of the connections between atmospheric new particle formation and organics at an urban site of Beijing

Z. B. Wang^{1,2}, M. Hu¹, Z. J. Wu^{1,2}, D. L. Yue¹, J. Zheng³, R. Y. Zhang^{1,3}, X. Y. Pei⁴, P. Paasonen^{4,5}, M. Dal Maso⁴, M. Boy⁴, and A. Wiedensohler²

¹State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

²Leibniz Institute for Tropospheric Research, Permoserstr. 15, 04318 Leipzig, Germany

³Department of Atmospheric Science, Texas A&M University, College Station, Texas 77843, USA

⁴Department of Physics, University of Helsinki, P.O. Box 64, 00014 Helsinki, Finland

⁵International Institute for Applied Systems Analysis, Schlossplatz 1, 2361 Laxenburg, Austria

Received: 30 December 2012 – Accepted: 2 January 2013 – Published: 6 February 2013

Correspondence to: M. Hu (minhu@pku.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

3419

Abstract

The role of low-volatility organic vapors in atmospheric new particle formation has been studied based on a data set of 17 nucleation events observed during the CAREBeijing 2008 campaign. The particle formation rates show good correlations with sulfuric acid and organic vapors implying that both play an important role in the atmospheric new particle formation. High correlation coefficients are observed in all investigated nucleation mechanisms. The best fit ($R = 0.73$, slope = 1.1) between the observed and modelled particle formation rates is achieved with the homogenous nucleation theory of sulfuric acid (both homomolecularly and heteromolecularly) with separate coefficients in $J = K_{SA1}[\text{H}_2\text{SO}_4]^2 + K_{SA2}[\text{H}_2\text{SO}_4][\text{Org}]$. The contributions of the sulfuric acid and the organics involving terms have been 43 % and 57 %, respectively. In addition, the higher particle formation rates are observed on polluted nucleation days, indicating the organic vapors should be involved in the new particle formation process in the polluted urban environment of Beijing with high background aerosol loading.

1 Introduction

New particle formation (NPF) in the atmosphere through nucleation of gas phase species and continuous condensational growth is a crucial secondary transformation process (Kulmala, 2003; Zhang et al., 2011). The freshly formed particles can grow into sizes where they can act as cloud condensation nuclei (CCN) and in this way influence cloud and climate-relevant properties. Model results show that nucleation is an important source of the total particle number concentration and CCN concentration on the global scale (Merikanto et al., 2009). Field observations also reveal the increasing CCN number concentrations originate from new particle formation (Kuang et al., 2009; Yue et al., 2011). Detailed knowledge on the formation mechanisms of new particles and the properties of their subsequent growth will lead to a better understanding on the various effects of atmospheric aerosols.

3420

NPF events have been observed in a variety of environments, including the remote atmosphere of the polar region, above and inside forests, coastal areas, rural and polluted urban regions (Kulmala et al., 2004; Holmes, 2007; Kulmala and Kerminen, 2008). Gaseous sulfuric acid has been identified as a key component in the nucleation process because of its low vapor pressure at typical atmospheric temperatures (Berndt et al., 2005; Sipila et al., 2010). Most studies reveal the relationship between observed nucleation rates and ambient gaseous sulfuric acid concentrations with the exponents in the range 1–2 (Sihto et al., 2006; Riipinen et al., 2007; Kuang et al., 2008; Nieminen et al., 2009; Paasonen et al., 2009, 2010). These exponent values are predicted with activation (exponent 1) and kinetic (exponent 2) nucleation theories (McMurry and Friedlander, 1979; Kulmala et al., 2006). Meanwhile, higher power values in Beijing have been reported recently (Wang et al., 2011), which may imply that thermodynamic processes seem to work better than the activation and kinetic nucleation theories or that sulfuric acid is the vapor mainly causing the very initial growth of particles (Paasonen et al., 2012).

Observation results, however, show large differences in the nucleation coefficients of kinetic nucleation in diverse ambient environments. The values vary in several orders of magnitude (Chen et al., 2012). The variability of the nucleation coefficients in different environments may be influenced by co-nucleation species such as low-volatility organic vapors (Paasonen et al., 2010). Recent results from laboratory studies and field observations imply that organic vapors participate in the nucleation process (Laaksonen et al., 2008; Zhang et al., 2009; Kerminen et al., 2010; Metzger et al., 2010). However, the identities of the specific organic species possibly directly involved in the atmospheric nucleation process are still not known.

A long term study on NPF in the urban of Beijing has been conducted since 2004 at Peking University (Wehner et al., 2004; Wu et al., 2007). Recent results suggest that sulfuric acid plays a dominant role in the nucleation process (Yue et al., 2010; Wang et al., 2011). In this paper, we apply the methods described by Paasonen et al. (2010) and present an analysis of the relationship between particle formation

3421

rates and precursor concentrations, taking into account the possible participation of organic vapors, which has not been discussed before. The role of organics in the nucleation process is evaluated and two typical nucleation event cases in the urban atmosphere of Beijing are discussed.

2 Measurements

2.1 Experimental method

The measurements were performed on the campus of Peking University (39.99° N, 116.31° E), which is located in the northwestern urban area of Beijing. The sampling site is located on the sixth floor of an academic building (about 20 m above the ground level). The local emissions are major contributed by the surrounding traffic. Detailed descriptions of the measurement site and surrounding environment can be found in Wu et al. (2008). Simultaneous measurements of particle number size distributions and gaseous sulfuric acid concentrations were conducted as part of the Campaign of Air Quality Research in Beijing and Surrounding areas in 2008 (CAREBeijing 2008) from 12 July to 25 September.

Number size distributions of atmospheric particles from 3 to 900 nm (mobility diameter) and ambient sulfuric acid concentrations were measured with TDMPMS (Twin Differential Mobility Particle Sizer) system and AP-ID-CIMS (atmospheric pressure ion drift – chemical ionization mass spectrometry), respectively. Detailed information on the instrumentations is given in previous publications (Birmili et al., 1999; Yue et al., 2010; Zheng et al., 2011). The 17 NPF event days investigated in this study are exactly the same as in the research by Wang et al. (2011), however, the data set of particle number size distribution is reevaluated including the additional correction for internal losses of TDMPMS as described in Wiedensohler et al. (2012).

3422

2.2 Data analysis

2.2.1 Particle formation rate J_3

The time evolution of the formation rate J_3 is calculated with the procedure described by Kulmala et al. (2012), recommended by Wang et al. (2011) and Vuollekoski et al. (2012):

$$J_3 = \frac{dN_{3-6}}{dt} + \text{CoagS}_4 \cdot N_{3-6} + \frac{1}{2nm} \text{GR}_6 \cdot N_{5-7} \quad (1)$$

Here, N_{3-6} refers to the particle number concentration in the size range 3–6 nm (covers the five lowest channels of the TDMPs) which are considered as the freshly nucleated particles. The value of GR_6 is assumed equal to the growth rate of the nucleation mode (GR_{3-7} in this study). CoagS_4 is the coagulation sink for the 4 nm particles which is calculated directly from the measured particle number size distribution (Kulmala et al., 2001). Hence, the right hand side of the Eq. (1) includes the measured change of the particle concentration per time interval (first term), the loss by coagulation scavenging (second term) and the growth out of the specific size range at 6 nm (third term).

2.2.2 Particle growth rate (GR) contributed by sulfuric acid

The growth rate of newly formed particles (GR_{3-7}) is estimated based on the particle number size distributions. The determination of the growth rate is made with the method presented by Lehtinen and Kulmala (2003). In this study, the method developed recently by Nieminen et al. (2010) is applied to calculate the vapor concentration required for growth rate of 1 nm h^{-1} in certain particle size ranges:

$$C_{\text{GR}=1 \text{ nm h}^{-1}} = \frac{2\rho_v d_v}{\gamma m_v \Delta t} \cdot \sqrt{\frac{\pi m_v}{8kT}} \cdot \left[\frac{2x_1 + 1}{x_1(x_1 + 1)} - \frac{2x_0 + 1}{x_0(x_0 + 1)} + 2 \ln \left(\frac{x_1(x_0 + 1)}{x_0(x_1 + 1)} \right) \right] \quad (2)$$

here x_0 and x_1 are the ratios of the vapour molecule diameter (d_v) to the initial and final particle diameter (3 nm and 7 nm in this study), respectively. The mass (m_v) and

3423

density (ρ_v) of sulfuric acid applied in this study are 135 amu and 1650 kg m^{-3} , respectively corresponding to hydrated sulfuric acid molecules (Kurten et al., 2007). However, the calculated $C_{\text{GR}=1 \text{ nm h}^{-1}, \text{H}_2\text{SO}_4}$ may be an underestimate because it is assumed that every sulfuric acid molecule colliding with the particle is attached to it which is not necessarily the case. Further descriptions of Eq. (2) can be found in Nieminen et al. (2010).

Then the growth rate contributed by sulfuric acid during the time period used for the determination of GR_{3-7} is calculated directly as:

$$\text{GR}_{\text{H}_2\text{SO}_4} = [\text{H}_2\text{SO}_4]_{\text{det}} / C_{\text{GR}=1 \text{ nm h}^{-1}, \text{H}_2\text{SO}_4} \quad (3)$$

where $[\text{H}_2\text{SO}_4]_{\text{det}}$ is the median value from the measured sulfuric acid concentration during the timeframe for the determination of GR_{3-7} .

2.2.3 Estimating concentration of low-volatility organic vapors

The sulfuric acid concentrations could not explain the observed growth rates of newly formed particles implying that the growth of aerosol particles is affected by other vapors (Boy et al., 2005; Fiedler et al., 2005; Kuang et al., 2010; Riipinen et al., 2011). Donahue et al., (2011) point out that organic condensation to freshly nucleated particles (below 10 nm) contributes substantially to their growth. Meanwhile, previous field observations (O'Dowd et al., 2002; Laaksonen et al., 2008; Riipinen et al., 2012) show the significant contribution of organic compounds to the growth of freshly formed particles which indicates that organic molecules play a crucial role in atmospheric new particle formation. However, the specific organic species in the nucleation and growth are still unclear. Therefore, we have to estimate their properties such as molecular formula, density and concentrations indirectly. For this, we use the following method presented by Paasonen et al. (2010).

We assume that a large gap between the observed growth rates and that due to sulfuric acid condensation is totally contributed by organic vapors, i.e.:

$$\text{GR}_{\text{Org}} = \text{GR}_{\text{det}} - \text{GR}_{\text{H}_2\text{SO}_4} \quad (4)$$

Apparently the determination of the GR_{Org} may cause an uncertainty and further influence the calculation of organic vapor concentration, the uncertainty analysis is presented in Sect. 3.3.

Field observations show evidence for the role of oxidation products of monoterpenes (MTO) in the nucleation and growth processes (Laaksonen et al., 2008). Here we assume the properties of these organic vapors as MTO with the chemical formula $C_{10}H_{16}O_2$ (Taipale et al., 2008). The corresponding molecular mass is 168 amu and the density is 1200 kg m^{-3} (Kannosto et al., 2008; Hallquist et al., 2009). Using Eq. (2) we calculate the potential organic vapor concentration $C_{GR=1 \text{ nm h}^{-1}, Org}$ leading to 1 nm h^{-1} growth, and further the GR_{Org} could be estimated with:

$$[Org]_{det} = GR_{Org} \cdot C_{GR=1 \text{ nm h}^{-1}, Org} \quad (5)$$

With the above calculations we can derive one value of the organic vapor concentration per day during the timeframe for determination of GR_{3-7} . The diurnal variations of organic vapor concentrations depend on both, their source (the oxidation rate of precursors) and sink (condensation on the preexisting particles). Because the concentrations of the precursors could not be achieved, we assume that the nucleation organic vapor concentration is determined by the concentration of hydroxyl radical (OH) and the condensation sink (CS). We approximate that the organic vapors condensing on the newly formed particles are produced via OH oxidation neglecting the other oxidation pathway such as ozonolysis in the daytime (Hao et al., 2009). Even though this assumption may not hold exactly, the diurnal cycle of OH is stronger than that of ozone, which may not have significant influence on the diurnal variation of organic vapors. Here global radiation (GlobRad) is used to make up the lack of the measured OH concentrations because it represents a good correlation with the UVB radiation which is the driving force of ozone photolysis to produce OH radicals. CS is referred to sink of organic vapors which could be obtained from particle number size distribution measurement. Hence, the estimated concentration of organic vapor could be expressed as:

3425

$$[Org](t) = [Org]_{det} \cdot \frac{GlobRad(t)/GlobRad_{det}}{CS(t)/CS_{det}} \quad (6)$$

where t means the certain time and subscript "det" refer to the median value during the timeframe of determination of the growth rate. However, it should be clarified that when we calculate the correlation between the nucleation rates and the precursor vapor concentrations, we only choose the data in the timeframe when strong nucleation event happening.

2.2.4 Nucleation mechanisms

The limitation of activation and kinetic nucleation mechanisms for the Beijing case have been discussed in a paper published by Wang et al. (2011). However these two mechanisms only consider sulfuric acid. In this study, we assume that those organic molecules which are responsible for the growth of the smallest particles are also involved in the nucleation process.

The following two mechanisms consider the organic species instead of sulfuric acid into the nucleation process via cluster activation:

$$J = A_{Org}[Org] \quad (7)$$

or assume both the sulfuric acid and organic vapor can activate clusters:

$$J = A_s([H_2SO_4] + [Org]) \quad (8)$$

Here, A_{Org} and A_s are the activation coefficients (in unit s^{-1}).

Meanwhile we can use organics instead of sulfuric acid in kinetic nucleation mechanism and can write:

$$J = K_{Org}[Org]^2 \quad (9)$$

3426

Also if we only use one organic replacing the sulfuric acid which means the homogeneous heteromolecular nucleation between sulfuric and organic vapor, the nucleation mechanism could be expressed as:

$$J = K_{\text{het}}[\text{H}_2\text{SO}_4][\text{Org}] \quad (10)$$

5 If we consider the homogeneous homomolecular nucleation of sulfuric acid and organic vapor to the heteromolecular nucleation, we obtain:

$$J = K_{\text{SA}} \left([\text{H}_2\text{SO}_4]^2 + [\text{H}_2\text{SO}_4][\text{Org}] \right) \quad (11)$$

$$J = K_{\text{S}} \left([\text{H}_2\text{SO}_4]^2 + [\text{H}_2\text{SO}_4][\text{Org}] + [\text{Org}]^2 \right) \quad (12)$$

10 where all the coefficients K refer to kinetic type nucleation in unit $\text{cm}^3 \text{s}^{-1}$.

In addition to the nucleation mechanisms with one coefficient, as described above, we discuss those mechanisms contain two terms (Eqs. 8 and 11) with separate nucleation coefficients. Then the nucleation mechanisms are rearranged as:

$$J = A_{s1}[\text{H}_2\text{SO}_4] + A_{s2}[\text{Org}] \quad (13)$$

15 and

$$J = K_{\text{SA}1}[\text{H}_2\text{SO}_4]^2 + K_{\text{SA}1}[\text{H}_2\text{SO}_4][\text{Org}] \quad (14)$$

The coefficients above are obtained by minimizing the sum of the squares of the differences between the observed and modelled nucleation rates in every data point.

Both the method to estimate the organic vapor concentration and proposed mechanisms have been discussed in detail in previous study which have been proved justified and reasonable. We refer the reader to Paasonen et al. (2010) for detailed material. Meanwhile, we should clarify that the correlation between formation rates (J_3) and vapor concentrations could only reflect the roles of vapors in atmospheric new particle formation, including the nucleation process and the initial growth of critical clusters from nucleation size to 3 nm.

3427

3 Results and discussion

3.1 Growth rate and organic vapor concentrations

The sulfuric acid concentration $C_{\text{GR}=1\text{nmh}^{-1},\text{H}_2\text{SO}_4}$ needed for 1nmh^{-1} growth varies between $1.1 \times 10^7 \text{cm}^{-3}$ and $1.5 \times 10^7 \text{cm}^{-3}$ due to the different ambient conditions.

5 The parameter Γ , which is defined as the ratio between the observed growth rate and the growth rate related to sulfuric acid, is shown in Fig. 1. The values vary from 6 to 44, which are consistent with previous studies (Stolzenburg et al., 2005; Wehner et al., 2005; Kuang et al., 2010). Hence, the contributions of gaseous sulfuric acid to growth rate are in the range of 2% to 16%, with the mean value of 5%. Since only a minor fraction of the observed, the huge difference between the observed growth rate and that which is explained by sulfuric acid indicates the important role of organics in the growth of newly formed particles.

10 The median organic vapor concentration calculated by the method presented in Sect. 2.2 is $5.7 \times 10^7 \text{moleculescm}^{-3}$ for the whole data set. This value is higher than estimated results for Hyytiälä ($1.6 \times 10^7 \text{moleculescm}^{-3}$), Melpitz ($1.6 \times 10^7 \text{moleculescm}^{-3}$) and Hohenpeissenberg (HPB) ($4.8 \times 10^7 \text{moleculescm}^{-3}$). They are however lower than that observed for San Pietro Capofiume (SPC) ($6.3 \times 10^7 \text{moleculescm}^{-3}$) (Paasonen et al., 2010). The daily maximum concentrations are in the range of $4.6\text{--}42.2 \times 10^7 \text{moleculescm}^{-3}$ with a mean value of $1.5 \pm 1.0 \times 10^8 \text{moleculescm}^{-3}$.

3.2 Correlation between particle formation rate J_3 and vapor concentration in different nucleation mechanisms

3.2.1 Organic vapor nucleation

15 Strong correlations ($R = 0.67$) between the particle formation rate J_3 and organic vapor concentration is observed in Beijing site (see Fig. 2). The same correlation coefficient

3428

R in activation nucleation and kinetic nucleation is due to that R is calculated by the logarithmic values of the observed and modeled particle formation rate, then the exponent of the vapor concentration does not effect on R . $V_{90/10}$ is the ratio of 90th and 10th percentiles of the specific nucleation coefficient, which is calculated through the division between observed nucleation rate and the applied parametrization. The variation of $V_{90/10}$ reflects the dispersion of the nucleation coefficient. For example, the $V_{90/10}$ equal to 10 means 80% of the specific nucleation coefficients are located within one order of magnitude. However, the variation $V_{90/10}$ is sensible to the order of the vapor concentration. Hence, the slightly higher $V_{90/10}$ values for the mechanisms involving the second order vapor concentration (exponent 2 or term $[\text{H}_2\text{SO}_4][\text{Org}]$) are found compared with those related to first order mechanisms. The values of $V_{90/10}$ are 10 and 25 in organic activation and kinetic nucleation mechanisms, respectively (the 10th and 90 percentils of coefficients in different nucleation mechanisms are also shown in Fig. 2). The median values of the nucleation coefficients in organic vapor activation and kinetic nucleation mechanisms are $4.4 \times 10^{-8} \text{ s}^{-1}$ and $6.6 \times 10^{-16} \text{ cm}^3 \text{ s}^{-1}$, respectively.

The comparisons of the correlation coefficient R with other sites in the nucleation mechanisms only involving one species are shown in Table 1. At the comparison sites, only one of the two studied precursors correlates with the particle formation rate: sulfuric acid at Hyytiälä, Melpitz, SPC and organic vapor at HPB. Instead, at Beijing site, J_3 correlates well, both with the sulfuric acid and organic vapor concentrations. The correlation coefficients are 0.66 and 0.67, respectively. This phenomenon implies that the sulfuric acid and low-volatility organic vapors are both have great contribution in the atmospheric new particle formation for Beijing case, compared with the other sites, which are dominated by a single precursor.

3.2.2 Nucleation of sulfuric acid and organic vapor

Since both the sulfuric acid and organic vapor play crucial roles in the new particle formation process at the Beijing site, fair correlation coefficients in the nucleation mechanisms involving both species are achieved as expected (see Table 2). The correlation

3429

coefficients vary from 0.67 to 0.73, which are higher in all proposed nucleation mechanisms compared with the other sites (Paasonen et al., 2010). Highest correlation coefficient ($R = 0.73$) for the entire data set is achieved with the model involving the sum of sulfuric acid homogenous heteromolecular nucleation and homogenous homomolecular nucleation, $J = K_{\text{SA}}([\text{H}_2\text{SO}_4]^2 + [\text{H}_2\text{SO}_4][\text{Org}])$, as shown in Fig. 3a. It should be noticed that the correlation coefficient become weaker ($R = 0.68$) if we consider the term squared organic vapor in the nucleation mechanism. This difference in correlation coefficient is rather small, when taking into account the considerable uncertainties in estimating the organic vapor concentration. These uncertainties, being much larger than those in the sulfuric acid concentration, accumulated in the term including squared organic vapor concentrations, thus automatically weakening the corresponding correlations. Meanwhile, it is also the reason that causes the larger $V_{90/10}$ in Eqs. (9) and (12) than that in Eqs. (10) and (11). Smallest variation, $V_{90/10} = 9.4$ is found in the model involving the sum of the vapor concentration with one nucleation coefficient A_s (see Fig. 3b). On the contrary, the largest variation, $V_{90/10} = 26.9$, results from the organics homogenous homomolecular nucleation ($J = K_{\text{Org}}[\text{Org}]^2$).

3.2.3 Optimized separate nucleation coefficients

The estimated organic vapor concentration is approximately one order of magnitude higher than the observed sulfuric acid concentration, implying that nucleation may rely more on organic compounds. Hence, in this study, the optimized separate nucleation coefficients are evaluated by minimizing the sum of the squares of the differences between the observed and modelled nucleation rates. The values are listed in Table 3. A lower correlation coefficient corresponding to the organic vapor compared to that of sulfuric acid is shown as expected. High correlation coefficient ($R = 0.73$) is achieved with the homogenous nucleation model for sulfuric acid both homomolecularly and hetermolecularly with separate coefficients $K_{s1} = 5.3 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1}$ and $K_{s2} = 5.8 \times 10^{-15} \text{ cm}^3 \text{ s}^{-1}$.

3430

The observed formation rates are plotted versus the modeled formation rates applying the optimized nucleation coefficients in Fig. 4. Even though the correlation coefficient could reach to 0.72 in the mechanism involving the sum of the vapor concentration with separate sulfuric acid and organic activation coefficients ($A_{s1} = 2.0 \times 10^{-7} \text{ s}^{-1}$ and $A_{s2} = 2.8 \times 10^{-8} \text{ s}^{-1}$), the slope between the observed and modelled formation rates is 1.7, which means that the overall modelled values underestimate the real formation rates by almost 50% (see Fig. 4a). On the other hand, the slope of Eq. (14) is 1.1, which is reasonable to explain the observed data.

The formation rate of each term could be calculated using the median values of observed sulfuric acid concentration ($4.9 \times 10^6 \text{ cm}^{-3}$) and estimated organic vapor concentration ($5.7 \times 10^7 \text{ cm}^{-3}$) of the whole investigated data set to multiply the corresponding nucleation coefficients. The contributions of the term only including the sulfuric acid ($[\text{H}_2\text{SO}_4]^2$) and the term involving organics ($[\text{H}_2\text{SO}_4][\text{Org}]$) are 43% and 57%, respectively. Although these values could not quantitative the real contribution of organics exactly, it may reflect its important role in the atmospheric new particle formation process.

3.3 Sensitivity analysis of the related parameters

Several assumptions are made, estimating the low-volatility organic vapor concentrations. Firstly, the remaining growth rate is assumed to be attributed to the organic vapor, which might overestimate its contribution. Secondly, the properties of organic vapors participating in the nucleation process are still unclear. Meanwhile the determination of the growth rate might also cause a double deviation (Kulmala et al., 2012). Hence, the sensitivities of the correlation coefficients to the growth rate, mass and density of the condensation organic vapor are shown in Table 4. The obvious changes of the correlation coefficients are not shown and the error is smaller than 0.1, indicating the change of assumptions could not have a great influence on the final results.

3431

3.4 Case studies

Two cases of the NPF events have been classified based on the correlations between particle formation rate and sulfuric acid concentration in Beijing (Wang et al., 2011): Case I: the exponent in the correlation of the formation rate and sulfuric acid between 1 and 2, which implies that activation or kinetic nucleation is the possible nucleation mechanism; Case II: the exponent is larger than 2.5, which indicates that thermodynamic nucleation is the dominant mechanism. And more than half of the NPF events (10 in 17 investigated days) belong to the second case which is rarely observed in clean environments.

The mean CS (during 08:00–11:00 LT) on the Case II ($0.024 \pm 0.012 \text{ s}^{-1}$) days is 1.7 times higher than that on Case I days ($0.014 \pm 0.006 \text{ s}^{-1}$), which indicates more polluted situation on those days. Hence the potential higher newly formed particles (presented as the ratio of sulfuric acid and CS) in Case II is lower than that in Case I as expected (see Fig. 5). However, the mean particle formation rate J_3 in Case II ($7.5 \pm 11.2 \text{ cm}^{-3} \text{ s}^{-1}$) is typically significant higher than that in Case I ($3.6 \pm 2.1 \text{ cm}^{-3} \text{ s}^{-1}$). A formation rate could be simply expressed as the variation rate of the newly nucleated particles within the certain time period. The opposite trend between the newly formed particle concentration and particle formation rate indicates that on Case II days, more precursor such as organic vapors should be involved in the new particle formation process to “sharp” the variation of newly formed particles.

Here we explore the roles of organic vapor in two kinds of NPF events. Figure 6 presents the mean of daily correlation coefficients in different nucleation mechanisms. It is clear that the mean correlation coefficients in Case II are higher than that in Case I in all nucleation mechanisms involving the organic vapor. This phenomenon implies that the more important role of organic vapor in the new particle formation process on polluted days.

3432

4 Conclusions

Although sulfuric acid has been identified as the key component in new particle formation in the polluted environment of Beijing, the growth rate of newly formed particles related to sulfuric acid could not explain the observed growth rates. Hence, the low-volatility organic vapor is needed to explain the observed growth. The involvement of organic vapors in the nucleation process was estimated in this study. Several nucleation mechanisms involving these organics vapors were tested and the relationships between observed particle formation rates J_3 and vapor concentrations were investigated using the data set of CAREBeijing 2008 campaign.

High correlation coefficient of $R = 0.67$ between the particle formation rate and organic vapor concentration was determined for the Beijing site. This is even higher than that between the formation rate and sulfuric acid concentration ($R = 0.66$). The formation rate correlates well with both the sulfuric acid and organic vapor concentrations, implying that both of them participate in the new particle formation process in Beijing. This is contrary to the clean environment sites, which are only dominated by a single precursor.

In the nucleation mechanisms with single nucleation coefficient (Eqs. 7–12), the highest correlation coefficient ($R = 0.73$) was achieved with the model describing the sum of sulfuric acid homogenous heteromolecular nucleation and homogenous homomolecular nucleation ($J = K_{SA}([H_2SO_4]^2 + [H_2SO_4][Org])$). The 10th and 90th percentiles of the nucleation coefficient K_{SA} are $3.3 \times 10^{-15} \text{ cm}^3 \text{ s}^{-1}$ and $3.6 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1}$, respectively, with the value of $V_{90/10}$ is close to one order of magnitude. Meanwhile, the largest variations of $V_{90/10}$ and smallest correlation coefficient R are obtained in the nucleation mechanism involving the organics homogenous homomolecular nucleation ($J = K_{Org}[Org]^2$), which may due to the uncertainty of the estimated organic vapor concentration.

High correlation coefficient ($R = 0.73$) was achieved with the model homogenous nucleation of sulfuric acid both homomolecularly and heteromolecularly with separate

3433

coefficients ($J = K_{s1}[H_2SO_4]^2 + K_{s2}[H_2SO_4][Org]$), where $K_{s1} = 5.3 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1}$ and $K_{s2} = 5.8 \times 10^{-15} \text{ cm}^3 \text{ s}^{-1}$. The contributions of the sulfuric acid and the organics involving terms have been 43% and 57%, respectively.

Although large uncertainties exist in determining the growth rate and estimating the organic vapor concentration, the sensitivity analysis showed that the assumptions won't have the obvious influence to the correlation coefficients, which indicates the results were reasonable in this study.

The roles of organic vapor in two kinds of NPF events which are distinguished by the exponent between particle formation rate and sulfuric acid concentration have been investigated. Higher correlation coefficients between particle formation rate and organic vapor concentrations in Case II ($n > 2.5$) than that in Case I ($n < 2.5$) suggest the important role of organic vapor in the new particle formation process during pollution days.

Even though Beijing results suggest that the organic vapor play important roles in the atmospheric new particle formation, the specific species of organics is still unknown. Further investigations on the roles of precursor in nucleation process are needed, including the measurements of the chemical compositions of nano-size particles.

Acknowledgements. This work is supported by the National Natural Science Foundation of China (21025728, 20977001, 21190052), the China Ministry of Environmental Protection's Special Funds for Scientific Research on Public Welfare (201009002).

References

- Berndt, T., Boge, O., Stratmann, F., Heintzenberg, J., and Kulmala, M.: Rapid formation of sulfuric acid particles at near-atmospheric conditions, *Science*, 307, 698–700, 2005.
- Birmili, W., Stratmann, F., and Wiedensohler, A.: Design of a DMA-based size spectrometer for a large particle size range and stable operation, *J. Aerosol Sci.*, 30, 549–553, 1999.
- Boy, M., Kulmala, M., Ruuskanen, T. M., Pihlatie, M., Reissell, A., Aalto, P. P., Keronen, P., Dal Maso, M., Hellen, H., Hakola, H., Jansson, R., Hanke, M., and Arnold, F.: Sulphuric

3434

- acid closure and contribution to nucleation mode particle growth, *Atmos. Chem. Phys.*, 5, 863–878, doi:10.5194/acp-5-863-2005, 2005.
- Chen, M., Titcombe, M., Jiang, J., Jen, C., Kuang, C., Fischer, M. L., Eisele, F. L., Siepmann, J. I., Hanson, D. R., Zhao, J., and McMurry, P. H.: Acid–base chemical reaction model for nucleation rates in the polluted atmospheric boundary layer, *P. Natl. Acad. Sci. USA*, 109, 18713–18718, 2012.
- Donahue, N. M., Trump, E. R., Pierce, J. R., and Riipinen, I.: Theoretical constraints on pure vapor-pressure driven condensation of organics to ultrafine particles, *Geophys. Res. Lett.*, 38, L16801, doi:10.1029/2011gl048115, 2011.
- Fiedler, V., Dal Maso, M., Boy, M., Aufmhoff, H., Hoffmann, J., Schuck, T., Birmili, W., Hanke, M., Uecker, J., Arnold, F., and Kulmala, M.: The contribution of sulphuric acid to atmospheric particle formation and growth: a comparison between boundary layers in Northern and Central Europe, *Atmos. Chem. Phys.*, 5, 1773–1785, doi:10.5194/acp-5-1773-2005, 2005.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- Hao, L. Q., Yli-Pirilä, P., Tiitta, P., Romakkaniemi, S., Vaattovaara, P., Kajos, M. K., Rinne, J., Heijari, J., Kortelainen, A., Miettinen, P., Kroll, J. H., Holopainen, J. K., Smith, J. N., Joutsensaari, J., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: New particle formation from the oxidation of direct emissions of pine seedlings, *Atmos. Chem. Phys.*, 9, 8121–8137, doi:10.5194/acp-9-8121-2009, 2009.
- Holmes, N. S.: A review of particle formation events and growth in the atmosphere in the various environments and discussion of mechanistic implications, *Atmos. Environ.*, 41, 2183–2201, 2007.
- Kannosto, J., Virtanen, A., Lemmetty, M., Mäkelä, J. M., Keskinen, J., Junninen, H., Hussein, T., Aalto, P., and Kulmala, M.: Mode resolved density of atmospheric aerosol particles, *Atmos. Chem. Phys.*, 8, 5327–5337, doi:10.5194/acp-8-5327-2008, 2008.
- Kerminen, V.-M., Petäjä, T., Manninen, H. E., Paasonen, P., Nieminen, T., Sipilä, M., Junninen, H., Ehn, M., Gagné, S., Laakso, L., Riipinen, I., Vehkamäki, H., Kurten, T., Ortega, I. K.,

- Dal Maso, M., Brus, D., Hyvärinen, A., Lihavainen, H., Leppä, J., Lehtinen, K. E. J., Mirme, A., Mirme, S., Hörrak, U., Berndt, T., Stratmann, F., Birmili, W., Wiedensohler, A., Metzger, A., Dommen, J., Baltensperger, U., Kiendler-Scharr, A., Mentel, T. F., Wildt, J., Winkler, P. M., Wagner, P. E., Petzold, A., Minikin, A., Plass-Dülmer, C., Pöschl, U., Laaksonen, A., and Kulmala, M.: Atmospheric nucleation: highlights of the EUCAARI project and future directions, *Atmos. Chem. Phys.*, 10, 10829–10848, doi:10.5194/acp-10-10829-2010, 2010.
- Kuang, C., McMurry, P. H., McCormick, A. V., and Eisele, F. L.: Dependence of nucleation rates on sulfuric acid vapor concentration in diverse atmospheric locations, *J. Geophys. Res.*, 113, D10209, doi:10.1029/2007jd009253, 2008.
- Kuang, C., McMurry, P. H., and McCormick, A. V.: Determination of cloud condensation nuclei production from measured new particle formation events, *Geophys. Res. Lett.*, 36, L09822, doi:10.1029/2009gl037584, 2009.
- Kuang, C., Riipinen, I., Sihto, S.-L., Kulmala, M., McCormick, A. V., and McMurry, P. H.: An improved criterion for new particle formation in diverse atmospheric environments, *Atmos. Chem. Phys.*, 10, 8469–8480, doi:10.5194/acp-10-8469-2010, 2010.
- Kulmala, M.: How particles nucleate and grow, *Science*, 302, 1000–1001, 2003.
- Kulmala, M. and Kerminen, V. M.: On the formation and growth of atmospheric nanoparticles, *Atmos. Res.*, 90, 132–150, 2008.
- Kulmala, M., Dal Maso, M., Makela, J. M., Pirjola, L., Vakeva, M., Aalto, P., Miikkulainen, P., Hameri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus B*, 53, 479–490, 2001.
- Kulmala, M., Vehkamäki, H., Petaja, T., Dal Maso, M., Lauri, A., Kerminen, V. M., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35, 143–176, 2004.
- Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster activation theory as an explanation of the linear dependence between formation rate of 3 nm particles and sulphuric acid concentration, *Atmos. Chem. Phys.*, 6, 787–793, doi:10.5194/acp-6-787-2006, 2006.
- Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement of the nucleation of atmospheric aerosol particles, *Nat. Protoc.*, 7, 1651–1667, 2012.
- Kurten, T., Torpo, L., Ding, C. G., Vehkamäki, H., Sundberg, M. R., Laasonen, K., and Kulmala, M.: A density functional study on water-sulfuric acid-ammonia clusters and implica-

- tions for atmospheric cluster formation, *J. Geophys. Res.*, 112, doi:10.1029/2006jd007391, 2007.
- Laaksonen, A., Kulmala, M., O'Dowd, C. D., Joutsensaari, J., Vaattovaara, P., Mikkonen, S., Lehtinen, K. E. J., Sogacheva, L., Dal Maso, M., Aalto, P., Petäjä, T., Sogachev, A., Yoon, Y. J., Lihavainen, H., Nilsson, D., Facchini, M. C., Cavalli, F., Fuzzi, S., Hoffmann, T., Arnold, F., Hanke, M., Sellegri, K., Umann, B., Junkermann, W., Coe, H., Allan, J. D., Alfarra, M. R., Worsnop, D. R., Riekkola, M.-L., Hyötyläinen, T., and Viisanen, Y.: The role of VOC oxidation products in continental new particle formation, *Atmos. Chem. Phys.*, 8, 2657–2665, doi:10.5194/acp-8-2657-2008, 2008.
- Lehtinen, K. E. J. and Kulmala, M.: A model for particle formation and growth in the atmosphere with molecular resolution in size, *Atmos. Chem. Phys.*, 3, 251–257, doi:10.5194/acp-3-251-2003, 2003.
- McMurry, P. H. and Friedlander, S. K.: New particle formation in the presence of an aerosol, *Atmos. Environ.*, 13, 1635–1651, 1979.
- Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, *Atmos. Chem. Phys.*, 9, 8601–8616, doi:10.5194/acp-9-8601-2009, 2009.
- Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions, *P. Natl. Acad. Sci. USA*, 107, 6646–6651, 2010.
- Nieminen, T., Manninen, H. E., Sihto, S. L., Yli-Juuti, T., Mauldin, R. L., Petaja, T., Riipinen, I., Kerminen, V. M., and Kulmala, M.: Connection of sulfuric acid to atmospheric nucleation in boreal forest, *Environ. Sci. Technol.*, 43, 4715–4721, 2009.
- Nieminen, T., Lehtinen, K. E. J., and Kulmala, M.: Sub-10 nm particle growth by vapor condensation – effects of vapor molecule size and particle thermal speed, *Atmos. Chem. Phys.*, 10, 9773–9779, doi:10.5194/acp-10-9773-2010, 2010.
- O'Dowd, C. D., Aalto, P., Hameri, K., Kulmala, M., and Hoffmann, T.: Aerosol formation – atmospheric particles from organic vapours, *Nature*, 416, 497–498, 2002.
- Paasonen, P., Sihto, S. L., Nieminen, T., Vuollekoski, H., Riipinen, I., Plass-Dulmer, C., Berresheim, H., Birmili, W., and Kulmala, M.: Connection between new particle formation and sulphuric acid at Hohenpeissenberg (Germany) including the influence of organic compounds, *Boreal Environ. Res.*, 14, 616–629, 2009.

3437

- Paasonen, P., Nieminen, T., Asmi, E., Manninen, H. E., Petäjä, T., Plass-Dülmer, C., Flen-tje, H., Birmili, W., Wiedensohler, A., Hõrrak, U., Metzger, A., Hamed, A., Laaksonen, A., Facchini, M. C., Kerminen, V.-M., and Kulmala, M.: On the roles of sulphuric acid and low-volatility organic vapours in the initial steps of atmospheric new particle formation, *Atmos. Chem. Phys.*, 10, 11223–11242, doi:10.5194/acp-10-11223-2010, 2010.
- Paasonen, P., Olenius, T., Kupiainen, O., Kurtén, T., Petäjä, T., Birmili, W., Hamed, A., Hu, M., Huey, L. G., Plass-Duelmer, C., Smith, J. N., Wiedensohler, A., Loukonen, V., McGrath, M. J., Ortega, I. K., Laaksonen, A., Vehkamäki, H., Kerminen, V.-M., and Kulmala, M.: On the formation of sulphuric acid – amine clusters in varying atmospheric conditions and its influence on atmospheric new particle formation, *Atmos. Chem. Phys.*, 12, 9113–9133, doi:10.5194/acp-12-9113-2012, 2012.
- Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III–IV campaigns in Heidelberg and Hyytiälä, *Atmos. Chem. Phys.*, 7, 1899–1914, doi:10.5194/acp-7-1899-2007, 2007.
- Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V.-M., Worsnop, D. R., Pandis, S. N., Donahue, N. M., and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei (CCN) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, doi:10.5194/acp-11-3865-2011, 2011.
- Riipinen, I., Yli-Juuti, T., Pierce, J. R., Petaja, T., Worsnop, D. R., Kulmala, M., and Donahue, N. M.: The contribution of organics to atmospheric nanoparticle growth, *Nat. Geosci.*, 5, 453–458, 2012.
- Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms, *Atmos. Chem. Phys.*, 6, 4079–4091, doi:10.5194/acp-6-4079-2006, 2006.
- Sipila, M., Berndt, T., Petaja, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin, R. L., Hyvarinen, A. P., Lihavainen, H., and Kulmala, M.: The role of sulfuric acid in atmospheric nucleation, *Science*, 327, 1243–1246, 2010.

3438

- Stolzenburg, M. R., McMurry, P. H., Sakurai, H., Smith, J. N., Mauldin, R. L., Eisele, F. L., and Clement, C. F.: Growth rates of freshly nucleated atmospheric particles in Atlanta, J. Geophys. Res., 110, D22S05, doi:10.1029/2005jd005935, 2005.
- Taipale, R., Ruuskanen, T. M., Rinne, J., Kajos, M. K., Hakola, H., Pohja, T., and Kulmala, M.: Technical Note: Quantitative long-term measurements of VOC concentrations by PTR-MS – measurement, calibration, and volume mixing ratio calculation methods, Atmos. Chem. Phys., 8, 6681–6698, doi:10.5194/acp-8-6681-2008, 2008.
- Vuollekoski, H., Sihto, S.-L., Kerminen, V.-M., Kulmala, M., and Lehtinen, K. E. J.: A numerical comparison of different methods for determining the particle formation rate, Atmos. Chem. Phys., 12, 2289–2295, doi:10.5194/acp-12-2289-2012, 2012.
- Wang, Z. B., Hu, M., Yue, D. L., Zheng, J., Zhang, R. Y., Wiedensohler, A., Wu, Z. J., Nieminen, T., and Boy, M.: Evaluation on the role of sulfuric acid in the mechanisms of new particle formation for Beijing case, Atmos. Chem. Phys., 11, 12663–12671, doi:10.5194/acp-11-12663-2011, 2011.
- Wehner, B., Wiedensohler, A., Tuch, T. M., Wu, Z. J., Hu, M., Slanina, J., and Kiang, C. S.: Variability of the aerosol number size distribution in Beijing, China: new particle formation, dust storms, and high continental background, Geophys. Res. Lett., 31, L22108, doi:10.1029/2004gl021596, 2004.
- Wehner, B., Petäjä, T., Boy, M., Engler, C., Birmili, W., Tuch, T., Wiedensohler, A., and Kulmala, M.: The contribution of sulfuric acid and non-volatile compounds on the growth of freshly formed atmospheric aerosols, Geophys. Res. Lett., 32, L16801, doi:10.1029/2005gl023827, 2005.
- Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Gröning, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H.-G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, Atmos. Meas. Tech., 5, 657–685, doi:10.5194/amt-5-657-2012, 2012.

3439

- Wu, Z. J., Hu, M., Liu, S., Wehner, B., Bauer, S., Maßling, A., Wiedensohler, A., Petäjä, T., Dal Maso, M., and Kulmala, M.: New particle formation in Beijing, China: statistical analysis of a 1-year data set, J. Geophys. Res., 112, D09209, doi:10.1029/2006jd007406, 2007.
- Wu, Z. J., Hu, M., Lin, P., Liu, S., Wehner, B., and Wiedensohler, A.: Particle number size distribution in the urban atmosphere of Beijing, China, Atmos. Environ., 42, 7967–7980, 2008.
- Yue, D. L., Hu, M., Zhang, R. Y., Wang, Z. B., Zheng, J., Wu, Z. J., Wiedensohler, A., He, L. Y., Huang, X. F., and Zhu, T.: The roles of sulfuric acid in new particle formation and growth in the mega-city of Beijing, Atmos. Chem. Phys., 10, 4953–4960, doi:10.5194/acp-10-4953-2010, 2010.
- Yue, D. L., Hu, M., Zhang, R. Y., Wu, Z. J., Su, H., Wang, Z. B., Peng, J. F., He, L. Y., Huang, X. F., Gong, Y. G. and Wiedensohler, A.: Potential contribution of new particle formation to cloud condensation nuclei in Beijing, Atmos. Environ., 45, 6070–6077, 2011.
- Zhang, R., Khalizov, A., Wang, L., Hu, M., and Xu, W.: Nucleation and growth of nanoparticles in the atmosphere, Chem. Rev., 112, 1957–2011, doi:10.1021/cr2001756, 2011.
- Zhang, R. Y., Wang, L., Khalizov, A. F., Zhao, J., Zheng, J., McGraw, R. L., and Molina, L. T.: Formation of nanoparticles of blue haze enhanced by anthropogenic pollution, P. Natl. Acad. Sci. USA, 106, 17650–17654, 2009.
- Zheng, J., Hu, M., Zhang, R., Yue, D., Wang, Z., Guo, S., Li, X., Bohn, B., Shao, M., He, L., Huang, X., Wiedensohler, A., and Zhu, T.: Measurements of gaseous H₂SO₄ by AP-ID-CIMS during CAREBeijing 2008 Campaign, Atmos. Chem. Phys., 11, 7755–7765, doi:10.5194/acp-11-7755-2011, 2011.

3440

Table 1. Comparisons of the correlation coefficient R in different nucleation mechanisms with other studies.

	Beijing	Hyytiälä	HPB	Melpitz	SPC
Air mass type	Urban	Forest	Mountain	Rural	Rural
# Event	17	10	15	8	3
Sulfuric acid	0.66	0.58	0.17	0.68	0.58
Organic vapor	0.67	0.29	0.61	-0.23	0.07
Ref.	This study	Paasonen et al. (2010)			

Sulfuric acid means the nucleation mechanisms $J = A[\text{H}_2\text{SO}_4]$ and $J = K[\text{H}_2\text{SO}_4]^2$.
Organic vapor means the nucleation mechanisms $J = A_{\text{Org}}[\text{Org}]$ and $J = K_{\text{Org}}[\text{Org}]^2$.

3441

Table 2. Percentile (10th, 25th, 75th and 90th), median and mean values of the nucleation coefficients of the whole dataset. $V_{90/10}$ is the ratio of 90th to 10th percentile values of the ratio between the observed and modeled particle formation rate. Correlation coefficients between observed and modeled particle formation rates are also presented.

Nucleation coefficient		Percentiles				Median	Mean	$V_{90/10}$	R
		10th	25th	75th	90th				
A_{Org} (10^{-8} s^{-1})	Eq. (7)	1.6	2.6	9.0	15.3	4.4	6.7	9.8	0.67
A_{s} (10^{-8} s^{-1})	Eq. (8)	1.5	2.3	8.1	13.6	4.1	6.0	9.4	0.69
K_{Org} ($10^{-16} \text{ cm}^3 \text{ s}^{-1}$)	Eq. (9)	2.0	3.7	19.6	54.6	6.6	23.8	26.9	0.67
K_{het} ($10^{-15} \text{ cm}^3 \text{ s}^{-1}$)	Eq. (10)	3.6	5.5	19.5	41.5	9.6	17.0	11.4	0.73
K_{SA} ($10^{-15} \text{ cm}^3 \text{ s}^{-1}$)	Eq. (11)	3.3	5.1	17.6	35.9	9.0	15.0	10.8	0.73
K_{s} ($10^{-16} \text{ cm}^3 \text{ s}^{-1}$)	Eq. (12)	1.9	3.4	17.6	43.9	6.0	18.8	23.4	0.68

3442

Table 3. Values of optimized separate nucleation coefficients and the corresponding $V_{90/10}$ and R .

Nucleation mechanism	A_{s1}	A_{s2}	K_{s1}	K_{s2}	$V_{90/10}$	R
	A (10^8 s^{-1})		K ($10^{-15} \text{ cm}^3 \text{ s}^{-1}$)			
$J = A_{s1}[\text{H}_2\text{SO}_4] + A_{s2}[\text{Org}]$	19.7	2.8	–	–	9.3	0.72
$J = K_{s1}[\text{H}_2\text{SO}_4]^2 + K_{s2}[\text{H}_2\text{SO}_4][\text{Org}]$	–	–	53.0	5.8	9.9	0.73

3443

Table 4. Correlation coefficients in different assumptions.

Nucleation mechanism	correlation coefficient R				
	GR	GR/2 ^a	2GR ^b	amu = 84 $\rho = 500 \text{ kg m}^{-3}$	amu = 336 $\rho = 1900 \text{ kg m}^{-3}$
$J = A_{\text{Org}}[\text{Org}]$	0.67	0.67	0.67	0.66	0.66
$J = K_{\text{Org}}[\text{Org}]^2$	0.67	0.67	0.67	0.66	0.66
$J = A_s([\text{H}_2\text{SO}_4] + [\text{Org}])$	0.69	0.70	0.68	0.68	0.68
$J = K_{\text{net}}[\text{H}_2\text{SO}_4][\text{Org}]$	0.73	0.73	0.73	0.72	0.72
$J = K_{\text{SA}}([\text{H}_2\text{SO}_4]^2 + [\text{H}_2\text{SO}_4][\text{Org}])$	0.73	0.74	0.73	0.73	0.72
$J = K_s([\text{H}_2\text{SO}_4]^2 + [\text{H}_2\text{SO}_4][\text{Org}] + [\text{Org}]^2)$	0.73	0.73	0.73	0.72	0.72
$J = A_{s1}[\text{H}_2\text{SO}_4] + A_{s2}[\text{Org}]$	0.72	0.73	0.72	0.72	0.72
$J = K_{s1}[\text{H}_2\text{SO}_4]^2 + K_{s2}[\text{H}_2\text{SO}_4][\text{Org}]$	0.73	0.74	0.73	0.73	0.72

^a GR/2 means the organic vapor only contribute 50 % of the remaining growth rate.^b 2GR means the possible error made in determination the growth rate, the real growth rate is two times higher than the observed growth rate.

3444




Fig. 1. Ratios between the observed growth rate from the particle number size distribution and the calculated growth rate contributed by sulfuric acid, as a function of the ambient sulfuric acid concentrations.

3445




Fig. 2. The particle formation rate J_3 versus (a) organic vapor concentrations and (b) squared organic vapor concentrations. The solid and dash lines present the 90th and 10th percentile values of the nucleation coefficients, respectively.

3446




Fig. 3. The particle formation rate J_3 versus (a) the sum of the squared sulfuric acid concentrations and the product of sulfuric acid and organic vapor concentrations and (b) the sum of the sulfuric acid and organic vapor concentrations, (b) the product of sulfuric acid and organic vapor concentrations. The solid and dash lines present the 90th and 10th percentile values of the nucleation coefficients, respectively.

3447




Fig. 4. The observed particle formation rate J_3 versus (a) the sum of the sulfuric acid and organic vapor concentrations multiplied with separate nucleation coefficients and (b) the sum of the squared sulfuric acid and the product of sulfuric acid and organic vapor concentrations multiplied with separate nucleation coefficients. The dash line presents the 1 : 1 line.

3448




Fig. 5. Comparisons of different parameters between two kinds of NPF events days. Red ones indicate the days with the exponent smaller than 2.5, blue ones indicate the days with the exponent larger than 2.5.

3449




Fig. 6. Comparisons of correlation coefficients in different nucleation mechanisms between two kind of NPF events days. Red ones indicate days with the exponent smaller than 2.5, blue ones indicate the days with the exponent larger than 2.5.

3450