

[Interactive  
Comment](#)

## ***Interactive comment on “Stratospheric SO<sub>2</sub> and sulphate aerosol, model simulations and satellite observations” by C. Brühl et al.***

**R. Saunders**

rws\_brancaaster@hotmail.com

Received and published: 20 May 2013

I was interested to read the paper and its implication of a meteoric dust sink to explain sulphur levels in the upper stratosphere. I'd like to bring to the authors attention, the paper by myself and colleagues (ACP, 12, 4387-, 2012) in which we modelled sulphuric acid loss to meteoric smoke (nano) particles in an effort to explain the low acid levels above 40 km, where a small number of measurements have been reported. We found that an uptake (sticking) coefficient of 0.01 or greater (section 3.4) was required for this heterogeneous uptake mechanism to account for the measured data i.e. the value used by the authors for this parameter in their treatment (0.01) is consistent with what we reported. As encouraging as such inter-model corroboration is, laboratory-measured

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



verification of this value using realistic meteoritic material and particle size is now vital.

---

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 11395, 2013.

ACPD

13, C2532–C2533, 2013

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C2533

