



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

A molecular-level approach for characterizing water-insoluble components of ambient organic aerosol particulates using ultra-high resolution mass spectrometry

A. S. Willoughby, A. S. Wozniak, and P. G. Hatcher

Department of Chemistry & Biochemistry, Old Dominion University, Norfolk, VA 23529, USA

Received: 2 April 2014 – Accepted: 9 April 2014 – Published: 24 April 2014

Correspondence to: P. G. Hatcher (phatcher@odu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The chemical composition of organic aerosols in the atmosphere is strongly influenced by human emissions, and the effect these have on the environment, human health, and climate change is determined by the molecular nature of these chemical species. The complexity of organic aerosol samples limits the ability to study the chemical composition, and, therefore, the associated properties and the impacts they have. Many studies address the water-soluble fraction of organic aerosols, and have had much success in identifying specific molecular formulas for thousands of compounds present. However, little attention is given to the water-insoluble portion, which can contain most of the fossil material that is emitted through human activity. Here we compare the organic aerosols present in water extracts and organic solvent extracts (pyridine and acetonitrile) of an ambient aerosol sample collected in a rural location that is impacted by natural and anthropogenic emission sources. A semi-quantitative method was developed using proton nuclear magnetic resonance spectroscopy to determine that the amount of organic matter extracted by pyridine is comparable to that of water. Electro spray ionization Fourier transform ion cyclotron resonance mass spectra show that pyridine extracts a molecularly unique fraction of organic matter compared to water or acetonitrile, which extract chemically similar organic matter components. The molecular formulas unique to pyridine were less polar, more aliphatic, and reveal formulas containing sulfur to be an important component of insoluble aerosol organic matter.

1 Introduction

The introduction and improvement of advanced spectroscopic methods has provided a wealth of new chemical information regarding organic aerosols in the last few decades. Establishing the chemical identity of the individual components that make up organic aerosols has remained an important goal in order to understand the relationships between their sources, transport, molecular identities and transformations

ACPD

14, 10393–10427, 2014

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and their impacts and fates in the environment. The importance of organic aerosols to climate and biogeochemical cycling has been well documented (Andreae and Crutzen, 1997; Andreae and Gelencsér, 2006; Bond et al., 2013; Booth et al., 2012; Crutzen and Andreae, 1990; Jacobson, 2000; Ramanathan et al., 2001). The specific molecular nature of organic aerosols directly influences the impact they have on the environment. For example, light absorption is caused by electronic transition associated with molecular double bonds and the number of double bonds and aromatic rings present within the structure has been linked to the ability of that molecule to absorb ultraviolet or visible light (Andreae and Gelencsér, 2006). Given the connections between chemical characteristics and environmental impact, establishing the relationship between aerosol OM source and associated chemical characteristics is important for modeling and predicting the net impact they have on environmental systems. To date, uncertainties in the molecular makeup of organic aerosols limit our ability to make these linkages between aerosol OM source, chemical characteristics, and environmental impact.

Between 10–90 % of total aerosol mass is comprised of OM (Jimenez et al., 2009; Kanakidou et al., 2005; and references within), and 10–70 % of that OM is water-soluble (Decesari et al., 2007; Kleefeld et al., 2002; Sullivan et al., 2004; Zappoli et al., 1999) depending on its physical and chemical composition. A suite of analytical techniques have provided bulk and specific chemical information about carbonaceous species present within fractions of atmospheric OM samples (Hallquist et al., 2009), including ambient aerosols, fog water and rainwater, and laboratory generated secondary organic aerosols (SOA). These efforts have shown aerosol OM to be made up of a highly diverse suite of oxygenated compounds that include aliphatic and conjugated functional groups (e.g., Decesari et al., 2007; Jimenez et al., 2009; Mayol-Bracero et al., 2002; Wozniak et al., 2008), which influence the water-solubility as well as the light-absorbing capacity of the compounds (Andreae and Gelencsér, 2006; Robertson and O'Reilly, 1987). Recently, ultra-high resolution mass spectrometry has shown an extraordinary capacity for characterizing aerosol OM, and is the only current technique able to provide elemental formula information for the thousands of compounds present

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

within individual ambient aerosols (Lin et al., 2012; Mazzoleni et al., 2012; Reemtsma et al., 2006; Schmitt-Kopplin et al., 2010; Wozniak et al., 2008) and laboratory generated aerosols (Bateman et al., 2010; Heaton et al., 2009; Laskin et al., 2010; Reinhardt et al., 2007) without prior chromatographic separation. In addition to oxygenated compounds, many other functionalized species, including those containing nitrogen and sulfur, have been identified in ambient aerosols, fog water (LeClair et al., 2012; Mazzoleni et al., 2010), and rainwater (Altieri et al., 2009, 2012; Mitra et al., 2013) from various sources.

Numerous publications characterize some fraction of aerosol OM using various solvent systems, ionization sources, and mass analyzers. To date, much of the work has focused on the water-soluble fraction, leaving little information regarding the water-insoluble fraction that comprises up to 90 % of the OM. Radiocarbon data for organic aerosols collected in rural environments suggests that much of the water-insoluble carbon is fossil-derived and from anthropogenic sources, whereas the water-soluble carbon is contemporary and biogenically-derived (Szidat et al., 2004; Wozniak et al., 2012a, b). Limiting molecular analysis to the water-soluble OM (WSOM) means that a quantitatively important component of organic aerosols is missing. Studies have examined the methanol-soluble (Heaton et al., 2009) and acetonitrile-soluble (Bateman et al., 2010; Heaton et al., 2009) fractions of SOAs and other laboratory generated aerosols. Studies of SOA, which are thought to be highly water-soluble, have shown that the water and acetonitrile extracted OM fractions are extremely similar at the molecular level (Bateman et al., 2010; Heaton et al., 2009). However, Bateman et al. (2010) examined a laboratory-generated biomass burning aerosol and determined that the acetonitrile-soluble component of aerosol OM had characteristically lower oxygen to carbon (O/C) ratios than the water-soluble component, suggesting molecular differences between water-soluble and solvent-soluble components of some organic aerosols. These differences imply that the water-insoluble materials in aerosol OM have molecular characteristics distinct from water-soluble OM. However, the differ-

ences in water-insoluble and water-soluble aerosol OM have still not been extensively explored.

In the present study we evaluate the specific molecular composition of the water-insoluble fraction of ambient aerosol OM. We employ Fourier transform ion cyclotron resonance mass spectrometry with negative electrospray ionization (ESI-FTICR-MS) for molecular characterization and NMR spectroscopy for additional structural characterization. We select pyridine and acetonitrile, two solvents that exhibit different polarities than water, have different solvating characteristics, and are compatible with ESI-FTICR-MS and NMR.

Three ambient aerosol particulate samples were collected from a coastal Virginia (USA) site influenced by a mixture of anthropogenic and biogenic aerosol OM sources typical to the east coast of the United States. The suitability of pyridine and acetonitrile as solvents for characterizing aerosol water-insoluble organic matter (WIOM) by ESI-FTICR-MS was established by comparing the molecular formulas assigned to the pyridine-soluble organic matter (PSOM), acetonitrile-soluble organic matter (ASOM), and WSOM extracts of these ambient aerosol particulate samples.

2 Methods

2.1 Sample collection

Three ambient aerosol total suspended particulate (TSP) samples were collected during the summer (16–17 August 2011; 24–25 June 2013; and 25–26 June 2013) at the Virginia Institute of Marine Science in Gloucester Point, VA (37.2482° N, 76.5005° W). Air was drawn through a pre-combusted (4 h, 475 °C) quartz microfiber filter (Whatman QM/A, 20.3 × 25.4 cm, 419 cm² exposed area, 0.6 μm effective pore size) using a TSP high volume air sampler (Model GS2310, Thermo Andersen, Smyrna, GA) at an average flow rate of 0.81 m³ min⁻¹. Air particles were collected for 24 h with total air volumes ranging between 1124–1169 m³, and are expected to have contributions from several

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nearby biogenic and anthropogenic sources (e.g., estuarine waters, natural vegetation, light vehicle traffic, oil refinery, coal-fired power plant) as is typical of the eastern United States. The samples were transferred to a pre-combusted foil pouch immediately after collection and stored at -8°C until analysis. A new and pre-combusted QM/A filter blank was stored under identical conditions as the sample, and was analyzed as a storage filter blank for the 2011 sample. A new and pre-combusted QM/A filter was attached to the air sampler immediately prior to the 2013 sampling (24 June 2013) and then stored under identical conditions as the samples, and was analyzed as a field filter blank for the 2013 samples.

2.2 Aerosol mass and carbon measurements

The QM/A filters were weighed before and after sampling to determine the total aerosol mass (24.3–29.7 mg TSP) and concentration ($20.8\text{--}26.2\ \mu\text{g m}^{-3}$). Triplicate aerosol core plug ($2.84\ \text{cm}^2$ area) samples were acidified to remove inorganic carbon by treating the plugs with 1 M HCl followed by drying in an oven (4 h, 80°C). Acidified filter plugs were evaluated for total carbon (C = organic + elemental/black – inorganic carbonates) and nitrogen (N) content using a Flash EA 1112 elemental analyzer (ThermoFinnigan), where quantification was achieved using an aspartic acid standard calibration curve. Acidified blank QM/A filters were evaluated for blank subtraction; however, C and N quantities for all of the blank QM/A filter plugs were below the detection levels of the instrument.

2.3 Solvent extractions

Replicate solvent extracts of the aerosols and the respective filter blank were obtained by combining three aerosol plugs or blank plugs (3 aerosol plugs contain between 0.116 and 0.147 mg C) with either 15 mL water (Millipore Synergy Ultrapure Water System), 5 mL pyridine (Acros, 99+%), or 5 mL acetonitrile (Fisher Optima, 99.9%) and thoroughly mixed on an orbital shaker table (150 RPM, 4 h, 21°C). Soluble particles

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were isolated using a syringe filter with a pre-combusted glass fiber filter (0.7 μm pore size). The water extracts were then desalted and concentrated using an established procedure for PPL solid-phase extraction cartridges (Dittmar et al., 2008). The desalted sample was eluted in 5 mL of methanol (WSOM, Acros, 99.9%). The pyridine filtrate (PSOM) and acetonitrile filtrate (ASOM) did not require further manipulation, and all three extracts were thus prepared so that the final solutions had the same filter plug to solvent volume ratio (3 plugs in 5 mL of solvent). Samples were stored at -8°C until FTICR-MS analysis, typically within 24 h of preparation.

2.4 Extraction efficiency determinations

The filtrates from the water-extracts were evaluated for non-purgeable organic carbon using a Shimadzu TOC-VCPH analyzer to measure water soluble organic carbon (WSOC) content (Wozniak et al., 2008, 2012a). The WSOC content was then compared with the total aerosol organic carbon content to find that 54–60% of the total aerosol organic carbon was WSOC.

An important aspect of choosing a suitable solvent for WIOM characterization is its extraction efficiency which could not be accurately determined for acetonitrile and pyridine using standard TOC analysis. Upon evaporation of solvent extracts, these organic solvents adhere to natural organic matter artificially inflating the carbon content in acetonitrile and pyridine extracts. Therefore, a method was developed based on ^1H NMR using glucose (98%, Acros Organics) as a standard, which is 100% soluble in pyridine at low concentrations ($< 1\text{ mg mL}^{-1}$). We determined that acetonitrile interferes greatly with our measurement strategy and so we abandoned attempts to determine extraction efficiency in this solvent. A known mass of glucose was dissolved into pyridine- D_5 (100% atom D, Acros Organics) providing known carbon and hydrogen concentrations to compare to ^1H NMR spectra. Aerosol plugs of known OC masses were each dissolved into pyridine- D_5 and water, separately. The WSOM and PSOM samples were then subjected to ^1H NMR spectroscopy using a Bruker Daltonics 400 MHz NMR with a BBI probe. WSOM samples were diluted using D_2O (100% atom D, Acros Organics)

at a ratio of 90 : 10 WSOM : D₂O. PSOM samples were analyzed in a solvent system of 100 % pyridine-D₅. All samples were analyzed for 4000 scans using a standard Bruker water-suppression pulse program, where the 90° pulse and the transmitter offset were optimized individually for each sample.

The signals obtained from ¹H NMR spectra were integrated over the range of 0.1 to 4.4 ppm to get a total signal response, and also integrated over three ranges of chemical shifts to determine contributions from the major proton types (Moretti et al., 2008; Shakya et al., 2012), aliphatic, unsaturated alkyl, and oxygenated aliphatics. Aliphatic hydrogen (H-C, 0.6–1.8 ppm for WSOM and 0.7–1.95 ppm for PSOM), unsaturated alkyl or hydrogen α to unsaturated carbons (H-C-C=, 1.8–3.2 ppm for WSOM and 1.95–3.2 ppm for PSOM), and oxygenated aliphatic hydrogen including alcohol, ether, and ester function groups (H-C-O, 3.2–4.4 ppm in WSOM and PSOM). Though aromatic protons (6.5–8.2 ppm) are used in other studies, we omit them from analysis due to the interference of exchanged proton in the pyridine-D₅ solvent. The region of 0.1–4.4 ppm is appropriate for this study because it is free of signals from the solvent and contains the majority of signal for these types of samples. The signal response for each proton region was normalized to the total signal response between 0.1–4.4 to establish the percent contribution for each proton type. These percentages were used to calculate the average H/C ratio for each sample to be used to convert hydrogen to carbon content (see Table S1). An H/C ratio of 2 is used for the aliphatic hydrogen and unsaturated alkyl hydrogen, and an H/C ratio of 1.1 is used for the oxygenated aliphatics (Decesari et al., 2007; Moretti et al., 2008; Shakya et al., 2012).

The signal obtained for glucose protons dissolved in pyridine-D₅ were used to establish a glucose response factor (area per mass unit hydrogen). The total area between 0.1–4.4 ppm in each sample was converted into a mass of dissolved hydrogen using the glucose response factor. The dissolved hydrogen mass was converted to dissolved carbon mass using the calculated average H/C ratio (1.94–1.98). Comparison of the calculated dissolved carbon amount to the starting mass of carbon gives a relative percentage of extractable organic matter. Details on these values are provided in Table S2.

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.5 ESI-FTICR-MS

Immediately prior to analysis, the WSOM replicates (in methanol) and filter blank extract were diluted by 2 using LC/MS grade water with a small amount (< 0.1 % total volume) of ammonium hydroxide to enhance ionization efficiency. The PSOM, ASOM, and respective filter blank extracts were diluted by 2 using methanol with a small amount (< 0.1 % total volume) of ammonium hydroxide. Prior to sample analysis, the instrument was externally calibrated using a polyethylene glycol standard. Each of the samples was introduced to an Apollo II electrospray ionization source (negative ion mode) at a flow rate of 120 nLmin⁻¹ on a Bruker Daltonics 12 Tesla Apex Qe FTICR-MS housed at the College of Sciences Major Instrumentation Cluster at Old Dominion University. Spray voltages were optimized for each sample. Ions were accumulated in the hexapole for 0.5–2.0 s before transfer into the ICR cell, where exactly 300 transients were co-added. Experimental duplicates were evaluated for each aerosol sample and solvent mixture to ensure good experimental and instrumental reproducibility.

2.6 Data processing

Each spectrum was calibrated internally using naturally occurring molecules (fatty acids, dicarboxylic acids, and other homologous series with only carbon, hydrogen, and oxygen in the molecular formula) within the sample (Sleighter et al., 2008). Salt peaks (mass defect 0.4–0.98 for $m/z < 400$, and mass defect 0.6–0.97 for $m/z < 400$), blank peaks (those present in the respective QM/A filter blank), and ¹³C isotopologue peaks were removed from each mass list prior to formula assignments. Additionally, each set of duplicates was evaluated for common m/z , where only common m/z were used for molecular formula assignments. Each set of duplicates was threshold-corrected for peaks that were below the S/N 3 threshold, but above S/N 2.5 (for example, if a peak was present at S/N 3.1 in one sample, and S/N 2.9 in the second sample, then it was considered common) (Sleighter et al., 2012). This corrects for minor differences in peak magnitude that may cause a peak to go undetected, when it is present at a magnitude

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



slightly below the method detection limit. Each set of duplicates had more than 67 % of the peaks (300–600 m/z) in common, indicating good instrumental and experimental reproducibility (Sleighter et al., 2012).

Molecular formulas were assigned to m/z common to duplicates using an in-house MatLab (The Math Works, Inc., Natick, MA) code according to the criteria $^{12}\text{C}_{5-50}$, $^1\text{H}_{5-100}$, $^{14}\text{N}_{0-5}$, $^{16}\text{O}_{1-30}$, $^{32}\text{S}_{0-2}$, and $^{31}\text{P}_{0-2}$, where the subscripts indicate the range of atoms allowed in a formula. The assigned formulas were screened to eliminate any chemically unreasonable formulas for naturally occurring organic compounds. The criteria for formula assignments are consistent with published procedures (Stubbins et al., 2010; Wozniak et al., 2008). Most (79–96 %) of the common peaks between 200–800 m/z could be assigned a formula. A large majority (> 90 %) of the formulas are within 0.5 ppm agreement of the measured m/z , and all formulas are within 1 ppm error.

3 Results and discussion

3.1 NMR spectroscopy

One of the initial concerns for this comparison was to determine whether the organic solvents chosen were as efficacious as water for removing OM from aerosol particulates. We resorted to ^1H NMR spectroscopy to evaluate the relative proportion of OM extracted by water and pyridine. Pyridine has been demonstrated to extract significant quantities of macromolecular OM from natural samples that have distinctly low water-solubility (e.g., coals, soils, kerogen, etc.) (McKee and Hatcher, 2010; Salmon et al., 2011; Wu et al., 2003). Extracts of pyridine lend themselves well to recovery estimates using ^1H NMR. By integrating the peaks in the main resonance absorption region of the spectrum of aerosol PSOM, between 0.1 and 4.4 ppm, we determined the area response from the sample and used the glucose response factor (area per mass unit of hydrogen) to determine the amount of carbon present in the PSOM. The starting mass of carbon for each aerosol is known, and the percent of extractable carbon, and there-

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fore extractable OM, could be determined. We determined that 31–59 % of the aerosol OM is soluble in pyridine, which is comparable to what is found for WSOM (54–60 %). We recognize that assumptions were made in order to determine these values, and that OM solubility in pyridine will vary with sample type; however, we are confident in these values due to the matched signal response in both PSOM and WSOM samples.

Acetonitrile has been used in the characterization of laboratory-generated aerosol OM in previous studies (Bateman et al., 2008, 2010; Heaton et al., 2009; Laskin et al., 2010; Reinhardt et al., 2007) and is examined by FTICR-MS in this study for this reason. We did not evaluate the acetonitrile extraction efficiency by NMR because this solvent displays its main signal in the 0.1 to 4.4 ppm region and a comparison like the one made with pyridine was not feasible. However, we can speculate that the efficiency is comparable to that of water and pyridine, considering its relative polarity as a solvent.

While the primary motivation for obtaining ^1H NMR spectra was to evaluate extraction efficiencies, the information contained therein is valuable for bulk characterization. A more detailed structure characterization is beyond the scope of this manuscript but will be the subject of future work. Figure 1 shows the ^1H NMR spectra for the WSOM and PSOM fractions for one of the aerosol samples (collected 25–26 June 2013) and the table inset gives the chemical shifts and average relative intensities (for all three aerosol samples) and standard deviations. Both spectra are dominated by aliphatic signals if one does not consider the strong resonances in the aromatic region of the PSOM spectrum assigned to pyridine protons. The peak positions are different for PSOM giving the impression that functionalized structures in WSOM are different from those of PSOM. For example, the methylenic peak (CH_2) in WSOM spectrum falls between 1.1–1.2 ppm, and is between 1.2–1.5 in the PSOM spectrum. However, it is likely that we are observing peak shifts due to solvent interactions, and some of the peaks in the PSOM spectrum are shifted downfield with respect to the WSOM spectrum. A majority of the signal is due to aliphatic hydrogen in both spectra. However, this group is proportionally of greater intensity (73.9 % of spectral intensity) in the PSOM spectrum relative to all other resonances. Methyl protons (CH_3 at 0.7–0.8 ppm for WSOM and

0.8–0.9 ppm for PSOM) are nearly in the same relative proportions to methylenic (CH₂ at 1.1–1.2 ppm for WSOM and 1.2–1.5 ppm for PSOM) protons in both extracts (a ratio of 4 for both WSOM and PSOM). This suggests aliphatic structures of similar chain lengths.

Additional differences between the WSOM and PSOM spectra are found in the regions downfield of 1.7 ppm, regions typically associated with protons near electron withdrawing functional groups. The WSOM extract shows higher relative peak areas in this region, evidence for a greater relative abundance of these types of resonances. Protons found in this region make up 41.8% of the spectral area for WSOM and 25.4% of the spectral area for PSOM. The peaks between 2.0 and 2.8 ppm in both spectra are characteristic of hydrogen attached to a carbon alpha to an unsaturated carbon (H-C-C=), which represent alkenes and carbonyl groups. The relative abundance is higher in the WSOM than PSOM (33.5% and 21.6%, respectively). This higher abundance suggests that WSOM is more selective for unsaturated compounds, which absorb light (Andreae and Gelencsér, 2006), and WSOM may contribute more light absorption on a per carbon basis than PSOM. The broad peak between 3.2 and 4.0 ppm in the WSOM spectrum is attributed to hydrogen adjacent to carbons attached to oxygen and nitrogen which could represent polyols, carbohydrate-like materials, or peptide-like materials. Their relative abundance in the PSOM spectrum is less than in the WSOM spectrum (3.8% vs. 8.3% spectral intensity) demonstrating that the two solvents extract chemically distinct portions of aerosol OM. The NMR spectral information thus show a greater preponderance of signals from protons near functionalized structures and unsaturated carbons in the WSOM and a higher relative signal for aliphatic protons in the PSOM.

3.2 Mass spectra and molecular formula characteristics

The ESI-FTICR mass spectra for ambient aerosol WSOM are comparable to previously published FTICR-MS spectra of aerosol WSOM extracts (Lin et al., 2012; Mazzoleni et al., 2012; Schmitt-Kopplin et al., 2010; Wozniak et al., 2008) averaging thousands of peaks across a broad range of 200–800 *m/z*; the PSOM and ASOM also contained

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thousands of peaks across the same mass range. There are clear spectral differences between the WSOM and the solvent extracts (i.e., PSOM and ASOM), including differences in the presence of some peaks, as well as the relative magnitudes of common peaks (Fig. 2).

Figure 2 shows a WSOM, PSOM, and ASOM spectra for one of the aerosol samples (collected 16–17 August 2011) expanded at a single nominal mass ($m/z = 427$). This distribution of peak intensities is representative of odd nominal masses across the spectral range. Each of the peaks with S/N of at least 3 is labeled with colored shapes to denote the elemental makeup of the assigned molecular formula. There are a few peaks that were not assigned a molecular formula because no chemically reasonable formula was possible using the given criteria suggesting these molecular formulas contain elements other than C, H, N, O, S, and/or P. One of the striking differences is the presence of more peaks at the low mass defect region (below 427.1 m/z) in the PSOM (Fig. 2b). Peaks in this region are either deficient in hydrogen or rich in oxygen, which suggests that PSOM contains either more unsaturated formulas or more highly oxygenated species than does the WSOM. Those specific peaks at 427.05294 m/z and 427.05518 m/z represent CHOS compounds with high O/C ratios (> 0.4) and moderate H/C ratios (1.0–1.5), indicating that the selectivity is likely due to the compounds being more highly oxygenated rather than hydrogen deficient. In addition to peak presence and absence, there is a difference in the peak intensity distributions between the three solvents. The most intense peak in the WSOM (Fig. 2a) is in the center of all peaks for that nominal mass at 427.19723 m/z , and has been assigned as a formula containing only CHO ($C_{21}H_{32}O_9$). This peak is present in both ASOM and PSOM, but the relative intensity is considerably lower. The most intense peak (427.09168 m/z) in the ASOM and PSOM is a CHOS formula ($C_{15}H_{24}O_{12}S$) located at a lower mass defect than the most intense peak in the WSOM spectrum. This peak is also present in WSOM, but at a much lower relative intensity. These differences in relative magnitude of peaks and the presence and absence of some peaks reflect the differences in the ability of these three solvents to extract and detect certain compounds. Differences in

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



solvent composition have not been evaluated to determine how the matrix affects ESI efficiency, so magnitude-weighted contributions are not considered in this study.

A majority of the detected peaks in each extract were assigned molecular formulas within 0.5 ppm error. All of the assigned formulas present in any of the three aerosol samples were combined for each of the three solvents to generate master lists containing every molecular formula assigned to each WSOM, PSOM, and ASOM. Additionally, the master lists for the three solvents were combined to evaluate the molecular properties of the aerosol sample as a whole. The master lists for each solvent were compared for distinct molecular characteristics. The molecular characteristics for each solvent are arranged by their elemental makeup (e.g., number and percentage of CHO formulas) and numerical averages for O/C, H/C and a modified aromaticity index (AI_{mod} , Eq. 1), shown in Table 1. For simplicity and due to low number frequency, all formulas containing phosphorus (CHOP, CHONP, and CHOSP) have been grouped together and reported as CHOP(N,S). AI_{mod} is a metric established by Koch and Dittmar (2006) that estimates the degree of aromaticity of an organic compound using only its molecular formula. AI_{mod} is calculated as follows:

$$AI_{\text{mod}} = (1 + C - 0.5O - S - 0.5H)/(C - 0.5O - S - N - P), \quad (1)$$

for any molecular formula comprised of C, H, N, O, S, and P. AI_{mod} is similar to other aromaticity metrics, such as carbon normalized double bond equivalent (Hockaday et al., 2006), and it includes heteroatoms as points of unsaturation and conservatively assumes that half of all O participate in a double bond, such as that of a carbonyl group. A low AI_{mod} indicates a low degree of aromaticity where a value of zero is an aliphatic compound, a value between 0–0.5 is representative of olefinic compounds (containing at least one double bond) and includes alicyclic molecules. A high AI_{mod} indicates a higher degree of aromaticity where a compound having a value above 0.5 is aromatic, and a value of 0.67 or higher indicates condensed aromatic compounds (fused aromatic rings). Aromatic and condensed aromatic compounds play an important role in the light-absorbing ability of organic aerosols.

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Before analyzing the molecular properties of the individual solvents, every molecular formula (5106 formulas total) assigned to any of the three solvents was combined to evaluate the aerosol samples as a whole. The Venn diagram shown in Fig. 3a shows the percentage of the total formulas unique to each of the solvents, and the ones found in one or more of the solvents. Of all the formulas, WSOM analyzes the largest fraction (3396, 67 %). There are 3152 formulas (62 %) identified in ASOM, and 2397 formulas (47 %) found in PSOM. There are many formulas common between WSOM and the organic solvent extracts, where 1077 formulas (21 %) are found in all three solvents, 1367 (27 %) are common with PSOM, and 2072 (41 %) are common with ASOM. There are 1710 formulas (33 %) found in the PSOM and/or ASOM spectra that are not present in WSOM spectra (Fig. 3a). The formulas present in ASOM and/or PSOM, but not in WSOM, are representative of compounds that are either (1) not water-soluble or (2) outside the analytical window of WSOM using ESI-FTICR-MS. Because of this, we refer to these compounds as “water-insoluble” organic matter (WIOM). One thing to note is that 67 % of the formulas are found in WSOM suggesting that WSOM is more complex molecularly than WIOM despite extracting similar amounts of material. This is likely due, in part, to the poor ionization efficiency of aliphatic material and compounds low in oxygen. The ^1H NMR of the PSOM shows us that pyridine is selective for more aliphatic compounds and compounds that are not extensively substituted with polar functional groups.

The 5106 individual formulas assigned to the three solvents are made up of 2051 CHO (40 %), 1472 CHOS (29 %), 857 CHON (17 %), 599 CHONS (12 %), and 127 CHOP(N,S) (2.5 %, Fig. 3b). Most (> 95 %) of the detected compounds are classified as either aliphatic (40 %, 2043 formulas) or olefinic (56 %, 2837 formulas) based on their low Al_{mod} values (Fig. 3c). Both anthropogenic and biogenic emission sources release OM that fall under the classification of aliphatic and olefinic such as alkanes, alkenes, alkanolic acids, alkenolic acids, alkanals, alkenals, and terpenes (Rogge et al., 1993a, b, c), which can act as precursors to SOA. The polar compounds are formed through atmospheric processing reactions (i.e., photooxidation or reaction with NO_x

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and SO_x) with these biogenically and anthropogenically released precursor molecules (Alfarra et al., 2006; Andreae and Gelencsér, 2006; Jacobson, 2000). Less than 5% of all formulas are classified as aromatic (3.6%, 183 formulas) or condensed aromatic (0.8%, 43 formulas) based on their high AI_{mod} . The low abundance of these aromatic species identified in these samples suggests only small contributions from combustion sources, which are known sources of carbonaceous aromatic compounds such as black carbon, oxygenated aromatic compounds, and polycyclic aromatic hydrocarbons. The sampling site from which these aerosols were collected is influenced by local combustion sources (i.e., coal-fired power plant and light vehicular traffic), but not to a large extent. This observation is supported by a previous study of aerosol OM from a similar rural site located 36 km north of our sampling site, and they show that aromatic compounds made up a quantitatively small (less than 1% on average) component of the total organic carbon and that only half of that aromatic OM was derived from fossil sources (Wozniak et al., 2012b). The small amount of aromatic material in the WSOM sample detected by FTICR-MS is supported by the lack of intensity ($< 3\%$ of total spectral intensity) of peaks that represent aromatic protons (chemical shift 6.5–8.2 ppm) in the ^1H NMR spectra (Fig. S1a). Unfortunately, aromatic protons from the aerosol OM are not distinguishable from the exchanged protons in the pyridine- D_5 ^1H NMR spectra (Fig. S1b); therefore, no conclusion can be made about the amount of aromatic compounds in the PSOM based on ^1H NMR.

The WSOM mass spectra contain the greatest number of molecular formulas (3396), which are dominated by 1563 CHO (46%) and 868 CHOS (26%) formulas followed by CHON (671 formulas, 20%), CHONS (214 formulas, 6.3%), and CHOP(N,S) (80 formulas, 2.4%), as displayed in Fig. 3b. This distribution of molecular formulas, specifically the dominance by CHO formulas, is consistent with other ambient aerosol samples collected in the eastern United States (Wozniak et al., 2008). There are more CHOS than CHON formulas, suggesting that sulfur species (e.g., SO_4^{2-}) were important to the atmospheric processes in this region at the time of sampling.



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The PSOM contains the fewest number of total formulas (2397); however, the molecular characteristics of PSOM are distinct from that of WSOM or ASOM. PSOM contains mostly CHOS (976 formulas, 41 %) and CHO (613 formulas, 26 %) formulas, followed by CHONS (489 formulas, 20 %), CHON (294 formulas, 12 %), and CHOP(N,S) (25 formulas, 1.0 %) formulas (Fig. 3b). Nearly half of the PSOM formulas (1030 formulas, 43 %) are not found in WSOM (PSOM-WIOM), indicating that they are either truly water-insoluble compounds or outside the analytical window of ESI-FTICR-MS, possibly due to poor ionization efficiency. The PSOM-WIOM is dominated by sulfur-containing species, 442 CHOS (43 %) and 311 CHONS (30 %) formulas, with smaller contributions from CHO (155 formulas, 15 %), CHON (101 formulas, 9.8 %), and CHOP(N,S) (21 formulas, 2.0 %) compounds (Table 1). This sulfur predominance suggests that pyridine is more selective towards organic compounds with sulfur as part of its molecular structure. A large majority (88 %) of the sulfur formulas in the PSOM-WIOM have O/S ratios of at least 4, suggesting that many of these formulas represent organosulfates and nitrooxyorganosulfates.

The ASOM fraction contains 3152 formulas, and 1080 of those formulas were not assigned to peaks in the WSOM spectra. Like WSOM, the whole ASOM sample is dominated by CHO (1367 formulas, 43 %) and CHOS (1038 formulas, 33 %) compounds, followed by CHON (419 formulas, 13 %), CHONS (257 formulas, 8.2 %) and CHOP(N,S) (71 formulas, 2.3 %, Fig. 3b). The formulas present in ASOM, but not in WSOM (i.e., ASOM-WIOM), shows a similar atomic distribution (Table 1) with CHO (428 formulas, 40 %) and CHOS (387 formulas, 36 %) formulas being most abundant, but there are more CHONS (136 formulas, 13 %) than CHON (101 formulas, 9.4 %) formulas.

The relative distributions of formulas for each compound class based on molecular structure (as determined by AI_{mod} values as described above) is shown in Fig. 3c. WSOM and ASOM contain mostly olefinic compounds (60 % in each case), and have significant contributions from aliphatic compounds (36 % and 37 %, respectively). Contrarily, PSOM is dominated by aliphatic formulas (58 %) with a smaller amount of olefinic

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0.2 and 0.8 and H/C ratios between 1.00 and 1.75. The formulas in this region above an H/C ratio of 1.50 have previously been attributed to peptides; however, 97 % of these formulas have only one nitrogen and cannot truly be peptides because multiple nitrogen atoms would be required. Additionally, all of these CHON formulas have an O/N ratio of at least 3, which suggests that the functionality of the nitrogen may be that of a nitrate group (NO₃). We also recognize the possibility of reduced nitrogen functional groups (e.g., amines and azo compounds) present in molecules containing other ionizable functional groups (e.g., carboxylic acid), but we are unable to differentiate the two possibilities with this method of analysis. WSOM formulas containing both sulfur and nitrogen (CHONS, Fig. 4c) are predominantly aliphatic with relatively high O/C ratios (> 0.5). Most (93 %) of these formulas have sufficient oxygen atoms (≥ 7) to contain at least one nitrate and one sulfate functional group, indicating that these formulas could be nitrooxyorganosulfates that have been identified in previous ambient aerosol samples (e.g., Surratt et al., 2007). CHOS formulas are the second most dominant formula type in the WSOM, and these formulas separate into two distinct regions of the van Krevelen diagram (Fig. 4d). One region contains mostly aliphatic and some olefinic formulas with O/C ratios greater than 0.25 and H/C ratios greater than 1.3. All of the CHOS formulas in this region have O/S of at least 4, suggesting that they are saturated organosulfates or organosulfates with a few double bonds. The other distinct CHOS region contains olefinic and aromatic formulas with a low O/C (< 0.4) and H/C ratios less than 1.4. Most (70 %) of these formulas have O/S ratios of at least 4, indicating that they could be aromatic organosulfates or organosulfates with multiple unsaturations. The CHOS formulas that do not have sufficient O atoms (O/S < 4) to be organosulfates must contain a reduced sulfur functional group (e.g. sulfonates and thiols). Organic compounds with reduced forms of sulfur, specifically thia arenes, have been identified in anthropogenic aerosol emissions sources and are known to be toxic (Eastmond et al., 1984; Rogge et al., 1993c). Sulfonates are ubiquitous in detergents and personal care products (Debelius et al., 2008; Lara-Martín et al., 2006), and have been previously identified in atmospheric OM (Altieri et al., 2009).

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As stated previously, PSOM is able to dissolve or detect 1030 compounds that WSOM does not (Table 1). The characteristics for all formulas in PSOM-WIOM indicate a lower average O/C (0.29) than WSOM (0.46), a higher average H/C (1.54) than WSOM (1.45), and lower average AI_{mod} (0.11) than WSOM (0.15, Table 1). The CHO formulas found in the PSOM-WIOM plot in a region of higher H/C and lower O/C than those identified in the WSOM (Fig. 4e), implying that they are, on average, more aliphatic and less oxidized than the CHO compounds found in WSOM. The CHON compounds in PSOM-WIOM show variable H/C and O/C ratios (Fig. 4f). Most of these compounds are either aliphatic or olefinic with O/N ratios of at least 3, indicating they may be organonitrates which have been identified previously in aerosol OM (e.g., LeClair et al., 2012). The CHONS formulas make up a large portion of the PSOM-WIOM formulas (30%) and they plot into two distinct regions on the van Krevelen diagram (Fig. 4g). The first region contains a majority of the formulas, and they are largely aliphatic molecules with O/C ratios greater than 0.5 and H/C ratios greater than 1.25, and the second region contains mostly aromatic formulas with O/C ratios less than 0.4 and H/C ratios less than 1.25. Many (70%) of the CHONS formulas contain enough oxygen atoms to contain at least one sulfate and one nitrate functional group, and are potentially nitrooxyorganosulfates. Most (81%) of the compounds in the high O/C and high H/C region contain sufficient oxygen, but only a few (6%) of the formulas in the low O/C and low H/C region have sufficient oxygen to be nitrooxyorganosulfates. The CHONS formulas that do not have sufficient oxygen to form nitrooxyorganosulfates must contain at least one sulfur or nitrogen atom present in a reduced form. CHOS formulas make up the largest fraction (43%) of PSOM-WIOM, and the bulk (96%) of those formulas are aliphatic or olefinic. Like WSOM, the CHOS formulas in PSOM-WIOM plot in two distinct regions of the van Krevelen diagram (Fig. 4h). The first region contains mostly aliphatic formulas spanning the entire O/C range between 0.03 and 1.2 and have H/C ratios greater than 1.2. The second region contains mostly olefinic and aromatic compounds with O/C ratios less than 0.5 and H/C ratios less than 1.5. The average O/C ratio for CHOS compounds in PSOM-WIOM is nearly the same as

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that of the WSOM (0.49 ± 0.31 and 0.47 ± 0.23 , respectively), but the standard deviation for PSOM-WIOM is greater. This larger deviation indicates that the PSOM-WIOM CHOS compounds are more diverse than those in WSOM. Most (80 %) of the CHOS formulas in the PSOM-WIOM have O/S ratios of at least 4, suggesting organosulfates.

Formulas with O/S ratios less than 4 indicate the presence of a more reduced form of sulfur within the structure. The average AI_{mod} of compounds with more reduced forms of sulfur is much greater than that of compounds with O/S sufficient to form organosulfates (0.38 and 0.04, respectively). The major difference between PSOM-WIOM and WSOM is the increased detection of organosulfates and nitrooxyorganosulfates in the PSOM-WIOM, suggesting that pyridine is a more suitable solvent than water for detecting these compounds in aerosol OM using ESI-FTICR-MS.

In spite of having many general molecular formula similarities to WSOM, the characteristics for all formulas in ASOM-WIOM (1080) indicate a lower average O/C (0.39) than WSOM (0.43), a higher average H/C (1.53) than WSOM (1.45), and an identical average AI_{mod} (0.15). These characteristics suggest that the formulas unique to ASOM are less oxygenated and/or have longer carbon chains. This is clearly shown by the many formulas that plot on the left side (low O/C) of the van Krevelen diagram for ASOM-WIOM (Fig. 3i–l). The average number of carbon atoms per formula is slightly larger at 24 carbon atoms for ASOM-WIOM and 22 carbon atoms for WSOM. Both ASOM-WIOM and WSOM are dominated by CHO formulas, but the CHO compounds in the ASOM-WIOM are localized to the upper left corner (low O/C and high H/C) of the diagram (Fig. 4i). Essentially all (99 %) of these formulas are aliphatic or olefinic. There are a small number of CHON formulas (101) found in ASOM-WIOM, and most of these formulas have higher H/C (mean value of 1.66), and are split between a lower O/C (< 0.6) and a high O/C (> 0.8). Nearly all (97 %) of the CHON formulas have an O/N ratio greater than 3 indicating the possibility of these compounds to contain nitrate as part of their structure. The CHONS formulas are also relatively scarce, and they plot in two separate areas on the van Krevelen diagram (Fig. 4k). Roughly half of the formulas plot below O/C ratio of 0.4 and the other half above 0.4. All of the formulas with the

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



higher O/C have sufficient oxygen atoms (at least 7) to form at least one nitrate and one sulfate group. However, only 24 % (17 of 70) of the CHONS formulas in the lower O/C region have enough oxygen atoms to form nitrooxyorganosulfates, which indicates the presence of more reduced N- or S-containing functional groups. Like WSOM, CHOS formulas are the second most abundant molecule type in the ASOM-WIOM comprising 387 of the 1080 formulas (36 %). These formulas are split into two regions of the diagram (Fig. 4I), where one region is dominated by lower O/C and H/C ratios and contains mostly olefinic and aromatic compounds. The other region is predominantly aliphatic compounds covering the full range of O/C ratios. A majority (76 %) of all of the CHOS formulas have sufficient oxygen to form organosulfates ($O/S \geq 4$). All of the formulas in the higher H/C and aliphatic region contain sufficient oxygen to form organosulfates, but more than half (55 %) of the formulas in the low H/C and low O/C region have O/S ratios less than 4. These formulas in the low O/C and low H/C region have multiple unsaturations and have more reduced forms of sulfur in their structure. The major differences between ASOM-WIOM and WSOM are in the high H/C and low O/C region, suggesting that acetonitrile is a more suitable solvent than water for detecting less polar aerosol OM compounds (i.e., fewer oxygen and heteroatoms and/or larger carbon chains) by ESI-FTICR-MS.

The WIOM compounds have chemical characteristics distinct from those in WSOM. Both organic solvents extracted compounds that were more aliphatic in nature than those found in the WSOM, as indicated by the lower O/C ratios and longer carbon chain lengths (for ASOM) of the CHO formulas. Previous work shows that aliphatic components of aerosol WSOM are quantitatively small, implying that they are, as expected, largely water-insoluble (Wozniak et al., 2012b). Radiocarbon signatures indicate that the water-insoluble components are mostly fossil-derived (Wozniak et al., 2012b), but can also come from contemporary plant material (Rogge et al., 1993b). The aliphatic and olefinic hydrocarbon material that is released during fossil fuel combustion can be functionalized through various atmospheric oxidation reactions, thus, increasing its polarity and water-solubility.

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nearly 75 % of the formulas unique to PSOM include sulfur as part of the molecular formula, indicating a potential selectivity for organosulfates and nitrooxyorganosulfates. Sulfur species, especially SO_x , are well known to play important role in atmospheric aging reactions. Organosulfates are formed in the atmosphere through the acid-catalyzed ring-opening reaction of epoxides in the presence of acidic sulfate seed aerosols (Minerath and Elrod, 2009), and these organosulfates can undergo nighttime NO_3 radical oxidation and photo-oxidation to form nitrooxyorganosulfates (Surratt et al., 2008). These compounds have been identified in ambient atmospheric OM (e.g., Altieri et al., 2012; Mazzoleni et al., 2012; Mitra et al., 2013; Reemtsma et al., 2006; Schmitt-Kopplin et al., 2010; Wozniak et al., 2008). Sulfate is emitted from numerous anthropogenic and biogenic sources and is ubiquitous in the atmosphere. The aerosol OM at this sampling site was in proximity to a known SO_x emission source (coal-fired power plant), providing ample opportunity for atmospheric aging reactions with sulfate to occur and form the observed organosulfate compounds. Organosulfates are very polar in nature which may increase the ability of aerosol OM to act as cloud condensation nuclei and, therefore, have an indirect radiative effect (Hallquist et al., 2009).

4 Conclusions

Pyridine and acetonitrile are suitable solvents for analyzing organic aerosols using ESI-FTICR-MS and identify a molecularly-unique portion of aerosol OM when compared to the water-soluble fraction. While the water-soluble fraction is of paramount importance due to the implications those compounds have on environmental processes, such as cloud condensation nuclei formation and mobility in watersheds, analysis of only the WSOM omits a large portion of OM (as much as 90 %). The PSOM and ASOM fractions are expected to be more lipophilic and are likely to be more interactive with biological tissues in the environment. Acetonitrile was able to analyze several unique water-insoluble molecular formulas; however, the elemental distributions and formula types (e.g., aliphatic) of these compounds were very similar to those of WSOM. Pyridine elu-

cidated a molecularly unique and, therefore, complementary set of chemical formulas than those in either ASOM or WSOM. PSOM has a stronger preference for extracting or analyzing sulfur-containing formulas, which are important aerosol components due to their abundance and hygroscopicity which allows them to act as cloud condensation nuclei and impact climate via the indirect effect. Because WIOM has been found to contain the majority of fossil fuel derived OM, analysis of the WIOM, such as OM found in ASOM and PSOM, provides clues to the molecular composition of the fossil material present in aerosols emitted from anthropogenic sources and therefore, insights into its potential fates and impacts in the environment.

Supplementary material related to this article is available online at <http://www.atmos-chem-phys-discuss.net/14/10393/2014/acpd-14-10393-2014-supplement.pdf>.

Acknowledgements. The work presented in this manuscript was funded by the Frank Batten endowment to P.G. Hatcher. A.S. Willoughby was funded by the US EPA STAR graduate fellowship program, Grand FP 91736001. The EPA has not officially endorsed this publication and the views expressed herein may not reflect the views of the EPA. The authors acknowledge assistance from the ODU COSMIC facility for FTICR-MS and NMR analyses. We thank Rachel Sleighter for useful discussions about the data.

References

Alfarra, M. R., Paulsen, D., Gysel, M., Garforth, A. A., Dommen, J., Prévôt, A. S. H., Worsnop, D. R., Baltensperger, U., and Coe, H.: A mass spectrometric study of secondary organic aerosols formed from the photooxidation of anthropogenic and biogenic precursors in a reaction chamber, *Atmos. Chem. Phys.*, 6, 5279–5293, doi:10.5194/acp-6-5279-2006, 2006.

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Altieri, K. E., Turpin, B. J., and Seitzinger, S. P.: Oligomers, organosulfates, and nitrooxy organosulfates in rainwater identified by ultra-high resolution electrospray ionization FT-ICR mass spectrometry, *Atmos. Chem. Phys.*, 9, 2533–2542, doi:10.5194/acp-9-2533-2009, 2009.
- 5 Altieri, K. E., Hastings, M. G., Peters, A. J., and Sigman, D. M.: Molecular characterization of water soluble organic nitrogen in marine rainwater by ultra-high resolution electrospray ionization mass spectrometry, *Atmos. Chem. Phys.*, 12, 3557–3571, doi:10.5194/acp-12-3557-2012, 2012.
- Andreae, M. and Crutzen, P.: Atmospheric aerosols: biogeochemical sources and role in atmospheric chemistry, *Science*, 276, 1052–1058, 1997.
- 10 Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, doi:10.5194/acp-6-3131-2006, 2006.
- Bateman, A. P., Walser, M. L., Desyaterik, Y., Laskin, J., Laskin, A., and Nizkorodov, S. A.: The effect of solvent on the analysis of secondary organic aerosol using electrospray ionization mass spectrometry, *Environ. Sci. Technol.*, 42, 7341–7346, 2008.
- 15 Bateman, A. P., Nizkorodov, S., Laskin, J., and Laskin, A.: High-resolution electrospray ionization mass spectrometry analysis of water-soluble organic aerosols collected with a particle into liquid sampler, *Anal. Chem.*, 82, 8010–8016, 2010.
- 20 Bond, T., Doherty, S., Fahey, D., Forster, P., Berntsen, T., DeAngelo, B., Flanner, M., Ghan, S., Kärcher, B., and Koch, D.: Bounding the role of black carbon in the climate system: a scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 1–173, 2013.
- Booth, B. B., Dunstone, N. J., Halloran, P. R., Andrews, T., and Bellouin, N.: Aerosols implicated as a prime driver of twentieth-century North Atlantic climate variability, *Nature*, 484, 228–232, 2012.
- 25 Crutzen, P. J. and Andreae, M. O.: Biomass burning in the tropics: impact on atmospheric chemistry and biogeochemical cycles, *Science*, 250, 1669–1678, 1990.
- Debelius, B., Forja, J., Del Valls, A., and Lubián, L.: Effect of linear alkylbenzene sulfonate (LAS) and atrazine on marine microalgae, *Mar. Pollut. Bull.*, 57, 559–568, 2008.
- 30 Decesari, S., Mircea, M., Cavalli, F., Fuzzi, S., Moretti, F., Tagliavini, E., and Facchini, M.: Source attribution of water-soluble organic aerosol by nuclear magnetic resonance spectroscopy, *Environ. Sci. Technol.*, 41, 2479–2484, 2007.

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Dittmar, T., Koch, B., Hertkorn, N., and Kattner, G.: A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater, *Limnol. Oceanogr.-Meth.*, 6, 230–235, 2008.
- 5 Eastmond, D. A., Booth, G. M., and Lee, M. L.: Toxicity, accumulation, and elimination of polycyclic aromatic sulfur heterocycles in *Daphnia magna*, *Arch. Environ. Con. Tox.*, 13, 105–111, 1984.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- 10 Heaton, K., Sleighter, R., Hatcher, P., Hall IV, W., and Johnston, M.: Composition domains in monoterpene secondary organic aerosol, *Environ. Sci. Technol.*, 43, 6950–6955, 2009.
- Hockaday, W., Grannas, A., Kim, S., and Hatcher, P.: Direct molecular evidence for the degradation and mobility of black carbon in soils from ultrahigh-resolution mass spectral analysis of dissolved organic matter from a fire-impacted forest soil, *Org. Geochem.*, 37, 501–510, 2006.
- 20 Jacobson, M.: Physically-based treatment of elemental carbon optics: implications for global direct forcing of aerosols, *Geophys. Res. Lett.*, 27, 217–220, 2000.
- Jimenez, J., Canagaratna, M., Donahue, N., Prevot, A., Zhang, Q., Kroll, J., DeCarlo, P., Allan, J., Coe, H., and Ng, N.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.
- 25 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- 30 Kleefeld, S., Hoffer, A., Krivacsy, Z., and Jennings, S.: Importance of organic and black carbon in atmospheric aerosols at Mace Head, on the West Coast of Ireland (53°19' N, 9°54' W), *Atmos. Environ.*, 36, 4479–4490, 2002.

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Koch, B. and Dittmar, T.: From mass to structure: an aromaticity index for high-resolution mass data of natural organic matter, *Rapid Commun. Mass Sp.*, 20, 926–932, 2006.
- Lara-Martín, P. A., Gómez-Parra, A., and González-Mazo, E.: Development of a method for the simultaneous analysis of anionic and non-ionic surfactants and their carboxylated metabolites in environmental samples by mixed-mode liquid chromatography–mass spectrometry, *J. Chromatogr. A*, 1137, 188–197, 2006.
- Laskin, J., Laskin, A., Roach, P., Slysz, G., Anderson, G., Nizkorodov, S., Bones, D., and Nguyen, L.: High-resolution desorption electrospray ionization mass spectrometry for chemical characterization of organic aerosols, *Anal. Chem.*, 82, 2048–2058, 2010.
- LeClair, J. P., Collett, J. L., and Mazzoleni, L. R.: Fragmentation analysis of water-soluble atmospheric organic matter using ultrahigh-resolution FT-ICR mass spectrometry, *Environ. Sci. Technol.*, 46, 4312–4322, 2012.
- Lin, P., Rincon, A. G., Kalberer, M., and Yu, J. Z.: Elemental composition of HULIS in the Pearl River Delta region, China: results Inferred from positive and negative electrospray high resolution mass spectrometric data, *Environ. Sci. Technol.*, 46, 7454–7462, 2012.
- Mayol-Bracero, O., Guyon, P., Graham, B., Roberts, G., Andreae, M., Decesari, S., Facchini, M., Fuzzi, S., and Artaxo, P.: Water-soluble organic compounds in biomass burning aerosols over Amazonia 2. Apportionment of the chemical composition and importance of the polyacidic fraction, *J. Geophys. Res.*, 107, 8091, doi:10.1029/2001JD000522, 2002.
- Mazzoleni, L. R., Ehrmann, B., Shen, X., Marshall, A., and Collett Jr, J.: Water-soluble atmospheric organic matter in fog: exact masses and chemical formula identification by ultrahigh-resolution Fourier transform ion cyclotron resonance mass spectrometry, *Environ. Sci. Technol.*, 44, 3690–3697, 2010.
- Mazzoleni, L. R., Saranjampour, P., Dalbec, M. M., Samburova, V., Hallar, A. G., Zielinska, B., Lowenthal, D. H., and Kohl, S.: Identification of water-soluble organic carbon in non-urban aerosols using ultrahigh-resolution FT-ICR mass spectrometry: organic anions, *Environ. Chem.*, 9, 285–297, 2012.
- McKee, G. A. and Hatcher, P. G.: Alkyl amides in two organic-rich anoxic sediments: a possible new abiotic route for N sequestration, *Geochim. Cosmochim. Ac.*, 74, 6436–6450, 2010.
- Minerath, E. C. and Elrod, M. J.: Assessing the potential for diol and hydroxy sulfate ester formation from the reaction of epoxides in tropospheric aerosols, *Environ. Sci. Technol.*, 43, 1386–1392, 2009.



**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

troscopy: organosulfates as photochemical smog constituents, *Anal. Chem.*, 82, 8017–8026, 2010.

Shakya, K. M., Place, P. F., Griffin, R. J., and Talbot, R. W.: Carbonaceous content and water-soluble organic functionality of atmospheric aerosols at a semi-rural New England location, *J. Geophys. Res.-Atmos.*, 117, D03301, doi:10.1029/2011JD016113, 2012.

Sleighter, R. L. and Hatcher, P.: The application of electrospray ionization coupled to ultrahigh resolution mass spectrometry for the molecular characterization of natural organic matter, *J. Mass Spectrom.*, 42, 559–574, 2007.

Sleighter, R. L., McKee, G., Liu, Z., and Hatcher, P.: Naturally present fatty acids as internal calibrants for Fourier transform mass spectra of dissolved organic matter, *Limnol. Oceanogr.-Meth.*, 6, 246–253, 2008.

Sleighter, R. L., Chen, H., Wozniak, A. S., Willoughby, A. S., Caricasole, P., and Hatcher, P. G.: Establishing a measure of reproducibility of ultrahigh-resolution mass spectra for complex mixtures of natural organic matter, *Anal. Chem.*, 84, 9184–9191, 2012.

Stubbins, A., Raitos, D. E., Hinrichsen, H. H., Douglass, J. G., Rose, A. L., Hedley, J., Finlay, K., Dong, H. P., Archer, S. D., and Blukacz, E.: Illuminated darkness: molecular signatures of Congo River dissolved organic matter and its photochemical alteration as revealed by ultrahigh precision mass spectrometry, *Limnol. Oceanogr.*, 55, 1467–1477, 2010.

Sullivan, A., Weber, R., Clements, A., Turner, J., Bae, M., and Schauer, J.: A method for on-line measurement of water-soluble organic carbon in ambient aerosol particles: results from an urban site, *Geophys. Res. Lett.*, 31, L13105, doi:10.1029/2004GL019681, 2004.

Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., and Jaoui, M.: Evidence for organosulfates in secondary organic aerosol, *Environ. Sci. Technol.*, 41, 517–527, 2007.

Surratt, J. D., Gómez-González, Y., Chan, A. W. H., Vermeylen, R., Shahgholi, M., Kleindienst, T. E., Edney, E. O., Offenberg, J. H., Lewandowski, M., and Jaoui, M.: Organosulfate formation in biogenic secondary organic aerosol, *J. Phys. Chem. A*, 112, 8345–8378, 2008.

Szidat, S., Jenk, T., Gaggeler, H., Synal, H., Fisseha, R., Baltensperger, U., Kalberer, K., Samburova, V., Wacker, L., and Saurer, M.: Source apportionment of aerosols by ¹⁴C measurements in different carbonaceous particle fractions, *Radiocarbon*, 46, 475–484, 2004.

Wozniak, A. S., Bauer, J. E., Sleighter, R. L., Dickhut, R. M., and Hatcher, P. G.: Technical Note: Molecular characterization of aerosol-derived water soluble organic carbon using ultrahigh

**Molecular
characterization of
aerosol WIOM**

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resolution electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry, *Atmos. Chem. Phys.*, 8, 5099–5111, doi:10.5194/acp-8-5099-2008, 2008.

Wozniak, A. S., Bauer, J. E., and Dickhut, R. M.: Characteristics of water-soluble organic carbon associated with aerosol particles in the eastern United States, *Atmos. Environ.*, 46, 181–188, 2012a.

Wozniak, A. S., Bauer, J. E., Dickhut, R. M., Xu, L., and McNichol, A. P.: Isotopic characterization of aerosol organic carbon components over the eastern United States, *J. Geophys. Res.*, 117, D13303, doi:10.1029/2011JD017153, 2012b.

Wu, Z., Jernström, S., Hughey, C. A., Rodgers, R. P., and Marshall, A. G.: Resolution of 10 000 compositionally distinct components in polar coal extracts by negative-ion electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry, *Energ. Fuel*, 17, 946–953, 2003.

Zappoli, S., Andracchio, A., Fuzzi, S., Facchini, M., Gelencser, A., Kiss, G., Krivacsy, Z., Molnar, A., Meszaros, E., and Hansson, H. C.: Inorganic, organic and macromolecular components of fine aerosol in different areas of Europe in relation to their water solubility, *Atmos. Environ.*, 33, 2733–2743, 1999.

Molecular
characterization of
aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Molecular formula distributions for each solvent extract based on elemental composition with number average characteristics. The values for PSOM and ASOM do not include formulas common with WSOM, thus represent characteristics for WIOM.

Solvent Extract	Elemental Composition	# of Formulas	% of Formulas	Number Averages		
				O/C	H/C	Al _{mod}
WSOM	CHO	1563	46%	0.43 ± 0.15	1.43 ± 0.24	0.17 ± 0.14
	CHON	671	20%	0.47 ± 0.23	1.41 ± 0.20	0.16 ± 0.16
	CHONS	214	6.3%	0.71 ± 0.21	1.65 ± 0.20	0.02 ± 0.10
	CHOS	868	26%	0.47 ± 0.23	1.46 ± 0.35	0.15 ± 0.21
	CHOP(N,S)	80	2.4%	0.39 ± 0.09	1.54 ± 0.14	0.09 ± 0.10
	Total	3396	100%	0.46 ± 0.19	1.45 ± 0.27	0.15 ± 0.17
PSOM (WIOM)	CHO	155	15%	0.29 ± 0.20	1.59 ± 0.36	0.13 ± 0.17
	CHON	101	9.8%	0.54 ± 0.32	1.53 ± 0.42	0.19 ± 0.21
	CHONS	311	30%	0.64 ± 0.23	1.52 ± 0.28	0.07 ± 0.18
	CHOS	442	43%	0.49 ± 0.31	1.54 ± 0.38	0.10 ± 0.17
	CHOP(N,S)	21	2.0%	0.49 ± 0.18	1.43 ± 0.50	0.23 ± 0.29
	Total	1030	100%	0.51 ± 0.29	1.54 ± 0.35	0.11 ± 0.18
ASOM (WIOM)	CHO	428	40%	0.25 ± 0.16	1.67 ± 0.24	0.10 ± 0.12
	CHON	101	9.4%	0.48 ± 0.35	1.66 ± 0.42	0.12 ± 0.22
	CHONS	136	13%	0.45 ± 0.25	1.27 ± 0.29	0.25 ± 0.25
	CHOS	387	36%	0.49 ± 0.32	1.42 ± 0.36	0.16 ± 0.20
	CHOP(N,S)	28	2.6%	0.42 ± 0.20	1.49 ± 0.25	0.14 ± 0.19
	Total	1080	100%	0.39 ± 0.28	1.53 ± 0.35	0.15 ± 0.19

Molecular characterization of aerosol WIOM

A. S. Willoughby et al.




Fig. 1. Expanded ^1H NMR spectra between 0–5 ppm for **(a)** WSOM and **(b)** PSOM for the aerosol particulate sample collected 25–26 June 2013. The table insets give the chemical shifts and average relative intensities (normalized to total intensity between 0 and 4.4 ppm) and standard deviations for the three aerosol samples in the major proton regions, including aliphatic (H-C), unsaturated (H-C-C=), and oxygenated aliphatics (H-C-O).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Molecular
characterization of
aerosol WIOM

A. S. Willoughby et al.




Fig. 2. FTICR mass spectra expanded for m/z at a nominal mass of 427 for **(a)** WSOM, **(b)** PSOM and **(c)** ASOM. Peaks with $S/N \geq 3$ have a colored shape above the peak to indicate the elemental makeup of the assigned molecular formula. Blue circles represent CHO formulas, green triangles represent CHOS formulas, orange diamonds represent CHON formulas, red squares represent CHONS formulas, and a black “x” denotes m/z that did not have a formula assigned under the chosen criteria.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Molecular
characterization of
aerosol WIOM

A. S. Willoughby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. The **(a)** Venn diagram showing the relative distributions of all molecular formulas (5106 total) present in any of the three solvents. Percentages in areas of overlap are percentages of molecular formulas that appear in both (or all three) of those samples. Percentages in areas of no overlap are molecular formulas that are unique to that individual sample. The **(b)** histogram of the fractional contributions (%) of molecular formulas from various elemental combinations to the total for all formulas (5106 total), WSOM formulas (3396 total), PSOM formulas (2397 total), or ASOM formulas (3152 total). The **(c)** pie charts showing the fractional contributions (%) of molecular structure classes as determined by AI_{mod} calculations.

Molecular
characterization of
aerosol WIOM

A. S. Willoughby et al.




Fig. 4. van Krevelen diagrams for molecular formulas assigned to the (a–d) WSOM, (e–h) PSOM-WIOM, and (i–l) ASOM-WIOM extracts. Any formula present in WSOM has been removed from the PSOM-WIOM and ASOM-WIOM plots. Each diagram is plotted based on elemental content of each molecular formula (CHO, CHON, CHONS, and CHOS). The labeled regions in (a) WSOM CHO formulas correspond to their aromaticity based on AI_{mod} and these regions are consistent for all of the diagrams. Formulas above the solid line are aliphatic, just below the solid line are olefinic, below the dashed line are aromatic, and below the dotted line are condensed aromatic.