



Using the chemical equilibrium partitioning space

F. Wania et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Using the chemical equilibrium partitioning space to explore factors influencing the phase distribution of compounds involved in secondary organic aerosol formation

F. Wania^{1,2}, Y. D. Lei^{1,2,3}, C. Wang^{1,2}, J. P. D. Abbatt², and K.-U. Goss^{4,5}

¹Department of Physical and Environmental Sciences, University of Toronto Scarborough, 1265 Military Trail, Toronto, Ontario, M1C 1A4, Canada

²Department of Chemistry, University of Toronto, Toronto, Ontario, M1C 1A4, Canada

³Department of Chemical Engineering and Applied Chemistry, University of Toronto, Toronto, Ontario, M1C 1A4, Canada

⁴Department of Analytical Environmental Chemistry, Centre for Environmental Research UFZ Leipzig-Halle, Permoserstraße 15, 04318 Leipzig, Germany

⁵Institute of Chemistry, University of Halle-Wittenberg, Kurt-Mothes-Straße 2, 06120 Halle, Germany

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 22 September 2014 – Accepted: 2 October 2014 – Published: 23 October 2014

Correspondence to: F. Wania (frank.wania@utoronto.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

14, 26545–26590, 2014

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Many atmospheric and chemical variables influence the partitioning equilibrium between gas phase and condensed phases of compounds implicated in the formation of secondary organic aerosol (SOA). The large number of factors and their interaction makes it often difficult to assess their relative importance and concerted impact. Here we introduce a two-dimensional space, which maps regions of dominant atmospheric phase distribution within a coordinate system defined by equilibrium partitioning coefficients between the gas phase, an aqueous phase and a water insoluble organic matter (WIOM) phase. Placing compounds formed from the oxidation of *n*-alkanes, terpenes and mono-aromatic hydrocarbons on the maps based on their predicted partitioning properties allows for a simple graphical assessment of their equilibrium phase distribution behaviour. Specifically, it allows for the simultaneous visualization and quantitative comparison of the impact on phase distribution of changes in atmospheric parameters (such as temperature, salinity, WIOM phase polarity, organic aerosol load, and liquid water content), and chemical properties (such as oxidation state, molecular size, functionalization, and dimerisation). The graphical analysis reveals that the addition of hydroxyl, carbonyl and carboxyl groups increases the affinity of aliphatic, alicyclic and aromatic hydrocarbons for the aqueous phase more rapidly than their affinity for WIOM, suggesting that the aqueous phase may often be relevant even for substances that are considerably larger than the C2 and C3 compounds that are typically believed to be associated with aqueous SOA. In particular, the maps identify some compounds that contribute to SOA formation if partitioning to both WIOM and aqueous phase is considered, but would remain in the gas phase if either condensed phase were neglected. For example, many semi-volatile α -pinene oxidation products will contribute to aqueous SOA under the high liquid water content conditions encountered in clouds, but would remain vapours in wet aerosol. It is conceivable to develop parameterizations of “partitioning basis sets” that group compounds with comparable partitioning properties,

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which – when combined with data on the abundance of those groups of compounds – could serve in the simulation of SOA formation.

1 Introduction

Many atmospheric and chemical variables influence the partitioning equilibrium between gas phase and condensed phases of compounds implicated in the formation of secondary organic aerosol (SOA) (Hallquist et al., 2009). For example, the molecular size of the compounds and the functional groups they contain have a strong influence on their preference for either gas phase or the condensed phase(s) making up an aerosol. Atmospheric temperature strongly impacts that phase equilibrium. The concentration and the polarity of the organic particulate matter present in the atmosphere play a role, as do relative humidity and the resultant liquid water content of the atmosphere. The presence of electrolytes such as ammonium sulphate and sodium chloride further modifies the gas phase-aqueous phase equilibrium. How all these variables impact gas/particle partitioning is reasonably well understood and in most cases can even be quantified with some confidence (e.g. Zuend and Seinfeld, 2012; Shiraiwa et al., 2013). Nevertheless, due to the large number of the variables and their interaction, it is often difficult to assess the relative importance of each of these variables and to fully comprehend the concerted impact of all of them acting together. In fact, different variables will be important for different chemicals and under different atmospheric circumstances. For example, lower temperatures shift the gas/aqueous phase equilibrium of volatile substances, such as α -pinene and n -alkanes, to the gas phase, but have the opposite effect for less volatile substances, such as the oxidation products formed from those volatile substances.

Graphical methods illustrating the effect of multiple variables within a common framework can be a powerful tool for understanding their relative importance and concerted impact (e.g. Wong and Wania, 2011). They can also aid in the identification and prioritization of data gaps by indicating for which compounds and which circumstances

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a more precise quantification of the effect of a variable on phase partitioning is required and when approximate knowledge can be tolerated.

Ten years ago, we introduced the chemical partitioning space as a graphical method for furthering the understanding of the concerted impact of multiple variables on the phase distribution of semi-volatile organic trace contaminants, such as polycyclic aromatic hydrocarbons and polychlorinated biphenyls, in the atmosphere (Lei and Wania, 2004). The basic idea of the chemical partitioning space is to calculate and display regions of dominant phase distribution within a coordinate system defined by equilibrium phase partitioning coefficients, and then to place substances on the resulting maps based on their partitioning properties under certain prevailing conditions (e.g. of temperature). This allows for a simple graphical assessment of the substances' equilibrium phase distribution behaviour and in turn of their depositional and reactive fate (Lei and Wania, 2004; Westgate and Wania, 2013). We believe, and seek to demonstrate here, that the chemical partitioning space could also be used to gain a deeper appreciation of the impact of various chemical and atmospheric variables on the phase distribution behaviour of SOA compounds. Because SOA formation strongly depends on the phase distribution behaviour of those substances, this also amounts to an exploration of the impact of those variables on the likelihood and the extent of SOA formation, as well as on the nature of the SOA formed.

The atmospheric chemistry community has a long tradition of using graphical tools to illustrate and advance understanding of atmospheric processes (e.g. Heald et al., 2010; Kroll et al., 2011) and there are precedents of displaying the phase distribution behaviour of SOA compounds as a function of chemical equilibrium partitioning parameters. The Volatility Basis Set (VBS) approach by Donahue and co-workers (2006, 2011, 2012) in both its one-dimensional and two-dimensional manifestations, is sometimes used to indicate the distribution between gas and aerosol phase as a function of organic aerosol load and a compound's saturation concentration, which has been shown to be the reciprocal of a gas-particle partitioning coefficient (Donahue et al., 2006). Similarly, Volkamer et al. (2009) displayed the distribution of glyoxal between

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gas and aqueous phase as a function of an effective Henry's law constant and the atmospheric liquid water content. However, to the best of our knowledge, no attempt has been made so far to similarly display graphically the distribution behaviour of SOA compounds between the gas phase, the atmospheric liquid water phase, and the organic aerosol phase at the same time. Notably, all two-dimensional representations of organic compounds used in the SOA community employ a chemical equilibrium partitioning parameter for at most one of the two dimensions and thus do not have a sole focus on phase distribution. Even if they do consider phase distribution behaviour (e.g. the 2-D VBS, Donahue et al., 2011, 2012), the existence of a liquid water phase in the atmosphere is ignored.

After first introducing the chemical partitioning space for the atmosphere, we explain how to place SOA compounds within the space and how to account for the influence of various variables on that placement. Using the partitioning space we then illustrate the impact of prediction uncertainty, temperature, salting-out effect, phase polarity, liquid water content, organic aerosol load, and chemical ageing (i.e. functionalization and oligomerisation) on the phase distribution of selected oxidation products of *n*-alkanes, biogenic terpenes and aromatic hydrocarbons.

2 Methods

2.1 Defining the chemical partitioning space

The equilibrium partition coefficient of a chemical between two phases X and Y is defined as:

$$K_{X/Y} = \frac{C_X}{C_Y} \quad (1)$$

where C_X and C_Y are the concentrations in phase X and Y in units of mol m^{-3} of phase. $K_{X/Y}$ thus m^3 of phase Y per m^3 of phase X. In a system consisting of the gas phase

(G), an aqueous phase (W) and a water insoluble organic matter phase (WIOM) there are three partition coefficients, which are related to each other according to:

$$K_{W/G} = \frac{K_{W/WIOM}}{K_{G/WIOM}} \quad (2)$$

Two of the three coefficients are therefore sufficient to describe a chemical's relative preferences for the three phases. Two coefficients in logarithmic form can be used to define a two dimensional coordinate system, which we call the chemical partitioning space. In Fig. 1a, $\log K_{W/G}$ and $\log K_{WIOM/G}$ are used to define such a space.

The equilibrium partition coefficient between WIOM and the gas phase ($K_{WIOM/G}$) on the x axis is related to the gas-particle partition coefficient K' in units of $\text{m}^3 \mu\text{g}^{-1}$ (Pankow, 2003) through:

$$K_{WIOM/G} = K' \delta_{WIOM} \frac{10^6 \mu\text{g}}{\text{g}} \quad (3)$$

where δ_{WIOM} is the density of WIOM in units of g m^{-3} . The equilibrium partition coefficient between water and gas phase $K_{W/G}$ on the y axis is related to the Henry's Law constants H in units of $\text{mol Pa}^{-1} \text{m}^{-3}$, or to H' in the non-SI units of M atm^{-1} favoured by the atmospheric chemistry community, through:

$$K_{W/G} = HRT = H'RT \times \frac{1 \text{ atm L}}{101.325 \text{ Pa m}^3} \quad (4)$$

where R is the ideal gas constant ($8.314 \text{ Pa m}^3 \text{ K}^{-1} \text{ mol}^{-1}$) and T is absolute temperature in K. A chemical can be placed in such a coordinate system based on its equilibrium partition coefficients.

Using the chemical equilibrium partitioning space

F. Wania et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.2 Displaying chemical phase distribution in the partitioning space

If a three-phase system consisting of gas, water and WIOM is at chemical equilibrium, the fraction ϕ_X of the total amount of a chemical that is in one of the three phases can be calculated using (Lei and Wania, 2004):

$$\phi_G = 1 / \left(1 + K_{W/G} \frac{V_W}{V_G} + K_{WIOM/G} \frac{V_{WIOM}}{V_G} \right) \quad (5)$$

$$\phi_W = 1 / \left(1 + K_{G/W} \frac{V_G}{V_W} + K_{WIOM/W} \frac{V_{WIOM}}{V_W} \right) \quad (6)$$

$$\phi_{WIOM} = 1 / \left(1 + K_{G/WIOM} \frac{V_G}{V_{WIOM}} + K_{W/WIOM} \frac{V_W}{V_{WIOM}} \right) \quad (7)$$

This means that the chemical's phase distribution depends only on its equilibrium partitioning coefficients and the phase volumes V_X in m^3 . In the atmosphere V_W and V_{WIOM} are negligibly small compared to V_G and the volume ratios V_W/V_G and V_{WIOM}/V_G can be calculated from the liquid water content (LWC) and the organic aerosol load (OAL) in units of $\mu g m^{-3}$ using:

$$\frac{V_W}{V_G} = \text{LWC in } \mu g m^{-3} \times 10^{-6} \frac{g}{\mu g} \times 10^{-6} \frac{m^3}{g} \quad (8)$$

$$\frac{V_{WIOM}}{V_G} = \text{OAL in } \mu g m^{-3} \times 10^{-6} \frac{g}{\mu g} \times 10^{-6} \frac{m^3}{g} \quad (9)$$

We thereby assume that water and WIOM have a density of $10^6 g m^{-3}$.

If we now assume LWC and OAL to be fixed, the phase distribution as calculated in Eqs. (5)–(7) only depends on the partition coefficients and we can display the values of ϕ_G , ϕ_W , and ϕ_{WIOM} within the partitioning space. In Fig. 1a this has been done assuming a LWC of $10 \mu g m^{-3}$ and an OAL of $10 \mu g m^{-3}$. In the red-coloured, upper

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



left of the partitioning space are chemicals with low $K_{W/G}$ and low $K_{WIOM/G}$, i.e. chemical favouring the gas phase over the aqueous phase and WIOM. Accordingly they are predominately in the gas phase (different shades of red are used to indicate areas of the space where ϕ_G exceeds 50, 90 and 99 %). In the blue-coloured lower left of the space are chemicals with high $K_{W/G}$ and low $K_{WIOM/G}$, i.e. highly water-soluble chemicals that favour the gas phase over the WIOM phase and thus partition predominantly into the aqueous phase (different shades of blue indicate areas of the space where ϕ_W exceeds 50, 90 and 99 %). Finally, in the green-coloured upper right of the space are chemicals with low $K_{W/G}$ and high $K_{WIOM/G}$, i.e. sparingly water-soluble substances of low volatility, that partition predominantly into the WIOM phase (different shades of green indicate areas of the space where ϕ_{WIOM} exceeds 50, 90 and 99 %).

The lines between the fields coloured in red, blue and green ($\phi_G = \phi_W = \phi_{WIOM} = 50\%$) indicate the $K_{W/G}$ and $K_{WIOM/G}$ thresholds, where the transition of chemicals from mostly in the gas phase to mostly in the aqueous or WIOM phase occurs (Fig. 1a). The vertical threshold between predominant partitioning into gas and WIOM phase occurs at $\log K_{WIOM/G} = \log(V_G/V_{WIOM})$ and the horizontal threshold between predominant partitioning into gas and aqueous phase at $\log K_{W/G} = \log(V_G/V_W)$. At the assumed LWC and OAL of $10 \mu\text{g m}^{-3}$ this corresponds to $\log K_{W/G}$ and $\log K_{WIOM/G}$ values of 11.

Because of the thermodynamic relationship of Eq. (2), $K_{WIOM/W}$ equals $K_{WIOM/G}$ divided by $K_{W/G}$ and diagonal lines from the upper left to the lower right in the partitioning space designate lines of equal $K_{WIOM/W}$. The diagonal lines in the lower right of Fig. 1a therefore show where the transition from dominant partitioning into WIOM and aqueous phase takes place. If liquid water and WIOM are equally abundant (LWC and OAL is $10 \mu\text{g m}^{-3}$) the threshold falls at $\log K_{WIOM/G} = \log(V_G/V_{WIOM}) = \log 1 = 0$. This threshold could obviously also be calculated from the other two thresholds ($\log K_{WIOM/G} - \log K_{W/G} = 11 - 11 = 0 = \log K_{WIOM/G}$). Finally, the white triangle in the centre of the three coloured areas designates the partitioning properties of chemicals that partition to none of the three phases to more than 50 %, i.e. they are present in equal proportions in all three phases.

Using the chemical equilibrium partitioning space

F. Wania et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Of course, the thresholds are not fixed, but depend on the size of the aqueous and WIOM phase in the atmosphere. OAL in the atmosphere ranges between 0.1 and $100 \mu\text{g m}^{-3}$, with most measurements yielding OAL between 1 and $10 \mu\text{g m}^{-3}$ (Jimenez et al., 2009). In Fig. 1b, broken vertical lines indicate where the threshold between predominant partitioning into gas and WIOM phase would fall, if OALs of 0.1, 1, and $100 \mu\text{g m}^{-3}$ prevailed. With each order of magnitude increase in the OAL, the threshold moves one log unit to the left, i.e. more volatile chemicals (i.e. with lower $\log K_{\text{WIOM/G}}$) will shift their equilibrium phase distribution from the gas to the WIOM phase. Similarly, the broken horizontal lines in Fig. 1b indicate the threshold between predominant partitioning into gas and aqueous phase, if larger or lower volumes of water were associated with the aerosol, i.e. at LWCs of 100 or $1 \mu\text{g m}^{-3}$, respectively. With each order of magnitude increase in the LWC, the threshold moves one log unit up, i.e. more volatile chemicals (i.e. with lower $\log K_{\text{W/G}}$) will shift their phase distribution from the gas to the aqueous phase.

The amount of liquid water in an aerosol depends on temperature, relative humidity and on the aerosol concentration and type, i.e. its hygroscopicity. A range of LWC between 1 and $100 \mu\text{g m}^{-3}$ was deemed plausible based on calculations with the E-AIM model (Wexler and Clegg, 2002). At high relative humidity and if the aerosol can act as cloud condensation nuclei, water starts to condense to a much greater extent and the LWC will adopt values that are more than 5 orders of magnitude higher. Figure 1c thus maps ϕ_{G} , ϕ_{W} , and ϕ_{WIOM} for a LWC of 0.3g m^{-3} ($V_{\text{W}}/V_{\text{G}} = 3 \times 10^{-7} \text{m}^3 \text{m}^{-3}$), which is typical for conditions within a cloud. The horizontal threshold moves sharply higher to a $\log K_{\text{W/G}}$ of 6.5, which means that at equilibrium many more chemicals will partition into the aqueous phase of a cloud, than in an aerosol under clear sky conditions (Fig. 1b). We note that the left side of the partitioning space plots is similar to a figure illustrating the partitioning of water-soluble molecules to cloud, fog and aerosol liquid water presented previously (Fig. 8 in Volkamer et al., 2009).

In atmospheric chamber experiments conducted at low relative humidity, the liquid water content of aerosol may be negligibly low. In that case the fraction in the aqueous

phase ϕ_W is also negligible, and the blue field disappears from the partitioning space altogether (Fig. 1d), or rather the vertical threshold moves so far down to no longer fall within the range of displayed $K_{W/G}$ values. During some chamber experiments, OAL exceed levels observed in ambient air, (e.g. $1000 \mu\text{g m}^{-3}$) and the vertical threshold can move far to the left, i.e. to a $\log K_{W_{IOM}/G}$ of around 9 (Fig. 1d).

We should stress that while Fig. 1b–d only shows the threshold for shifts in dominant phase partitioning, i.e. the lines where 50 % of a chemical is in one phase, even very small fractions of a chemical partitioning into a phase could be important, for example, if they have an efficient reactive or depositional sink within that phase or if they have reached high concentrations in their preferred phases.

2.3 Estimating the equilibrium partitioning coefficients for SOA compounds

A chemical occupies a place within the chemical partitioning space based on its partitioning coefficients $K_{W/G}$ and $K_{W_{IOM}/G}$. Ideally, one would use measured partition coefficients to locate a substance in the partitioning space (Lei and Wania, 2004). However, because such data are not available for many, if not most of the substances involved in SOA formation, we need to rely on prediction methods when trying to locate SOA compounds in the chemical partitioning space. We recently applied three such methods to the prediction of $K_{W_{IOM}/G}$ for substances involved in SOA formation (Wania et al., 2014). These methods, which were first applied to the issue of gas-particle partitioning by Arp et al. (2008, 2009), are poly-parameter linear free energy relationships (ppLFERs) calibrated with empirical aerosol sorption data, the solvation models implemented in SPARC Performs Automated Reasoning in Chemistry (SPARC), and the quantum-chemical software COSMOtherm (Version C30_1401 with BP_TZVP_C30_1401 parameterization, COSMOlogic GmbH & Co. KG, Leverkusen, Germany, 2014), which predict solvation equilibria from molecular structure alone. The same three methods can also be used to predict $K_{W/G}$.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We compiled the chemical identity of compounds believed to form during the oxidation of normal alkanes (Jordan et al., 2008), terpenes (Chen and Griffin, 2005) and aromatic hydrocarbons (Forstner et al., 1997). Additional oxidation products of α -pinene were taken from Camredon et al. (2010), Shilling et al. (2009) and Valorso et al. (2011).

Molecular structures of a total of more than 350 molecules were drawn with ChemSketch (ACD/Labs, Advanced Chemistry Development Inc. Toronto, Canada), which yields each molecule's SMILES (simplified molecular-input line-entry system) string.

For the ppLFER approach, the molecules' solute descriptors were predicted with ABSOLV, which relies on a group contribution quantitative structure property relationship (ACD/Labs, Toronto, Canada). $K_{W/G}$ at different temperatures were calculated directly using the ppLFER equations given by Goss (2006). $K_{WIOM/G}$ at 15 °C for four different aerosols (sampled from urban, suburban, rural and coastal sites during different seasons) were calculated using ppLFER equations by Arp et al. (2008b), and adjusted for temperature using an enthalpy of phase partitioning estimated using the approach described in Wania et al. (2014). The temperature-adjusted $\log K_{WIOM/G}$ for the four aerosol samples was averaged.

SPARC directly predicts $K_{W/G}$ and $K_{WIOM/G}$ as a function of temperature when provided with a SMILES code. For the $K_{WIOM/G}$ prediction, SPARC requires a solvent to be specified, which should have solvation properties that resemble those of WIOM in SOA (Wania et al., 2014). As a default choice we used a relatively non-polar, hypothetical structure termed *B* proposed to be an oligomerisation product of the oxidation of volatile aromatics (Kalberer et al., 2004; Arp and Goss, 2009; Wania et al., 2014). For the estimation of $K_{WIOM/G}$ of the oxidation products of α -pinene, we additionally used two molecules as surrogate solvents that are themselves oxidation products of α -pinene: a dimer formed by esterification of pinic acid and 10-hydroxy-pinonic acid, referred to as ESTER_dimer by Zuend and Seinfeld (2012), and 5-hydroperoxy-4-(hydroxymethyl)-5-methyl-2-oxohexanoic acid, designated as C813OOH in the Master Chemical Mechanism model. The latter chemical is highly oxygenated and polar.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wania et al. (2014) describe the prediction of $K_{\text{WIOM}/\text{G}}$ for the n -alkane oxidation products using COSMOtherm and the surrogate solvent B mentioned above. COSMOtherm was similarly used to calculate $K_{\text{W}/\text{G}}$ for the n -alkane oxidation products. COSMOtherm was also applied to estimate $K_{\text{W}/\text{G}}$ and $K_{\text{WIOM}/\text{G}}$ at selected temperatures for 22 α -pinene oxidation products, using either structure B , ESTER_dimer and C813OOH as surrogate solvents to represent WIOM.

The salting-out effect on aqueous phase/gas phase partitioning was quantified by estimating a compound specific Setschenow coefficient K_{S} in ammonium sulphate solution using a ppLFER by Wang et al. (2014) and then calculating a salt solution/gas phase equilibrium partitioning coefficient $K_{\text{S}/\text{G}}$ using:

$$\log K_{\text{S}/\text{G}} = \log K_{\text{W}/\text{G}} - [\text{salt}]K_{\text{S}} \quad (10)$$

where [salt] is the molar concentration of ammonium sulphate in aqueous salt solution, which was varied between 0 and 6.6 M, the latter corresponding to a supersaturated solution as may be encountered in aqueous aerosol (Zuend and Seinfeld, 2012).

All estimated partitioning data are given in an Excel[®] file in the Supplement.

2.4 Major assumptions underlying the chemical partitioning space approach

Before proceeding to consider and interpret chemical partitioning space plots for SOA compounds, it is worthwhile to discuss two major assumptions that are made when drawing these diagrams. The first assumption is that chemicals achieve quasi-instantaneous equilibrium between the phases present in the atmosphere. The second is that the condensed phase separates into a largely aqueous phase and a phase largely made up of water-insoluble organic matter. In particular, we do not consider the possibility of the simultaneous presence of two immiscible, largely non-aqueous organic phases, e.g. a non-polar hydrocarbon-like phase separated from a polar, highly oxidized, non-aqueous phase. (Such a phase separation is for example predicted by the 2-D VBS approach (Donahue et al., 2012), but this approach also disregards the

presence of water in the condensed phase.) Because of the equilibrium assumption, particle morphology, i.e. the spatial arrangement of the two condensed phases relative to each other, is of no concern in our analysis (Shiraiwa et al., 2013).

While for a long time, the time scale of chemical diffusion in largely liquid-like atmospheric aerosol phases was judged sufficiently short to assume that equilibrium partitioning prevails most of the time, more recently evidence is emerging that suggests that the physical state of organic aerosol may at times be semi-solid or glassy (Mikhailov et al., 2009; Virtanen et al., 2010) and the rate of mass transfer in such phases is too slow to assure equilibrium distribution at atmospheric time scales (e.g. Koop et al., 2011; Perraud et al., 2012). However, this view is not yet universally acknowledged and there is similarly experimental evidence of the validity of the quasi-instantaneous equilibrium assumption (e.g. Saleh et al., 2013). In any case, even if chemical equilibrium is not always achieved the equilibrium distribution is still of paramount importance as the state that the system strives to reach.

Support for separation into predominantly organic and aqueous phases in aerosol stems from laboratory (e.g. Song et al., 2012), field (You et al., 2012) and theoretical investigations (e.g. Bertram et al., 2011; Zuend and Seinfeld, 2012). These same investigations also suggest the possibility of the formation of a single condensed aqueous phase at high relative humidity and if the organic material is highly oxidized. Incidentally, even though we assume the presence of a WIOM phase separate from an aqueous phase when drawing the partitioning space map, the map still can indicate when such phase separation is unlikely. Namely, if most of the organic compounds present in an aerosol fall in the part of the partitioning space indicating predominant partitioning into the aqueous phase and few fall into the part corresponding to partitioning into WIOM, it stands to reason that no WIOM phase will form and that instead all of the organic material present will be in a single, largely aqueous condensed phase.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

3.1 Placing the SOA compounds in the chemical partitioning space

Figure 2 shows the location of the SOA compounds in the partitioning space. The chemicals are located based on their $K_{W/G}$ and $K_{WIOM/G}$ at 15°C estimated with SPARC using structure *B* as a surrogate for WIOM and assuming no salting out effect. Differently coloured markers designate compounds derived from different precursors.

Comparing a chemical's location relative to the thresholds for dominant phase distribution, that were introduced in Fig. 1 and are integrated into Fig. 2, allows us to graphically deduce its phase distribution behaviour. It is apparent that the oxidation products of *n*-alkanes (blue dots in Fig. 2) generally have higher $\log K_{WIOM/G}$ and lower $K_{W/G}$ than the oxidation products of terpenes and aromatic hydrocarbons and therefore tend to have a higher affinity for the WIOM phase than the aqueous phase when they become insufficiently volatile to remain in the gas phase. Lower $K_{W/G}$ values of the products formed from α - and β -pinene and d-limonene (green and red dots in Fig. 2, respectively) means that they cluster around the line separating predominant partitioning to WIOM and aqueous phase. The graphical analysis also suggests that most of the oxidation products of the mono-aromatic hydrocarbons identified by Forstner et al. (1997) (yellow dots in Fig. 2) are too volatile to notably contribute to organic aerosol formation under most atmospheric conditions. An exception is a cloud, where many oxidation products from all types of precursors could be expected to partition substantially into cloud water.

Figure 2 does not show the position of small water-soluble compounds, such as the C2 and C3 compounds typically associated with aqueous SOA (e.g. glyoxal, methylglyoxal, oxalic acid, etc.) (Lim et al., 2010). The distribution of these compounds between gas and aerosol phase is often strongly influenced by hydration and possibly other reactions in the aqueous phase (Ip et al., 2009; Volkamer et al., 2009; Kampf et al., 2013) leading to "effective" partition coefficients that are orders of magnitude higher than estimated ones (e.g. Rossignol et al., 2012). Furthermore, many smaller acids

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are expected to ionize substantially under the pH condition prevailing in aerosol water and the $K_{W/G}$ predicted for the neutral form may often underestimate the preference for the aqueous phase. Following Volkamer et al. (2009), it would be possible to locate these compounds in the partitioning space based on their “effective” partitioning coefficients, as long as these can be reliably derived.

Figure 3 is identical to Fig. 2, except that now the markers designating the SOA compound are coloured based on their average carbon oxidation state (OS_C) and number of carbons (n_C). Figure S1 in the Supplement displays a similar plot based on the compounds’ oxygen : carbon ratio (O : C). These plots allow us to relate a chemical’s location in the partitioning space with its molecular size and oxidation state. An increase in n_C corresponds to a shift to the right in the chemical partitioning space, i.e. larger chemicals have a higher affinity for the WIOM phase relative to the gas phase. Increasing n_C in the form of methylene groups, on the other hand, hardly affects a chemical’s relative affinity for water and the gas phase, i.e. an additional $-CH_2-$ decreases vapour pressure and water solubility almost to the same extent. n -alkane oxidation products that only deviate in their chain length are arranged along lines that are parallel to the x axis (Fig. 3a). If a series of compound varies in n_C because of additional aromatic carbons, such as the polycyclic aromatic hydrocarbons, the increase in $K_{WIOM/G}$ is accompanied by a smaller increase in $K_{W/G}$ (Lei and Wania, 2014). An increase in a compound’s OS_C is reflected by a shift down and slightly to the right in the partitioning space, i.e. more oxidized species have a higher relative affinity for the aqueous than the WIOM and gas phase. SOA compounds’ OS_C increases largely because of the addition of oxygen containing functional group, which leads to an increase in $K_{W/G}$. The O : C ratio shows similar but less clear trends as the OS_C (Fig. S1).

3.2 Categorizing compounds based on their atmospheric phase distribution behaviour

We can now classify organic compounds based on their atmospheric phase distribution behaviour. Figure 4 assigns labels to different zones of the chemical partitioning

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



space. Based on their tendency to partition to condensed phases, compounds are classified as volatile (V), semi-volatile (SV), low volatile (LV), and extremely low volatile (ELV). Based on their preferred condensed phase, compounds are further categorised as hydrocarbon-like (HOC, preference for WIOM), oxygenated (OOC, no obvious preferred condensed phase), and water-soluble (WSOC, preference for aqueous phase).

At equilibrium less than 1 % of a VOC ($\log K_{\text{WIOM/G}}$ and $\log K_{\text{W/G}} < 8$) is predicted to be in the condensed phase even under high OAL conditions ($100 \mu\text{g m}^{-3}$). A chemical is presumed to be a LVOC ($\log K_{\text{WIOM/G}}$ and $\log K_{\text{W/G}} > 14$), if less than 1 % is predicted to be in the gas phase even under low OAL conditions ($1 \mu\text{g m}^{-3}$). SVOCs ($8 < \log K_{\text{WIOM/G}}$ or $\log K_{\text{W/G}} < 14$) are in between VOCs and LVOCs. Similarly, at equilibrium less than 1 % of a HOC ($\log K_{\text{WIOM/W}} > 3$) partitions into the aqueous phase, even under low OAL conditions, and less than 1 % of a WSOC ($\log K_{\text{WIOM/W}} < -3$) partitions into the WIOM phase, even if OAL is high. OOCs fall in between HOCs and WSOCs ($-3 < \log K_{\text{WIOM/W}} < 3$). Because of the widely different LWC, SV-WSOCs occupy a different part of the partitioning space under clear conditions and in a cloud. Whereas chemicals with a $\log K_{\text{W/G}} < 8$ are VOCs at low RH, that threshold shifts to a $\log K_{\text{W/G}}$ of 3.5 within a cloud.

The nomenclature in Fig. 4 builds upon previous classifications of SOA compounds (Kroll et al., 2011; Donahue et al., 2012). In particular, the aerosol mass spectrometry community uses HOA and OOA to refer to hydrocarbon-like and oxygenated organic aerosol. However, the use of the terms here may not always be entirely congruous with previous efforts. For example, our definition of SVOCs comprises some of the compounds that would be called intermediate volatility organic compounds (IVOCs) and LVOCs by Donahue et al. (2012).

So far, we have used a single point to designate a chemical's location in the partitioning space, although this location is variable for two primary reasons: prediction uncertainty and environmental variability. In the next two sections, we will explore these sources of variability in more detail.

3.3 Influence of prediction uncertainty on placement of *n*-alkane oxidation products in the partitioning space

In order to evaluate the uncertainty of the placement of chemicals in the partitioning space, we sought to quantify the confidence of the $K_{W_{IOM}/G}$ and $K_{W/G}$ predictions by comparing the results of different prediction methods. The values obtained for the partitioning coefficients of three *n*-alkanes (C9, C13, C17) and their oxidation products at 25 °C by ppLFER (average for four aerosol), SPARC (using structure *B* for WIOM), and COSMOtherm (using structure *B* for WIOM) are displayed in Fig. 5. The $K_{W_{IOM}/G}$ predictions had previously been compared in Wania et al. (2014). We use arrows to illustrate the displacement of a chemical in the partitioning space when we apply different prediction methods: the origin of each arrow is the location obtained by SPARC predictions. The end of the blue and red arrow is the location obtained using ppLFRs and COSMOtherm, respectively. The length of the arrow is thus a measure of the uncertainty of the prediction (the horizontal and vertical displacement indicating the discrepancy in $K_{W_{IOM}/G}$ and $K_{W/G}$ predictions, respectively) and arrows consistently pointing in a particular direction indicate a bias of one method relative to another.

The arrows in Fig. 5 tend to get longer when moving from the upper left to the lower right of the partitioning space, suggesting that the discrepancy between predictions becomes larger as chemicals become less volatile. This is expected as predictions for these less volatile, multi-functional chemicals involve an extrapolation from the type of chemicals for which empirical partitioning data are available and which are therefore the basis of the calibration of the ppLFER and also the SPARC method. The vertical displacements in Fig. 5 tend to be larger than the horizontal ones, indicating that the uncertainty of the $K_{W/G}$ prediction is higher than that of $K_{W_{IOM}/G}$. Most arrows are pointing to the left, which means that SPARC predicts generally higher $K_{W_{IOM}/G}$ than the other two methods. Most red arrows are pointing upwards, indicating that SPARC also predicts higher $K_{W/G}$ than COSMOtherm.

Using the chemical equilibrium partitioning space

F. Wania et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The uncertainty in the placement of chemicals falling into the area close to the phase transition between gas and WIOM phase is generally less than an order of magnitude for $K_{\text{WIOM}/\text{G}}$ and less than two orders of magnitude for $K_{\text{W}/\text{G}}$ (Fig. 5). The latter is generally of little consequence because the oxidation products of the n -alkanes are not sufficiently polar to partition to the aqueous aerosol phase, unless the LWC is very high (i.e. in a cloud). The uncertainty of $K_{\text{WIOM}/\text{G}}$ on the other hand is sufficiently large to significantly change the predicted partitioning behaviour of SV-HOCs with a log $K_{\text{WIOM}/\text{G}}$ ranging from 10 to 12. It is worthwhile, however, to reiterate Wania et al. (2014)'s observation that the prediction uncertainty in $K_{\text{WIOM}/\text{G}}$ for the n -alkane oxidation products is much smaller than the prediction uncertainty of saturation vapour pressure normally used to predict the phase distribution behaviour of SOA compounds.

3.4 Influence of environmental variability on placement of α -pinene oxidation products in the partitioning space

We use α -pinene oxidation products to illustrate the effect of temperature, aqueous phase salinity and WIOM phase polarity on the placement of chemicals in the partitioning space. Because it is difficult to discuss the impact of environmental variability without also considering the prediction uncertainty discussed in Sect. 3.3, we produced multiple plots using different predictions. Figure 6a and b is based on SPARC and COSMOtherm-predicted partition coefficients, respectively, and Fig. S2 in the Supplement is based on ppLFER predictions. The number of compounds with COSMOtherm predictions is smaller than for the other two methods. In these plots, three intersecting lines represent each chemical. The diagonal line indicates the variability in the partitioning coefficients caused by temperature (upper left of the line is for 30 °C, lower right is for 0 °C). The vertical line designates the salting-out effect, which yields a decrease in $K_{\text{W}/\text{G}}$. The uppermost value on that line indicates the $K_{\text{W}/\text{G}}$ in a 6.6 M supersaturated solution of $(\text{NH}_4)_2\text{SO}_4$. The horizontal line indicates the range of $K_{\text{WIOM}/\text{G}}$ obtained by using different surrogate molecules to represent the WIOM phase (Fig. 6). In the case of the ppLFER-based plot (Fig. S2), the horizontal line indicates the range of values ob-

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and is due to the well-known linear correlation between the logarithm of a partition coefficient and the enthalpy of phase transfer (Goss and Schwarzenbach, 1999). The vertical lines, on the other hand, decrease in length with increasing oxygenation, i.e., the salting-out effect is weaker for the higher generation oxidation products. This is because Setschenow coefficients are smaller for more polar substances (Endo et al., 2012; Wang et al., 2014).

By comparing the placement of chemicals in Fig. 6 with the boundaries of dominant phase distribution introduced in Fig. 1, we can graphically predict the phase distribution behaviour of α -pinene oxidation products under different conditions (Fig. 6a and b displays the boundaries for low and high LWC, respectively). At low LWC (Fig. 6a) products with four oxygen atoms are starting to contribute to the condensed phase, especially at higher OALs and lower temperature. Products with five and more oxygen atoms are already predicted to have only minor fractions in the gas phase, falling close to the boundary between aqueous and WIOM phase. The salting-out effect likely assures that those α -pinene oxidation products would be found mostly in the WIOM phase. Products with seven and more oxygen atoms (i.e. the ELVOCs observed as major products of α -pinene oxidation in the field by Ehn et al., 2014) would clearly no longer be in the gas phase under any conditions.

The partitioning space analysis also suggests that under high LWC conditions (i.e. in a cloud, Fig. 6b) oxidation products of α -pinene with three and even only two oxygen would partition strongly into the aqueous condensed phase, also because the salting-out effect would not longer be active. This incidentally highlights why an analysis of phase partitioning that ignores liquid water in the atmosphere can lead to erroneous conclusions. The red outlines in Fig. 6a and b designate the region where chemicals fall whose contribution to SOA formation would not be taken into account, if the aqueous phase were ignored. At low LWC (solid red outline), this applies to few, if any, oxidation products, but at high LWC (dashed red outline), a very large number of oxidation products would remain in the gas phase. This appears to be in agreement with Pankow

et al. (2013), who predicts “significant RH-induced enhancement of organic particulate matter condensation” in the α -pinene/ozone system.

Incidentally, none of the α -pinene oxidation products are predicted to partition notably into WIOM at high LWC (Fig. 6b). This probably means that unless there is another source of WIOM to the atmosphere, a single aqueous aerosol phase would form under such conditions.

We stress that there are other reasons why a chemical’s location may depend on environmental conditions. For example, some SOA compounds are acids with pK_A values in the pH range encountered in the aqueous phase of aerosol and cloud droplets, i.e. they can be expected to be ionized to some degree. Ionization increases $K_{W/G}$ relative to the value that is estimated by the prediction methods for the protonated form of the acids. It also means that $K_{W/G}$, and therefore the placement in the partitioning space depends on the pH of the aqueous phase.

3.5 Illustrating displacement within the partitioning space caused by chemical transformations

So far, the SOA compounds have been considered in isolation and not as part of a chemical reaction sequence. Figure 7 shows the oxidation products of the C17 n -alkane linked by arrows designating the chemical reactions elucidated by Lim and Ziemann (2005). Like the α -pinene oxidation products in Fig. 6, the oxidation products fall along a band that starts at the precursor VOC in the upper left and extends to the lower right of the partitioning space. Remarkably, the slope of that band and even the extent of its broadening toward the lower right is very similar for C17 (and by inference all n -alkanes) and α -pinene. This similarity is not really surprising, as in both cases hydroxyl, carbonyl, peroxy, nitrate, and carboxylic acid functional groups are being added to an aliphatic compound.

The slope of these bands of compounds within the partitioning space suggests that on average such functionalization roughly increases the $K_{W/G}$ by two log units for every log unit increase in $\log K_{WIOM/G}$, i.e. the addition of functional groups increases

26567

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the affinity for the aqueous phase more rapidly than the affinity for the WIOM phase. Whether the functionalization of a precursor VOC ends up generating compounds that prefer the WIOM or the aerosol aqueous phase thus depends to some extent on the location of that precursor in the partitioning space. The lower the $K_{W/G}$ and the higher the $K_{WIOM/G}$ of a precursor, the more likely it is to generate oxidation products that favour the WIOM phase.

The broadening of the band is due to different changes in $K_{W/G}$ and $K_{WIOM/G}$ occurring upon the addition of different functional groups. According to Lim and Ziemann (2005), the first few steps of the oxidation of *n*-alkanes by OH in the presence of NO_x, involves primarily five types of transformations. The displacement within the partitioning space caused by these transformations is depicted in Fig. 8.

As expected from the general trend toward the lower right seen in Fig. 7, the addition of functional groups to the aliphatic chain increases both $K_{WIOM/G}$ and $K_{W/G}$, i.e., arrows pointing to the lower right represent the reaction steps. The exception is the formation of cyclical dihydrofurans from δ -hydroxycarbonyls, which leads to a displacement in the opposite direction, i.e. to a more volatile substance. Size and direction of the displacement in the partitioning space depends on the functional groups being added. The addition of oxygen-containing groups such as hydroxyl, carbonyl, and carboxylester, increases $K_{W/G}$ more than $K_{WIOM/G}$, i.e. the mean vertical displacement is generally more than double the mean horizontal displacement. The addition of nitrate, on the other hand, leads to about equal displacement in the vertical and horizontal direction, i.e., causes $K_{W/G}$ and $K_{WIOM/G}$ to increase by the same extent. This incidentally explains why nitrated compounds (represented by broken lines in Fig. 6a) and especially di-nitrated compounds (UR37, UR38, UR39, UR41, UR42 in Fig. 7) tend to fall on the right side of the band of oxidation products, whereas those without nitrate substitutions (RP20, RP21, UR36 in Fig. 7) on the left. This implies that nitration increases the affinity for WIOM more than the addition of carbonyls, hydroxyls and carboxylester does.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To which compound a functional group is being added often does not seem to matter much (i.e. the arrows within each panel of Fig. 8 are of similar length and point in a similar direction). In particular the alkyl-chain length plays a limited role (blue and green arrows are very similar). However, the displacement caused by the addition of nitrate is different for different compounds. For example, nitration of the dihydrofuran leads to a larger change in $K_{W_{IOM}/G}$ and $K_{W/G}$ (longer arrows) than nitration of the aliphatic chain. The addition of a second nitrate leads to a smaller increase in $K_{W/G}$ (arrows point more to the right) than the first nitrate.

To obtain a broader set of displacements caused by chemical transformations, we also looked at the functionalization occurring during oxidation of aromatic hydrocarbons (Fig. S5). This analysis confirmed that the addition of oxygen-containing functional groups again increases both $K_{W_{IOM}/G}$ and $K_{W/G}$, with the latter increasing more rapidly, i.e., the vertical displacement being slightly larger than the horizontal one. The displacement is increasing from carbonyl to hydroxyl to carboxyl groups (i.e. arrows are getting longer in Fig. S5), whereby the sum of the displacements caused by a carbonyl and a hydroxyl group is larger than the displacement caused by the addition of a carboxyl group. Similar to the addition of a nitrate group to the *n*-alkane oxidation products (Fig. 8), the addition of a nitro-group to a mono-aromatic hydrocarbon increases $K_{W_{IOM}/G}$ and $K_{W/G}$ to the same extent, i.e., horizontal and vertical displacements are very similar (Fig. S5). Increasing the size of a mono-aromatic compound through the addition of methyl- and methylene-groups has only a very minor effect on the placement in the partitioning space, increasing $K_{W_{IOM}/G}$ only slightly and $K_{W/G}$ hardly at all (Fig. S5).

In the case of $-OH$ and $-NO_2$ addition to a mono-aromatic hydrocarbon, the displacement in the partitioning space can be strongly dependent on the type of functional groups already present in the molecule. The difference in the change in $K_{W_{IOM}/G}$ and $K_{W/G}$ occurring upon the addition of $-OH$ directly to the aromatic ring or to an existing carbonyl group (i.e. conversion of benzaldehyde to benzoic acid) is relatively small (Fig. 9a). On the other hand, the effect on partitioning upon the addition of $-OH$

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tors on SOA formation. For example, it allows for a direct comparison of the uncertainty of the prediction of partitioning coefficients with the impact of atmospheric variability on these coefficients. It facilitates a comparison of the direction and size of the effects of temperature, salinity and organic phase polarity on the partitioning behaviour of different SOA compounds. The change in partitioning behaviour occurring upon dimerisation can be compared with that of adding a hydroxyl or a nitrate functional group. Because everything is displayed within the same two-dimensional framework, even very different factors can be assessed comparatively: prediction uncertainty can be compared with the effect of a change in OAL or LWC on partitioning, the effect of a chemical reaction, or the effect of a change in temperature and ammonium sulphate concentration.

This suitability for comparative visualisation can serve to aid in the understanding of dynamic processes, where several of the factors affecting phase distribution are changing at the same time. For example, during a photochemical smog episode, or during a chamber experiment, the precursor VOCs become oxidized and with increasing functionalization – and eventually also dimerisation – the oxidation products partition increasingly into the condensed phase (i.e. they move to the lower right of the partitioning space) (Fig. 11). As a result, the OAL is increasing causing the vertical threshold separating dominant phase distribution into gas and WIOM phase to move to the left (Fig. 11). This implies that with ongoing SOA formation, more and more volatile chemicals become available for partitioning to the WIOM. Overall, the changes in partitioning behaviour caused by the chemical processes are considerably larger than the shift in the threshold caused by the increase in OAL (Fig. 11).

Similarly, during cloud droplet formation a large upward shift in the horizontal threshold between dominant phase distribution into water and gas phase coincides with a downward shift in the position of SOA compounds in the partitioning space, as salts become diluted and the salting out effect becomes negligible. Another example is the dilution of tail pipe emissions, which is accompanied by a rapid decrease in both OAL and temperature, having opposing effects on the partitioning between gas and WIOM phase (May et al., 2013).

centrations (Carlton et al., 2013). Therefore, partitioning between the two condensed phases and the value of $K_{W/WIOM}$ is important for predicting SOA formation, composition and yield. Also, the occurrence and kinetics of condensed phase reactions that oxidation products may undergo, such as dimerisations or reactions with sulphate are likely to be different in the WIOM and aqueous phase.

Several 2-dimensional property spaces have been advocated for describing organic aerosol characteristics (Pankow and Brasanti, 2009; Kroll et al., 2011; Cappa and Wilson, 2012), the 2-D VBS by Donahue et al. (2011, 2012) using saturation concentration and O : C ratio (or OS_C) being the most conspicuous. One of the key shortcomings of the 2-D VBS approach is, that it “does not allow consideration of water uptake or variable partition coefficients” (Pankow et al., 2013). Donahue et al. (2012) actually mention this when introducing the 2-D-VBS: “The formulation so far does not address the role of water. [. . .] However, because water can be a dominant component, and therefore its unique properties are more likely to challenge the mean-field assumption, it is likely that explicit treatment of humidity effects will be required.” Here we have presented what essentially amounts to an alternative 2-dimensional approach to SOA characterization based on partitioning coefficients from gas phase into WIOM and water. This approach includes partitioning into an aqueous phase and directly addresses one of the key issues that atmospheric modelers of SOA formation aim to address: which fraction of the products formed from the oxidation of VOCs will end up in the SOA after equilibration under different atmospheric conditions. We could imagine that it is possible to develop parameterizations of the chemical partitioning space introduced here, which – combined with gas phase concentrations of sets of compounds with comparable partitioning properties – are suitable for implementation in atmospheric models seeking to explicitly describe the formation of SOA.

**The Supplement related to this article is available online at
doi:10.5194/acpd-14-26545-2014-supplement.**

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. The authors acknowledge financial support from the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- 5 Arp, H. P. H. and Goss, K. U.: Ambient gas/particle partitioning, 3. Estimating partition coefficients of apolar, polar, and ionizable organic compounds by their molecular structure, *Environ. Sci. Technol.*, 43, 1923–1929, 2009.
- Arp, H. P. H., Schwarzenbach, R. P., and Goss, K. U.: Ambient gas/particle partitioning, 1. Sorption mechanisms of apolar, polar, and ionizable organic compounds, *Environ. Sci. Technol.*, 42, 5541–5547, 2008.
- 10 Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A., You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase separation, efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the organic component, *Atmos. Chem. Phys.*, 11, 10995–11006, doi:10.5194/acp-11-10995-2011, 2011.
- 15 Camredon, M., Hamilton, J. F., Alam, M. S., Wyche, K. P., Carr, T., White, I. R., Monks, P. S., Rickard, A. R., and Bloss, W. J.: Distribution of gaseous and particulate organic composition during dark α -pinene ozonolysis, *Atmos. Chem. Phys.*, 10, 2893–2917, doi:10.5194/acp-10-2893-2010, 2010.
- 20 Cappa, C. D. and Wilson, K. R.: Multi-generation gas-phase oxidation, equilibrium partitioning, and the formation and evolution of secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 9505–9528, doi:10.5194/acp-12-9505-2012, 2012.
- Carlton, A. G. and Turpin, B. J.: Particle partitioning potential of organic compounds is highest in the Eastern US and driven by anthropogenic water, *Atmos. Chem. Phys.*, 13, 10203–10214, doi:10.5194/acp-13-10203-2013, 2013.
- 25 Chen, J. J. and Griffin, R. J.: Modeling secondary organic aerosol formation from oxidation of alpha-pinene, beta-pinene, and d-limonene, *Atmos. Environ.*, 39, 7731–7744, 2005.
- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T.,
- 30

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Using the chemical
equilibrium
partitioning space**

F. Wania et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Dal Maso, M., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary aerosol, *Nature*, 506, 476–479, 2014.
- Endo, S., Pfennigsdorff, A., and Goss, K.-U.: Salting-out effect in aqueous NaCl solutions: trends with size and polarity of solute molecules, *Environ. Sci. Technol.*, 46, 1496–1503, 2012.
- Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, 2006.
- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11, 3303–3318, doi:10.5194/acp-11-3303-2011, 2011.
- Donahue, N. M., Kroll, J. H., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set – Part 2: Diagnostics of organic-aerosol evolution, *Atmos. Chem. Phys.*, 12, 615–634, doi:10.5194/acp-12-615-2012, 2012.
- Forstner, H. J. L., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol from the photooxidation of aromatic hydrocarbons: molecular composition, *Environ. Sci. Technol.*, 31, 1345–1358, 1997.
- Goss, K.-U.: Prediction of the temperature dependency of Henry's law constant using poly-parameter linear free energy relationships, *Chemosphere*, 64, 1369–1374, 2006.
- Goss, K.-U. and Schwarzenbach, R. P.: Empirical prediction of heats of vaporization and heats of adsorption of organic compounds, *Environ. Sci. Technol.*, 33, 3390–3393, 1999.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q., Martin, S. T., Farmer, D. K., and Artaxo, P.: A simplified description of the evolution of organic aerosol composition in the atmosphere, *Geophys. Res. Lett.*, 37, L08803, doi:10.1029/2010GL042737, 2010.

Using the chemical
equilibrium
partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ip, H. S. S., Huang, X. H. H., and Yu, J. Z.: Effective Henry's law constants of glyoxal, glyoxylic acid, and glycolic acid, *Geophys. Res. Lett.*, 36, L01802, doi:10.1029/2008GL036212, 2009.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S., Zhang, Q., Kroll, J. H., De Carlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.
- Jordan, C. E., Ziemann, P. J., Griffin, R. J., Lim, Y. B., Atkinson, R., and Arey, J.: Modeling SOA formation from OH reactions with C-8-C-17 *n*-alkanes, *Atmos. Environ.*, 42, 8015–8026, 2008.
- Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004.
- Kampf, C. J., Waxman, E. M., Slowik, J. G., Dommen, J., Pfaffenberger, L., Praplan, A. P., Prevot, A. S. H., Baltensperger, U., Hoffmann, T., and Volkamer, R.: Effective Henry's law partitioning and the salting constant of glyoxal in aerosols containing sulfate, *Environ. Sci. Technol.*, 47, 4236–4244, 2013.
- Koop, T., Bookhold, J., Shiraiwa, M., and Poeschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238–19255, 2011.
- Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R., Altieri, K. E., Mazzoleni, L. R., Wozniak, A. S., Bluhm, H., Mysak, E. R., Smith, J. D., Kolb, C. E., and Worsnop, D. R.: Carbon oxidation state as a metric for describing the chemistry of atmospheric organic aerosol, *Nature Chem.*, 3, 133–139, 2011.
- Lei, Y. D. and Wania, F.: Is rain or snow a more efficient scavenger of organic chemicals?, *Atmos. Environ.*, 38, 3557–3571, 2004.

**Using the chemical
equilibrium
partitioning space**

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Lim, Y. B. and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from reactions of *n*-alkanes with OH radicals in the presence of NO_x, Environ. Sci. Technol., 39, 9229–9236, 2005.

Lim, Y. B., Tan, Y., Perri, M. J., Seitzinger, S. P., and Turpin, B. J.: Aqueous chemistry and its role in secondary organic aerosol (SOA) formation, Atmos. Chem. Phys., 10, 10521–10539, doi:10.5194/acp-10-10521-2010, 2010.

May, A. A., Presto, A. A., Hennigan, C. J., Nguyen, N. T., Gordon, T. D., and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: 1. gasoline vehicle exhaust, Atmos. Environ., 77, 128–139, 2013.

Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, Atmos. Chem. Phys., 9, 9491–9522, doi:10.5194/acp-9-9491-2009, 2009.

Pankow, J. F.: Gas/particle partitioning of neutral and ionizing compounds to single and multi-phase aerosol particles, 1. Unified modeling framework, Atmos. Environ., 37, 3323–3333, 2003.

Pankow, J. F. and Barsanti, K. C.: The carbon number-polarity grid: a means to manage the complexity of the mix of organic compounds when modeling atmospheric organic particulate matter, Atmos. Environ., 43, 2829–2835, 2009.

Pankow, J. F., Niakan, N., and Asher, W. E.: Combinatorial variation of structure in considerations of compound lumping in one- and two-dimensional property representations of condensable atmospheric organic compounds, 1. Lumping by 1-D volatility with *n_C* fixed, Atmos. Environ., 80, 172–183, 2013.

Perraud, V., Bruns, E. A., Ezell, M. J., Johnson, S. N., Yu, Y., Alexander, M. L., Zelenyuk, A., Imre, D., Chang, W. L., Dabdub, D., Pankow, J. F., and Finlayson-Pitts, B. J.: Nonequilibrium atmospheric secondary organic aerosol formation and growth, P. Natl. Acad. Sci. USA, 109, 2836–2841, 2012.

Rossignol, S., Chiappini, L., Perraudin, E., Rio, C., Fable, S., Valorso, R., and Doussin, J. F.: Development of a parallel sampling and analysis method for the elucidation of gas/particle partitioning of oxygenated semi-volatile organics: a limonene ozonolysis study, Atmos. Meas. Tech., 5, 1459–1489, doi:10.5194/amt-5-1459-2012, 2012.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Saleh, R., Donahue, N. M., and Robinson, A. L.: Time scales for gas-particle partitioning equilibration of secondary organic aerosol formed from alpha-pinene ozonolysis, *Environ. Sci. Technol.*, 47, 5588–5594, 2013.
- Shilling, J. E., Chen, Q., King, S. M., Rosenoern, T., Kroll, J. H., Worsnop, D. R., DeCarlo, P. F., Aiken, A. C., Sueper, D., Jimenez, J. L., and Martin, S. T.: Loading-dependent elemental composition of α -pinene SOA particles, *Atmos. Chem. Phys.*, 9, 771–782, doi:10.5194/acp-9-771-2009, 2009.
- Shiraiwa, M., Zuend, A., Bertram, A. K., and Seinfeld, J. H.: Gas–particle partitioning of atmospheric aerosols: interplay of physical state, non-ideal mixing and morphology, *Phys. Chem. Chem. Phys.*, 15, 11441–11453, 2013.
- Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation and morphology of internally mixed dicarboxylic acids/ammonium sulfate/water particles, *Atmos. Chem. Phys.*, 12, 2691–2712, doi:10.5194/acp-12-2691-2012, 2012.
- Stenzel, A., Goss, K.-U., and Endo, S.: Prediction of partition coefficients for complex environmental contaminants: validation of COSMOtherm, ABSOLV, and SPARC, *Environ. Toxicol. Chem.*, 33, 1537–1543, 2014.
- Valorso, R., Aumont, B., Camredon, M., Raventos-Duran, T., Mouchel-Vallon, C., Ng, N. L., Seinfeld, J. H., Lee-Taylor, J., and Madronich, S.: Explicit modelling of SOA formation from α -pinene photooxidation: sensitivity to vapour pressure estimation, *Atmos. Chem. Phys.*, 11, 6895–6910, doi:10.5194/acp-11-6895-2011, 2011.
- Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirila, P., Leskinen, J., Makela, J. M., Holopainen, J. K., Poschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824–827, 2010.
- Volkamer, R., Ziemann, P. J., and Molina, M. J.: Secondary Organic Aerosol Formation from Acetylene (C_2H_2): seed effect on SOA yields due to organic photochemistry in the aerosol aqueous phase, *Atmos. Chem. Phys.*, 9, 1907–1928, doi:10.5194/acp-9-1907-2009, 2009.
- Wang, C., Lei, Y. D., Endo, S., and Wania, F.: Measuring and modeling the salting out effect in ammonium sulfate solutions, *Environ. Sci. Technol.*, accepted, doi:10.1021/es5035602, 2014.
- Wania, F., Lei, Y. D., Wang, C., Abbatt, J. P. D., and Goss, K.-U.: Novel methods for predicting gas-particle partitioning during the formation of secondary organic aerosol, *Atmos. Chem. Phys. Discuss.*, 14, 21341–21385, doi:10.5194/acpd-14-21341-2014, 2014.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- and H_2O , *J. Geophys. Res.*, 107, 4207, doi:10.1029/2001JD000451, 2002.

Wong, F. and Wania, F.: Visualising the equilibrium distribution and mobility of organic contaminants in soil using the chemical partitioning space, *J. Environ. Monitor.*, 13, 1569–1578, 2011.

Westgate, J. N. and Wania, F.: Model-based exploration of the drivers of mountain cold-trapping in soils, *Environ. Sci.: Processes Impacts*, 15, 2220–2232, 2013.

You, Y., Renbaum-Wolff, L., Carreras-Sospedra, M., Hanna, S. J., Hiranuma, N., Kamal, S., Smith, M. L., Zhang, X., Weber, R. J., Shilling, J. E., Dabdub, D., Martin, S. T., and Bertram, A. K.: Images reveal that atmospheric particles can undergo liquid–liquid phase separations, *P. Natl. Acad. Sci. USA*, 109, 13188–13193, 2012.

Zuend, A. and Seinfeld, J. H.: Modeling the gas-particle partitioning of secondary organic aerosol: the importance of liquid-liquid phase separation, *Atmos. Chem. Phys.*, 12, 3857–3882, doi:10.5194/acp-12-3857-2012, 2012.

Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 1. Different versions of the chemical partitioning space for the atmosphere, showing in red, blue and green the combinations of partitioning properties that lead to dominant equilibrium partitioning to the gas, aqueous, and water insoluble organic matter (WIOM) phase, respectively. **(a)** refers to typical atmospheric conditions of $10 \mu\text{g}$ of both WIOM and liquid water per m^3 of air and demonstrates that the transition from 1 (10) to 99 (90) % in a particular phase occurs over a four (two) order of magnitude range in a partition coefficient. A clear atmosphere **(a and b)**, a cloud **(c)**, and a dry chamber experiment **(d)** only differ in terms of the relative volume of the three phases present, leading to shifts in the thresholds indicating 50 % partitioning into any one phase. Dashed lines in **(b–d)** indicate the 50 % threshold for various loadings of WIOM and various liquid water contents.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. Placement of the selected oxidation products of *n*-alkanes, biogenic terpenes and mono-aromatic hydrocarbons in the partitioning space. $K_{WIOM/G}$ and $K_{W/G}$ at 15 °C and 0 M salt of the compounds was estimated using SPARC assuming the solvation properties of WIOM can be approximated by those of structure *B*.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 3. Same as Fig. 2, except that markers are coloured based on a chemical's number of carbon atoms (n_C) (left) or its average carbon oxidation state (OS_C).

Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 4. Same as Fig. 2, except that different regions of the chemical partitioning space have been named according to the type of compounds that fall into these regions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.



n-alkanes

(C9, C13, C17)

25 °C
0 M salt
WIOM B

Origin of arrow:
SPARC-prediction

— ppLFER
— COSMOtherm

Figure 5. Arrows indicating the uncertainty of chemical placement in the partitioning space based on the discrepancy of $K_{WIOM/G}$ and $K_{W/G}$ predictions made with SPARC (origin of arrows), ppLFRs (tip of blue arrows) and COSMOtherm (tip of red arrows). For some compounds, no COSMOtherm predictions are available.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 6. Placement of α -pinene and some of its oxidation products in the chemical partitioning space based on predictions with SPARC (a) and COSMOtherm (b). Each chemical is represented by three lines, which account for the influence of temperature variations between 0 and 30 °C (diagonal line), the salting-out effect by 6.6 M $(NH_4)_2SO_4$ (vertical line) and the polarity of the surrogate representing the solvation properties of WIOM (horizontal line). The chemicals are superimposed on the dominant phase distribution maps for wet aerosol (left) and a cloud (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 7. Oxidation products of the *n*-alkane with 17 carbon atoms placed within the chemical partitioning space based on partitioning coefficients at 25 °C predicted with pplFERS and connected by arrows based on sequence of reaction by which they are formed. The code names for the compounds are based on Jordan et al. (2008).

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 8. Displacement of chemicals in the partitioning space caused by transformations involved in the oxidation of *n*-alkanes with 10 (blue) and 17 (green) carbons using predictions of $K_{W/O/M/G}$ and $K_{W/G}$ at 25 °C obtained by ppLFER. Also given are the mean vertical displacement (MVD) and mean horizontal displacement (MHD).

Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 9. Displacement in the chemical partitioning space occurring upon the addition of a hydroxyl (a) or nitro-group (b) to a mono-aromatic hydrocarbon, as predicted by SPARC for 15 °C and using structure *B* for WIOM (VD: vertical displacement, HD: horizontal displacement).

Using the chemical equilibrium partitioning space

F. Wania et al.




Figure 10. Displacements in the chemical partitioning space due to various dimerisation reactions proposed for ozonolysis reaction products of α -pinene by Camredon et al. (2010). $K_{WIOM/G}$ and $K_{W/G}$ at 15 °C were estimates with SPARC using structure *B* as a surrogate for WIOM. Also shown is the categorisation into compounds with different atmospheric phase distribution behaviour introduced in Fig. 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 11. Visualisation in the partitioning space of processes occurring during the oxidation of VOC precursors in a chamber experiment or a photochemical smog episode (in the presence of water). While the addition of functional groups and dimerisation reactions lead to an increase in $K_{WIOM/G}$ and $K_{W/G}$ and therefore a gradual shift to the lower right, the attendant increase in the organic aerosol load shifts the threshold for the gas-WIOM transition to the left.

Using the chemical equilibrium partitioning space

F. Wania et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

