



PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Polycyclic aromatic hydrocarbons, polychlorinated biphenyls, and chlorinated pesticides in background air in central Europe – investigating parameters affecting wet scavenging of polycyclic aromatic hydrocarbons

P. Shahpoury¹, G. Lammel^{1,2}, A. Holubová Šmejkalová^{3,4}, J. Klánová²,
P. Příbylová², and M. Váňa^{3,4}

¹Max-Planck Institute for Chemistry, Mainz, Germany

²Research Centre for Toxic Compounds in the Environment, Masaryk University,
Brno, Czech Republic

³Czech Hydrometeorological Institute, Košetice, Czech Republic

⁴Global Change Research Centre, ASCR, Prague, Czech Republic

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 13 September 2014 – Accepted: 2 October 2014 – Published: 28 October 2014

Correspondence to: G. Lammel (lammel@recetox.muni.cz),
P. Shahpoury (p.shahpoury@mpic.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

14, 26939–26970, 2014

PAHs, PCBs, and CPs in background air in central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Concentrations of polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), and chlorinated pesticides (CPs) were measured in air and precipitation at a background site in central Europe. Σ PAH concentrations in air and rainwater ranged from 0.7 to 327.9 ng m⁻³ and below analytical method detection limit (< MDL) to 2.1 × 10³ ng L⁻¹. The concentrations of PCBs and CPs in rainwater were < MDL. Σ PCB and Σ CP concentrations in air ranged from < MDL to 44.6 and < MDL to 351.7 pg m⁻³, respectively. The potential relationships between PAH wet scavenging and particulate matter and rainwater properties were investigated. The concentrations of ionic species in particulate matter and rainwater were significantly correlated, highlighting the importance of particle scavenging process. Overall, higher scavenging efficiencies were found for relatively less volatile PAHs, underlining the effect of analyte gas-particle partitioning on scavenging process. The PAH wet scavenging was more effective when the concentrations of ionic species were high. In addition, the elemental and organic carbon contents of the particulate matter were found to influence the PAH scavenging.

1 Introduction

Semi-volatile organic compounds (SOCs), such as polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), and chlorinated pesticides (CPs), can reach remote and background regions through atmospheric transport and deposition processes (Blais, 2005; Hageman et al., 2006; Grimalt et al., 2001). Wet scavenging is an important mechanism through which atmospheric SOC_s enter terrestrial and aquatic ecosystems (van Ry et al., 2002). Non-reactive gaseous SOC_s are removed from the atmosphere by gas scavenging process, which is controlled by the SOC equilibrium partitioning with falling raindrops, as described by Henry's law. In contrast, particle-associated species are removed from the atmosphere mainly through particle scavenging (Ligocki et al., 1985), a process controlled by physical parameters including par-

ACPD

14, 26939–26970, 2014

PAHs, PCBs, and CPs in background air in central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 ticulate matter (PM) characteristics, cloud microphysics, and meteorological conditions (Poster and Baker, 1996). Therefore, factors that affect SOC gas-particle partitioning, such as ambient temperature, relative humidity, PM chemical composition and surface characteristics, and SOC vapor pressure (Lohmann and Lammer, 2004), may play an important role in determining the relative contributions of gas and particle scavenging processes. It was suggested that SOC scavenging efficiencies would be between one to three orders of magnitude higher in case of in-cloud scavenging compared to that of below-cloud scavenging (Ligocki et al., 1985). Potential variability in scavenging efficiencies has also been suggested between rain events originated from warm and cold clouds (Bidleman, 1988). A number of studies observed variations in scavenging efficiencies of SOCs between different precipitation events (Offenberg and Baker, 2002; Agrell et al., 2002; Liu et al., 2013; Kaupp and McLachlan, 2000; McLachlan and Sellström, 2009; Atlas and Giam, 1988). To this date, studies have mainly focused on determining the relative importance of SOC gas and particle scavenging processes. However, to the best of our knowledge, no comprehensive study has been conducted to understand the causes behind the variability in scavenging efficiencies. Therefore, in this study we aimed at filling this gap by investigating the relationships between a set of aerosol and rainwater properties and SOC scavenging. The present study was conducted at a background site in central Europe, as a follow up with a previously published research (Škrdlíková et al., 2011) in which PAH scavenging efficiencies were determined at that site. The specific objectives of the present study were to (1) measure the concentrations of PAHs, PCBs, and CPs in air and rainwater samples, (2) estimate wet depositional fluxes of target compounds in the study area, (3) determine wet scavenging efficiencies and scavenged mass fractions of the analytes, and (4) investigate the potential relationships between analyte scavenging and meteorological parameters as well as aerosol and rain characteristics. PAHs, PCBs, and CPs were selected because they are widespread atmospheric contaminants and have been found in air at background locations across Europe (Dvorská et al., 2009; Gioia et al., 2007; Roots et al., 2010; Halse et al., 2011). In addition, these chemicals can persist in the

environment due to their resistance to photolytic, chemical, and biological degradation (Yolsal et al., 2014).

2 Experimental

2.1 Sampling site

5 The sampling was conducted at Košetice observatory located in an open land about 70 km southeast of Prague in the Czech Republic (Fig. S1 in the Supplement). The observatory is classified as a European background site (Dvorská et al., 2008). It is managed by the Czech Hydrometeorological Institute (CHMI) and is part of the European Monitoring and Evaluation Program. The area is covered with snow two to three
10 months per year with the mean annual temperature and precipitation of 7.1 °C and 625 mm. The potential source of pollution in the region is the nearby highway (~ 7 km west of the observatory) connecting Prague to Brno. In addition, due to having dominant westerly winds, atmospherically transported contaminants from the industrial and urban areas of the Czech Republic, notably Prague, may contribute to the pollution in
15 the area.

2.2 Sampling procedure

Rainwater and air samples were collected during the period of December 2011 to January 2014 (Table S1 in the Supplement). A total of 231 rain samples were collected using an automatic precipitation sampler (Baghira, Czech Republic) equipped with a 1 m²
20 stainless steel collection funnel, a lid, and a 5 L glass collection bottle. The opening and closing of the lid is triggered by a precipitation sensor mounted on the sampler. During sampling, rain drains from the funnel and accumulates in the collection bottle. The collected sample volumes were between 0.5 and 1 L, depending on the amount of rainfall for each event. Larger or smaller volumes were discarded – the latter was not deemed
25 to provide enough analyte mass for trace analysis.

26943

PAHs, PCBs, and CPs in background air in central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air samples were collected on quartz fiber filters (QFF) (Grade QM-A, 10.1 cm ID, Whatman, UK) and in polyurethane foam (PUF) plugs (55 cm ID, 50 cm height, Organika, Poland) using a Graseby Andersen PS-1 high-volume sampler (GA, USA) with a flow rate of $17 \text{ m}^3 \text{ h}^{-1}$. For each sampling period, two consecutive PUF plugs and one filter paper were deployed. A total of 162 gaseous and particulate air samples were collected during the period of study. Nevertheless, only samples from 54 events were considered for determining scavenging ratios because only these were followed by rain events. The collection of these samples started between 6 to 35 h prior to the onset of rain (predicted by CHMI) and stopped when rainfall started (Table S2 in the Supplement). This sampling approach was chosen in order to sample rainwater from the same air mass as that of PM. In addition, PM has been suggested to be removed from a travelling air parcel by up to 8 and 23 % in the first 24 and 72 h after the onset of rain, respectively (Wiman et al., 1990; Škrdlíková et al., 2011). This means that prolonging the air sampling beyond the onset of rain would potentially lead to underestimating contaminant concentrations in air and, therefore, overestimating scavenging ratios and washed-out mass fractions. In addition, relative humidity substantially increases during rainfall and this would potentially affect gas-particle partitioning of SOCs in the air leading to underestimation or overestimation of gaseous and particulate fraction of contaminants.

Field blanks for air samples were prepared at the site following the standard protocol for mounting QFF and PUF plugs onto the sampler without turning on the sampler. No field blanks were generated for rain samples.

2.3 Meteorological and other supporting data

Cloud top heights were estimated from radiosonde measurements over Prague (Station ID. 11520) (IGRA, 2014) (Table S2 in the Supplement). Meteorological parameters – i.e. near-ground temperature, precipitation type and intensity, cloud base heights (determined through on-site ceilometer measurements), concentrations of $\text{PM}_{2.5}$ and PM_{10} , and their ionic species (i.e. SO_4^{2-} , NO_3^- , NH_4^+), elemental carbon (EC), and or-

ganic carbon (OC) contents – were obtained from CHMI (Tables S2 and S3 in the Supplement). Aerosol number size distribution data, used for calculating the aerosol surface area, was obtained from the Academy of Sciences of the Czech Republic. Analyte physico-chemical properties were obtained from Estimation Programs Interface Suite 4.11 (USEPA, 2012). The weather charts used to determine the occurrence of frontal passage over the study site were obtained from Berliner Wetterkarte (BWK, 2013).

2.4 Chemical analysis and quality control

QFF and PUF samples were extracted with dichloromethane using an automatic extraction system (Büchi B-811, Switzerland). PUF plugs ($n = 2$) related to each sampling period were extracted together. Field blanks were extracted along with each set of 10 samples. The extracts were concentrated under a gentle stream of nitrogen in ambient temperature and fractionated using a silica column for PAHs and a sulfuric acid modified silica column for PCBs and CPs. Analytes in rainwater were extracted using solid-phase extraction (C18 Speedisks, Bakerbond, the Netherlands). The analytes were later eluted using 40 mL of (1 : 1) dichloromethane: *n*-hexane, concentrated under a gentle stream of nitrogen, and passed through a glass column packed with anhydrous sodium sulfate (1 cm ID, 3 cm height) to remove residual water. Prior to extraction, all air and rainwater samples and blanks were spiked with a solution containing d_8 -naphthalene, d_{10} -phenanthrene, d_{12} -perylene, PCB 30, and PCB 185, which were used as recovery standards.

Samples were analyzed for 26 parent PAHs (i.e. naphthalene (NAP), acenaphthylene (ACY), acenaphthene (ACE), fluorene (FLN), phenanthrene (PHE), anthracene (ANT), fluoranthene (FLT), pyrene (PYR), benzo(*a*)anthracene (BAA), chrysene (CHR), benzo(*b*)fluoranthene (BBF), benzo(*k*)fluoranthene (BKF), benzo(*a*)pyrene (BAP), indeno(123-*cd*)pyrene (IPY), dibenz(*a,h*)anthracene (DHA), benzo(*g,h,i*)perylene (BPE), benzo(*b*)fluorene (BFN), benzo(*g,h,i*)fluoranthene (BGF), cyclopenta(*c,d*)pyrene (CPP), triphenylene (TPH), benzo(*j*)fluoranthene

**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(BJF), benzo(*e*)pyrene (BEP), perylene (PER), dibenz(*a,c*)anthracene (DCA), anthanthrene (ATT), coronene (COR)), one heterocyclic PAH (i.e. benzo-naphthothiophene (BNT)), and one alkylated PAH (i.e. retene (RET)) using a Hewlett-Packard gas chromatograph (GC 6890) interfaced to a Hewlett-Packard mass selective detector (MS 5973). Seven PCBs (i.e. PCB 28, PCB 52, PCB 101, PCB 118, PCB 153, PCB 138, PCB 180) and 13 CPs, namely α -hexachlorocyclohexane (HCH), β -HCH, γ -HCH, δ -HCH, ε -HCH, *o,p'*-dichlorodiphenyldichloroethylene (DDE), *p,p'*-DDE, *o,p'*-dichlorodiphenyldichloroethane (DDD), *p,p'*-DDD, *o,p'*-dichlorodiphenyltrichloroethane (DDT), *p,p'*-DDT, pentachlorobenzene (PeCB), hexachlorobenzene (HCB), were analyzed using an Agilent GC (7890) coupled with an Agilent Triple Quadrupole MS/MS (7000B). All analytes were separated on a J&W Scientific capillary column (HP-5ms, 0.25 mm ID, 0.25 μ m film thickness). D₁₄-*p*-terphenyl and PCB 121 were used as internal standards for PAH and PCB/CP analysis, respectively.

Limits of quantification (LOQ) for analytes were calculated based on instrument detection limits, which in turn are determined using three times the chromatogram baseline noise level. The median LOQs for individual PAHs and PCBs/CPs in air samples were 0.006 and 0.0005 ng m⁻³, respectively. The median LOQs in rainwater samples were 0.05 and 0.45 ng L⁻¹, respectively. LOQ values were used in cases where analyte concentrations in field blanks were < LOQ. The mean concentrations of analytes in five field blanks were subtracted from those in the corresponding samples. The concentrations that were lower than mean +3 standard deviations of those in field blanks were considered below method detection limit (< MDL) and were substituted with LOQ/2 for calculation of averages.

Analysis of ionic species in rainwater was done using a Hewlett-Packard capillary electrophoresis system (HP 3D CE). Measurements were performed using an uncoated silica capillary column (75 μ m ID, 70 cm length). The applied separation voltage was 10 kV with hydrodynamic injection of 50 mbar (20 s)⁻¹. Analytes were detected indirectly at 254 nm wavelength. The electrolyte was composed of 5 mmol sodium chro-

where, b_r and b_a are analyte burdens (ng m^{-2}) in rainwater and air, respectively, and h (m) is the height of the air column subject to precipitation, which corresponds to the cloud height from which precipitation originates.

3 Results and discussion

3.1 Concentrations and distribution of analytes

3.1.1 Concentrations in air

NAP was targeted for analysis but it did not pass our quality control criteria and, therefore, is not discussed hereafter. In addition, only analytes that were detected in $> 10\%$ of the samples are discussed herein (Tables 1 and 2). Σ PAH (i.e. the sum of 27 PAHs) concentrations in gas phase ranged from 0.6 to 138.6 (mean \pm standard deviation (SD): 11.3 ± 15.7) ng m^{-3} (Table 1). PAHs with molecular mass < 228 Da, namely ACY, ACE, FLN, PHE, ANT, FLT, and PYR, were predominant in gas phase and, on average, accounted for 93 % of the Σ PAH concentrations. PHE, FLN, and FLT showed the highest mean contributions accounting for 42, 24, and 12 % of the Σ PAH gaseous concentrations, respectively. BKF, BAP, IPY, DHA, BPE, BNT, CPP, BJF, BEP, PER, DCA, ATT, and COR were detected in $< 10\%$ of the gaseous samples ($n = 162$) (Table 1). The Σ PAH concentrations in the particulate phase ranged from 0.1 to 189.3 (mean \pm SD: 9.3 ± 21.2) ng m^{-3} (Table 1). In sum, PAHs with molecular mass > 228 Da were predominant in particulate phase and, on average, contributed to 67 % of the Σ PAH particulate concentrations. However, FLT and PYR showed the highest individual contributions and, on average, accounted for 13 and 11 % of the Σ PAH concentrations, respectively. The sum of the concentrations of gas and particulate phase PAHs ranged from 0.7 to 327.9 ng m^{-3} , with the mean \pm SD being 20.6 ± 35.9 ng m^{-3} . The PAH concentration ranges in gas and particulate phase are noticeably higher than those measured by Škrdlíková et al. (2011) (i.e. 0.9 to 34.6 ng m^{-3} and 0.1 to 16.8 ng m^{-3} , respectively) for

samples collected from the same site during 2007–2008 period, but are by far lower than those reported by Holoubek et al. (2007) for the period between 1996 and 2005 (i.e. 0.4 to 208 ng m⁻³ and 0.1 to 359 ng m⁻³, respectively).

Except for *p,p'*-DDE, the targeted chlorinated compounds were mainly detected in gas phase (i.e. < 10% in particulate phase) (Table 2). The concentrations of *p,p'*-DDE in particulate phase ranged from < MDL to 6.8 (mean ± SD: 0.4 ± 0.7) pg m⁻³. In gas phase, *β*-HCH, *δ*-HCH, *ε*-HCH, and *o,p'*-DDD were detected in < 10% of all samples (*n* = 162). The concentrations in gas phase ranged from < MDL to 43.5 (mean ± SD: 8.1 ± 7.2) pg m⁻³ for Σ PCBs (i.e. the sum of the concentrations of PCB 28, 52, 101, 118, 138, 153, 180), < MDL to 73.8 (11.0 ± 11.6) pg m⁻³ for Σ HCHs (i.e. the sum of the concentrations of *α*- and *γ*-HCH), < MDL to 104.3 (23.2 ± 19.8) pg m⁻³ for Σ DDTs (i.e. the sum of the concentrations of *o,p'*-DDE, *p,p'*-DDE, *p,p'*-DDD, *o,p'*-DDT, and *p,p'*-DDT) and < MDL to 332.3 (96.1 ± 42.0) pg m⁻³ for Σ CBs (i.e. the sum of the concentrations of PeCB and HCB) (Table 2). On average, PCB 28 (40%), *γ*-HCH (56%), *p,p'*-DDE (83%), and HCB (90%) predominated Σ PCB, Σ HCH, Σ DDT, and Σ CB concentrations, respectively. The sum of the concentrations of gas and particulate phase PCBs and CPs ranged from < MDL to 44.6 and < MDL to 351.7 pg m⁻³, respectively. The measured concentrations in the present study are considerably lower than those reported by Holoubek et al. (2007) for the samples collected from the same location between 1996 and 2005 (i.e. < MDL to 390 pg m⁻³ for Σ PCBs, < MDL to 771 pg m⁻³ for Σ HCHs, 1 to 207 pg m⁻³ for Σ DDTs, and < MDL to 831 pg m⁻³ for HCB). As production and use of these compounds are banned in Europe, the relatively low concentrations in the present study could indicate emission due to volatilization from contaminated soil around Košetice (Fig. S1). This argument is supported by significant correlations found between the ambient temperature at the sampling site and the measured concentrations of PCB 28 (*r* = 0.70, *P* < 0.05, *n* = 162), *γ*-HCH (*r* = 0.74, *P* < 0.05), and *p,p'*-DDE (*r* = 0.71, *P* < 0.05). Despite having the highest mean concentration among the other chlorinated analytes, relatively small correlation was found between the concentrations of HCB and ambient temperature changes (*r* = 0.25). This could be due to

PAHs, PCBs, and CPs in background air in central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the fact that this compound has different source pathways compared to all other chlorinated chemicals investigated in this study. Although banned, HCB could potentially be released to the environment as an unintended byproduct of organic solvent and aluminum manufacturing and waste burning (EPER, 2014). In addition, this compound is present in some pesticide formulations, such as chlorothalonil, which is currently registered for use in Europe.

3.1.2 Concentrations in rain

Σ PAH concentrations in rainwater ranged from < MDL to 2.1×10^3 (mean \pm SD: 173.3 ± 256.1) ng L^{-1} (Table 3), predominated (mean: 69%) by congeners with < 228 Da molecular mass. FLT, PHE, and PYR showed the highest individual contributions and, on average, accounted for 35, 14, and 10% of the Σ PAH concentrations. The Σ PAH concentration range in the present study was higher than the one reported by Škrdlíková et al. (2011) for Košetice (i.e. 7.1 to 485.9 ng L^{-1}) but noticeably lower than that measured by Holoubek et al. (2007) (i.e. 2.4 to 6310 ng L^{-1}). The concentrations of PCBs and CPs, targeted for analysis in rainwater samples, were below LOQs and, therefore, are not discussed hereafter.

3.2 Wet deposition fluxes

Σ PAH daily wet deposition fluxes ranged from < MDL to 5.5×10^3 (mean \pm SD: 632.9 ± 900.1) ng m^{-2} (Table 4). Σ PAH seasonal fluxes were noticeably higher during winter and spring compared to summer and autumn (Table 5), which is in agreement with previous observations in Europe (Kiss et al., 2001; Škrdlíková et al., 2011). Seasonal changes in SOC fluxes could be due to different factors, such as variations in atmospheric concentration, precipitation amount (providing that atmospheric concentrations do not change drastically between seasons), and SOC gas-particle partitioning behavior. The increase in PAH fluxes during winter and spring cannot be explained by changes in precipitation amounts, as indicated in Table 5. However, atmospheric con-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



centrations of PAHs were noticeably higher in winter than they were in other seasons (Fig. 1), in agreement with a previously published study in Europe (Lammel et al., 2011). This may be due to higher emission rate, lower boundary layer mixing height (Birgöl et al., 2011), and lower concentrations of OH radicals in winter (Halsall et al., 2001). This may partly explain the higher fluxes seen during winter periods. Finally, lower temperatures in winter could potentially shift partitioning of the more temperature sensitive SOCs towards particulate phase, and particle scavenging, as opposed to gas scavenging, was suggested to be the dominant wet scavenging mechanism for the removal of PAHs from the atmosphere (Bidleman, 1988; Poster and Baker, 1996; Offenberg and Baker, 2002). In the present study, significant negative regressions were found between near-ground temperature and θ for a number of PAHs including PYR ($r^2 = 0.72$, $P < 0.05$, $n = 150$), FLT ($r^2 = 0.69$, $n = 155$), RET ($r^2 = 0.75$, $n = 156$), and TPH ($r^2 = 0.73$, $n = 162$) (Table S4 in the Supplement), which could provide further explanation for higher fluxes during cold seasons.

3.3 Scavenging ratios

Scavenging ratios were only determined for those events with air and rainwater sampled from the same air mass ($n = 54$). The rejection criterion was that frontal passage did not occur over the site prior (± 1 h uncertainty) to the onset of rain. Out of 54 pair samples, only 32 met the abovementioned criterion (Table S2 in the Supplement) and, hence, W_t ratios were determined for 32 rain events.

The relative importance of gas and particle scavenging can be calculated theoretically using W_t and gas scavenging ratio obtained through dimensionless Henry's law constant (Offenberg and Baker, 2002; Poster and Baker, 1996). However, this approach was avoided in the present study due to the fact that the equilibrium partitioning of SOCs between gas phase and raindrops, and consequently gas scavenging ratios, is greatly affected by temperature, which may change drastically from cloud to ground level. This process is non-linear and a representative temperature is unknown. This means that Henry's law constants, which are corrected using ground temperatures,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



may not necessarily represent the true gas scavenging ratios related to falling rain-drops. One should also note that theoretical gas scavenging ratios for SOCs in gas phase, being the inverse of their Henry's law constants, may not entirely reflect the underlying mechanism of the compound removal from the atmosphere. For instance, PAHs with higher vapor pressure normally show higher Henry's law constants, as can be seen from experimental data (Bamford et al., 1999). This would result in relatively lower gas scavenging ratios for gaseous PAHs with higher vapor pressure and solubility, which contradicts the concept of SOC gas scavenging.

In the present study, the concentrations of ionic species (i.e. the sum of SO_4^{2-} , NO_3^-) in PM and those in adjacent rainwater samples were significantly correlated ($r = 0.71$, $P < 0.05$, $n = 23$). This indicates the efficient removal of PM by rainfall and highlights the importance of particle scavenging in the present study. In general, W_f ratios for individual PAHs ranged from 0 (in cases where concentrations in rainwater were $< \text{LOQ}$) to 3.5×10^5 , with the exception of FLT for which W_f ranged between 3.7×10^3 and 1.3×10^6 (median: 1.4×10^4) (Table 6). Overall, W_f values for relatively more volatile PAHs (log K_{oa} between 6.27 and 7.57), namely ACY, ACE, FLN, ANT, and PHE, were up to 10^4 , whereas the values for the rest of PAHs (log $K_{\text{oa}} > 8.70$) were up to 10^6 , but mainly dominated by the values close to 10^5 (Table 6). Taking into account that SOCs are more efficiently removed from the atmosphere by particle scavenging, the current results may imply that increase in affinity of individual PAHs towards organic phase would overall lead to higher scavenging efficiencies. This, together with other parameters that affect analyte gas-particle partitioning, namely temperature, aerosol surface area and chemical composition (e.g. EC and OC contents) (Junge, 1977; Pankow, 1987; Bidleman, 1988), and factors that affect particle removal rate including rain intensity, aerosol and raindrop size and collision efficiency (Mircea et al., 2000; Poster and Baker, 1996; Slinn et al., 1978), may contribute to the observed variability in measured scavenging ratios.

Since chlorinated compounds in the present study were mainly detected in gas phase in the air (Table 2), it is likely that their removal from the atmosphere was deter-

mined by gas scavenging. The fact that concentrations of these analytes in rainwater were below detection limits may provide further evidence for this argument, as the magnitude of gas scavenging is negligible.

The upper- and lower-bound scavenged mass fractions of target analytes corresponding to cloud base and top heights were determined for 32 events, except in one case for which cloud top height was not available (Table S2). Σ PAH upper-bound scavenged mass fractions ranged from 1.2×10^{-3} to 0.8 (median: 5.3×10^{-2}), whereas lower-bound mass fractions were between 5.2×10^{-4} and 0.1 (median: 9.7×10^{-3}) (Table 7). The median values for the analyte lower-bound mass fractions were nearly an order of magnitude different from those reported by Škrdlíková et al. (2011) for the same site, except for FLN and CHR, which were found to be in agreement (Table 7). This could potentially be related to the relatively small number of events considered in that study ($n = 10$). It should be noted that scavenged mass fractions were calculated assuming that analyte concentrations near ground represented those within the air column up to cloud top – i.e. assuming perfect mixing throughout the column. The true scavenged mass fractions could potentially be higher due to a negative vertical concentration gradient (Škrdlíková et al., 2011).

3.4 Factors affecting the scavenging ratios

As can be seen from Tables 6 and S3, W_t ratios for Σ PAHs in the present study ranged from 2.4×10^3 to 2.3×10^5 , with noticeable variability between the events (mean \pm SD: $1.7 \times 10^4 \pm 3.8 \times 10^4$). As discussed earlier, different parameters could contribute to the variability seen in scavenging ratios. In this section, we investigate potential relationships between PAH scavenging efficiencies and precipitation intensity, near-ground temperature, the concentrations of $\text{PM}_{2.5}$ and PM_{10} , PM ionic species (i.e. the sum of the concentrations of SO_4^{2-} , NO_3^- , and NH_4^+), EC and OC contents, PM surface area, and rainwater ionic species (i.e. the sum of SO_4^{2-} , NO_3^- , and Cl^-). As can be seen from Table S3, the highest Σ PAH W_t was found for the samples collected on 14 April 2012.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**PAHs, PCBs, and CPs
in background air in
central Europe**P. Shahpoury et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ratio was nearly two orders of magnitude higher compared to that obtained for another event with similar precipitation type and ambient temperature (i.e. 11–12 October 2012). Apart from the concentrations of PM_{10} and ionic species in PM and rainwater, the rest of the parameters were somewhat similar for these two events (Table S3).

The concentrations of PM_{10} as well as those for ionic species in PM and rainwater for the former event were 1.4, 1.8, and 11 times the values found for the latter event. This observation could suggest that PAH scavenging by rain is more efficient in conditions where PM and rainwater contain high quantities of ionic species.

In order to explore the relative importance of the abovementioned parameters in relation to scavenging ratios, correlation analysis was performed between these variables. Among all parameters, aerosol surface area was only available for 2012. In addition, aerosol EC and OC contents could only be obtained for 15 rain events out of 32, as they were recorded once every six days (Table S3). Owing to the fact that EC and OC have high importance in SOC gas-particle partitioning (Lohmann and Lammel, 2004), and in order to enhance the comparability of results, for all other parameters, correlation analysis was only performed for the corresponding rain events. The results showed that the concentrations of rainwater ionic species were significantly correlated with Σ PAH scavenging ratios ($r = 0.70$, $P < 0.05$, $n = 15$), whereas weak correlations were found for all other parameters, except the ratio of PM ionic species/ PM_{10} , for which a moderate correlation was found ($r = 0.34$, $P > 0.05$). However, when the data related to 14 April 2012 (with exceptionally high W_t ratio) was excluded from the analysis, the suggested importance of rainwater and PM ionic species decreased ($r = 0.14$ and 0.07 , respectively, $P > 0.05$, $n = 14$). This indicates that the initial analysis may have been overpowered by significantly high values of ionic species related to a single event. Without the data point related to 14 April 2012, moderate correlations were obtained between Σ PAH W_t ratios and EC/PM_{10} ($r = 0.34$, $P > 0.05$) and OC/PM_{10} ($r = 0.37$, $P > 0.05$). Although not statistically significant, the current results suggest that the aerosol EC and OC contents are important parameters in determining PAH scavenging efficiencies. This may reflect the affinity of PAHs towards EC and OC, rep-

representing adsorption and absorption processes (Lohmann and Lammel, 2004). However, the magnitude of these effects could be different when considering individual PAHs. For instance, the coefficients of correlation between W_t and EC/PM₁₀ ratios for PAHs with moderate to high particulate mass fractions, namely PYR (mean $\theta = 0.40$), CHR (mean $\theta = 0.74$), and BBF (mean $\theta = 0.97$), were 0.38, 0.20, and 0.33 ($n = 10$), whereas the coefficients for correlation with OC/PM₁₀ ratios were 0.51, 0.53, and 0.03, respectively. Despite not having statistical significance, the results may indicate that the aerosol OC content was more effective than EC for wet scavenging of PYR and CHR. It should be noted that precipitation type (i.e. snow vs. rain) was only available for eight events out of 15. These were mainly identified as rainfall (Table S3), which ruled out the potential impact of precipitation type on scavenging efficiencies. The lack of significant linear relationships between scavenging ratios and aerosol and rainwater properties indicates that the process is not controlled by a single factor but rather by a combination of parameters. The findings of the present study highlight the need for incorporating other PM components, such as EC and OC, in air pollution models for more accurate estimation of particulate PAH scavenging.

4 Conclusions

The results suggest that volatilization from contaminated soil was the main source of chlorinated compounds in the air over Košetice. Higher Σ PAH fluxes in winter were possibly related to higher PAH atmospheric concentrations and particulate mass fractions in that season. Particle scavenging was found to be significant in the present study, as indicated by the strong correlation between rainwater and PM ionic species. In sum, greater scavenging ratios were found for less volatile PAHs, highlighting the importance of SOC gas-particle partitioning in determining their scavenging efficiencies. The results indicated that PAH wet scavenging could be more efficient in cases where the concentrations of ionic species in PM and rainwater are high. In addition, the high affinity of PAHs towards EC and OC were found to be reflected in the scavenging effi-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ciency. In most model applications which study PAHs, particulates are scavenged either as in conventional air pollution modeling – i.e. accounting for PM ionic composition, but no other PM components (parameterization based on Köhler theory or empirical approaches based on cloud droplet number) (Abdul-Razzak and Ghan, 2000; Gong et al., 2003) – or with insoluble aerosols or aerosol modes (Sehili and Lammel, 2007; Friedman and Selin, 2012). Future studies would need to focus on in-depth analysis of PM chemical composition with the aim to include OC and EC in the parameterizations of particulate PAH wet deposition.

**The Supplement related to this article is available online at
doi:10.5194/acpd-14-26939-2014-supplement.**

Acknowledgements. We thank Miroslava Bittová and Jan Kuta (Masaryk University) for ion analyses, Jaroslav Pekarek (CHMI) for providing meteorological data, and Vladimir Zdimal (Institute for Chemical Process Fundamentals, the Academy of Sciences of the Czech Republic) for providing aerosol number size distribution data. This research was supported by the Granting Agency of the Czech Republic (GACR project No. P503/11/1230) and has been co-funded by the European Social Fund and the state budget of the Czech Republic. The research infrastructure (i.e. RECETOX) has been supported by the Czech Ministry of Education (LO1214 and LM2011028).

The service charges for this open access publication have been covered by the Max Planck Society.

References

Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation 2. Multiple aerosol types, *J. Geophys. Res.*, 105, 6837–6844, 2000.

**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Agrell, C., Larsson, P., Okla, L., and Agrell, J.: PCB congeners in precipitation, wash out ratios and depositional fluxes within the Baltic Sea region, Europe, Atmos. Environ., 36, 371–383, 2002.

Atlas, E. and Giam, C. S.: Ambient concentration and precipitation scavenging of atmospheric organic pollutants, Water. Air. Soil. Poll., 38, 19–36, 1988.

Bamford, H. A., Poster, D. L., and Baker, J. E.: Temperature dependence of Henry's law constants of thirteen polycyclic aromatic hydrocarbons between 4 °C and 31 °C, Environ. Toxicol. Chem., 18, 1905–1912, 1999.

Bidleman, T. F.: Atmospheric processes, Environ. Sci. Technol., 22, 361–367, 1988.

Birgül, A., Tasdemir, Y., and Cindoruk, S. S.: Atmospheric wet and dry deposition of polycyclic aromatic hydrocarbons (PAHs) determined using a modified sampler, Atmos. Res., 101, 341–353, 2011.

Blais, J. M.: Biogeochemistry of persistent bioaccumulative toxicants: Processes affecting the transport of contaminants to remote areas, Can. J. Fish. Aquat. Sci., 62, 236–243, 2005.

BWK: Verein Berliner Wetterkarte, e. V. (Ed.), www.berliner-wetterkarte.de (last access: 16 February 2014) 2013.

Dvorská, A., Lammel, G., Klánová, J., and Holoubek, I.: Košetice, Czech Republic – ten years of air pollution monitoring and four years of evaluating the origin of persistent organic pollutants, Environ. Pollut., 156, 403–408, 2008.

Dvorská, A., Lammel, G., and Holoubek, I.: Recent trends of persistent organic pollutants in air in central Europe – Air monitoring in combination with air mass trajectory statistics as a tool to study the effectivity of regional chemical policy, Atmos. Environ., 43, 1280–1287, 2009.

EPER: European Pollutant Emission Register, <http://www.eea.europa.eu/data-and-maps/data/eper-the-european-pollutant-emission-register-4> (last access: 04 March 2014), 2014.

Friedman, C. L. and Selin, N. E.: Long-range atmospheric transport of polycyclic aromatic hydrocarbons: A global 3-D model analysis including evaluation of arctic sources, Environ. Sci. Technol., 46, 9501–9510, 2012.

Gioia, R., Sweetman, A. J., and Jones, K. C.: Coupling passive air sampling with emission estimates and chemical fate modeling for persistent organic pollutants (POPs): a feasibility study for northern Europe, Environ. Sci. Technol., 41, 2165–2171, 2007.

Gong, S. L., Barrie, L. A., Blanchet, J. P., von Salzen, K., Lohmann, U., Lesins, G., Spacek, L., Zhang, L. M., Girard, E., Lin, H., Leaitch, R., Leighton, H., Chylek, P., and Huang, P.: Canadian Aerosol Module: A size-segregated simulation of atmospheric aerosol processes for

**PAHs, PCBs, and CPs
in background air in
central Europe**

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



climate and air quality models 1. Module development, *J. Geophys. Res.*, 108, AAC 3-1–AAC 3-16, 2003.

Grimalt, J. O., Fernandez, P., Berdie, L., Vilanova, R. M., Catalan, J., Psenner, R., Hofer, R., Appleby, P. G., Rosseland, B. O., Lien, L., Massabuau, J. C., and Battarbee, R. W.: Selective trapping of organochlorine compounds in mountain lakes of temperate areas, *Environ. Sci. Technol.*, 35, 2690–2697, 2001.

Hageman, K. J., Simonich, S. L., Campbell, D. H., Wilson, G. R., and Landers, D. H.: Atmospheric deposition of current-use and historic-use pesticides in snow at National Parks in the Western United States, *Environ. Sci. Technol.*, 40, 3174–3180, 2006.

Halsall, C. J., Sweetman, A. J., Barrie, L. A., and Jones, K. C.: Modelling the behaviour of PAHs during atmospheric transport from the UK to the Arctic, *Atmos. Environ.*, 35, 255–267, 2001.

Halse, A. K., Schlabach, M., Eckhardt, S., Sweetman, A., Jones, K. C., and Breivik, K.: Spatial variability of POPs in European background air, *Atmos. Chem. Phys.*, 11, 1549–1564, doi:10.5194/acp-11-1549-2011, 2011.

Holoubek, I., Klánová, J., Jarkovský, J., and Kohoutek, J.: Trends in background levels of persistent organic pollutants at Kosec observatory, Czech Republic. Part I. ambient air and wet deposition 1996–2005, *J. Environ. Monitor.*, 9, 557–563, 2007.

IGRA: Integrated Global Radiosonde Archive, <ftp://ftp.ncdc.noaa.gov/pub/data/igra> (last access: 23. February 2014), 2014.

Junge, C. E.: *Fate of Pollutants in the Air and Water Environments*, edited by: Suffett, I. H., John Wiley and Sons, New York, 1977.

Kaupp, H. and McLachlan, M. S.: Distribution of polychlorinated dibenzo-P-dioxins and dibenzofurans (PCDD/Fs) and polycyclic aromatic hydrocarbons (PAHs) within the full size range of atmospheric particles, *Atmos. Environ.*, 34, 73–83, 2000.

Kiss, G., Varga-Puchony, Z., Tolnai, B., Varga, B., Gelencsér, A., Krivácsy, Z., and Hlavay, J.: The seasonal changes in the concentration of polycyclic aromatic hydrocarbons in precipitation and aerosol near Lake Balaton, Hungary, *Environ. Pollut.*, 114, 55–61, 2001.

Lammel, G., Novák, J., Landlová, L., Dvorská, A., Klánová, J., Čupr, P., Kohoutek, J., Reimer, E., and Škrdlíková, L.: Sources and distributions of polycyclic aromatic hydrocarbons and toxicity of polluted atmosphere aerosols, in: *Urban Airborne Particulate Matter*, edited by: Zereini, F. and Wiseman, C. L. S., Springer, Berlin Heidelberg, 39–62, 2011.

Ligocki, M. P., Leuenberger, C., and Pankow, J. F.: Trace organic compounds in rain. III. particle scavenging of neutral organic compounds, *Atmos. Environ.*, 19, 1619–1626, 1985.

PAHs, PCBs, and CPs in background air in central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Liu, F., Xu, Y., Liu, J., Liu, D., Li, J., Zhang, G., Li, X., Zou, S., and Lai, S.: Atmospheric deposition of polycyclic aromatic hydrocarbons (PAHs) to a coastal site of Hong Kong, South China, *Atmos. Environ.*, 69, 265–272, 2013.

Lohmann, R. and Lammel, G.: Adsorptive and absorptive contributions to the gas-particle partitioning of polycyclic aromatic hydrocarbons: state of knowledge and recommended parametrization for modeling, *Environ. Sci. Technol.*, 38, 3793–3803, 2004.

McLachlan, M. S. and Sellström, U.: Precipitation scavenging of particle-bound contaminants – a case study of PCDD/Fs, *Atmos. Environ.*, 43, 6084–6090, 2009.

Mircea, M., Stefan, S., and Fuzzi, S.: Precipitation scavenging coefficient: influence of measured aerosol and raindrop size distributions, *Atmos. Environ.*, 34, 5169–5174, 2000.

Offenberg, J. H. and Baker, J. E.: Precipitation scavenging of polychlorinated biphenyls and polycyclic aromatic hydrocarbons along an urban to over-water transect, *Environ. Sci. Technol.*, 36, 3763–3771, 2002.

Pankow, J. F.: Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere, *Atmos. Environ.*, 21, 2275–2283, 1987.

Poster, D. L. and Baker, J. E.: Influence of submicron particles on hydrophobic organic contaminants in precipitation. 2. scavenging of polycyclic aromatic hydrocarbons by rain, *Environ. Sci. Technol.*, 30, 349–354, 1996.

Roots, O., Roose, A., Kull, A., Holoubek, I., Čupr, P., and Klánová, J.: Distribution pattern of PCBs, HCB and PeCB using passive air and soil sampling in Estonia, *Environ. Sci. Pollut. R.*, 17, 740–749, 2010.

Sehili, A. M. and Lammel, G.: Global fate and distribution of polycyclic aromatic hydrocarbons emitted from Europe and Russia, *Atmos. Environ.*, 41, 8301–8315, 2007.

Škrdlíková, L., Landlová, L., Klánová, J., and Lammel, G.: Wet deposition and scavenging efficiency of gaseous and particulate phase polycyclic aromatic compounds at a central European suburban site, *Atmos. Environ.*, 45, 4305–4312, 2011.

Slinn, W. G. N., Hasse, L., Hicks, B. B., Hogan, A. W., Lal, D., Liss, P. S., Munnich, K. O., Sehmel, G. A., and Vittori, O.: Some aspects of the transfer of atmospheric trace constituents past the air-sea interface, *Atmos. Environ.*, 12, 2055–2087, 1978.

USEPA: United States Environmental Protection Agency, Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.11, 2012.

PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** PAH concentrations and detection frequencies (DF) in air ($n = 162$).

	Gaseous Concentrations (ng m^{-3})			DF %	Particulate Concentrations (ng m^{-3})			DF %
	Min.	Max.	Mean \pm SD		Min.	Max.	Mean \pm SD	
ACY	< MDL	34.5	1.0 ± 3.0	87	< MDL	1.9	$7.8 \times 10^{-2} \pm 2.4 \times 10^{-1}$	53
ACE	< MDL	3.2	0.3 ± 0.5	90	< MDL	0.2	$1.1 \times 10^{-2} \pm 3.1 \times 10^{-2}$	22
FLN	< MDL	26.3	2.7 ± 4.0	98	< MDL	2.5	$1.1 \times 10^{-1} \pm 3.4 \times 10^{-1}$	49
PHE	< MDL	45.0	4.7 ± 5.9	98	< MDL	14.6	0.8 ± 2.2	72
ANT	< MDL	4.7	$1.3 \times 10^{-1} \pm 4.0 \times 10^{-1}$	81	< MDL	1.5	$6.3 \times 10^{-2} \pm 1.9 \times 10^{-1}$	53
FLT	7.0×10^{-2}	13.8	1.3 ± 1.6	100	3.0×10^{-3}	26.1	1.3 ± 3.3	99
PYR	3.9×10^{-3}	7.7	0.6 ± 0.8	93	3.5×10^{-3}	20.7	1.1 ± 2.5	99
BAA	< MDL	0.3	$1.4 \times 10^{-2} \pm 3.7 \times 10^{-2}$	38	< MDL	11.1	0.5 ± 1.2	87
CHR	< MDL	0.4	$6.0 \times 10^{-2} \pm 6.8 \times 10^{-2}$	95	< MDL	15.8	0.7 ± 1.7	97
BBF	< MDL	0.9	$1.3 \times 10^{-2} \pm 8.2 \times 10^{-2}$	14	< MDL	13.6	0.7 ± 1.5	96
BKF	< MDL	0.3	$6.2 \times 10^{-3} \pm 2.6 \times 10^{-2}$	4	< MDL	5.0	0.3 ± 0.6	88
BAP	< MDL	0.5	$8.7 \times 10^{-3} \pm 4.2 \times 10^{-2}$	3	< MDL	9.1	0.4 ± 0.9	83
IPY	< MDL	0.5	$7.2 \times 10^{-3} \pm 4.6 \times 10^{-2}$	3	< MDL	10.4	0.5 ± 1.0	81
DHA				2	< MDL	1.2	$4.1 \times 10^{-2} \pm 1.1 \times 10^{-1}$	56
BPE	< MDL	0.5	$7.4 \times 10^{-3} \pm 4.2 \times 10^{-2}$	3	< MDL	6.6	0.4 ± 0.7	90
RET	6.5×10^{-3}	5.0	0.2 ± 0.4	100	< MDL	4.0	0.2 ± 0.5	73
BFN	< MDL	0.8	$5.2 \times 10^{-2} \pm 8.2 \times 10^{-2}$	80	< MDL	4.7	0.2 ± 0.5	72
BNT	< MDL	2.1×10^{-2}	$2.4 \times 10^{-3} \pm 2.2 \times 10^{-3}$	6	< MDL	0.5	$2.3 \times 10^{-2} \pm 6.2 \times 10^{-2}$	46
BGF	< MDL	0.4	$5.1 \times 10^{-2} \pm 5.4 \times 10^{-2}$	93	< MDL	7.7	0.3 ± 0.8	92
CPP	< MDL	0.2	$5.0 \times 10^{-3} \pm 2.2 \times 10^{-2}$	3	< MDL	10.3	0.3 ± 0.9	73
TPH	< MDL	0.1	$1.5 \times 10^{-2} \pm 1.5 \times 10^{-2}$	87	< MDL	2.8	$1.4 \times 10^{-1} \pm 3.2 \times 10^{-1}$	86
BJF	< MDL	0.4	$8.4 \times 10^{-3} \pm 3.9 \times 10^{-2}$	6	< MDL	7.5	0.4 ± 0.9	91
BEP	< MDL	0.5	$1.0 \times 10^{-2} \pm 4.6 \times 10^{-2}$	5	< MDL	7.2	0.4 ± 0.8	87
PER				2	< MDL	1.5	$6.9 \times 10^{-2} \pm 1.5 \times 10^{-1}$	61
DCA				2	< MDL	0.6	$2.9 \times 10^{-2} \pm 6.7 \times 10^{-2}$	55
ATT				2	< MDL	1.7	$4.7 \times 10^{-2} \pm 1.5 \times 10^{-1}$	48
COR				2	< MDL	3.3	$1.3 \times 10^{-1} \pm 3.2 \times 10^{-1}$	64
Σ PAHs	0.6	138.6	11.3 ± 15.7		0.1	189.3	9.3 ± 21.2	

SD: standard deviation; < MDL: below method detection limit; Σ PAHs: the sum of the concentrations of individual PAHs with DF > 10% in either gas or particulate phase. Concentration ranges and means are only reported for analytes with DF > 2%.

PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** PCB and CP concentrations and detection frequencies (DF) in air ($n = 162$).

	Gaseous Concentrations ($\mu\text{g m}^{-3}$)			DF %	Particulate Concentrations ($\mu\text{g m}^{-3}$)			DF %
	Min.	Max.	Mean \pm SD		Min.	Max.	Mean \pm SD	
PCB 28	< MDL	14.3	3.2 ± 2.4	88				1
PCB 52	< MDL	8.4	1.5 ± 1.4	68				1
PCB 101	< MDL	13.9	1.2 ± 1.8	60				1
PCB 118	< MDL	4.0	0.3 ± 0.5	10				< MDL
PCB 138	< MDL	3.9	0.5 ± 0.5	24				2
PCB 153	< MDL	8.0	1.0 ± 1.2	62	< MDL	3.1	0.3 ± 0.3	4
PCB 180	< MDL	2.1	0.3 ± 0.2	14	< MDL	3.3	0.2 ± 0.3	6
α -HCH	< MDL	44.5	4.4 ± 6.0	57				1
β -HCH				2				< MDL
γ -HCH	< MDL	31.4	6.6 ± 6.5	75				1
δ -HCH	< MDL	2.0	0.4 ± 0.3	5				1
ε -HCH				1				< MDL
o, p' -DDE	< MDL	1.9	0.4 ± 0.3	27				< MDL
p, p' -DDE	< MDL	87.3	19.5 ± 15.9	98	< MDL	6.8	0.4 ± 0.7	31
o, p' -DDD	< MDL	2.2	0.3 ± 0.2	6				< MDL
p, p' -DDD	< MDL	2.4	0.4 ± 0.4	17				1
o, p' -DDT	< MDL	8.9	1.5 ± 1.9	45				< MDL
p, p' -DDT	< MDL	11.2	1.6 ± 2.3	36				1
PeCB	< MDL	52.8	9.6 ± 7.3	99				2
HCB	< MDL	279.5	86.5 ± 40.4	99	< MDL	62.7	0.9 ± 6.2	7
Σ PCBs	< MDL	43.5	8.1 ± 7.2					
Σ HCHs	< MDL	73.8	11.0 ± 11.6					
Σ DDTs	< MDL	104.3	23.2 ± 19.8					
Σ CBs	< MDL	332.3	96.1 ± 42.0					

SD: standard deviation; < MDL: below method detection limit; Σ PCBs: the sum of the concentrations of PCB 28, 52, 101, 118, 138, 153, and 180; Σ HCHs: the sum of the concentrations of α - and γ -HCH; Σ DDTs: the sum of the concentrations of o, p' -DDE, p, p' -DDE, o, p' -DDD, o, p' -DDT, and p, p' -DDT; Σ CBs: the sum of the concentrations of PeCB and HCB. Note that only analytes with DF > 10% in either gas or particulate phase were considered for calculating the sums. Concentration ranges and means are only reported for analytes with DF > 2%.

PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 3.** PAH concentrations and detection frequencies (DF) in rainwater ($n = 231$).

	Concentrations (ng L ⁻¹)			DF %
	Min.	Max.	Mean ± SD	
ACY	< MDL	23.7	2.1 ± 2.8	67
ACE	< MDL	7.0	1.0 ± 1.2	54
FLN	< MDL	41.6	6.0 ± 5.8	99
PHE	< MDL	268.0	23.5 ± 29.2	99
ANT	< MDL	41.2	1.1 ± 3.2	56
FLT	< MDL	1.1 × 10 ³	61.4 ± 144.6	99
PYR	< MDL	282.5	17.7 ± 27.0	91
BAA	< MDL	76.9	3.1 ± 6.7	63
CHR	< MDL	141.8	9.1 ± 14.9	75
BBF	< MDL	85.2	6.5 ± 13.0	69
BKF	< MDL	42.3	2.8 ± 4.8	63
BAP	< MDL	50.0	2.3 ± 4.9	55
IPY	< MDL	58.2	3.6 ± 7.1	60
DHA	< MDL	3.9	0.3 ± 0.4	25
BPE	< MDL	50.9	3.6 ± 6.5	59
RET	< MDL	44.0	3.1 ± 4.6	73
BFN	< MDL	67.2	3.1 ± 6.0	62
BNT	< MDL	6.7	0.5 ± 0.7	43
BGF	< MDL	70.8	4.9 ± 7.7	70
CPP	< MDL	17.4	1.1 ± 2.2	48
TPH	< MDL	44.2	3.5 ± 5.4	67
BJF	< MDL	63.0	5.6 ± 9.7	64
BEP	< MDL	60.0	4.8 ± 7.6	73
PER	< MDL	7.8	0.5 ± 0.9	39
DCA	< MDL	7.7	0.4 ± 0.7	27
ATT	< MDL	8.0	0.3 ± 0.7	18
COR	< MDL	22.1	1.5 ± 2.9	52
Σ PAHs	< MDL	2.1 × 10 ³	173.3 ± 256.1	

SD: standard deviation; < MDL: below method detection limit; Σ PAHs: the sum of the concentrations of 27 individual PAHs listed on the table.

PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 6.** PAH scavenging ratios (W_i).

	W_i			$\log K_{oa}$	$\log K_{aw}$
	Min.	Max.	Median		
ACY	0	6.9×10^4	4.7×10^3	6.27	-2.33
ACE	0	3.0×10^4	4.5×10^3	6.31	-2.12
FLN	863.9	5.0×10^4	4.8×10^3	6.79	-2.41
ANT	0	8.4×10^4	2.4×10^3	7.55	-2.64
PHE	2.2×10^3	2.5×10^4	5.4×10^3	7.57	-2.76
RET	0	3.2×10^4	6.2×10^3	8.70	-2.35
PYR	0	1.1×10^5	1.2×10^4	8.80	-3.31
FLT	3.7×10^3	1.3×10^6	1.4×10^4	8.88	-3.44
BAA	0	1.5×10^5	4.7×10^3	9.07	-3.31
CHR	0	1.6×10^5	1.3×10^4	9.48	-3.67
BFN	0	1.6×10^5	8.6×10^3	9.57	-3.80
BGF	0	1.4×10^5	1.4×10^4	9.78	-4.26
PER	0	1.5×10^5	1.8×10^3	10.08	-3.83
CPP	0	1.8×10^5	563	10.15	-4.45
BBF	0	1.4×10^5	7.5×10^3	10.35	-4.57
BJF	0	1.4×10^5	9.2×10^3	10.59	-4.48
TPH	0	2.0×10^5	1.9×10^4	10.69	-5.20
BKF	0	1.4×10^5	5.6×10^3	10.73	-4.62
BAP	0	9.1×10^4	1.6×10^3	10.86	-4.73
DCA	0	1.4×10^5	0	11.11	-4.70

K_{oa} : analyte octanol-air partitioning coefficient; K_{aw} : analyte air-water partitioning coefficient. For ease of interpretation, analytes are sorted based on their $\log K_{oa}$ values. $\log K_{oa}$ and $\log K_{aw}$ values were obtained from Estimation Programs Interface Suite 4.11 (USEPA, 2012); zero value indicates an event in which analyte concentration in rainwater was < LOQ.

PAHs, PCBs, and CPs
in background air in
central Europe

P. Shahpoury et al.

Table 7. Continued.

	^a Upper-bound ε			^b Lower-bound ε			^c Median
	Min.	Max.	Median	Min.	Max.	Median	
BEP	0	2.9	4.0×10^{-2}	0	7.5×10^{-2}	6.5×10^{-3}	NA
IPY	0	3.5	1.7×10^{-2}	0	0.3	1.8×10^{-3}	NA
DHA	0	3.1	0	0	3.8×10^{-2}	0	NA
BPE	0	3.6	2.3×10^{-2}	0	0.2	2.5×10^{-3}	NA
COR	0	4.8	2.2×10^{-3}	0	0.2	7.2×10^{-4}	NA
BNT	0	2.8	0	0	0.2	0	NA
ATT	0	3.8	0	0	4.7×10^{-2}	0	NA
Σ PAHs	1.2×10^{-3}	0.8	5.3×10^{-2}	5.2×10^{-4}	0.1	9.7×10^{-3}	NA

^a Upper- and ^b lower-bound ε correspond to cloud base and top height, respectively; ^c median ε values reported by Skrdlíková et al. (2011) for cloud top heights ($n = 10$); zero value indicates an event in which analyte concentration in rainwater was < LOQ; NA: not available. Analytes are sorted based on their log K_{oa} values.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



