

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

New emission factors for Australian vegetation fires measured using open-path Fourier transform infrared spectroscopy – Part 2: Australian tropical savanna fires

T. E. L. Smith¹, C. Paton-Walsh², C. P. Meyer³, G. D. Cook⁴, S. W. Maier⁵, J. Russell-Smith⁵, M. J. Wooster¹, and C. P. Yates⁵

¹King's College London, Earth and Environmental Dynamics Research Group, Department of Geography, London, UK

²University of Wollongong, School of Chemistry, Wollongong, New South Wales, Australia

³CSIRO Marine and Atmospheric Sciences, PMB 1, Aspendale, Victoria, Australia

⁴CSIRO Ecosystem Sciences Sustainable Agriculture Flagship, PMB 44, Winnellie, Northern Territory, Australia

⁵Charles Darwin University, Research Institute for the Environment and Livelihoods, Darwin, Northern Territory, Australia

ACPD
14, 6311–6360, 2014

New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.

Title Page
Abstract
Conclusions
Tables
Introduction
References
Figures

◀ ▶
◀ ▶
Back Close
Full Screen / Esc

Printer-friendly Version
Interactive Discussion



Received: 30 December 2013 – Accepted: 13 February 2014 – Published: 11 March 2014

Correspondence to: T. E. L. Smith (thomas.smith@kcl.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

14, 6311–6360, 2014

New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Abstract

Savanna fires contribute approximately 40–50 % of total global annual biomass burning carbon emissions. Recent comparisons of emission factors from different savanna regions have highlighted the need for a regional approach to emission factor development, and better assessment of the drivers of the temporal and spatial variation in emission factors. This paper describes the results of open-path Fourier Transform Infrared (OP-FTIR) spectroscopic field measurements at twenty-one fires occurring in the tropical savannas of the Northern Territory, Australia, within different vegetation assemblages and at different stages of the dry season. Spectra of infrared light passing through a long (22–70 m) open-path through ground-level smoke released from these fires were collected using an infrared lamp and a field-portable FTIR system. The IR spectra were used to retrieve the mole fractions of fourteen different gases present within the smoke, and these measurements used to calculate the emission ratios and emission factors of the various gases emitted by the burning. Only a handful of previous emission factor measures are available specifically for the tropical savannas of Australia and here we present the first reported emission factors for methanol, acetic acid, and formic acid for this biome. Given the relatively large sample size, it was possible to study the potential causes of the within-biome variation of the derived emission factors. We find that the emission factors vary substantially between different savanna vegetation assemblages; with a majority of this variation being mirrored by variations in the modified combustion efficiency (MCE) of different vegetation classes. We conclude that a significant majority of the variation in the emission factor for trace gases can be explained by MCE, irrespective of vegetation class, as illustrated by variations in the calculated methane emission factor for different vegetation classes using data subsetted by different combustion efficiencies. Therefore, the selection of emission factors for emissions modelling purposes need not necessarily require detailed fuel type information, if data on MCE (e.g. from future spaceborne total column measurements) or a correlated variable were available.

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



From measurements at twenty-one fires, we recommend the following emission factors for Australian tropical savanna fires (in grams of gas emitted per kilogram of dry fuel burned) which are our mean measured values: 1674 g kg⁻¹ of carbon dioxide; 87 g kg⁻¹ of carbon monoxide; 2.1 g kg⁻¹ of methane; 0.11 g kg⁻¹ of acetylene; 0.49 g kg⁻¹ of ethylene; 0.08 g kg⁻¹ of ethane; 1.57 g kg⁻¹ of formaldehyde; 1.06 g kg⁻¹ of methanol; 1.54 g kg⁻¹ of acetic acid; 0.16 g kg⁻¹ of formic acid; 0.53 g kg⁻¹ of hydrogen cyanide; and 0.70 g kg⁻¹ of ammonia.

1 Introduction

Estimates of mean annual carbon emissions from worldwide savanna and grassland fires amount to ~1 Pg C year⁻¹ (2000–2009), or approximately a mean of 40–50 % of total global annual biomass burning carbon emissions (van der Werf et al., 2010). In terms of the Australian savanna, the direct gross carbon emissions from fires have a similar, if not greater, magnitude than those from Australian fossil fuel combustion (Haverd et al., 2013). However, in tropical Australia, net carbon emissions from fires are considered negligible as the carbon lost to burning is approximately balanced by the carbon sequestered during regrowth (Haverd et al., 2013). Akagi et al. (2011) present summaries of emission factors (EF) that can be used to convert estimates of total biomass (or carbon) consumed in fires into estimates of individual trace gas species release. Emissions of non-CO₂ gases from savanna fires, such as nitrous oxide and methane, whose emissions are not sequestered by regrowth, comprise 2–4 % of total Australian national greenhouse gas emissions (DCCEE, 2010), and are included in the accountable greenhouse gas inventory. At state or territory scales, pyrogenic emissions of CO₂, CH₄, and N₂O can be far more significant, with emissions from fires comprising nearly 50 % of the Northern Territory's accountable carbon emissions (DCCEE, 2010).

A clear understanding of the emissions characteristics of Australia's tropical savannas is important for regional emissions modelling and for associated greenhouse gas accounting schemes. As with the fire regimes of other savanna regions, that in Aus-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

tralia's tropical savanna is heavily influenced by human fire use and fire management (Garde, 2009; Cook et al., 2010). Currently, policy in the Northern Territory of Australia aims to reduce overall savanna fire emissions by promoting early dry season (pre mid-July) fires which consume less fuel and create a network of fire breaks that reduce the extent of late dry season (post mid-July) fires, reducing overall fire frequency (Cook and Meyer, 2009; Russell-Smith et al., 2009). Given that the seasonality of the emission factor for methane in Australian tropical savannas has recently been shown to be different to that observed in African savannahs (Meyer et al., 2012), it is evidently important to make regionally-specific measurements of emission factors so that the most accurate assessments of the effectiveness of such policies, and the impact of fire management on greenhouse gas and reactive gas emissions, can be provided.

Despite the prevalence of savanna fires on the Australian continent, with approximately 400 000 km² of tropical savannas (~ 15 % of global tropical savannas) and 150 000 km² of arid savannas burning annually (Russell-Smith et al., 2007; Meyer et al., 2012), the focus of most savanna emissions research has been on Africa (e.g. Bertschi et al., 2003a; Yokelson et al., 2003; Sinha et al., 2003; Keane et al., 2006). Only Hurst et al. (1994a), Hurst et al. (1994b), and Shirai et al. (2003) have published emission factors from Australian savannas, with most measurements made using airborne campaigns. Most recently, Meyer et al. (2012) have published emission factors for methane (CH₄) and nitrous oxide (N₂O) from Australian savannas using ground-based sampling methods.

Careful interpretation of data collected from these various methods is required to ensure the accurate characterisation of "fire-averaged" emissions (Akagi et al., 2011). Beyond these methodological complications are the factors that lead to the natural variability of emissions (e.g. interactions between meterology, seasonality and combustion efficiency). As the recent study by Van Leeuwan and van der Werf (2011) shows, within-biome temporal and spatial variations in emission factors are substantial, and despite their attempts to explain these using relationships between emission factors and a number of environmental variables (tree cover density, vegetation greenness, temperature,



precipitation, length of dry season), correlations between CO, CH₄ and CO₂ EFs, and MCE, were rather low, with environmental variables explaining 33 %, 38 %, 19 %, and 34 % of the variation in the reported emissions metrics. Van Leeuwen and van der Werf (2011) posit some potential reasons for the poor explanation of within-biome variability, including different measurement techniques for EFs, assumptions used to weight EFs by the amount of flaming and smouldering combustion and uncertainties in the environmental variables themselves. Explaining the large natural variability of EFs that is so pronounced in the findings of Akagi et al. (2011) remains an important challenge for biomass burning emissions science.

Through measurements of a large number of fires during multiple campaigns at different stages of tropical Australia's dry season, this work aims to expand the current emission factors inventory for Australian savannas, whilst also exploring the nature and causes of the natural variability of fire emissions within this ecosystem. Long Open-Path Fourier Transform InfraRed (OP-FTIR) spectroscopy was used for field measurements of trace gases emitted from 21 different experimental fires or wildfires in the tropical savanna of the Northern Territory, Australia.

2 Methods

2.1 The open-path FTIR system and deployment

FTIR systems can simultaneously record infrared spectra at a high enough spectral resolution and across a wide wavenumber range, such that estimates of the path-averaged mole fraction of many of the trace gases present in the path can be retrieved via spectral analysis. The OP-FTIR system operated by King's College London consists of a MIDAC Corporation M2000 Series FTIR spectrometer (0.5 cm⁻¹ resolution) equipped with a Stirling-cycle cooled mercury-cadmium-telluride detector (MCT), and fitted with a MIDAC custom-built 76 mm Newtonian telescope. Unlike the mono-static pre-modulated source system used by Paton-Walsh et al. (2014), the MIDAC system

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

employed here operates using a bi-static configuration, where the FTIR spectrometer is operated in “active” mode, viewing a remotely located infrared source. Here, the infrared source consists of a 12 V silicon carbide glowbar operating at 1500 K fitted in front of a 20 cm diameter gold-plated aluminium collimator. With a maximum optical path distance of 2 cm, each scan of the spectrometer mirror takes ~1.2 s. To improve the signal-to-noise ratio, it is advantageous to average (co-add) a number of spectral measurements prior to analysis. The MIDAC system and its use to characterise biomass burning emissions via long OP-FTIR spectroscopy is detailed in Wooster et al. (2011), whilst an accuracy assessment with respect to the retrieval of greenhouse gas mole fractions from the collected spectra was undertaken by Smith et al. (2011).

A mixture of purposely lit “experimental” plots and wildfires were studied. At most plots, the OP-FTIR path was positioned on a road, track, or fire-break downwind of the fire plot with pathlengths ranging from 22 to 70 m (Table 1). There were, however, some exceptions. For some plots, where there was no safe location downwind of the fire plot, the OP-FTIR path was positioned over the plot, with smoke passing between the spectrometer and IR source before or after the fire front passed through this path. Spectra collected during the time in which the fire was burning within the open path were affected by infrared emission from the flames; this effect precludes the retrieval of gas mole fractions from spectra due to the superposition of emission and absorption lines in the spectral signatures (one disadvantage of the bi-static OP-FTIR configuration). Consequently, all spectra affected by thermal emission from hot flames were removed prior to the analysis stage. For other plots, and especially for the wildfire measurements, it was not possible to position a path beside or across the fire plot. In these situations, either the IR source, or both the spectrometer and the IR source, were positioned within the fire plot. To protect the equipment from the fire, the equipment was positioned within natural firebreaks (rocky outcrops, areas of low biomass) or artificial firebreaks (raked vegetation) with the OP-FTIR path penetrating into the fire plot, or being entirely enclosed by the fire plot (Fig. 1).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figure

1



Back

Close

Full Screen / Esc

Printer friendly Version

Interactive Discussion



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

For the experimental plots, it was possible to assemble the OP-FTIR equipment prior to each fire. Burn-offs and wildfires, by their nature, are unpredictable; identification of a position for the OP-FTIR path and assembly of the equipment took place whilst the fire was burning upwind. In one case (at the Adelaide River Station), it was possible to follow the fire along a track, assembling and reassembling paths as the fire front passed each path location. Given that most plots were small (< 1 ha) and in some cases burned quickly, spectra were collected using 4 co-added scans to increase signal-to-noise yet still allow for a high measurement frequency (generating approximately one measured spectrum every 5 s). Observations of atmospheric temperature and pressure (parameters required for the trace gas retrieval analysis) were made using a co-located WXT510 weather station (Vaisala Oyj, Helsinki, Finland) at the Arnhem Land campaigns. For the Darwin region campaign, temperature and pressure were taken from a Bureau of Meteorology (Australian Government) weather station in Batchelor (located within 60 km of all of the burn sites).

2.2 Quantitative analysis of infrared spectra

For measurements that use a bi-static approach, where the FTIR spectrometer is used to measure a remote collimated non-modulated IR source, as described here, the measured spectrum includes radiation that has originated from the IR source (I_{SOURCE}), but also from instrument self-emission and from other ambient temperature sources (I_{AMB}) within the field of view of the receiving telescope. At the higher wavenumber (shorter wavelength) regions used to retrieve gases such as CO₂, CO, CH₄, H₂CO (2000–3100 cm⁻¹, see Fig. 2), the ambient signal contribution is almost non-existent. However, at wavenumber regions below ~ 1500 cm⁻¹, where gases such as C₂H₄, HCOOH, CH₃OH and NH₃ are retrieved, the contribution from ambient sources becomes more important to account for. This ambient signal component introduces a zero-baseline offset to the measured spectra which needs to be removed prior to any trace gas retrieval as this component contains no information related to trace gas absorption (Müller et al., 1999). When I_{AMB} is small (at wavenumbers > 1500 cm⁻¹) the effect

on measured transmittance is relatively minor. However, higher levels of I_{AMB} at lower wavenumbers can significantly alter the apparent transmittance (T_{app}):

$$T_{\text{app}} = \frac{I}{I_{\text{SOURCE}} + I_{\text{AMB}}} \quad (1)$$

To correct for this zero-baseline offset, it is necessary to make a measurement of the ambient spectrum (e.g. a spectral measurement with the active IR source turned off) and subtract this spectrum from each measurement of the active IR source, making slight adjustments for any MCT detector non-linearities in accordance with Müller et al. (1999) to ensure that the resultant “difference” spectra exhibit zero signal in saturated spectral regions, such as the 668–670 cm⁻¹ CO₂ absorption band (see Fig. 4 in Wooster et al., 2011). Wooster et al. (2011), who follow this methodology, confirm that subtracting the ambient signal has a large effect on retrieved mole fractions of NH₃ (retrieved using a spectral window around 940 cm⁻¹); NH₃ mole fractions increased by a factor of 2 due to the reduced calculated transmittance (through Eq. 1). Wooster et al. (2011) also stress that the impact of ambient signal subtraction is much smaller for retrievals of CO₂ and CO at ~2000 cm⁻¹ (< 10 %). During measurements of a particular fire, some changes in ambient background emission might be expected (e.g. due to changes in air temperature). The majority of measurements presented in this work took place over relatively short-lived fires (< 1 h) and should therefore not be substantially influenced by changes in the ambient background spectrum. Nevertheless, measurements of ambient spectra were made both before and after a fire where possible, to check for any large changes in the ambient spectrum during the course of a fire.

Trace gas mole fractions were retrieved from all open-path FTIR spectra (after subtracting the background spectrum) using the Multiple Atmospheric Layer Transmission (MALT) model, as described in Part 1 of this paper (Paton-Walsh et al., 2014). MALT includes a forward model and a parameter optimisation routine, and has been used previously for retrieving the mole fraction of biomass burning compounds from open-path

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

measurements of smoke in Australia (Griffith et al., 1991; Meyer et al., 2012), and in the United States (Akagi et al., 2013). Working in African savannah, Wooster et al. (2011) used an alternative retrieval code that again used similar physical principles to MALT. A full description of MALT can be found in Griffith et al. (1996).

There are a few differences between the retrieval parameters used here and those used in Part 1 of this paper. Due to the higher spectral resolution of the MIDAC spectrometer, a resolution of 0.5 cm^{-1} (fixed) was used here with a field of view of 20 milli-radians (fixed); here we apply a triangle apodisation function. Phase error and wavenumber shifts were fitted in accordance with Paton-Walsh et al. (2014).

2.3 Spectral regions and parameters for quantitative spectral analysis

A set of standardised spectral windows and retrieval parameters are outlined in Part 1 of this paper (Table 1 in Paton-Walsh et al., 2014). The higher spectral resolution and greater wavenumber range ($700\text{--}4000\text{ cm}^{-1}$) of the MIDAC spectrometer and detector deployed for the measurements presented here allowed for the additional detection of HCN and C_2H_2 in the $710\text{--}760\text{ cm}^{-1}$ region, beyond what was detectable using the spectrometer deployed by Paton-Walsh et al. (2014). A full set of example spectral fits to our 0.5 cm^{-1} spectra is presented in Fig. 2. Due to the underlying spectral response of the MIDAC spectrometer, a higher-order background polynomial was required for the HCOOH retrieval window (Fig. 2f); here we use a 3rd-order polynomial, rather than a 2nd-order polynomial as indicated in Table 1 in Paton-Walsh et al. (2014).

2.4 Fire-averaged emission factors and probing natural variability

Individual fire emission factors were calculated using the method outlined in Part 1 (Paton-Walsh et al., 2014; Sect. 1.5). Emission factors for CO_2 and CO are calculated via the summation method, using background mole fractions for these gases measured before ignition, or upwind of the fires. Emission factors for the other gases are calculated using their respective emission ratios to either CO_2 or CO as also described in



Sect. 1.5 of Paton-Walsh et al. (2014). Given that the results presented here cover measurements at 21 different fires from distinct tropical savanna ecosystem vegetation assemblages, and measured at different stages of the dry season (Table 1), further analysis will include an assessment of the *within-ecosystem* variability in emission factors.

5

3 Site and fire descriptions

Savanna fire emissions were measured at experimental fires in July and September 2009 near Kulnguki on the Arnhem Land Plateau, Northern Territory. Further measurements of experimental fires, land management fires and wildfires were carried out
10 during August to October 2010 in the cities of Darwin and Palmerston, Litchfield Municipality, and Coomalie Shire of the Northern Territory. The flora of these areas is typical of eucalyptus-dominated savannas that comprise much of northern Australia, from the Kimberley region of Western Australia, through the Top End of the Northern Territory to Cape York in Queensland (Fig. 3).

15

3.1 Arnhem Land

Measurements in Arnhem Land were conducted during two field campaigns described at length by Meyer et al. (2012). Six experimental fires were measured during the early-dry season (EDS) between 2–7 July 2009, and a further five experimental fires were measured during the late-dry season (LDS) between 30 September and 4 October 2009.

20

The climate of West Arnhem Land is tropical monsoonal, and 97 % of the mean annual rainfall falls between November and April. The closest weather station, at Oenpelli ($12^{\circ}20' S$, $133^{\circ}03' E$), 100 km west-northwest of the experimental fire sites records a mean annual rainfall of 1417 mm, with a mean maximum temperature of $30^{\circ}C$ in July and $38^{\circ}C$ in October.

25

ACPD

14, 6311–6360, 2014

New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



New emission factors for Australian vegetation fires

T. E. L. Smith et al.

The experimental fire sites were located near Kulnguki, in West Arnhem Shire, Northern Territory (12°38' S, 133°55' E, Fig. 4d), 330 km east of Darwin. One set of fires were situating on the Queue landsystem (Lynch and Wilson, 1998; Meyer et al., 2012), a system of sandstone plains dominated by a semi-continuous canopy of *Eucalyptus* and *Corymbia* trees, with an *Acacia* mid-storey (Fig. 5a). The understorey consists of sorghum grasses with a number of shrub species to about one metre high. For consistency with Meyer et al. (2012), the vegetation of this system will be referred to as "tussock grass open woodland" (TGOW). Another set of fires were situated on the sandstone plateau of the Buldiva landsystem (Lynch and Wilson, 1998; Meyer et al., 2012). Here, the dominant trees are similar to that of the TGOW, whereas the understorey is mainly dominated by hummock grass (*Triodia*) with a variety of shrubs (Fig. 5c). The vegetation of this system will be referred to as "hummock grass open woodland" (HGOW). While tussock grasses can be found in savanna systems worldwide, the hummocky spinifex grasses are endemic to Australia (Meyer et al., 2012), adapted to the arid climate with a shrub-like growth form. In addition to the TGOW and HGOW vegetation classes, fires were measured in open woodland areas with very few trees, dominated by sorghum grasses (Fig. 5b); this vegetation class will be referred to as "sorghum open woodland" (SOW). Individual hummocks of spinifex grasses were also burned and measured for their emissions (Fig. 5d); this vegetation class will be referred to as "pure spinifex" (SPIN). An extended description of the particular species at these fire sites can be found in Meyer et al. (2012).

The tussock-grass open woodland and sorghum open woodland fires were conducted in a series of adjacent plots established prior to the first campaign along an access road. Each tussock-grass open woodland plot was approximately 500 m × 200 m in size, whilst each sorghum open woodland plot was approximately 100 m × 100 m in size. The dissected quartz sandstone landscape of the hummock-grass open woodland allowed for the identification of two isolated plots prior to the campaign, each with areas of 50–70 ha, surrounded by natural fire barriers. The plots were situated in isolated country accessible only by helicopter, approximately 6–10 km from the tussock-grass

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

Back

Close

Full Screen / Esc

Page 1 of 1



plots. These experimental plots were ignited by a line ignition pattern on the upwind side of the plot, with some plots having a back fire also ignited on the downwind side of the plot. The isolated hummocks of spinifex could be ignited individually without risk of establishing a fire front due to the surrounding rocky outcrop. For further descriptions of the Arnhem Land experimental fires, including fuel loads and fuel moisture content, refer to Meyer et al. (2012).

3.2 Darwin Region

A third measurement campaign to the Northern Territory was conducted during August to October in the dry season of 2010. Measurements were made at a number of wild-fires, as well as experimental fires, at various locations within a region 120 km to the south/southeast of Darwin (Fig. 4c). The climate of this areas is tropical monsoonal, with 95 % of mean annual rainfall falling between April and November. According to the weather station at Batchelor (within 60 km of all fire measurement sites), mean annual rainfall is 1528 mm, with a mean maximum temperature of 32 °C in July and 37 °C in October.

The opportunistic nature of this campaign allowed for emissions to be measured from fires in a number of different savanna land systems within the vicinity of Darwin. These included the gently undulating plains of the Koolpinyah land system (Christian et al., 1953), dominated by open eucalypt woodland with tussock grasses (Fig. 5e and f). This vegetation class is similar to the tussock grass open woodland of the Arnhem Land campaigns and will also be indicated using TGOW. Measurements were also made at fires on the flat perennially flooded plains of the Marakai land system (Christian et al., 1953), mainly consisting of tussock grasslands (Fig. 5 g). This vegetation class will be referred to as “tussock grass floodplain” (TGFP). Fires were also measured on the flat country of the Batchelor land system (Christian et al., 1953), dominated by tall closed woodland with tussock and gamba (*Andropogon gayanus*) grasses (Fig. 5h). This vegetation class will be referred to as “tussock and gamba grass woodland” (TGGW).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unlike the purely experimental fires of the Arnhem Land campaign, most of the fires measured for the Darwin region campaign were either wildfires or “burn-off”, where there was no control over the timing and pattern of ignition, and with many fires burning across many hectares. Locations of burn-off and wildfires were identified by regular monitoring of the NAFI website (www.firenorth.org.au), which shows the locations of satellite-detected fires; and by regular phone calls to Bushfires NT (Northern Territory Government), who are responsible for supporting landholders with fire mitigation.

Wildfires were measured at the Adelaide River Station on the Marrakai Road (TGFP) on 1 October 2010; near Lambells Lagoon off the Arnhem Highway (TGOW) on 1 October 2010; and at Berry Springs (TGOW) on 4 October 2010. Burn-offs were measured at the Gould airstrip near Batchelor (TGGW) on 11 September 2010 and at Southport (TGOW) on 6 October 2010. In addition to the wildfires and burn-offs, measurements were made at a set of three experimental fires in the Territory Wildlife Park (TGOW) near Berry Springs on 5 October 2010. Each of the experimental plots was approximately 130 m × 80 m in size, and the fires and fire conditions from this campaign are presented in Table 1.

4 Results

A time series of smoke tracer mole fractions retrieved from FTIR spectra collected near to the downwind boundary of one of the experimental fires is shown in Fig. 6, with CO₂, CO and MCE shown in detail in Fig. 7. In most of the experimental burns (with the exception of the hummock grass plots), the fires were ignited with a back-burning fire line on the down-wind boundary of the fire plot; this moves slowly against the wind, forming a fire break. The up-wind boundary is then ignited, with the majority of the plot burning with a head-fire driven by the prevailing wind. Generally, the smoke passing through the infrared beam will have first originated from the flaming combustion in the back-burn fire line (notice the high MCE in the first minute of the time series in Fig. 7). Immediately following the initial burst of smoke from the flaming combustion,

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the smoke from the flaming back-burn will be mixing with smouldering smoke, also originating from the back-burn, with increasing CO mole fraction and a corresponding decrease in MCE (until \sim 7 min in Fig. 7). As the back-burn fire line burns away from the OP-FTIR path, most of the smoke crossing the infrared beam will have been emitted

5 from the steadily increasing area of smouldering-phase combustion, with contributions from both the flaming combustion of the head-fire and the smouldering area behind the head-fire (towards the end of the time series in Fig. 7, as MCE decreases to \sim 90 % and below). This equates to the area of “flaming combustion” (i.e. the fireline) and the associated “smoking zone” of smouldering combustion located close behind the fire front,
10 as detailed in the OP-FTIR African savannah measurements of Wooster et al. (2011).

4.1 Emission ratios

The emission ratios of each trace gas relative to CO_2 and CO are calculated using the generalised least squares linear regression slope (this is the best fit to the points that minimises deviations from the line of fit in both the x- and y-axes, weighted by
15 uncertainties in both x and y) of the trace gas species mole fraction vs. the CO_2 or CO mole fraction. As an example, Figs. 8 and 9 show a series of trace gas mole fractions plotted against CO_2 and CO, respectively. The generalised least squares linear best fits are shown in each case, with the gradient taken as the emission ratio ($\text{ER}_{\text{x}/\text{CO}_2}$ or $\text{ER}_{\text{x}/\text{CO}}$). As is evident in both Figs. 8 and 9, it is clear that for all gases there is
20 a stronger correlation to CO than to CO_2 (due to their preferential production during smouldering-phase combustion, as is also the case for CO). For most gases (CH_4 , C_2H_2 , C_2H_4 , C_2H_6 , CH_3OH , CH_3COOH , HCOOH , NH_3 , HCN), their emission ratio to CO is used for the calculation of their respective emission factors. For H_2CO , the only
25 gas to show a better mean correlation with CO_2 (across all 21 fires), the emission ratio to CO_2 is used for the calculation of its emission factor (Paton-Walsh et al., 2014, also found C_2H_4 to be better correlated to CO_2). Following Sinha et al. (2003) and Wooster et al. (2011), the strength of the coefficient of determination (R^2) between the mole fractions of each target gas and that of CO or CO_2 is used to confirm whether

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

the emission ratio for the target gas is well determined. For gases with stronger IR absorption features (e.g. CO, CH₄, C₂H₄, NH₃), R^2 in the figures was generally higher than 0.8, whilst gases with weaker IR absorption (e.g. CH₃OH, HCN, HCOOH) showed poorer correlations to CO₂ and CO, due to greater uncertainties in the retrieval process.

5 Those species for which the coefficient of determination fell below 0.4 are assumed to have poorly determined emission ratios (Sinha et al., 2003; Wooster et al., 2011). Consequently, emission ratios where R^2 fell below 0.4 were not used in the calculation of emission factors.

Given that MCE is known to decrease as the fire progresses through the plot (Fig. 7),
10 with a greater contribution from the increasing area of smouldering combustion behind the fire front, it is assumed that measurements with the highest MCE are of smoke that has predominantly originated from flaming combustion, whilst those measurements with lower MCE are of smoke that has a higher proportion of smouldering combustion (Yokelson et al., 1996). Whilst the *within-fire* variation in emission ratios is quite limited,
15 as shown by the high value of R^2 on the linear least squares regression fits in the bivariate plots (Figs. 8 and 9), this ever-changing ratio of flaming-to-smouldering smoke does lead to some variation in the emission ratio of most of the gases to CO₂ (as evident in the deviation from the linear best-fit in most of the scatter plots shown in Fig. 8). To account for this, it is possible to calculate different emission ratios to CO₂ for the different
20 combustion stages of each fire. Conventionally, laboratory studies use observations of fire activity to determine the combustion stage of emissions (e.g. Yokelson et al., 1996) or an approach where the maximum rate of increase of CO emission is taken as the beginning of the smouldering stage (e.g. Lobert et al., 1991). The OP-FTIR study of Wooster et al. (2011) had access to simultaneous airborne imaging, so was able
25 to separate out times of different fire activity based on those thermal/optical airborne observations. Here, measured emissions cannot be ascribed to particular combustion stages using observations, and furthermore the relative contribution to measured emissions from flaming and smouldering combustion may vary depending on changes in the wind direction, ignition pattern of the fire, and heterogeneous vegetation cover

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and fuel moisture within the fire plot. Instead, the emissions measurements are split by MCE, so that measurements with high MCE (> 90 %) and those with low MCE (< 90 %) may be used separately to calculate emission ratios (Fig. 10). This approach has been used in the past for laboratory analysis, and the 90 % threshold was taken from studies

5 that have published MCE for flaming and smouldering combustion stages (e.g. Lobert et al., 1991; Yokelson et al., 1996; Bertschi et al., 2003b). Emission ratios of each target species to both CO_2 and CO were assessed in this way and are presented in Table 2 and Table 3, respectively, in addition to emission ratios derived from all measurements.

As discussed above, the *within-fire* variation in emission ratios is fairly limited, and
10 can be understood by splitting measurements by MCE as already described. The
between-fire variation, however, is larger, with the flaming emission ratio of CH_4 , for
example, varying from 0.0010 (for spinifex) to 0.0045 for one of the tussock grass open
woodland plots. Most of this variation is caused by comparison between the “land-
scape” experimental fires (those that incorporate a number of fuel species; e.g. TGOW,
15 HGOW, SOW) and the experimental fires that were conducted on the individual clumps
of spinifex. Given that the spinifex measured here appeared to burn with high combus-
tion efficiency (possibly due to measured high burn temperature, aerated structure, and
leaf resin content), producing little observable “smoke” (see Fig. 5d), it is perhaps no
surprise that its emissions characteristics are different from those from experimental
20 fires containing multiple species, where vegetation that burn at different combustion
efficiencies contribute to a plume-averaged efficiency (e.g. the tussock grass and hum-
mock grass open woodland landscapes consist of both fine and coarse fuels, each with
different structure and degrees of compaction). As the purpose of this work is to char-
acterise landscape-averaged emissions, the emission ratios and emission factors from
25 the spinifex fires have been excluded from calculating average emissions parameters.

The emission ratios presented in Table 2 ($\text{ER}_{\text{X}/\text{CO}_2}$) and Table 3 ($\text{ER}_{\text{X}/\text{CO}}$) show the mean emission ratios for the landscape fire measurements (including HGOW, SOW,
TGFP, TGGW, TGOW vegetation classes), excluding those derived from individual
species fires (SPIN). Table 2 confirms that the emission ratios to CO_2 (i.e. $\text{ER}_{\text{X}/\text{CO}_2}$),

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



for all of the gases investigated here, are higher in the measurements we class as “smouldering” (based on MCE < 90 % than in those measurements classed as “flaming” (MCE > 90 %). The close correlation between the emissions of gases and those of CO (Fig. 9) also explains why there is no significant difference in the emission ratios to CO ($ER_{X/CO}$) recorded during these “flaming” and “smouldering” classes (the difference between “flaming” and “smouldering” $ER_{X/CO}$ is smaller than the standard deviation of each ER). For formaldehyde (H_2CO), the higher emission ratio to CO during the “flaming” class might be due to the production of formaldehyde during the closely-related (in space and time) pyrolysis combustion phase (Lobert and Warnatz, 1993); with some of the pyrolysed gas escaping combustion due to advective or convective winds. Wooster et al. (2011) suggested pyrolysis was the cause of the similarly higher OP-FTIR derived emission ratios to CO for formaldehyde in the flaming rather than the residual smoldering combustion stage of southern African savannah fires.

4.2 Drivers of variations in emission ratios & emission factors

Figures 8 and 9 depict trace gas retrievals and emission ratios derived from data of a typical experimental fire. Given that a number of different vegetation types were sampled during the three Australian campaigns, with measurements made throughout the early- and late-dry season, and the likelihood that each experimental fire would have burned at different efficiencies depending on the fuel and meteorological conditions, it is important to assess the relative importance of each of these variables. Figure 11 includes all data from the 2009 campaigns, showing the scatter plot for CH_4 vs. CO_2 only, with each subplot (a–c) depicting the same data but with each data point (i.e. each individual retrieval from 4 co-added spectra collected over ~ 5 s) colour-coded depending on its MCE (Fig. 11a); the season of the experimental fire (Fig. 11b); and the predominant vegetation type for the burn (Fig. 11c). As discussed above, the between-fire variation in emission ratios can be primarily explained by a dependence on vegetation class (as is evident in Fig. 11c), with the spinifex (individual species) fires grouping towards the lower emission ratios and the sorghum dominated fires grouping towards the higher

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

emission ratios. Figure 11a clearly demonstrates that this grouping is mainly a result of differences in the combustion efficiencies of these different vegetation classes, with MCE appearing to be the major control on the emission ratio slope, despite differences in vegetation classes. There is no clear pattern to discern when the data are plotted by season (Fig. 11b).

By sub-sampling the data in Fig. 11a into 2 % bins of MCE (e.g. only measurements where $80\% < \text{MCE} \leq 82\%$), it is possible to calculate a range of emission ratios to CO_2 depending on MCE. Figure 12 depicts eleven different CH_4 emission ratios to CO_2 ($\text{ER}_{\text{CH}_4/\text{CO}_2}$) calculated in this way. This figure demonstrates the strong relationship between the emission ratio of methane and the modified combustion efficiency, which is unsurprising due to the well defined CO to CO_2 and CH_4 to CO emission ratios already shown in Fig. 9. A clear relationship between the slope of the linear least squares regression lines and MCE is evident as the slope increases systematically with decreasing MCE. This relationship applies for the other trace gases that also correlate well with carbon monoxide. These MCE-dependent emission ratios can then be used to calculate MCE-dependent emission factors, discussed further below.

4.3 Final emission factors

Emission factors for CO_2 and CO for each fire were calculated by the summation method as described in Paton-Walsh et al. (2014). Emission factors for all other gases were calculated from the CO_2 and CO emission factors for the relevant fires and the emission ratio to CO_2 or CO (given in Table 2 and Table 3). These emission factors are presented in Table 4. Mean emission factors calculated from the individual fire emission factors in Table 4 are listed in Table 5.

By combining measurements from *all* fires, it is possible to build a detailed understanding of the relationship between MCE and emission ratios (as depicted earlier in Fig. 11a and Fig. 12). MCE-dependent emission ratios can be used to calculate MCE-dependent emission factors. Using the same methodology applied to calculate emission factors for individual fires, it is possible to calculate emission factors for each

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

2 %‐MCE bin. Figure 13 shows the relationship between MCE and the emission factor of CH₄ (EF CH₄) calculated using the measurements from different vegetation classes and when using all measurements combined. This plot demonstrates that the majority of variation in the emission factor for CH₄ is determined by MCE irrespective of vegetation type. The uncertainty bounds of the CH₄ emission factor for any particular vegetation class lie within the bounds of the emission factor determined from all measurements for each MCE bin (except TGFP in the 86–88 % MCE bin).

5 Comparison to past ER and EF measures

The emission factors presented here are the first to be reported from OP‐FTIR measurements of tropical savannas in Australia and follow only a few ground-based (Hurst et al., 1994a) and airborne studies (Hurst et al., 1994b; Shirai et al., 2003) in this ecosystem. The emission factors for CH₃OH, CH₃COOH, and HCOOH are the first to be reported by any methodology for Australian savanna fires (Table 5). The emission factors presented here have been estimated from measurements at up to 21 separate fires from a variety of vegetation types and from both controlled experimental burns and wildfires (Table 4). Table 5 compares the emission factors reported here with those reported by previous studies of tropical savannas in Australia (listed above) and with those reported by the emission factor databases for global savanna and grasslands in Andreae and Merlet (2001) and global savannas in Akagi et al. (2011).

20 5.1 Carbonaceous species

A broad range of emission factors for carbon dioxide (EF CO₂) are reported in the literature from savanna ecosystems (Table 5). The emission factors reported here are higher than those reported by previous studies of northern Australian savanna fires, but fall within the standard deviation of measurements in northern Australia made by Hurst et al. (1994b) and Shirai et al. (2003). The EF CO₂ reported here agrees well with



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

those reported in Andreae and Merlet (2001 and subsequent updates in 2009), and Akagi et al. (2011), which draw on numerous studies in savanna ecosystems worldwide (the mean EF CO₂ reported here is only 0.7 % smaller than that reported by Akagi et al. (2011) for savanna ecosystems).

The carbon monoxide emission factor (EF CO) reported here agrees well with previous studies of biomass burning in Australian tropical savannas, which appear to be slightly higher than the global average for savannas (Akagi et al. (2011) report an EF CO of 63 g kg⁻¹, whilst the mean EF CO for the Australian studies, including this one, is 81 g kg⁻¹). This might indicate that the efficiency of burning in Australian tropical savannas might be lower than the global savanna average; indeed, the global ecosystem averages include emissions from grasslands, which are believed to burn with higher MCE due to the relatively higher abundance of finer fuels than is found in woodland savannas (van Leeuwen and van der Werf et al., 2011).

For the methane emission factor (EF CH₄), the EF reported here agrees with all of the previous Australian studies and the global savanna EF CH₄. The mean emission factors of C₂H₂ and C₂H₄ are both lower than those reported by Andreae and Merlet (2001) and Akagi et al. (2011), and are closer to those measured in the northern Australian studies (the mean EF C₂H₂ and EF C₂H₄ for the northern Australian studies, including this one, at 0.11 g kg⁻¹ and 0.44 g kg⁻¹ respectively, are both about half of those reported for global savannas).

The emission factor for formaldehyde (EF H₂CO) is six-times higher than the only previous study to report EF H₂CO for Australian savanna fires (Hurst et al., 1994a), but is in closer agreement to the estimates for global savannas (Andreae and Merlet, 2001; Akagi et al., 2011). Paton-Walsh et al. (2010) (who report emission ratios to CO, but not emission factors, for northern Australian fires from solar occultation FTIR measurements) also find that their emission ratio for H₂CO is far higher than that reported in Hurst et al. (1994a) and suggest that one reason for this discrepancy might be that the H₂CO measured by Hurst et al. (1994a), whose study was based on cryogenically stored samples, may have been subject to losses during sampling or storage. Both the

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



OP-FTIR methodology used here and the methodology of Paton-Walsh et al. (2010) use long open-path measurements and so do not suffer from such problems associated with losses to the walls of sampling or storage equipment. Wooster et al. (2011), who also use a long open-path FTIR methodology, also find their H₂CO emission factor for fires measured in Kruger National Park (South Africa) to be significantly elevated compared with literature values for African savannah fires.

The emission factors for CH₃OH, CH₃COOH and HCOOH are the first to be reported for Australian savannas. Globally, only two savanna studies report EFs for these gases in the database of Andreae and Merlet (2001, with 2009 updates), whilst the standard deviation for the EFs of these gases is high in Akagi et al. (2011) (ranging from 35–48 % of the mean EF). EF HCOOH and EF CH₃OH reported here lies closest to the latest estimates of Akagi et al. (2011) for global savannas, whilst EF CH₃COOH is somewhat lower than those reported in both Andreae and Merlet (2001) and Akagi et al. (2011). The number of studies that contribute to this “global” average, however, is limited (only two studies for EF CH₃COOH in Andreae and Merlet, 2001, with 2009 updates, and five studies in Akagi et al., 2011), and it is likely that the lower emission factor reported here is within the realms of natural variability for CH₃COOH emissions.

5.2 Nitrogenous species

There is a very clear discrepancy between the emission factors for HCN (EF HCN) reported here and those from Hurst et al. (1994a, b). EF HCN reported here is over an order of magnitude higher than those from the previous Australian savanna studies. As was the case for H₂CO, Paton-Walsh et al. (2010) also find their measured emission ratio of HCN to CO to be over 6 times higher and 3 times higher than that of Hurst et al. (1994a) and Hurst et al. (1994b) respectively. It is suggested that one reason for these discrepancies might be that the HCN measured by Hurst et al. (1994a) and Hurst et al. (1994b) may also have been subject to losses during sample handling and storage. The EF HCN reported here agrees with measurements from other savanna ecosystems (Table 5).

The emission factor for ammonia (EF NH₃) lies closest to that reported by Hurst et al. (1994b) for Australian savannas (Table 5), and shows no significant difference from that reported for global savannas in Andreae and Merlet (2001) and Akagi et al. (2011).

5 6 Summary and conclusions

Griffith (1991) first demonstrated the use of long open-path FTIR for vegetation fire emission ratio measurements, and this approach was then revisited by Wooster et al. (2011) where a methodology for deriving biomass burning trace gas emission factors from these long open-path FTIR measurements was developed and applied to fires in the savannahs of Kruger. That work focussed on five key gases retrieved in different parts of the infrared spectral region ($900\text{--}3100\text{ cm}^{-1}$). Most recently, Akagi et al. (2013) also deploy a long open-path FTIR methodology at fires in the United States, greatly expanding the number of gases retrieved using this approach. The current work and the companion paper by Paton-Walsh et al. (2013) have retrieved emission factors for fifteen different trace gases, has explored different ways of combining time-series of emission measurements to derived “whole-fire” emission factors, and has explored relationships between these emission factors and the measurements modified combustion efficiencies, seasonality and vegetation class. This work has applied these methods to a large number (21) of fires conducted in Australian savanna, including both prescribed/experimental burns and wildfires, and is one of only a few limited studies of biomass burning emissions in the Australian savannas, which is surprising given that Australian savanna burning is an important source of a number of climatologically important atmospheric trace gases at both a national and global level. Generally, the averaged emission factors presented here are in line with global savanna emission factors (e.g. Akagi et al., 2011; Table 5), with the exception of formaldehyde (H_2CO) that has an emission factor of around double of that reported by Andreae and

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

Tables Figures

Tables

Figures



Merlet (2001) and Akagi et al. (2011). This is in line with other studies that deploy open-path methodologies (Paton-Walsh et al., 2010; Wooster et al., 2011).

The emission factors for H₂CO and HCN are substantially larger than those measured in previous studies of Australian savannas (Hurst et al., 1994a, b). As suggested by Paton-Walsh et al. (2010), who also find discrepancies between their measurements of H₂CO and HCN in northern Australia and those of Hurst et al. (1994a) and Hurst et al. (1994b), one explanation for this underestimation by these earlier studies might be due to losses of H₂CO and HCN during sample handling and storage. For CH₃OH, CH₃COOH, and HCOOH, the emission factors reported here represent one of only a few studies to report EFs of these gases in savanna burning globally.

The reported variability in emission factors for individual fires (Table 4) and the standard deviation of the mean emission factor calculated from all of the measured fires is large, although it is in line with the variability reported by Akagi et al. (2011). Understanding the causes of this variability is one of the important scientific questions posed by those who wish to better model global biomass burning emissions (van Leeuwan and van der Werf, 2011). Here, and in Meyer et al. (2012), we find that the emission ratios and emission factors varied substantially for different vegetation classes (Fig. 11 and Table 4), but that the MCE of the burns can be used to explain the majority of this variation (Fig. 12 and Fig. 13).

Acknowledgements. We thank the North Australian Indigenous Land Sea Management Alliance (NAILSMA), Indigenous Rangers of Warrdeken Land Management Limited, the Mok Clan as traditional owners of the field sites. The 2009 Arnhem Land campaigns formed part of a collaborative project supported by the Australian Government Caring For Our Country Program through NAILSMA. We are also grateful for the technical support provided by Bushfires NT and CSIRO. We would like to thank Christopher MacLellan and Alasdair MacArthur of NERC FSF for their wide ranging support, including the loan of the FTIR spectrometer used here. Funding for the studentship of T. E. L. Smith comes from NERC/ESRC Studentship ES/F012551/1. The contribution of Martin Wooster was supported by the NERC National Centre for Earth Observation (NCEO) and NERC grant NE/J010502/1.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



References

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Garde, M.: The language of fire: seasonality, resources and landscape burning on the Arnhem Land Plateau, in: *Culture, Ecology and Economy of Savanna Fire Management in Northern Australia: Rekindling the Wurrk Tradition*, edited by: Russell-Smith, J., Whitehead, P., and Cooke, P., CSIRO Publishing, Collingwood, Victoria, 85–164, 2009.

5 Griffith, D. W. T.: Synthetic calibration and quantitative analysis of gas-phase FTIR spectra, *Appl. Spectrosc.*, 50, 59–70, 1996.

Griffith, D. W. T., Mankin, W. G., Coffey, M. T., Ward, D. E., and Riebau, A.: FTIR remote sensing of biomass burning emissions of CO₂, CO, CH₄, CH₂O, NO, NO₂, NH₃ and N₂O, in: *Global Biomass Burning*, edited by: Levine, J. S., The MIT Press, Cambridge, Massachusetts, 230–239, 1991.

10 Haverd, V., Raupach, M. R., Briggs, P. R., J. G. Canadell., Davis, S. J., Law, R. M., Meyer, C. P., Peters, G. P., Pickett-Heaps, C., and Sherman, B.: The Australian terrestrial carbon budget, *Biogeosciences*, 10, 851–869, doi:10.5194/bg-10-851-2013, 2013.

Hurst, D. F., Griffith, D. W. T., and Cook, G. D.: Trace gas emissions from biomass burning in tropical Australian savannas, *J. Geophys. Res.*, 99, 16441–16456, 1994a.

15 Hurst, D. F., Griffith, D. W. T., Carras, J. N., Williams, D. J., and Fraser, P. J.: Measurements of trace gases emitted by Australian savanna fires during the 1990 dry season, *J. Atmos. Chem.*, 18, 33–56, 1994b.

Keane, W. C., Lobert, J. M., Crutzen, P. J., Maben, J. R., Scharffe, D. H., Landmann, T., He'ly, C., and Brain, C.: Emissions of major gaseous and particulate species during experimental burns of southern African biomass, *J. Geophys. Res.*, 111, D04301, doi:10.1029/2005JD006319, 2006.

Lobert, J. M. and Warnatz, J.: Emissions from the combustion process in vegetation, In: *Fire in the Environment: the Ecological, Atmospheric, and Climatic Importance of Vegetation Fires*, edited by: Crutzen, P. J. and Goldammer, J. G., John Wiley, New York, 15–37, 1993.

25 Lobert, J. M., Scharffe, D. H., Hao, W. M., Kuhlbusch, T. A., Seuwen, R., Warneck, P., and Crutzen, P. J.: Experimental evaluation of biomass burning emissions: nitrogen and carbon containing compounds, in: *Global Biomass Burning: Atmospheric, Climatic and Biospheric Implications*, edited by: Levine, J. S., MIT Press, Cambridge, Massachusetts, 289–304, 1991.

30 Lynch, B. T. and Wilson, P. L.: Land systems of Arnhem Land, Natural Resources Division, Northern Territory Department of Lands, Planning and Environment, Palmerston, 168 pp., 1998.

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

- Meyer, C. P., Cook, G. D., Reisen, F., Smith, T. E. L., Tattaris, M., Russell-Smith, J., Maier, S. W., Yates, C. P., and Wooster, M. J.: Direct measurements of the seasonality of emission factors from savanna fires in northern Australia. *J. Geophys. Res.*, 117, D20305, doi:10.1029/2012JD017671, 2012.
- 5 Müller, U., Kurte, R., and Heise, H. M.: Investigation of photometric errors in FTIR-spectra obtained in open-path monitoring, *J. Mol. Struct.*, 482–483, 539–544, 1999.
- Parks Australia: Terrestrial Ecoregions in Australia, available at: <http://www.environment.gov.au/topics/land/nrs/science-maps-and-data/australias-bioregions-ibra/australias-ecoregions> (last access: 24 September 2012), 2010.
- 10 Paton-Walsh, C., Deutscher, N. M., Griffith, D. W. T., Forgan, B. W., Wilson, S. R., Jones, N. B., and Edwards, D. P.: Trace gas emissions from savanna fires in northern Australia. *J. Geophys. Res.*, 115, D16314, doi:10.1029/2009JD013309, 2010.
- 15 Paton-Walsh, C., Smith, T. E. L., Young, E. L., Griffith, D. W. T., and Guérette, É.-A.: New emission factors for Australian vegetation fires measured using open-path Fourier transform infrared spectroscopy – Part 1: methods and Australian temperate forest fires, *Atmos. Chem. Phys. Discuss.*, 14, 4327–4381, doi:10.5194/acpd-14-4327-2014, 2014.
- Russell-Smith, J., Yates, C. P., Whitehead, P. J., Smith, R., Craig, R., Allan, G. E., Thackway, R., Frakes, I., Cridland, S., Meyer, C. P., and Gill, A. M.: Bushfires “down under”: patterns and implications of contemporary Australian landscape burning, *Int. J. Wildland Fire*, 16, 361–377, 2007.
- 20 Russell-Smith, J., Murphy, B. P., Meyer, M. C. P., Cook, G. D., Maier, S., Edwards, A. C., Schatz, J., and Brocklehurst, P.: Improving estimates of savanna burning emissions for greenhouse accounting in northern Australia: limitations, challenges, applications, *Int. J. Wildland Fire*, 18, 1–18, 2009.
- 25 Shirai, T., Blake, D. R., Meinardi, S., Rowland, F. S., Russell-Smith, J., Edwards, A., Kondo, Y., Koike, M., Kita, K., Machida, T., Takegawa, N., Nishi, N., Kawakami, S., and Ogawa, T.: Emission estimates of selected volatile organic compounds from tropical savanna burning in northern Australia, *J. Geophys. Res.*, 108, D38406, doi:10.1029/2001JD000841, 2003.
- Sinha, P., Hobbs, P. V., Yokelson, R. J., Bertschi, I. T., Blake, D. R., Simpson, I. J., Gao, S., Kirchstetter, T. W., and Novakov, T.: Emissions of trace gases and particles from savanna fires in southern Africa, *J. Geophys. Res.*, 108, D138487, doi:10.1029/2002JD002325, 2003.
- 30 Smith, T. E. L., Wooster, M. J., Tattaris, M., and Griffith, D. W. T.: Absolute accuracy and sensitivity analysis of OP-FTIR retrievals of CO₂, CH₄ and CO over concentrations representative

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



of “clean air” and “polluted plumes”, *Atmos. Meas. Tech.*, 4, 97–116, doi:10.5194/amt-4-97-2011, 2011.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

van Leeuwen, T. T. and van der Werf, G. R.: Spatial and temporal variability in the ratio of trace gases emitted from biomass burning, *Atmos. Chem. Phys.*, 11, 3611–3629, doi:10.5194/acp-11-3611-2011, 2011.

10 Wooster, M. J., Freeborn, P. H., Archibald, S., Oppenheimer, C., Roberts, G. J., Smith, T. E. L., Govender, N., Burton, M., and Palumbo, I.: Field determination of biomass burning emission ratios and factors via open-path FTIR spectroscopy and fire radiative power assessment: headfire, backfire and residual smouldering combustion in African savannahs, *Atmos. Chem. Phys.*, 11, 11591–11615, doi:10.5194/acp-11-11591-2011, 2011.

15 Yokelson, R. J., Griffith, D. W. T., and Ward, D. E.: Open-path Fourier transform infrared studies of large-scale laboratory fires, *J. Geophys. Res.*, 101, 21067–21080, 1996.

Yokelson, R. J., Bertschi, I. T., Christian, T. J., Hobbs, P. V., Ward, D. E., and Hao, W. M.: Trace gas measurements in nascent, aged, and cloud-processed smoke from african savanna fires by airborne Fourier transform infrared spectroscopy (A-FTIR), *J. Geophys. Res.*, 108, D138478, doi:10.1029/2002JD002322, 2003.

New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



New emission factors for Australian vegetation fires

T. E. L. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 1. List of fires studied using OP-FTIR during the three Northern Territory measurements campaigns in Arnhem Land and the Darwin Region, with information about the plot location, OP-FTIR pathlength, and meteorological variables.

Date	Local time	Dur. ^a (min)	Site name	Fire type	Veg. type	Location		OP-FTIR		Weather		
						Latitude	Longitude	Path (m)	T (°C)	RH (%)	P (hPa)	U (ms ⁻¹)
Arnhem Land Campaign (early dry season)												
2 Jul 2009	1430	90	ED-TGOW-A	experiment	tgow	12°44.45' S	133°53.13' E	46.4	30.6	39	997	6.0
3 Jul 2009	1330	70	ED-TGOW-B	experiment	tgow	12°44.58' S	133°53.13' E	41.2	28.9	46	998	5.0
4 Jul 2009	1250	40	ED-SOW-A	experiment	sow	12°39.07' S	133°53.99' E	38.5	25.6	42	1000	4.0
4 Jul 2009	1400	35	ED-SOW-B	experiment	sow	12°39.12' S	133°53.95' E	38.5	26.3	41	999	4.0
5 Jul 2009	1430	35	ED-SPIN	experiment	spin	12°38.32' S	133°54.75' E	22–27 ^b	27.8	37	996	2.0
Arnhem Land Campaign (late dry season)												
30 Sep 2009	1400	45	LD-TGOW-A	experiment	tgow	12°44.76' S	133°53.17' E	40	34.7	34	993	4.8
1 Oct 2009	1400	40	LD-TGOW-B	experiment	tgow	12°44.89' S	133°53.17' E	48.5	32.1	46	994	1.6
2 Oct 2009	1430	30	LD-SOW-A	experiment	sow	12°44.92' S	133°53.13' E	51	34.3	35	995	3.2
2 Oct 2009	1530	60	LD-SPIN	experiment	spin	12°38.32' S	133°54.75' E	22–25 ^b	34.9	34	994	0.3
3 Oct 2009	1300	90	LD-HGOW	experiment	hgow	12°43.01' S	133°50.43' E	70.5	35.2	31	1001	11.5
Darwin Region Campaign												
11 Sep 2010	1430	90	BAT-TGGW	burn-off	tggw	13°04.95' S	131°02.75' E	40.0	36.2	31	1009	4.2
1 Oct 2010	1330	70	ARS-TGFP-A	wildfire	tgfp	12°48.93' S	131°20.05' E	30.5	36.9	19	1007	1.7
1 Oct 2010	1250	40	ARS-TGFP-B	wildfire	tgfp	12°48.93' S	131°20.15' E	34.8	36.9	19	1007	1.7
1 Oct 2010	1400	35	ARS-TGFP-C	wildfire	tgfp	12°48.93' S	131°20.25' E	40.8	36.9	19	1007	1.7
1 Oct 2010	1430	35	ARS-TGFP-D	wildfire	tgfp	12°48.93' S	131°20.35' E	38.0	36.9	19	1007	1.7
1 Oct 2010	1300	90	AHW-TGOW	wildfire	tgow	12°36.32' S	131°14.57' E	34.0	34	19	1007	1.7
4 Oct 2010	1300	25	BSP-TGOW	wildfire	tgow	12°42.19' S	131°01.43' E	29.4	36.9	31	1006	1.1
5 Oct 2010	1400	45	TWP-TGOW-A	experiment	tgow	12°41.37' S	130°58.80' E	35.0	29.8	65	1011	0.6
5 Oct 2010	1400	40	TWP-TGOW-B	experiment	tgow	12°41.69' S	130°58.80' E	40.0	29.8	65	1011	0.6
5 Oct 2010	1430	30	TWP-TGOW-C	experiment	tgow	12°42.02' S	130°58.80' E	38.5	29.8	65	1011	0.6
6 Oct 2010	1530	60	SPT-TGOW	burn-off	tgow	12°43.63' S	130°56.71' E	46.0	29.8	65	1011	0.1

^a Duration for the wildfires and burn-off fires refers to the length of the dataset.

^b Multiple OP-FTIR paths were used for the spinifex fires.

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

Table 2. Mean emission ratios for compound Y with respect to CO₂ (ER_{Y/CO_2}) calculated from OP-FTIR measurements at 21 tropical savanna fires over the three campaigns to the Northern Territory. The standard deviation of the emission ratios is also included. The “Flaming” and “Smouldering” classes are defined on the basis of an MCE threshold of 90 %, as detailed in the main text.

Gas	All measurements $ER_{Y/CO_2} \pm SD$	Flaming $ER_{Y/CO_2} \pm SD$	Smouldering $ER_{Y/CO_2} \pm SD$
CO	0.0949 ± 0.0351	0.0814 ± 0.0262	0.1353 ± 0.0396
CH ₄	0.00281 ± 0.00183	0.00213 ± 0.00100	0.00483 ± 0.00142
C ₂ H ₂	0.00010 ± 0.00004	0.00009 ± 0.00004	0.00016 ± 0.00013
C ₂ H ₄	0.00046 ± 0.00021	0.00039 ± 0.00022	0.00061 ± 0.00032
C ₂ H ₆	0.00017 ± 0.00014	0.00010 ± 0.00002	0.00018 ± 0.00009
H ₂ CO	0.00139 ± 0.00044	0.00119 ± 0.00026	0.00186 ± 0.00078
CH ₃ OH	0.00062 ± 0.00057	0.00039 ± 0.00021	0.00087 ± 0.00059
CH ₃ COOH	0.00055 ± 0.00023	0.00046 ± 0.00021	0.00103 ± 0.00037
HCOOH	0.00010 ± 0.00006	0.00010 ± 0.00007	0.00010 ± 0.00003
HCN	0.00048 ± 0.00025	0.00037 ± 0.00019	0.00067 ± 0.00024
NH ₃	0.00091 ± 0.00058	0.00077 ± 0.00042	0.00170 ± 0.00069

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

Table 3. Same as Table 2, but for mean emission ratios for compound Y with respect to CO ($ER_{Y/CO}$).

Gas	All measurements $ER_{Y/CO} \pm SD$	Flaming $ER_{Y/CO} \pm SD$	Smouldering $ER_{Y/CO} \pm SD$
CO	12.07 ± 4.62	13.39 ± 3.96	7.79 ± 1.57
CH_4	0.0399 ± 0.0072	0.0384 ± 0.0051	0.0402 ± 0.0092
C_2H_2	0.0011 ± 0.0006	0.0014 ± 0.0007	0.0012 ± 0.0009
C_2H_4	0.0057 ± 0.0018	0.0058 ± 0.0026	0.0048 ± 0.0025
C_2H_6	0.0014 ± 0.0010	0.0010 ± 0.0001	0.0014 ± 0.0006
H_2CO	0.0182 ± 0.0028	0.0202 ± 0.0044	0.0148 ± 0.0055
CH_3OH	0.0083 ± 0.0038	0.0074 ± 0.0021	0.0080 ± 0.0050
CH_3COOH	0.0081 ± 0.0014	0.0082 ± 0.0016	0.0079 ± 0.0035
$HCOOH$	0.0008 ± 0.0002	0.0010 ± 0.0003	0.0008 ± 0.0002
HCN	0.0053 ± 0.0024	0.0048 ± 0.0023	0.0051 ± 0.0021
NH_3	0.0129 ± 0.0042	0.0128 ± 0.0041	0.0128 ± 0.0055

New emission factors for Australian vegetation fires

T. E. L. Smith et al.

Table 4. Emission factors (g kg^{-1} of dry fuel burned) for each individual fire, calculated using the standardised method outlined in Paton-Walsh et al. (2014). Uncertainties were calculated in quadrature from those associated with the trace gas emission ratios and a $\pm 10\%$ uncertainty in the assumed fuel carbon.

	CO_2 EF	CO_2 unc	CO EF	CO unc	CH_4 EF	CH_4 unc	C_2H_2 EF	C_2H_2 unc	C_2H_4 EF	C_2H_4 unc	C_2H_6 EF	C_2H_6 unc	CH_2O EF	CH_2O unc	CH_3OH EF	CH_3OH unc	CH_3COOH EF	CH_3COOH unc	HCN EF	HCN unc	NH_3 EF	NH_3 unc
LD-TGOW-A	1638 \pm 164	110 \pm 18	2.56 \pm 0.41	0.16 \pm 0.03	0.82 \pm 0.08	nr \pm nr	1.76 \pm 0.18	1.15 \pm 0.18	1.97 \pm 0.32	0.14 \pm 0.02	0.60 \pm 0.10	0.91 \pm 0.15										
LD-TGOW-B	1603 \pm 160	134 \pm 21	3.22 \pm 0.51	0.11 \pm 0.02	0.64 \pm 0.06	nr \pm nr	2.08 \pm 0.21	1.20 \pm 0.19	1.96 \pm 0.31	0.12 \pm 0.02	0.49 \pm 0.08	0.82 \pm 0.13										
LD-SOW-A	1528 \pm 153	169 \pm 27	5.94 \pm 0.95	0.18 \pm 0.03	0.75 \pm 0.08	nr \pm nr	2.43 \pm 0.24	3.76 \pm 0.60	3.28 \pm 0.52	0.31 \pm 0.05	1.12 \pm 0.18	1.63 \pm 0.26										
LD-SPIN	1716 \pm 172	75 \pm 12	1.28 \pm 0.20	0.18 \pm 0.03	0.11 \pm 0.01	nr \pm nr	0.53 \pm 0.05	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	0.36 \pm 0.06	0.61 \pm 0.10									
LD-HGOW	1662 \pm 166	100 \pm 16	2.01 \pm 0.32	0.09 \pm 0.01	0.24 \pm 0.02	nr \pm nr	0.67 \pm 0.11	1.57 \pm 0.25	nr \pm nr	nr \pm nr	0.66 \pm 0.11	0.51 \pm 0.08										
ED-TGOW-A	1669 \pm 167	96 \pm 15	1.92 \pm 0.31	nr \pm nr	0.16 \pm 0.02	0.04 \pm 0.01	1.44 \pm 0.14	nr \pm nr	0.96 \pm 0.15	nr \pm nr	nr \pm nr	0.11 \pm 0.02										
ED-TGOW-B	1679 \pm 168	87 \pm 14	1.81 \pm 0.29	nr \pm nr	0.74 \pm 0.07	nr \pm nr	0.81 \pm 0.13	1.64 \pm 0.26	nr \pm nr	nr \pm nr	nr \pm nr	0.60 \pm 0.10										
ED-SOW-A	1670 \pm 167	92 \pm 15	2.16 \pm 0.35	nr \pm nr	0.52 \pm 0.05	nr \pm nr	1.54 \pm 0.15	0.99 \pm 0.16	1.92 \pm 0.31	0.17 \pm 0.03	nr \pm nr	0.58 \pm 0.09										
ED-SOW-B	1624 \pm 162	118 \pm 19	2.87 \pm 0.46	0.07 \pm 0.01	0.56 \pm 0.06	nr \pm nr	1.39 \pm 0.14	1.48 \pm 0.24	2.64 \pm 0.42	nr \pm nr	0.17 \pm 0.03	0.97 \pm 0.16										
ED-SPIN	1763 \pm 176	39 \pm 6	0.71 \pm 0.11	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr										
BAT-TGGW	1713 \pm 171	60 \pm 10	1.65 \pm 0.26	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	1.02 \pm 0.16	nr \pm nr	nr \pm nr	0.42 \pm 0.07									
ARS-TGFP-A	1746 \pm 175	45 \pm 7	1.11 \pm 0.18	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	1.35 \pm 0.13	nr \pm nr	0.89 \pm 0.14	nr \pm nr	nr \pm nr	0.40 \pm 0.06									
ARS-TGFP-B	1676 \pm 168	86 \pm 14	1.61 \pm 0.26	0.13 \pm 0.02	nr \pm nr	0.07 \pm 0.01	1.62 \pm 0.16	0.69 \pm 0.11	1.30 \pm 0.21	nr \pm nr	0.32 \pm 0.05	0.62 \pm 0.10										
ARS-TGFP-C	1695 \pm 169	76 \pm 12	1.45 \pm 0.23	nr \pm nr	0.34 \pm 0.03	nr \pm nr	1.34 \pm 0.13	0.62 \pm 0.10	1.39 \pm 0.22	nr \pm nr	nr \pm nr	0.51 \pm 0.08										
ARS-TGFP-D	1746 \pm 175	46 \pm 7	0.84 \pm 0.13	0.10 \pm 0.02	0.29 \pm 0.03	nr \pm nr	1.36 \pm 0.14	nr \pm nr	0.87 \pm 0.14	0.07 \pm 0.01	0.38 \pm 0.06	0.35 \pm 0.06										
ARN-TGFP	1736 \pm 174	49 \pm 8	1.09 \pm 0.17	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	0.39 \pm 0.06	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr										
BSP-TGOW	1695 \pm 170	76 \pm 12	1.79 \pm 0.29	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	1.09 \pm 0.11	0.70 \pm 0.11	1.15 \pm 0.18	nr \pm nr	nr \pm nr	0.84 \pm 0.14									
TWP-TGOW-A	1740 \pm 174	49 \pm 8	1.15 \pm 0.18	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	1.04 \pm 0.10	0.44 \pm 0.07	0.94 \pm 0.15	nr \pm nr	nr \pm nr	0.54 \pm 0.09									
TWP-TGOW-B	1655 \pm 165	96 \pm 15	2.60 \pm 0.42	nr \pm nr	nr \pm nr	nr \pm nr	nr \pm nr	1.77 \pm 0.18	nr \pm nr	1.56 \pm 0.25	0.14 \pm 0.02	nr \pm nr	1.24 \pm 0.20									
TWP-TGOW-C	1627 \pm 163	114 \pm 18	3.06 \pm 0.49	0.07 \pm 0.01	0.46 \pm 0.05	0.13 \pm 0.02	2.32 \pm 0.23	0.87 \pm 0.14	1.54 \pm 0.25	0.14 \pm 0.02	nr \pm nr	1.04 \pm 0.17										
SPT-TGOW	1721 \pm 172	59 \pm 9	1.23 \pm 0.20	nr \pm nr	0.31 \pm 0.03	nr \pm nr	0.99 \pm 0.10	nr \pm nr	1.21 \pm 0.19	nr \pm nr	nr \pm nr	0.49 \pm 0.08										



New emission factors for Australian vegetation fires

T. E. L. Smith et al.

Table 5. Emission factors (g kg^{-1} dry fuel burned) reported by this study and those for the same trace gases reported by previous Australian savanna studies (Hurst et al., 1994a, b; Shirai et al., 2003) and by Andreae and Merlet (2001) and Akagi et al. (2011) who estimate global savanna emission factors by amalgamating suite of emissions studies. For this study and the Australian savanna studies, the mean and standard deviation of the emission factor is calculated from individual fires. The emission factor reported by Andreae and Merlet (2001) is the mean emission factor of all of the contributing studies, and the standard deviation is that of the studies (unless only two studies were used for a particular emission factor, in which case, the uncertainty is represented by a range). The emission factor reported by Akagi et al. (2011) is the mean of the contributing studies, whilst the uncertainty is an indication of the natural variability of the emission factor estimated from the reported variability in the contributing studies.

	Australian Tropical Savanna						All Savanna					
	This Study		Hu1994		Hu1994		Sh2003		AM2001		Ak2011	
	EF	SD	EF	SD	EF	SD	EF	SD	EF	SD	EF	NV
CO_2	1674 ± 56		1595 ± 121		1646 ± 106		1613 ± 111		1646 ± 99		1686 ± 38	
CO	87 ± 33		91.4 ± 27		58.9 ± 20		88 ± 8		61 ± 16		63 ± 17	
CH_4	2.10 ± 1.16		2.3 ± 1.29		2.3 ± 0.79		2.2 ± 0.4		2.2 ± 0.8		1.94 ± 0.85	
C_2H_2	0.11 ± 0.04		0.12 ± 0.08		0.10 ± 0.05		0.12 ± 0.02		0.27 ± 0.22		0.24 ± 0.1	
C_2H_4	0.49 ± 0.22		nr ± nr		0.53 ± 0.24		0.31 ± 0.04		0.84 ± 0.46		0.82 ± 0.35	
C_2H_6	0.08 ± 0.05		0.05 ± 0.03		0.07 ± 0.02		0.53 ± 0.07		0.32 ± 0.15		0.66 ± 0.41	
CH_2O	1.57 ± 0.44		0.26 ± 0.16		nr ± nr		nr ± nr		0.71 ± 0.42		0.73 ± 0.62	
CH_3OH	1.06 ± 0.87		nr ± nr		nr ± nr		nr ± nr		$1.47 \pm 1.20\text{--}1.74$		1.18 ± 0.41	
CH_3COOH	1.54 ± 0.64		nr ± nr		nr ± nr		nr ± nr		$2.61 \pm 2.40\text{--}2.81$		3.55 ± 1.47	
HCOOH	0.16 ± 0.07		nr ± nr		nr ± nr		nr ± nr		$0.63 \pm 0.62\text{--}0.63$		0.21 ± 0.10	
HCN	0.53 ± 0.31		0.03 ± 0.02		0.03 ± 0.02		nr ± nr		0.23 ± 0.55		0.41 ± 0.15	
NH_3	0.70 ± 0.36		1.53 ± 0.86		0.61 ± 0.39		nr ± nr		0.74 ± 0.25		0.52 ± 0.35	



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.




Fig. 1. Photographs showing the use of natural and artificial firebreaks to protect the OP-FTIR equipment during fires in which the open path between the spectrometer and IR source lay within the fire plots. **(a)** The FTIR spectrometer positioned on a rocky outcrop, with the fire being allowed to pass around the rocks; **(b)** the IR lamp protected by raked vegetation, notice the distinction between the burnt and unburned area.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

New emission factors for Australian vegetation fires

T. E. L. Smith et al.



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 2. Typical retrieval spectral windows for each of the trace gases retrieved for this paper: **(a)** CO₂ and CO; **(b)** CH₄; **(c)** C₂H₄ and NH₃; **(d)** H₂CO; **(e)** CH₃OH; **(f)** HCOOH; **(g)** CH₃COOH; **(h)** C₂H₆; and **(i)** C₂H₂ and HCN. The measured spectrum (black line) and the best-fit modelled spectrum (dashed red line) are shown along with each of the individually modelled trace gas spectra (multiple colours). The modelled spectrum is the sum of each of the individual trace gas spectra convolved to the background polynomial (see Table 1 in Paton-Walsh et al., 2014).

**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.




Fig. 3. Map of Australia, showing the two field campaign locations. Both the Arnhem Land and Darwin Region campaigns are situated in the “tropical and subtropical grasslands, savannas and shrublands” terrestrial ecoregion (adapted from Parks Australia, 2010).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Maps showing the location of each of the fire sites discussed in this paper (see Table 1 for full site descriptions).


**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.



Fig. 5. Photographs depicting the various savanna fire classes experienced during both the Arnhem Land (Arnhem) and Darwin region (Darwin) campaigns. The abbreviations in parentheses are used throughout this paper, more details about specific fire plots are given in Table 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



6350

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Fig. 6. Time series of path-averaged trace gas mole fractions (in ppm) for plot LD-TGOW-A (30 September 2009) during the late-dry season Arnhem Land campaign. Trace gas path-averaged mole fractions were retrieved from the OP-FTIR spectra using MALT. An example smoke spectrum with the best-fit modelled spectrum for each of the spectral windows used to retrieve the mole fraction of each of the trace gases presented in this figure can be found in Fig. 2. Gaps in the time series of specific trace gases are due to periods of low signal-to-noise within the spectral window used for the retrieval of that particular gas.



New emission factors
for Australian
vegetation fires

T. E. L. Smith et al.





Fig. 7. (top) Time series of retrieved CO₂ and CO pathlength averaged mole fractions, and (bottom) Modified Combustion Efficiency (MCE). Only the first ~60 s following ignition are dominated by smoke from purely flaming combustion, with increasing contribution from the smouldering-phase combustion zone towards the end of this time series (as reflected by the decreasing MCE throughout). These data are taken from LD-TGOW-A (30 September 2009). The full suite of gas mole fractions are presented in Fig. 6.

New emission factors for Australian vegetation fires

T. E. L. Smith et al.



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 8. Example scatter plots of the measured trace gas pathlength-averaged mole fractions used to calculate emission ratios. The data presented here are from LD-TGOW-A (30 September 2009). The slope of the least squares linear best fit to these data (thick red line) is used to derive the relevant emission ratio (ER_{X/CO_2}). For each of the ten trace gases shown above, the emission ratio and the R^2 is given towards the top of each plot. The equivalent plots for emission ratios to CO are presented in Fig. 9.

New emission factors for Australian vegetation fires

T. E. L. Smith et al.




Fig. 9. Same as Fig. 8, but with relation to CO. Notice the higher correlation (R^2) between each of the trace gases and carbon monoxide, than between each of the trace gases and carbon dioxide (Fig. 8).

New emission factors for Australian vegetation fires

T. E. L. Smith et al.





Fig. 10. An example scatter plot of the measured pathlength-averaged mole fractions of CO_2 and CH_4 (as in Fig. 8, top centre). Here, the data have been split using MCE, with measurements having an MCE greater than 90 % (red) being considered separately to those with a lower MCE (black). Different emission ratios and emission factors for flaming and smouldering emissions can then be determined (also shown on the plot in red and black text respectively, the green text indicates the emission ratio and factors determined using all data).



6357

Title Page

Abstract Introduction

Inclusions References

Tables Figures

|◀ ▶|

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Fig. 11. Scatter plots of the measured mole fractions of CO₂ and CH₄, including all measurements made during the two Arnhem Land campaigns in 2009. All three plots show the same data, but are colour-coded differently using **(a)** MCE; **(b)** season of measurement; and **(c)** vegetation class.

**New emission factors
for Australian
vegetation fires**

T. E. L. Smith et al.




Fig. 12. Same as Fig. 11a, but including linear best-fit lines that have been calculated using measurements subsampled by narrow (2 %) bins of MCE. Each line represents the least squares linear best fit to the subsampled emissions measurements (the MCE bin for each line is indicated on the plot).

New emission factors for Australian vegetation fires

T. E. L. Smith et al.




Fig. 13. The emission factor for CH_4 (EF CH_4) grouped by MCE, calculated using the emission ratios determined from 2 bins for different vegetation classes, and for all measurements combined (error bars indicate the estimated uncertainty calculated in accordance with Paton-Walsh et al., 2014).