



## Abstract

At the East Antarctic deep ice core drilling site Dome C, daily precipitation measurements have been initiated in 2006 and are being continued until today. The amounts and stable isotope ratios of the precipitation samples as well as crystal types are determined. Within the measuring period, the two years 2009 and 2010 showed striking contrasting temperature and precipitation anomalies, particularly in the winter seasons. The reasons for these anomalies and their relation to stable isotope ratios are analysed using data from the mesoscale atmospheric model WRF (Weather Research and Forecasting Model) run under the Antarctic Mesoscale Prediction System (AMPS). 2009 was relatively warm and moist due to frequent warm air intrusions connected to amplification of Rossby waves in the circumpolar westerlies, whereas the winter of 2010 was extremely dry and cold. It is shown that while in 2010 a strong zonal atmospheric flow was dominant, in 2009 an enhanced meridional flow prevailed, which increased the meridional transport of heat and moisture onto the East Antarctic plateau and led to a number of high-precipitation/warming events at Dome C. This was also evident in a positive (negative) SAM index and a negative (positive) ZW3 index during the winter months of 2010 (2009). Changes in the frequency or seasonality of such event-type precipitation can lead to a strong bias in the air temperature derived from stable water isotopes in ice cores.

## 1 Introduction

Although Antarctic precipitation has been studied for approximately half a century (see e.g. Bromwich, 1988), a number of open questions remain. There are two key motivations for studying Antarctic precipitation. The first is that precipitation/snowfall is the most important positive component of the mass balance of Antarctica. This is receiving increasing attention in discussions of climate change since the mass balance response to global warming can considerably influence sea level change. A possible

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





the Southern Ocean, the  $^{18}\text{O}$  values are lower during cold periods (winter/glacial) than during warm periods (summer/interglacial), which leads to clear seasonal variations and likewise large differences between glacial and interglacial periods in the stable isotope ratios measured in the ice core.

This spatially derived linear relationship has been found not to hold temporally, however (Jouzel et al., 2003). Apart from air temperature, several other factors influence the stable isotope ratio, such as seasonality of precipitation, location of and conditions at the moisture sources and conditions along moisture transport paths (e.g. Jouzel et al., 2003; Noone et al., 1999; Schlosser, 1999). Thus, for a correct interpretation of the ice core data a thorough understanding of the atmospheric processes responsible for the precipitation is needed, as it was the precipitation that ultimately formed the glacier ice investigated in the cores. In particular, information about moisture sources, moisture transport paths, and atmospheric conditions at the final deposition site is required.

Measuring Antarctic precipitation is a challenge, not only due to the remoteness and extreme climate of the continent, but also due to difficulties in distinguishing between drifting/blowing snow and falling precipitation. The latter is due to the high wind speeds that typically accompany precipitation events in coastal areas. In the interior of the continent, while wind speeds are lower than at the coast, the threshold for drifting snow is often lower due to lower snow densities as well. Measurements are also complicated by the extremely small amounts of precipitation produced in the cold and dry air. Precipitation measurements with optical devices may hold some hope for improved data in the future, but these instruments are currently in the testing phase in Antarctica (S. Colwell, personal communication, 2015). In light of the lack of observations, atmospheric models have become increasingly useful tools to investigate Antarctic precipitation (Bromwich et al., 2004; Noone et al., 1999; Schlosser et al., 2008, 2010a, b).

In this study, observational precipitation and stable isotope data from the deep-drilling site Dome C, East Antarctica, combined with data from a mesoscale atmospheric model are used to investigate the precipitation regime and its relation to stable

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



isotope ratios of the precipitation. The Dome C precipitation series is the first and so far only multi-year precipitation/stable isotope series at an Antarctic deep ice core drilling site.

This investigation concentrates on the years 2009 and 2010. These years were chosen because they showed striking contrasting temperature and precipitation anomalies, particularly in the winter seasons. Fogt (2010) reports that temperatures in the Antarctic were persistently above average in the mid-to-lower troposphere during the winter of 2009. The positive surface temperature anomalies were most marked in East Antarctica. In 2010, the picture was very different from 2009, with generally below-average temperatures on the East Antarctic plateau in winter and spring (Fogt, 2011).

## 2 Study site

Dome C (75.106° S, 123.346° E, elevation 3233 m) is one of the major domes on the East Antarctic ice sheet. Its mean annual temperature is  $-54.5^{\circ}\text{C}$ , and the mean annual accumulation derived from ice cores amounts to  $25\text{ mm w.e. yr}^{-1}$ . Several deep ice cores have been retrieved at Dome C, the first one in 1977/78, reaching a depth of 906 m, corresponding to an age of approximately 32 000 yr. The thermally drilled core was retrieved during the International Antarctic Glaciological Project (Lorius, 1979).

The oldest ice to date has been obtained at Dome C through the European deep drilling project EPICA (European Project for Ice Coring in Antarctica). The drilling was completed in January 2006; at the base of the 2774.15 m long ice core the age of the ice was estimated to be 800 000 yr, thus covering eight glacial cycles (EPICA community members, 2004). To support the EPICA drilling operation, the French-Italian Antarctic wintering base Dome Concordia became operational in 2005.

### Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Previous work

#### 3.1 Synoptics and precipitation

Precipitation conditions in the interior of Antarctica are very different from those in coastal areas. Whereas precipitation at the coast is usually caused by frontal systems of passing cyclones that form in the circumpolar trough (e.g. Simmonds et al., 2002), in the interior different precipitation mechanisms are at play. On the majority of days, only diamond dust, also called clear-sky precipitation, is observed. It forms due to radiative cooling in a nearly saturated air mass. Although diamond dust is predominant temporally, it does not necessarily account for the largest fraction of the total yearly precipitation. It has been shown that a few snowfall events per year can bring up to 50 % of the total annual precipitation (Braaten et al., 2000; Reijmer and Van den Broeke, 2003; Schlosser et al., 2010a; Fujita and Abe, 2006). Those events are due to amplification of Rossby waves in the circumpolar westerlies, which increases the meridional transport of heat and moisture polewards. In extreme cases this can even mean a transport from the Atlantic sector across the continent to the Pacific side (Sinclair, 1981; Schlosser et al., 2015b) The relatively moist and warm air is orographically lifted over the ice sheet, followed by cloud formation and/or precipitation (Noone et al., 1999; Massom et al., 2004; Birnbaum et al., 2006; Schlosser et al., 2010a, b). Except for the study by Fujita and Abe (2006), all of these investigations were based on model and AWS data, rather than daily precipitation measurements.

For a long time it was believed in ice core studies that precipitation represented in Antarctic ice cores is formed close to the upper boundary of the temperature inversion layer assuming that the largest moisture amounts are found where the air temperature is highest (Jouzel and Merlivat, 1984). However, more recent studies have shown that humidity inversions are parallel to the temperature inversion only in 50 % of the cases, and often multiple humidity inversions occur (Nygard et al., 2013). In particular, the local cycle of sublimation and re-sublimation (deposition) is poorly known, but it is important for both mass balance and isotope fractionation studies.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



poral variations of  $\delta^{18}\text{O}$  were highly correlated with air temperature. The lowest  $\delta^{18}\text{O}$  value measured was  $-81.9\text{‰}$ , which is the isotopically lightest water ever collected on the Earth's surface. Half of the annual precipitation resulted from only 11 events (18 days), without showing any seasonality. The other half was due to diamond dust. Similar results were found in studies by Schlosser et al. (2010a), at Kohlen Station (see Fig. 1) and by Reijmer and Van den Broeke (2003), who used data from automatic weather stations in Dronning Maud Land. The precipitation-weighted temperature was significantly higher than the mean annual surface temperature because the precipitation events were related to warm-air advection. Recently, Dittmann et al. (2015) investigated the stable isotope data obtained by Fujita and Abe (2006) at Dome Fuji for all days with dynamically caused snowfall in a combined approach of synoptic analysis and isotope modelling. They found that, for single events, the relationship between deuterium excess and atmospheric conditions at the moisture source used in ice core studies was not existent.

## 4 Data and methods

### 4.1 Precipitation and isotopes

Daily precipitation measurements were initiated at Dome C in 2006, and have, with some interruptions, been continued until today. Daily precipitation amounts are measured using a wooden platform set up at a distance of 800 m from the main station, at a height of 1 m above the snow surface to avoid contributions from low drifting snow. For the same reason, the platform is surrounded by a rail of approximately 8 cm height. The measurements include precipitation sampling and analysis of stable water isotopes ( $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ) of the samples. Additionally the crystal structure of the precipitation is analysed in order to distinguish between diamond dust, snowfall, and drift snow. Diamond dust consists of extremely fine ice needles whereas synoptic snowfall shows various types of regular snow crystals, which tend to be broken in case of drifting/blowing snow.

The snow crystal type depends on air temperature during formation in the cloud. Samples of mixed crystal types can also occur.

While errors of the precipitation measurements cannot be quantified, it is understood that they can exceed 100 % given the extremely small precipitation amounts.

The snow samples were sent to the Geochemistry Laboratory of the University of Trieste, where they were melted and stored in freezers at approximately  $-20^{\circ}\text{C}$  until, provided the precipitation amount was sufficient, they were analysed using a mass spectrometer (Thermo-Fisher Delta Plus XP). Very small samples were analysed using a Picarro I1102-I cavity-ringdown spectroscopy (CRDS) analyser. The precision of the Picarro I1102-I is 0.1 ‰ for  $\delta^{18}\text{O}$  and 0.5 ‰ for  $\delta\text{D}$  (Stenni et al., 2015). Details of the measurements and an extensive discussion of the full data set can be found in Stenni et al. (2015).

## 4.2 AWS data

The Antarctic Meteorological Research Center (AMRC) and Automatic Weather Station (AWS) Program are sister projects of the University of Wisconsin-Madison funded under the United States Antarctic Program (USAP) that focus on data for Antarctic research support, providing real-time and archived weather observations and satellite measurements and supporting a network of automatic weather stations across Antarctica.

The current AWS at Dome C was set up by the AMRC, in December 1995. The station measures the standard meteorological variables of air temperature, pressure, wind speed, wind direction, and humidity. Data can be obtained from <http://amrc.ssec.wisc.edu>. Note that an initial AWS (named Dome C) had been set up in 1985, however, at a distance of about 70 km from the current site. Thus, only data from the new station (Dome C II) are used in the present study.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 4.3 WRF Model output from the AMPS archive

In addition to the observations described above, this study uses numerical weather prediction (NWP) model output for analysis of the synoptic environments of the target years, of precipitation processes, and of events. The output is from forecasts of the Weather Research and Forecasting (WRF) Model (Skamarock et al., 2008) run under the Antarctic Mesoscale Prediction System (AMPS) (Powers et al., 2003, 2012), a real-time NWP capability that supports the weather forecasting for the United States Antarctic Program (USAP). The (US) National Center for Atmospheric Research (NCAR) has run AMPS since 2000 to produce twice-daily forecasts covering Antarctica with model grids of varying resolutions. The AMPS WRF forecasts have been stored in the AMPS Archive and used extensively in studies (e.g. Monaghan et al., 2005; Seefeldt and Cassano, 2008; Schlosser et al., 2008; Seefeldt and Cassano, 2012). For 2009 and 2010, the WRF output over the Dome C region reflects a forecast domain with a horizontal grid spacing of 15 km, employing 44 vertical levels between the surface and 10 hPa. This 15 km grid was nested within a 45 km grid covering the Southern Ocean, and Fig. 2 shows these domains.

Model output from AMPS has been verified through various means over the years. Multi-year AMPS forecast evaluations have been conducted (Bromwich et al., 2005), and WRF's ability for the Antarctic in particular has been confirmed (Bromwich et al., 2013). AMPS's and WRF's Antarctic performance has also been documented in a number of case and process studies (e.g. Bromwich et al., 2013; Powers, 2007; Nigro et al., 2011, 2012). For model development within AMPS, verification for both warm and cold season periods is performed prior to changes in model versions or configurations (Powers et al., 2012). The reliability of AMPS WRF forecasts is also reflected in their demand from international Antarctic operations and field campaign forecasting efforts (see e.g. Powers et al., 2012). Lastly, similarly to how it is used here, AMPS output has been a key tool in previous published studies of Antarctic precipitation related to ice core analyses (Schlosser et al., 2008, 2010a, b).

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





–30 °C can be seen. Minimum temperatures barely exceed –70 °C whereas in 2010, minima are close to –80 °C. The highest temperature in the winter of 2010 was only slightly above –50 °C.

A very high precipitation value of 1.36 mm on 9 February 2010, followed by 0.67 mm on 10 February, both classified as diamond dust, stems from only one event around 9 February. These values should be considered with care given the high error possibilities of the measurements. Considering the extremely low density of diamond dust, a diamond dust amount of more than 1 mm day<sup>-1</sup> seems to be unlikely. However, the model data do show a precipitation event connected to warm air advection from the north (see below) for this day, which would indicate the occurrence of snowfall rather than diamond dust. Most likely a mixture of crystal types was found during this event. The precipitation totals for May to September are 12.0 mm w.e. for 2009 and 4.3 mm w.e. for 2010. Daily sums exceed 0.25 mm only three times in 2010, but 16 times in 2009. Usually, high daily precipitation amounts are associated with relative maxima in air temperature. In general, the winter of 2010 was cold and dry, whereas 2009 was relatively warm and moist compared to the long-term average.

Figure 4a shows monthly precipitation amounts for 2009 and 2010, distinguishing between diamond dust, hoar frost, and snowfall; Fig. 4b gives the relative frequencies of the three different observed types of precipitation for both years. Again, large differences between 2009 and 2010 are found. While approximately half of the precipitation fell as snow in 2009, less than a quarter of the total precipitation stemmed from snowfall in 2010, when mostly diamond dust was observed. As seen before, the winter months of May to September exhibit the largest differences. In particular, the extremely “warm” July of 2009 brought high amounts of snowfall. The lowest amounts of precipitation are seen in austral summer 2009/10, with no precipitation observed in November and only very small amounts in December and January.

The total amount of precipitation measured on the raised platform is 16.5 mm w.e. for 2009 and 13.4 mm w.e. for 2010, compared to the mean annual accumulation of 25 mm w.e. derived from firn core and stake measurements (Frezzotti et al., 2005).

Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



From the given data it cannot be determined whether the difference is due to snow removed from the measuring platform by wind or sublimation or snow added to the snow surface by wind (blowing or drifting snow) or deposition (re-sublimation).

## 5.2 Atmospheric flow conditions

### 5.2.1 Synoptic analyses with AMPS archive data

The synoptic situations that caused precipitation at Dome C were analysed using WRF output data from the AMPS archive. In particular, fields of 500hpa geopotential height and 24 h precipitation were used. For the 500 hPa geopotential height information the 12 h forecast was utilized. For 24 h precipitation, the 12–36 h forecast sums of precipitation (rather than 0–24 h) were used to allow for model spin up of clouds and micro-physical fields. This is considered long enough for moist process spin-up, but avoids error growth reflected in longer forecast times (Bromwich et al., 2005).

For all precipitation events with observed daily sums exceeding 0.2 mm, the synoptic situations that caused the precipitation were investigated. In total, 29 events were studied, 20 in 2009 and 9 in 2010. For 2009 (2010), the model showed precipitation at Dome C in 44 (50 %) of the studied cases and precipitation in the vicinity in 33 (25) % of the cases; no precipitation was shown in the model in 22 (25) % of the cases. In total, approximately half of the precipitation events were represented well by the model, one quarter showed synoptic events that did not bring precipitation exactly at the location and time of the measurements, and one quarter of the cases were not forecast by the model at all.

Generally, snowfall events were found to be associated with an amplification of the Rossby waves in the circumpolar westerlies, which causes a northerly flow across the Dome C region between a trough to the west and an upper-level ridge to the east of Dome C. This northerly flow brings relatively warm and moist air from as far as 35–40° S to the East Antarctic plateau, leading to orographic precipitation when it is forced to ascend on the way from the coast to the high-altitude interior. Variations of this general

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ture source. (The 500 hPa trajectory seems to have some inconsistencies on the 5th day which should not be over-interpreted). Whereas it is not possible to exactly determine the moisture source (under the simplifying assumption of a single moisture source), the information is sufficient to distinguish between a source in the Southern Ocean and one at middle latitudes, which is most important for ice core interpretation. A frequent occurrence of the synoptic situation described (as it was the case in 2009) means a more northern mean moisture source than on average, which has to be taken into account for deriving air temperature from stable isotopes. (A detailed study using trajectory calculations for all observed precipitation events at Dome C is ongoing.) It was also found to be typical for precipitation events at Dome C that the main flow is split into a meandering branch and a branch that remains zonal. This is observed more often at Dome C than at Dome F (Dittmann et al., 2015) or at Kohlen Station (Schlosser et al., 2010a).

Figure 6 presents an example for a case with no precipitation in the model, but relatively large observed precipitation amounts. The 500 hPa geopotential height field (Fig. 6a) shows a cutoff-high west of Dome C on the day after the precipitation event shown in Fig. 5. The remaining atmospheric moisture is not sufficient to produce precipitation in the model (Fig. 6b), but it does lead to remarkably high amounts of diamond dust and/or hoar frost (0.7 mm observed during this event). This synoptic situation was also found by Hirasawa et al. (2013) in a detailed study of the synoptics and precipitation during and after a blocking event at Dome Fuji. (Note that neither diamond dust nor hoar frost formation is parameterized in the model.) In 2010, the flow was mainly zonal and the synoptic situations described above were much less frequent than in 2009 and not as strongly developed.

Using the WRF output, monthly composite fields of 500 hPa geopotential height were calculated to compare the general flow conditions in 2009 and 2010. Figure 7 shows the composite mean 500 hPa geopotential height for July 2009 and 2010, respectively. Even in the monthly mean, the distinct upper-level ridge in 2009 that projects onto the

East Antarctic plateau and leads to warm air advection and increased precipitation at Dome C is clearly seen.

In 2010, in the monthly average, the flow was mainly zonal, which reduced the meridional exchange of heat and moisture, thus leading to lower temperatures and less precipitation in the interior of the Antarctic continent.

## 5.2.2 Southern Annular Mode

The occurrence of high-precipitation events on the Antarctic plateau due to amplification of Rossby waves is often connected to a strongly positive phase of the Southern Annular Mode (SAM). The SAM is the dominant mode of atmospheric variability in the extratropical Southern Hemisphere. It is revealed as the leading empirical orthogonal function in many atmospheric fields (e.g. Thompson and Wallace, 2000), such as surface pressure, geopotential height, surface temperature, and zonal wind (Marshall, 2003). Since pressure fields from global reanalyses commonly used to study the SAM are known to have relatively large errors in the polar regions, Marshall (2003) defined an SAM index based on surface observations. He calculated the pressure differences between 40 and 65° S using data from six mid-latitude stations and six Antarctic coastal stations to calculate the corresponding zonal means. A large meridional pressure gradient corresponds to a positive SAM index and vice versa. The positive index means strong, mostly zonal westerlies and comparatively little exchange of moisture and energy between middle and high latitudes, which leads to a general cooling of Antarctica, except for the Antarctic Peninsula that projects into the westerlies. A negative SAM index is associated with weaker westerlies and a larger meridional flow component.

Figure 8 shows the monthly mean SAM index for 2009 and 2010 (data can be found at <http://www.nerc-bas.ac.uk/icd/gjma/sam.html>). Whereas in the winter months (May to September) of 2009 the SAM index was generally negative (with the exception of a weakly positive value in June), 2010 has positive indices from April to August, with strongly positive values in June and July, and only a weakly negative index in September. This is consistent with the pattern of a strong zonal flow with few precipitation

events at Dome C due to amplified ridges in the winter of 2010, with the opposite situation holding in 2009. The highest SAM index is found in November 2010; however, in austral summer the relationship between the SAM index and precipitation seems to be less straightforward. Furthermore, it should be kept in mind that SAM explains only about one third of the atmospheric variability in the Southern Hemisphere (Marshall, 2007).

### 5.2.3 Zonal Wave Number 3

Another method to investigate the general atmospheric flow conditions is to analyse spatial and temporal variations of the quasi-stationary zonal waves in the Southern Hemisphere. In this study Zonal Wave Number 3 (ZW3) is used. While the atmospheric circulation in the Southern Hemisphere appears strongly zonal (or symmetric), there is a significant non-zonal (asymmetric) component and ZW3 represents a significant proportion of this asymmetry. It is a dominant feature of the circulation on a number of different time scales (e.g. Karoly, 1989), is responsible for 8 % of the spatial variance in the field (Van Loon and Jenne, 1972), and contributes significantly to monthly and interannual circulation variability (e.g. Trenberth, 1990; Trenberth and Mo, 1985). The asymmetry is revealed when the zonal mean is subtracted from the geopotential height field thereby creating a coherent pattern of zonal anomalies, with the flow associated with these patterns becoming apparent. ZW3 has preferred regions of meridional flow, which influence the meridional transport of heat and moisture into and out of the Antarctic. Raphael (2004) defined an index of ZW3 based on its amplitude (effectively the size of the zonal anomaly) at 50° S showing that ZW3 has identifiable positive and negative phases associated with the meridionality of the flow. A positive value for this index indicates more meridional flow (large zonal anomaly) and a negative value more zonal flow (small zonal anomaly). Figure 9a shows the monthly mean ZW3 index for the period 2009–2010. From June to September 2009 the ZW3 index was largely positive except for a comparatively small negative excursion in July. On the contrary, from June to September 2010 it was negative. The asymmetry in the circulation suggested

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by the index is shown in Fig. 9b (July 2009) and 9c (July 2010). This figure was created by subtracting the long-term zonal mean at each latitude, from the mean 500 hPa geopotential height field in July 2009 and 2010, respectively. The flow onto Dome C suggested by the alternating negative and positive anomalies is northerly in July 2009, but has a strong zonal component in July 2010. This information given by the ZW3 index and the patterns of zonal anomalies is consistent with that suggested by the SAM.

### 5.3 Stable isotopes

Since the main motivation of the presented precipitation study is the improvement of the climatic interpretation of stable isotope data, in Fig. 10 the daily mean temperature and the measured stable isotope ratios of the precipitation samples, namely  $\delta^{18}\text{O}$  and the second-order parameter deuterium excess  $d$  ( $d = \delta\text{D} - 8 \delta^{18}\text{O}$ , globally averaged), are displayed for 2009 and 2010. As expected,  $\delta^{18}\text{O}$  and air temperature exhibit a similar annual cycle, with high values in summer and the lowest values in the winter months. Consistent with the unusually “warm” winter of 2009, also the  $\delta^{18}\text{O}$  reaches higher values in winter 2009 than in winter 2010. Because of the more meridional flow and thus more northerly (and warmer) oceanic moisture source, the initial  $\delta^{18}\text{O}$  is already higher than on average and the condensation temperature at Dome C is above-average during the precipitation events as well. In addition to the warm-air advection, the existing near-surface temperature inversion layer is often removed because of increased wind speed and increased cloud cover, the latter causing a change in the radiation balance, namely increased down-welling long-wave radiation. In contrast to  $\delta^{18}\text{O}$ , the deuterium excess shows maxima in winter and minima in summer. In winter 2010, the deuterium excess is clearly higher than in 2009; the difference between the maxima in 2009 and 2010 amounts to 20 ‰. A comprehensive analysis of the full stable isotope data set of Dome C can be found in a companion paper by Stenni et al. (2015).

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 6 Discussion and conclusion

Precipitation and stable water isotope data from two contrasting years 2009 and 2010 at the Antarctic deep-drilling site Dome C, on the East Antarctic Plateau were investigated in the present study. The observations from Dome C are the first and only multi-year series of daily precipitation/stable isotope measurements at a deep-drilling site, even though “multi” means only nine years in this case. The differences between the two years 2009 and 2010 were most striking in winter. Whereas 2009 was relatively warm and moist due to frequent warm air intrusions connected to amplification of Rossby waves in the circumpolar westerlies, the winter of 2010 was extremely cold and dry, with the lowest monthly mean July temperature observed since the beginning of the AWS measurements in 1996. This can be explained by the prevailing strong zonal flow in the winter of 2010, related to a strongly positive SAM index and a negative ZW3 index. Also, the frequency distribution of the various precipitation types was largely different in 2009 and 2010, with snowfall prevailing in 2009 whereas diamond dust was dominant in 2010.

Similarly striking differences in weather conditions of 2009 and 2010 were seen in other parts of East Antarctica. Gorodetskaya et al. (2013) found that accumulation in 2009 was eight times higher than in 2010 at the Belgian year-round station “Princess Elisabeth”. At this location, the temperature was also higher in 2009 than in 2010, particularly in fall/early winter. The findings are supported by Boening et al. (2012), who used observations from GRACE (Gravity Recovery And Climate Experiment) and found an abrupt mass increase on the East Antarctic ice sheet in the period 2009–2011. Similarly, Lenaerts et al. (2013) investigated snowfall anomalies in Dronning Maud Land, East Antarctica. They state that the large positive anomalies of accumulation found in 2009 and 2011 stand out in the past approximately 60 years although comparable anomalies are found further back in time.

Distinguishing between the different forms of precipitation, namely diamond dust, hoar frost and dynamically caused snowfall, is important for both mass balance and

### Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







ANT-12456663). We thank Gareth Marshall for providing the SAM indices online. We thank all winterers at Dome C, who did the precipitation sampling.

## References

- Birnbaum, G., Brauner, R., and Ries, H.: Synoptic situations causing high precipitation rates on the Antarctic plateau: observations from Kohnen Station, Dronning Maud Land, *Antarct. Sci.*, 18, 279–288, doi:10.1017/S0954102006000320, 2006.
- Boening, C., Lebsack, M., Landerer, F., and Stephens, G.: Snowfall-driven mass change on the East Antarctic ice sheet, *Geophys. Res. Lett.*, 39, L21501, doi:10.1029/2012GL053316, 2012.
- Braaten, D. A.: Direct measurements of episodic snow accumulation on the Antarctic polar plateau, *J. Geophysic. Res.*, 105, 10119–10128, 2000.
- Bromwich, D. H.: Snowfall in high southern latitudes, *Rev. Geophys.*, 26, 149–168, 1988.
- Bromwich, D. H., Guo, Z., Bai, L., and Chen, Q.: Modeled Antarctic precipitation. Part I: Spatial and temporal variability, *J. Climate*, 17, 427–447, 2004.
- Bromwich, D. H., Monaghan, A. J., Manning, K. W., and Powers, J. G.: Real-time forecasting for the Antarctic: an evaluation of the Antarctic mesoscale prediction system (AMPS), *Mon. Weather Rev.*, 133, 579–603, 2005.
- Bromwich, D. H., Otieno, F. O., Hines, K. M., Manning, K. W., and Shilo, E.: Comprehensive evaluation of polar weather research and forecasting performance in the Antarctic, *J. Geophys. Res.*, 118, 274–292, doi:10.1029/2012JD018139, 2013.
- Church, J. A., Gregory, J. M., Huybrechts, P., et al.: Sea level change, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 881 pp., 2013.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus XVI*, 4, 436–468, 1964.
- Dittmann, A., Schlosser, E., Fujita, K., Manning, K. W., Powers, J. G., Duda, M. G., Masson-Delmotte, V., and Werner, M.: Precipitation regime and stable isotopes at Dome Fuji, Antarctica, in preparation, 2015.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Enomoto, H., Motoyama, H., Shiraiwa, T., Saito, T., Kameda, T., Furukawa, T., Takahashi, S., Kodama, Y., and Watanabe, O.: Winter warming over Dome Fuji, East Antarctica and semi-annual oscillation in the atmospheric circulation, *J. Geophys. Res.*, 103, 23103–23111, 1998.

5 EPICA community members: 8 Glacial cycles from an Antarctic ice core, *Nature*, 429, 623–628, 2004.

Fogt, R. L.: State of the climate in 2009, 6. Antarctica, *B. Am. Meteorol. Soc. (special supplement)*, 91, 125–134, 2010.

10 Fogt, R. L.: State of the climate in 2010, 6. Antarctica, *B. Am. Meteorol. Soc. (special supplement)*, 92, 161–172, 2011.

Frezzotti, M., Pourchet, M., Onelio, F., Gandolfi, S., Gay, M., Urbini, S., Vincent, C., Becagli, S., Gragnani, R., Proposito, M., Severi, M., Traversi, R., Udisti, R., and Fily, M.: Spatial and temporal variability of snow accumulation in East Antarctica from traverse data, *J. Glaciol.*, 51, 113–123, 2005.

15 Frieler, K., Clark, P. U., He, F., Buizert, C., Reese, R., Ligtenberg, S. R. M., Van den Broeke, M. R., Winkelmann, R., and Levermann, A.: Consistent evidence of increasing Antarctic accumulation with warming, *Nature Climate Change*, 5, 348–352, doi:10.1038/NCLIMATE2574, 2015.

20 Fujita, K. and Abe, O.: Stable isotopes in daily precipitation at Dome Fuji, East Antarctica, *Geophys. Res. Lett.*, 33, L18503, doi:10.1029/2006GL026936, 2006.

Godfred-Spenning, C. and Simmonds, I.: An analysis of Antarctic sea-ice and extratropical cyclone associations, *Int. J. Climatol.*, 16, 1315–1332, 1996.

25 Gorodetskaya, I. V., Van Lipzig, N. P. M., Van den Broeke, M. R., Mangold, A., Boot, W., and Reijmer, C. H.: Meteorological regimes and accumulation patterns at Utsteinen, Dronning Maud Land, East Antarctica: analysis of two contrasting years, *J. Geophys. Res.*, 118, 1–16, doi:10.1002/jgrd.50177, 2013.

Hirasawa, N., Nakamura, H., and Yamanouchi, T.: Abrupt changes in meteorological conditions observed at an inland Antarctic station in association with wintertime blocking, *Geophys. Res. Lett.*, 27, 1911–1914, 2000.

30 Hirasawa, N., Nakamura, H., Motoyama, H., Hayashi, M., and Yamanouchi, T.: The role of synoptic-scale features and advection in a prolonged warming and generation of different forms of precipitation at dome Fuji station, Antarctica, following a prominent blocking event, *J. Geophys. Res.*, 118, 6916–6928, doi:10.1002/jgrd.50532, 2013.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jouzel, J., Vimeux, F., Caillon, N., Delaygue, G., Hoffmann, G., Masson-Delmotte, V., and Parrenin, F.: Magnitude of isotope/temperature scaling for interpretation of central Antarctic ice cores, *J. Geophys. Res.*, 108, 4361, doi:10.1029/2002JD002677, 2003.

Karoly, D. J.: Southern Hemisphere circulation features associated with El Nino-Southern Oscillation events, *J. Climate*, 2, 1239–1251, 1989.

Lenaerts, J. T. M., Van Meijgaard, E., Van den Broeke, M. R., Ligtenberg, S. R. M., Horwarth, M., and Isaksson, E.: Recent snowfall anomalies in Dronning Maud Land, East Antarctica, in a historical and future climate perspective, *Geophys. Res. Lett.*, 40, 2684–2688, doi:10.1002/grl.50559, 2013.

Lorius, C., Merlivat, L., Jouzel, J., and Pourchet, M.: A 30 000 years isotope climatic record from Antarctic ice, *Nature*, 280, 644–647, 1979.

Marshall, G. J.: Trends in the Southern Annular Mode from observations and reanalyses, *J. Climatol.*, 16, 4134–4143, 2003.

Marshall, G. J.: Half-century seasonal relationship between the Southern Annular Mode and Antarctic temperatures, *Int. J. Climatol.*, 27, 373–383, 2007.

Massom, R., Pook, M. J., Comiso, J. C., Adams, N., Turner, J., Lachlan-Cope, T., and Gibson, T.: Precipitation over the interior East Antarctic ice sheet related to midlatitude blocking-high activity, *J. Climate*, 17, 1914–1928, 2004.

Monaghan, A. J., Bromwich, D. H., Powers, J. G., and Manning, K. W.: The climate of McMurdo, Antarctica, region as represented by one year of forecasts from the Antarctic mesoscale prediction system, *J. Climate*, 18, 1174–1189, 2005.

Monaghan, A. J., Bromwich, D. H., Fogt, R. L., Wang, S., Mayewski, P. A., Dixon, D. A., Ekaykin, A., Frezzotti, M., Goodwin, I., Isaksson, E., Kaspari, S. D., Morgan, V. I., Oeter, H., Van Ommen, T. D., Van der Veen, C. J., and Wen, J.: Insignificant change in Antarctic snowfall since the international geophysical year, *Science*, 313, 827–831, doi:10.1126/science.1128243, 2006.

Nicolas, J. P. and Bromwich, D. H.: Climate of West Antarctica and influence of marine air intrusions, *J. Climate*, 24, 49–67, doi:10.1175/2010JCLI3522.1, 2011.

Nigro, M. A., Cassano, J. J., and Seefeldt, M. W.: A weather pattern-based approach to evaluate the Antarctic mesoscale prediction system (AMPS) forecasts: comparison to automatic weather station observations, *Weather Forecast.*, 26, 184–198, doi:10.1175/2010WAF2222444.1, 2011.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Nigro, M. A., Cassano, J. J., and Knuth, S. L.: Evaluation of Antarctic mesoscale prediction system (AMPS) cyclone forecasts using infrared satellite imagery, *Antarct. Sci.*, 24, 183–192, doi:10.1017/S0954102011000745, 2012.
- Noone, D., Turner, J., and Mulvaney, R.: Atmospheric signals and characteristics of accumulation in Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 104, 19191–19211, 1999.
- Nygard, T., Valkonen, T., and Vihma, T.: Antarctic low-tropopause humidity inversions: 10 yr climatology, *J. Climate*, 26, 5205–5219, doi:10.1175/JCLI-D-12-00446.1, 2013.
- Powers, J. G.: Numerical prediction of an Antarctic severe wind event with the weather research and forecasting (WRF) model, *Mon. Weather Rev.*, 135, 3134–3157, 2007.
- Powers, J. G., Monaghan, A. J., Cayette, A. M., Bromwich, D. H., Kuo, Y., and Manning, K. W.: Real-time mesoscale modeling over Antarctica. The Antarctic mesoscale prediction system, *B. Am. Meteorol. Soc.*, 84, 1522–1545, 2003.
- Powers, J. G., Manning, K. W., Bromwich, D. H., Cassano, J. J., and Cayette, A. M.: A decade of Antarctic science support through AMPS, *B. Am. Meteorol. Soc.*, 93, 1699–1712, 2012.
- Raphael, M. N.: A zonal wave 3 index for the Southern hemisphere, *Geophys. Res. Lett.*, 31, L23212, doi:10.1029/2004GL020365, 2004.
- Reijmer, C. H. and Van den Broeke, M. R.: Temporal and spatial variability of the surface mass balance in Dronning Maud Land, Antarctica, *J. Glaciol.*, 49, 512–520, 2003.
- Reijmer, C. H., Van den Broeke, M. R., and Scheele, M. P.: Air parcel trajectories and snowfall related to five deep drilling locations in Antarctica based on the ERA-15 dataset, *J. Climate*, 15, 1957–1968, 2002.
- Ritter, F., Steen-Larsen, H. C., Kipfstuhl, J., Orsi, A., Behrens, M., and Masson-Delmotte, V.: First continuous measurements of water vapor isotopes on the Antarctic Plateau, *Geophys. Res. Abstr.*, 16, EGU2014-9721, 2014.
- Sarchilli, C., Frezzotti, M., and Ruti, P. M.: Snow precipitation at four ice core sites in East Antarctica: provenance, seasonality and blocking factors, *Clim. Dynam.*, 37, 2107–2125, doi:10.1007/s00382-010-0946-4, 2010.
- Schlosser, E.: Effects of seasonal variability of accumulation on yearly mean  $\delta^{18}\text{O}$  values in Antarctic snow, *J. Glaciol.*, 45, 463–468, 1999.
- Schlosser, E., Duda, M. G., Powers, J. G., and Manning, K. W.: The precipitation regime of Dronning Maud Land, Antarctica, derived from AMPS (Antarctic mesoscale prediction system) Archive Data, *J. Geophys. Res.*, 113, D24108, doi:10.1029/2008JD009968, 2008.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schlosser, E., Manning, K. W., Powers, J. G., Duda, M. G., Birnbaum, G., and Fujita, K.: Characteristics of high-precipitation events in Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 115, D14107, doi:10.1029/2009JD013410, 2010a.

Schlosser, E., Powers, J. G., Duda, M. G., Manning, K. W., Reijmer, C. H., and Van den Broeke, M.: An extreme precipitation event in Dronning Maud Land, Antarctica – a case study using AMPS (Antarctic mesoscale prediction system) archive data, *Polar Res.*, 29, 330–344, doi:10.1111/j.1751-8369.2010.00164.x, 2010b.

Schlosser, E., Manning, K. W., Powers, J. G., Gillmeier, S., and Duda, M. G.: An extreme precipitation/warming event in Antarctica – a study with Polar WRF, in preparation, 2015.

Seefeldt, M. W. and Cassano, J. J.: An analysis of low-level jets in the greater Ross Ice Shelf region based on numerical simulations, *Mon. Weather Rev.*, 136, 4188–4205, doi:10.1175/2007JAMC1442.1, 2008.

Seefeldt, M. W. and Cassano, J. J.: A description of the Ross ice shelf air stream (RAS) through the use of self-organizing maps (SOMs), *J. Geophys. Res.*, 117, D09112, doi:10.1029/2011JD016857, 2012.

Simmonds, I., Keay, K., and Lim, E.: Synoptic activity in the seas around Antarctica, *Mon. Weather Rev.*, 131, 272–288, 2002.

Sinclair, M. R.: Record-high temperatures in the Antarctic – a synoptic case study, *Mon. Weather Rev.*, 109, 2234–2242, 1981.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, NCAR/TN 475+STR, Nat. Cent. for Atmos. Res., Boulder, Co, USA, 125 pp., 2008.

Steen-Larsen, H. C., Johnsen, S. J., Masson-Delmotte, V., Stenni, B., Risi, C., Sodemann, H., Balslev-Clausen, D., Blunier, T., Dahl-Jensen, D., Ellehøj, M. D., Falourd, S., Grindsted, A., Gkinis, V., Jouzel, J., Popp, T., Sheldon, S., Simonsen, S. B., Sjolte, J., Steffensen, J. P., Sperlich, P., Sveinbjörnsdóttir, A. E., Vinther, B. M., and White, J. W. C.: Continuous monitoring of summer surface water vapor isotopic composition above the Greenland Ice Sheet, *Atmos. Chem. Phys.*, 13, 4815–4828, doi:10.5194/acp-13-4815-2013, 2013.

Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E., Roethlisberger, R., and Selmo, E.: An oceanic cold reversal during the last deglaciation, *Science*, 293, 2074–2077, 2001.

Stenni, B., Masson-Delmotte, V., Selmo, E., Oerter, H., Meyer, H., Roethlisberger, R., Jouzel, J., Cattani, O., Falourd, S., Fischer, H., Hoffmann, G., Iacumin, P., Johnsen, S. F., Minster, B.,

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Udisti, R.: The deuterium excess records of EPICA Dome C and Dronning Maud Land ice cores (East Antarctica), *Quat. Scie. Rev.*, 29, 146–159, 2010.

Stenni, B., Bonazza, M., Cagnati, A., Cattani, F., Dreossi, G., Frezzotti, M., Frosini, D., Grigioni, P., Karlicek, D., Masson-Delmotte, V., Scarchilli, C., Risi, C., Schlosser, E., Udisti, R., and Valt, M.: Three year monitoring of stable isotopes of precipitation at Dome Concordia, Antarctica, in preparation, 2015.

Stoelinga, M. T.: A users guide to RIP Version 4.5: a program for visualizing mesoscale model output, NCAR online document, University of Washington, available at: [http://www2.mmm.ucar.edu/mm5/documents/ripug\\_V4.html](http://www2.mmm.ucar.edu/mm5/documents/ripug_V4.html) (last access: 3 November 2015), 2009.

Suzuki, K., Yamanouchi, T., and Motoyama, H.: Moisture transport to Syowa and Dome Fuji stations in Antarctica, *J. Geophys. Res.*, 113, D24114, doi:10.1029/2008JD009794, 2008.

Thompson, D. W. J. and Wallace, J. M.: Annular modes in the extratropical circulation. Part 1: Month-to-month variability, *J. Climate*, 13, 1000–1016, 2000.

Trenberth, K. E. and Mo, K. C.: Blocking in the Southern Hemisphere, *Mon. Weather Rev.*, 133, 38–53, 1985.

Van Loon, H.: The half-yearly oscillation in middle and high southern latitudes and the coreless winter, *J. Atmos. Sci.*, 24, 472–486, 1967.

Van Loon, H. and Jenne, R. L.: The zonal harmonic standing waves in the Southern Hemisphere, *J. Geophys. Res.*, 77, 992–1003, 1972.

Winkelmann, R., Levermann, A., Martin, M. A., and Frieler, K.: Increased future ice discharge from Antarctica owing to higher snowfall, *Nature*, 492, 239–242, 2012.

**Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica**

E. Schlosser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Figure 1.** Map of Antarctica indicating Dome C and other important deep-drilling sites in Antarctica.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.




**Figure 2.** AMPS domains used for model output analysis in this study.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.




**Figure 3.** (a) Mean monthly temperatures for 2009 and 2010 at Dome C AWS. Daily precipitation and daily mean temperature at Dome C for (b) 2009 and (c) 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica


E. Schlosser et al.



**Figure 4.** Monthly precipitation at Dome C **(a)** 2009 and **(b)** 2010, distinguishing three different types of precipitation: diamond dust, hoarfrost, and snowfall. Relative frequency of diamond dust, hoarfrost, and snowfall for **(c)** 2009 and **(d)** 2010. The types were determined from photos of the crystals on the platforms by the Avalanche Research Institute, Arabba, Italy.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica


E. Schlosser et al.



**Figure 5.** (a) 500 hPa geopotential height from AMPS archive data (Domain 1) 13 September 2009 00Z (The axis of the upper-level ridge mentioned in the text is marked by a bold black line.), (b) 24 h precipitation from AMPS 13 September 2009 00:00 to 24:00 GMT, (c) 5 day back-trajectories for parcels arriving at Dome C at 00:00 UTC 12 September 2009. Trajectories for three arrival levels are shown: (1) 600 hPa, (2) 500 hPa, (3) 300 hPa.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica


E. Schlosser et al.



**Figure 6.** Example for synoptic situation, during which precipitation is observed at Dome C, but not forecast by WRF in AMPS. **(a)** 500 hPa geopotential height, Domain 2. **(b)** 24 h precipitation total (mm) from AMPS.

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.




**Figure 7.** Mean July – 500 hPa geopotential height based on AMPS archive model output for 2009 and 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica**

E. Schlosser et al.

**Figure 8.** Mean monthly SAM index for 2009 and 2010 (after Marshal, 2003).[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.



**Figure 9.** (a) Monthly mean Zonal Wave Number 3 (ZW3) index for 2009–2010, (b) July 2009 500 hPa geopotential height anomaly: mean July 2009 height minus long-term zonal mean height, (c) July 2010 500 hPa geopotential height anomaly: mean July 2009 height minus long-term zonal mean height.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Precipitation regime and stable oxygen isotopes at Dome C, East Antarctica

E. Schlosser et al.



**Figure 10.** Daily mean air temperatures at Dome C 2009 and 2010 from AWS and stable isotopes ( $\delta^{18}\text{O}$  and deuterium excess) of corresponding precipitation samples.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

