



Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz¹, S. R. Proud^{1,2}, S. S. Seppäläinen¹, and D. J. Cziczo¹

¹Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

²University of Copenhagen, Copenhagen, Denmark

Received: 2 January 2015 – Accepted: 12 February 2015 – Published: 5 March 2015

Correspondence to: D. J. Cziczo (djciczo@mit.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Atmospheric aerosol particles are often partially or completely composed of inorganic salts, such as ammonium sulfate and sodium chloride, and therefore exhibit hygroscopic properties. Many inorganic salts have well-defined deliquescence and efflorescence points at which they take up and lose water, respectively. Deliquescence and efflorescence of simple inorganic salt particles have been investigated by a variety of methods, such as IR spectroscopy, tandem mobility analysis and electrodynamic balance. Field measurements have shown that atmospheric aerosols are not typically pure inorganic salt, instead they often also contain organic species. There is ample evidence from laboratory studies that suggests that mixed particles exist in a phase-separated state, with an aqueous inorganic core and organic shell. Although phase separation has not been measured in situ, there is no reason it would not also take place in the atmosphere. Many recent studies have focused on microscopy techniques that require deposition of the aerosol on a glass slide, possibly changing its surface properties. Here, we investigate the deliquescence and efflorescence points, phase separation and ability to exchange gas-phase components of mixed organic and inorganic aerosol using a flow tube coupled with FTIR spectroscopy. Ammonium sulfate aerosol mixed with organic polyols with different O : C ratios, including 1,4-butanediol, glycerol, 1,2,6-hexanetriol, 1,2-hexanediol, and 1,5-pentanediol have been investigated. Those constituents correspond to materials found in the atmosphere in great abundance, and therefore, particles prepared in this study should mimic atmospheric mixed phase aerosol particles. The results of this study tend to be in agreement with previous microscopy experiments, with several key differences, which possibly reveal a size-dependent effect on phase separation in organic/inorganic aerosol particles.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Organic-containing tropospheric aerosol particles can exist as solids or liquids with a range of viscosities, depending on temperature and relative humidity conditions. The chemical properties of those states are often radically different. For example, reactions of nitric acid with sea salt aerosols (SSA) are overwhelmingly faster in the aqueous phase (Liu et al., 2008; Tolocka et al., 2004) and the oxidation of N_2O_5 proceeds readily on surfaces of liquid particles, but not solid ones (Fried et al., 1994; Hu and Abbatt, 1997; Hallquist et al., 2003; Grassian, 2001). A particle's phase can also influence its radiative properties; it has been shown that liquid particles scatter radiation more than their solid counterparts with otherwise similar chemical composition (Martin, 2000). Additionally, there is evidence that organic aerosols can be efficient CCN, but also that the nature of the organic can alter the CCN properties (Clegg et al., 2001; Cruz and Pandis, 1998, 2000). Glassy organic coatings can even change ice nucleation properties of common atmospheric IN particles such as mineral dust (Schill et al., 2014).

This work uses bulk spectroscopic methods to investigate phase transitions of ammonium sulfate as a result of water uptake from the gas phase and the impact of organic constituents. Water uptake from the gaseous phase by aerosols whose organic and inorganic constituents are phase-separated is also determined. The phase transitions of pure ammonium sulfate have been well-characterized, but the particles in the atmosphere are rarely comprised of pure constituents. At mid-latitudes, organic material contributes as much as 20–50 % of the total fine aerosol mass (Parsons et al., 2006; Cruz and Pandis, 1998; Saxena et al., 1995; Zhang et al., 2007; Murphy et al., 1998). A study of several European urban and rural field sites indicated that the water-soluble fraction of fine aerosol was 65–70 %, 20–50 % of the total soluble aerosol material were organics and 70 % of the total organics present in the aerosol were also polar (Zappoli et al., 2007). Russell et al. (2002) used X-ray spectromicroscopy to analyze organic

ACPD

15, 6537–6566, 2015

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

coatings on SSA particles revealing a diversity of functional groups and a high degree of complexity of mixtures in those systems.

To investigate the phase changes of sub-micrometer diameter aerosol particles, Fourier transform IR (FTIR) spectroscopy in the mid-infra red (MIR) range was used because it can distinguish and quantify gas and condensed-phase water (Cziczko et al., 1997). The measurements were performed on a flowing aerosol sample exposed to variable humidity air. In this manner solid aerosols were observed to become droplets as they took up water from the gas phase (deliquescence) or liquid particles became solid as they lost water (efflorescence). Other measurement techniques that have been used to investigate aerosol phase changes include single particle levitation coupled with spectroscopy (Raman, Mie resonance, micro-FTIR) as well as different types of microscopy on a substrate (Liu et al., 2008; Zhao et al., 2008; Parsons et al., 2006; Bertram et al., 2011). Many microscopy techniques require depositing particles on a hydrophobic slide, which represents a possible surface for heterogeneous phase transition. Levitation techniques, such as electrodynamic balance, acoustic suspension and light pressure suspension coupled with spectroscopy are well suited for studying condensation and freezing events on single particles but not the properties of a multi-particle flow.

In the case of inorganic aerosol, droplets form when crystalline aerosol is exposed to humidified air. For pure salts, such as sodium chloride or ammonium sulfate, the water uptake starts rapidly within a very narrow range of RH. This corresponds to the deliquescence RH (DRH). After deliquescence, the particle enters the hygroscopic growth regime. When an aqueous particle is exposed to decreasing RH its water content decreases but it often does not return to zero at the DRH; instead it experiences hysteresis in the loss of condensed phase water. The particle exists in a supersaturated aqueous state until the efflorescence RH (ERH) at which point it rapidly crystallizes to an anhydrous solid. DRH and ERH for pure ammonium sulfate have been characterized extensively at room temperature to be, respectively, 85 and 30–35 %, the latter with a dependence on particle volume (Liu et al., 2008; Cziczko and Abbatt, 2000; Cziczko

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 1997). DRH and ERH do not change significantly with a decrease in temperature (Cziczo and Abbatt, 2000; Koop et al., 2000; Braban et al., 2001).

The ERH and DRH of $(\text{NH}_4)_2\text{SO}_4$ /organic/water mixtures are more complicated than that of solutions devoid of organics. This study focuses on phase transitions of such ternary systems. The organic compounds used here are diols and triols, such as glycerol, 1,4-butanediol and 1,2,6-hexanetriol. Organics were found to either suppress, enhance or have no effect on ERH and DRH of inorganic aerosol depending on the type of organic and its concentration relative to the concentration of the inorganic salt, $(\text{NH}_4)_2\text{SO}_4$ in this case, consistent with the data available in the literature (Bertram et al., 2011; Smith et al., 2011; Parsons et al., 2006; Cruz and Pandis, 1998; Robinson et al., 2014). The process is usually characterized chemically in terms of the salting-out effect, described by the Setchenov equation (Bertram et al., 2011; Desnoyers and Ichhaporla, 1969). Organics whose O : C ratio is bigger than ~ 0.7 , such as glycerol, were found by Bertram et al. (2011), in a study using optical microscopy, to lower the ERH and DRH of ammonium sulfate and suppress it completely at high concentrations. Efflorescence, considered a kinetic process where the free energy barrier for crystal formation must be overcome, is often described by homogeneous nucleation theory (Parsons et al., 2006). Crystalline ammonium sulfate has been shown to be a poor nucleus for subsequent heterogeneous nucleation of organics, such as malonic acid, which may serve to explain why certain organics suppress ERH of ammonium sulfate (Ciobanu et al., 2009; Parsons et al., 2006; Smith et al., 2011; Braban and Abbatt, 2004). Bertram et al. (2011) suggested that the Gibbs–Duhem relation, which implies that an increase of organic to sulfate ratio increases the chemical potential of organic but decreases the chemical potential of ammonium sulfate, explains the decrease in DRH and ERH values with the increase of organic concentration for certain organics. Thus, as the concentration of organic increases, DRH decreases to maintain unity solution saturation with respect to ammonium sulfate and ERH decreases to maintain critical supersaturation (Bertram et al., 2011).

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

ERH and DRH depression by certain organics, but not others, can be explained by liquid–liquid phase separation of the aerosol (Bertram et al., 2011). Phase separation occurs when the organic fraction of the aerosol does not mix with the aqueous inorganic fraction. This results in a particle whose core is aqueous inorganic and outer shell is organic. Therefore, ERH for the phase-separated core of such a particle would be close to the ERH of pure ammonium sulfate (Bertram et al., 2011; Buajarern et al., 2007). For systems in which phase separation does not occur, organics disrupt the nucleation resulting in ERH/DRH depression as described above (Bertram et al., 2011; Ciobanu et al., 2009; Smith et al., 2011). According to previous studies, particles whose organic fraction has O : C ratio greater than ~ 0.7 do not undergo phase separation but those that contain organics with O : C ratio less than ~ 0.7 do (Bertram et al., 2011; Song et al., 2012). In this study, $(\text{NH}_4)_2\text{SO}_4$ /glycerol system is the only system expected not to undergo phase separation according to the previous results.

Ciobanu et al. (2009) studied the liquid–liquid phase separation processes in millimeter-sized droplets (12–67 μm dry diameter) composed of PEG-400 and ammonium sulfate using Raman microscopy. Based on this work, there are two distinct mechanisms of phase separation, one based on classical nucleation theory and the other based on dispersed cluster growth and coalescence (Ciobanu et al., 2009). Liquid–liquid phase separation was observed for particles of all sizes investigated, but the mechanisms varied depending on particle composition (Ciobanu et al., 2009). A computer model of liquid–liquid phase separation in binary, ternary and multicomponent mixtures has been developed by Zuend et al. (2010) based on those results.

The evidence for liquid–liquid phase separation comes almost solely from microscopy and EDB studies (Bertram et al., 2011; You et al., 2013; Ciobanu et al., 2009; Choi and Chan, 2002; Marcolli and Krieger, 2006). As a limit of the resolution the particles studied are large, on the order of 1 μm diameter or greater (Bertram et al., 2011; Ciobanu et al., 2009; Marcolli and Krieger, 2006). None of those studies have reported a dependence of phase separation of particle size, but there is now evidence from cryo-transmission electron microscopy of smaller particles that liquid–liquid phase

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

separation is highly dependent on particle size in sub- μm diameter particles (Veghte et al., 2013). Veghte et al. (2013) found that for $(\text{NH}_4)_2\text{SO}_4$ /succinic acid system, no phase separation occurred below 170 nm and for $(\text{NH}_4)_2\text{SO}_4$ /pimelic acid system, no phase separation occurred below 270 nm. The technique used in this study, FTIR coupled with a flow tube set-up, can access sizes smaller than 1 μm diameter, and therefore can be used to study liquid–liquid phase separation in small particles relevant to the atmosphere. To study the effects of phase separation on the uptake of water vapor, the particles are exposed to a flow of D_2O vapor after given time to fully phase separate to investigate whether the organic layer formed on the particle surface inhibits the transfer of water vapor to the aqueous core.

2 Experimental

Polydisperse droplets were generated with an atomizer. An aqueous solution of Milli-Q water (18.2 $\text{M}\Omega\text{cm}$) and 10 weight % (wt%) reagent grade ammonium sulfate (99 %, Sigma Aldrich, St. Louis, MO) was used to produce inorganic aerosols. For organics, glycerol, 1,4-butanediol, 1,2,6-hexanetriol, 1,2-hexanediol, and 1,5-pentanediol were used (98 %, reagent grade, Sigma Aldrich). All pure organics were produced from 10 wt% solutions but the ratio of ammonium sulfate and organics in the ternary solutions was variable. A custom constant output atomizer was used which had an orifice diameter of 0.030 cm. This atomizer was designed to produce a 0.1 slpm flow, which was then diluted with 0–0.9 slpm dry nitrogen. The custom atomizer aerosol output was centered at ~ 120 nm, determined with a BMI, Inc. (Hayward, CA) Differential Mobility Analyzer (DMA).

For deliquescence experiments a dried aerosol flow was then mixed with a humidified nitrogen flow, which was passed through a bubbler (1.0–1.5 slpm; for conditions up to 80 % RH) or a Nafion tube surrounded by water (for conditions to ~ 100 %). The flow was then passed into a 9 L glass volume, which allowed for a ~ 1 min interaction time between vapor and particles, and to buffer any particle production variability before

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

moving into the flow tube. The RH was controlled by varying the ratio of the aerosol flow, which was diluted with dry gas and passed through a dryer, and the humidified flow through the water bubbler or Nafion tube. In some experiments, described in detail later, the aerosol flow was also mixed with a nitrogen flow passed through a D₂O bubbler.

For measurement of efflorescence the dryer was bypassed so that the aerosol flow remained at ~ 100 % RH. A dry flow of gas was then added to the mixing volume to lower the RH with higher flows corresponding to increasingly lower RH.

The RH was measured in two ways: with an E+E Elektronik EE08 (Engerwitzdorf, Austria) humidity/temperature sensor located at the inlet of the flow tube (Fig. 1) and by quantifying the amount of gas-phase water present in the FTIR spectra. The humidity sensor, with ±1 % quoted error, was verified using saturated solutions of salts. RH was directly determined from the spectra by integrating gas-phase water lines from 1874 to 1855 cm⁻¹. The total water vapor content obtained by integration was calibrated using the RH sensor for flows of varying RH. The value obtained by direct integration of the FTIR spectrum is used as the *x* axis in the figures in this paper.

FTIR has been a valuable tool in studying model atmospheric aerosol (Cziczo et al., 1997; Cziczo and Abbatt, 1999, 2000; Braban et al., 2003; Braban and Abbatt, 2004; Earle et al., 2010). A flow tube system coupled to an FTIR spectrometer, shown schematically in Fig. 1, was used in these studies. The flow tube used for the majority of experiments was glass, 80 cm in length and 49 mm in inner diameter. Flow tube windows were 3 mm thick, uncoated polished ZnSe crystals. ZnSe has a nearly flat transmittance over the range of primary interest (500 to 5000 cm⁻¹) and is both chemically inert and non-hygroscopic. The windows were joined to the glass by EpoTek (Billerica, MA) 353ND high-performance epoxy. Windows were mounted at a slight angle, ~ 5°, to prevent interference within the crystal. A Bruker (Billerica, MA) Tensor 37 FTIR spectrometer with MIR beamsplitter coupled to a liquid-nitrogen cooled external MCT/A detector was used to collect aerosol spectra in the range 500 to 8000 cm⁻¹ at 4 cm⁻¹ resolution. The MCT detector was enclosed in an acrylic box through which

a flow of dry nitrogen was maintained at ~ 1 slpm throughout the experiment in order to minimize interference of ambient water vapor in the spectra.

FTIR spectra acquired in this experiment include absorptions from water vapor, condensed water, ammonium sulfate and alcohol (Fig. 2, Table 1). Scattering from aerosol particles is indicated by the baseline rise toward higher wavenumber. Narrow water vapor lines are centered at 3750 and 1500 cm^{-1} . Broad condensed water bands are present at 3450, 1640, and 650 cm^{-1} , corresponding to OH stretch, HOH bend, and H-bonding, respectively. In Fig. 2, the HOH bend feature is not apparent because it coincides with the water vapor lines. The ammonium sulfate features are apparent at $2800\text{--}3300\text{ cm}^{-1}$ due to N-H stretch mode, $1420\text{--}1450\text{ cm}^{-1}$ due to NH_4^+ deformation mode and at 1115 and 620 cm^{-1} both due to sulfate. The NH_4^+ deformation mode is indistinguishable in the presence of the water vapor lines in Fig. 2. In Fig. 3 the gas water spectrum has been subtracted, and the NH_4^+ deformation mode at $1420\text{--}1450\text{ cm}^{-1}$, as well as the HOH bend at 1640 cm^{-1} (in some spectra) are apparent. The amount of liquid-phase water was quantified by integrating the band at 1640 cm^{-1} after gas-phase water subtraction. Figure 3 also indicates the FTIR spectral response during an effluorescence experiment as aerosol particles uptake water: the top panel is anhydrous ammonium sulfate at 30 % RH and condensed-phase water features are absent. Evidence of liquid water is apparent at 60 % as the condensed water spectral features. Upon dissolution, ammonium sulfate absorption bands change slightly. The 1115 cm^{-1} sulfate band broadens and shifts towards lower wavenumbers, and the $1420\text{--}1450\text{ cm}^{-1}$ band broadens and shifts towards higher wavenumbers. See Cziczko et al. (1997) for details.

Note that the polyols used in these experiments, unlike carboxylic acids, do not have visible IR absorption features. The C-H and O-H stretches are obscured by water features. The middle panel at Fig. 3 shows the same sequence of spectra for $(\text{NH}_4)_2\text{SO}_4/1,2,6\text{-hexanetriol}$ mixture, and the bottom panel shows spectra for $(\text{NH}_4)_2\text{SO}_4/\text{glycerol}$ mixture. The $2800\text{--}3300\text{ cm}^{-1}$ N-H stretch feature exhibits the presence of an alcohol. The HOH bend remains an indication of the presence of condensed-phase water in all spectra.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Phase separation is not directly observed with FTIR spectroscopy. However, gas-phase exchange of water across the organic boundary can be investigated by using D_2O vapor in place of water. Figure 4 is a set of spectra produced when different aerosol is exposed to D_2O . The spectra are from aerosol from an aqueous $(NH_4)_2SO_4/1,2,6$ -hexanetriol solution in 1 : 1 ratio by mass, exposed to D_2O vapor at increasing RH (from bottom to top). Gas-phase lines at 2600 – 2800 cm^{-1} correspond to D_2O and DOH. The broad absorption at 2500 cm^{-1} corresponds to condensed-phase DOH that forms when vapor phase D_2O exchanges with liquid water in the aerosol droplets. Condensed-phase DOH feature grow in response to increasing concentration of D_2O vapor and accessible condensed-phase H_2O . This growth was measured in a manner similar to the HOH bend proxy for condensed-phase H_2O ; integration of the absorption at 2500 cm^{-1} . The amount of D_2O/DOH vapor can be qualified in a manner similar to the RH calibration by integrating a gas-phase line; for this work 2870 – 2880 cm^{-1} .

3 Results and discussion

3.1 ERH and DRH properties of ternary solutions

Initial experiments were conducted on the binary $(NH_4)_2SO_4$ /water system to determine DRH and ERH for comparison to the literature. Experimental data of the condensed phase water peak are plotted in the upper left panel of Fig. 5. The plotted error in the condensed water band area was quantified by averaging the deviation and noise in the 0–30 % RH range, where no liquid water exists, after subtraction of water vapor features, and is shown for clarity in only the left-most data point. For pure ammonium sulfate, efflorescence was observed at $\sim 35\%$ RH and deliquescence at $\sim 80\%$, consistent with the literature (Cziczko et al., 1997).

Experiments were subsequently conducted on the more complex ternary solutions. The results of the efflorescence and deliquescence experiments on

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(NH₄)₂SO₄/1,4-butanediol mixtures are shown in the other panels of Fig. 5. Using quantification of the HOH bend area, the deliquescence point was ~ 80 %. There is evidence of liquid water, a relatively small HOH feature, below the ERH of the binary (NH₄)₂SO₄/water solution and the presence of the condensed phase is supported by D₂O experiments, discussed in a subsequent section. According to Bertram et al. (2011), there is evidence of phase separation in these solutions. Our results indicate some inhibition of ERH due to the presence of organics.

Experiments on (NH₄)₂SO₄/glycerol mixtures are detailed in Fig. 6. The presence of glycerol affects the DRH and ERH of the mixture. In the 1 : 3 glycerol/(NH₄)₂SO₄ solution the ERH is ~ 30 % and the DRH is ~ 40 %. In the 1 : 2 glycerol/(NH₄)₂SO₄ and 1 : 1 glycerol/(NH₄)₂SO₄ solutions there is no evidence of efflorescence and, consequently, deliquescence cannot be observed. This behavior suggests no phase separation has occurred (Bertram et al., 2011).

Figure 7 contains spectra from experiments on the (NH₄)₂SO₄/1,2,6-hexanetriol system. Mixtures of this system again show evidence of suppression of ERH and DRH. In the 1 : 3 1,2,6-hexanetriol/(NH₄)₂SO₄ mixture, the ERH appears unaffected by the organic but the deliquescence curve shows evidence of liquid water below 80 % RH, until about 60 % RH. In the 1 : 2 and 1 : 1 1,2,6-hexanetriol/(NH₄)₂SO₄ mixtures, organics appear to completely suppress efflorescence (e.g., panel d). These data suggest no occurrence of phase separation in 1,2,6-hexanetriol/(NH₄)₂SO₄ mixture.

The set of experiments conducted here are listed and compared to the literature in Table 2. The results of the efflorescence and deliquescence experiments are generally consistent with previous work that shows suppression of ERH in ternary organic/(NH₄)₂SO₄/water solutions with the extent of this effect dependent on specific organic (Bertram et al., 2011; You et al., 2013; Marcolli and Krieger, 2006). If complete suppression of ERH and DRH is taken to be evidence of no phase separation then the (NH₄)₂SO₄/glycerol and (NH₄)₂SO₄/1,2,6-hexanetriol systems showed no phase separation. For the (NH₄)₂SO₄/glycerol system, this is consistent with the literature, as glycerol has been shown not to undergo phase separation when mixed with inorganics

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Bertram et al., 2011; You et al., 2013). For the $(\text{NH}_4)_2\text{SO}_4/1,2,6$ -hexanetriol system, the data is inconsistent with the literature, as this system was shown to undergo phase separation at 76.7 % RH (You et al., 2013). Our data are consistent with phase separation in the $(\text{NH}_4)_2\text{SO}_4/1,4$ -butanediol system and the observed unaltered ERH and DRH points agree with previous studies (Marcolli and Krieger, 2006).

These data suggest the $(\text{NH}_4)_2\text{SO}_4$ /glycerol and $(\text{NH}_4)_2\text{SO}_4/1,2,6$ -hexanetriol systems exhibit a partial or complete suppression of ERH and DRH. This contradicts the literature for the $(\text{NH}_4)_2\text{SO}_4/1,2,6$ -hexanetriol system which does not report suppression. This result is important because it suggests potential differences between FTIR and microscopy techniques. Using electron microscopy techniques, Veghte et al. (2013) found that, for some ternary systems, sub-200 nm diameter organic/inorganic particles do not undergo phase separation. Aerosols in this size range cannot be observed by optical microscopy techniques used by Bertram et al. (2011), as only particles on the order of ~ 1000 nm and greater can be studied with this technique. FTIR, however, is a bulk technique, which studies the composite aerosol within the sample volume. As described in the Experimental section, the constant output atomizer used in this work produces a polydisperse particle distribution across this size range (i.e., from sub-200 nm to super-micrometer diameter). It is therefore likely that a combination of large phase separated and small non-phase separated particles were present in the flow tube. Indication of inhibition of ERH and DRH could have come from a subset of the overall population, in this case due to the small aerosol, as suggested by Veghte et al. (2013).

3.2 Heterogeneous chemical potential of ternary solutions

Cziczko et al. (1997) showed that D_2O vapor remained in the gas phase when only crystalline aerosol was present, whereas gas- and condensed-phase DOH was rapidly formed in the presence of condensed phase H_2O . For a comparison to the literature, initial experiments were conducted by adding an increasing flow of D_2O vapor to a anhydrous, sub-30 % RH, flow of $(\text{NH}_4)_2\text{SO}_4$ aerosol (i.e., aerosol below its ERH). These

data are plotted as the lower-most curve of Fig. 8 and indicate, consistent with the literature, that DOH is not formed in the presence of crystalline aerosol (Cziczo et al., 1997).

Using this methodology, the possible uptake of water by ternary aerosol was investigated by exposing particles to an increasing flow of D₂O. The formation of condensed-phase DOH in a 1 : 1 (NH₄)₂SO₄/1,2,6-hexanetriol mixture at variable RH is shown in Fig. 4. The area of the condensed phase DOH feature is proportional to the amount of liquid water in the spectrum and, assuming a relatively constant aerosol content, it can be compared between spectra. Assuming ammonium sulfate and 1,2,6-hexanetriol phase separates below ~ 78 % RH (Table 2), separation should have occurred for the 30 and 60 % RH points (the blue and green traces in Fig. 4). The data show evidence that DOH forms even in a system that should have phase separated.

Figure 8 is a composite plot of gas-phase exchange experiments, including the spectra in Fig. 4, on different organic/inorganic mixtures under three RH conditions: 80 % (top), 70 % (middle) and 60 % RH (bottom); this RH range was chosen because the organic/inorganic aerosols undergo phase separation within it (Table 2). Data for anhydrous ammonium sulfate (at 30 % RH) is shown for reference in each figure. Note that a lack of D₂O exchange is also exhibited by the 1 : 1 1,4-butanediol/(NH₄)₂SO₄ mixture below its efflorescence point (at 20 % RH), likely indicative of a phase-separated aerosol where the inorganic core has undergone efflorescence. These two curves, “negative control experiments”, are shown for reference in all three panels.

The other organic/inorganic solutions, none of which exhibited efflorescence, show gas-phase exchange at all RH conditions. The intensity of the exchange signal is largest in the (NH₄)₂SO₄/glycerol system, consistent with a mixture that does not phase separate or effloresce (i.e., liquid water is always expected to be present at the surface). Under the RH conditions studied, (NH₄)₂SO₄/1,4-butanediol, (NH₄)₂SO₄/1,2-hexanediol, and (NH₄)₂SO₄/1,5-pentanediol super-micrometer diameter particles are all expected to phase separate (Bertram et al., 2011). In these three systems sub-micrometer particles have not been studied to date and we can

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

not preclude that they remain in a non-phase separated state per the finding of Veghte et al. (2013). This is a possible indication of non-phase separated particles of small size remaining chemically active. Atmospherically, this is important because the results indicate that sub-micrometer aerosol that does not phase separate remains in a chemically active state with surface water present.

Noteworthy are the $(\text{NH}_4)_2\text{SO}_4/1,2,6\text{-hexanetriol}$ and $(\text{NH}_4)_2\text{SO}_4/1,4\text{-butanediol}$ systems which may phase separate regardless of size (M. Freedman, personal communication, 2014). The data collected here support a lack of phase separation for the former because of a lack of observed efflorescence while the latter may phase separate since it is observed to effloresce. The data in Fig. 8 indicate that if phase separation does proceed below $\sim 78\%$ RH for either system they continue to exchange D_2O with the gas phase (in the case of $(\text{NH}_4)_2\text{SO}_4/1,4\text{-butanediol}$ until efflorescence of the inorganic core). Therefore, if present, an organic shell does not appear to present a barrier for vapor exchange to the core material and these particles remain chemically active.

Shorter residence time experiments were also performed. For these measurements the 9 L mixing volume was removed to reduce the time during which phase separation could occur, before mixing with D_2O , to seconds. There was no evidence of exchange stopping at the shorter residence time. This reinforces (1) rapid phase separation and (2) no limitation of mass accommodation of water on this timescale (A. Bertram, personal communication, 2013).

4 Conclusions

This experiment examined hygroscopic properties of three mixed organic/inorganic aerosols: $(\text{NH}_4)_2\text{SO}_4/\text{glycerol}$, $(\text{NH}_4)_2\text{SO}_4/1,4\text{-butanediol}$ and $(\text{NH}_4)_2\text{SO}_4/1,2,6\text{-hexanetriol}$. Liquid-liquid phase separation is expected to occur in these systems if the O:C ratio of the organic is below ~ 0.7 (Bertram et al., 2011; You et al., 2013; Song et al., 2012). When liquid-liquid separation occurs organics have no effect on hygroscopic properties and efflorescence and deliquescence of

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

the inorganic proceeds as it would in a $(\text{NH}_4)_2\text{SO}_4$ -only solution (You et al., 2013). When liquid–liquid separation does not occur the organic can disrupt nucleation in the inorganic solution and the ERH and DRH can be inhibited (You et al., 2013). Given the O : C ratios used in this experiment (see Table 2), only the $(\text{NH}_4)_2\text{SO}_4$ /glycerol mixture was not expected to phase separate. ERH and DRH inhibition was observed in this system, consistent with the literature. Liquid–liquid phase separation should have occurred in the $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol and $(\text{NH}_4)_2\text{SO}_4$ /1,2,6-hexanetriol mixtures. The hygroscopic behavior of the $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol mixture did not exhibit inhibition of ERH and DRH, consistent with phase separation and the literature. Contrary to theory, the $(\text{NH}_4)_2\text{SO}_4$ /1,2,6-hexanetriol mixture behaved like the $(\text{NH}_4)_2\text{SO}_4$ /glycerol mixture: ERH and DRH were inhibited as would be the case for a non-phase separated aerosol. This supports the observation of Veghte et al. (2013) that sub-micrometer aerosol particles do not phase separate despite the requisite O : C ratio.

The effect of liquid–liquid phase separation on heterogeneous chemistry was also studied. The approach taken in this study was to determine water vapor exchange across the organic barrier formed on the aerosol surface in a phase separated system. After minutes of time in a mixing volume for phase separation to take place, the organic/inorganic aerosols were mixed with D_2O vapor. DOH was produced, an indication of accessibility of the core H_2O , in all systems except those that had no water in them: anhydrous ammonium sulfate and $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol below the ERH of ammonium sulfate. The $(\text{NH}_4)_2\text{SO}_4$ /glycerol system was expected not to phase separate and the exchange observation is consistent with this phase. The $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol, $(\text{NH}_4)_2\text{SO}_4$ /1,2-hexanediol, and $(\text{NH}_4)_2\text{SO}_4$ /1,5-pentanediol systems are expected to phase separate, at least at super-micrometer diameter sizes, and the $(\text{NH}_4)_2\text{SO}_4$ /1,2,6-hexanetriol system is expected to phase separate at all particle diameters. In these cases D_2O exchange is observed in all cases. This indicates that atmospherically-relevant sub-micrometer diameter particles either do not phase separate and/or an organic shell does not inhibit the ex-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

change of vapor species such as water. In either case, organics do not appear capable of inhibiting heterogeneous chemical reactions requiring an aqueous inorganic component.

We suggest a next step in these experiments would be to consider size-selected aerosol. This would allow further consideration of the Veghte et al. (2013) finding that small aerosol particles (< 200 nm) do not experience phase separation. Such experiments would require increased sensitivity to the lower aerosol concentrations required when only a size-selected portion of the population is used.

Acknowledgements. This work was supported by NASA Earth and Space Science Fellowship, NASA grant NNX13AO15G and NOAA grant NA11OAR4310159. The authors would like to thank Allan K. Bertram and Miriam A. Freedman for useful discussions.

References

- Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO₃, O₃ and NO₂ with deliquescent NaCl aerosols at room temperature, *J. Phys. Chem. A*, 102, 3719–3725, 1998.
- Bertram, A. K., Patterson, D. D., and Sloan, J. J.: Mechanisms and temperatures for the freezing of sulfuric acid aerosols measured by FTIR extinction spectroscopy, *J. Phys. Chem.*, 100, 2376–2383, 1996.
- Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A., You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase separation, efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the organic component, *Atmos. Chem. Phys.*, 11, 10995–11006, doi:10.5194/acp-11-10995-2011, 2011.
- Braban, C. F. and Abbatt, J. P. D.: A study of the phase transition behavior of internally mixed ammonium sulfate - malonic acid aerosols, *Atmos. Chem. Phys.*, 4, 1451–1459, doi:10.5194/acp-4-1451-2004, 2004.
- Braban, C. F., Abbatt, J. P. D., and Cziczo, D. J.: Deliquescence of ammonium sulfate particles at sub-eutectic temperatures, *Geophys. Res. Lett.*, 28, 3879–3882, 2001.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Braban, C. F., Carroll, M. F., Styler, S. A., and Abbatt, J. P. D.: Phase transitions of malonic and oxalic acid aerosols, *J. Phys. Chem. A*, 107, 6594–6602, 2003.
- Buajareern, J., Mitchem, L., and Reid, J. P.: Characterizing multiphase organic/inorganic/aqueous aerosol droplets, *J. Phys. Chem. A*, 111, 9054–9061, 2007.
- 5 Choi, M. Y. and Chan, C. K.: The effects of organic species on the hygroscopic behaviors of inorganic aerosols, *Environ. Sci. Technol.*, 36, 2422–2428, 2002.
- Ciobanu, V. G., Marcolli, C., Krieger, U. K., Weers, U., and Peter, T.: Liquid–liquid phase separation in mixed organic/inorganic aerosol particles, *J. Phys. Chem. A*, 113, 10966–10978, 2009.
- 10 Clegg, S. L., Seinfeld, J. H., and Brimblecombe, P.: Thermodynamic modeling of aqueous aerosols containing electrolytes and dissolved organic compounds, *J. Aerosol Sci.*, 32, 713–738, 2001.
- Cruz, C. N. and Pandis, S. N.: The effect of organic coatings on the cloud condensation nuclei activation of inorganic atmospheric aerosol, *J. Geophys. Res.*, 103, 13111–13123, 1998.
- 15 Cruz, C. N. and Pandis, S. N.: Deliquescence and hygroscopic growth of mixed inorganic–organic atmospheric aerosol, *Environ. Sci. Technol.*, 34, 4313–4319, 2000.
- Cziczo, D. J. and Abbatt, J. P. D.: Deliquescence, efflorescence, and supercooling of ammonium sulfate aerosols at low temperature: implications for cirrus cloud formation and aerosol phase in the atmosphere, *J. Geophys. Res.*, 104, 13781–13790, 1999.
- 20 Cziczo, D. J. and Abbatt, J. P. D.: Infrared observations of the response of NaCl, MgCl₂, NH₄HSO₄ and NH₄NO₃ aerosols to changes in relative humidity from 298 to 238 K, *J. Phys. Chem. A*, 104, 2038–2047, 2000.
- Cziczo, D. J. and Abbatt, J. P. D.: Ice nucleation in NH₄HSO₄, NH₄NO₃ and H₂SO₄ aqueous particles: implications for cirrus cloud formation, *Geophys. Res. Lett.*, 28, 963–966, 2001.
- 25 Cziczo, D. J., Nowak, J. B., Hu, J. H., and Abbatt, J. P. D.: Infrared spectroscopy of model tropospheric aerosols as a function of relative humidity: observation of deliquescence and crystallization, *J. Geophys. Res.*, 102, 18843–18850, 1997.
- Desnoyers, J. E. and Ichhaporia, F. M.: Salting-in and salting-out of polar nonelectrolytes, *Can. J. Chem.*, 47, 4639–4643, 1969.
- 30 Earle, M. E., Kuhn, T., Khalizov, A. F., and Sloan, J. J.: Volume nucleation rates for homogeneous freezing in supercooled water microdroplets: results from a combined experimental and modelling approach, *Atmos. Chem. Phys.*, 10, 7945–7961, doi:10.5194/acp-10-7945-2010, 2010.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Fried, A., Henry, B. E., and Calvert, J. G.: The reaction probability of N_2O_5 with sulfuric acid aerosols at stratospheric temperatures and compositions, *J. Geophys. Res.*, 99, 3517–3532, 1994.
- Grassian, V. H.: Heterogeneous uptake and reaction of nitrogen oxides and volatile organic compounds on the surface of atmospheric particles including oxides, carbonates, soot and mineral dust: implications for the chemical balance of the troposphere, *Int. Rev. Phys. Chem.*, 20, 467–548, 2001.
- Hallquist, M., Stewart, D. J., Stephenson, S. K., and Cox, R. A.: Hydrolysis of N_2O_5 on sub-micron sulfate aerosols, *Phys. Chem. Chem. Phys.*, 5, 3453–3463, 2003.
- Hu, J. H. and Abbatt, J. P. D.: Reaction probabilities for N_2O_5 hydrolysis on sulfuric acid and ammonium sulfate aerosols at room temperature, *J. Phys. Chem. A*, 101, 871–878, 1997.
- Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures at low temperatures: implications for ozone chemistry in the polar marine boundary layer, *J. Geophys. Res.*, 105, 26393–26402, 2000.
- Liu, Y., Yang, Z., Desyaterik, Y., Gassman, P. L., Wang, H., and Laskin, A.: Hygroscopic behavior of substrate-deposited particles studied by micro-FT-IR spectroscopy and complementary methods of particle analysis, *Anal. Chem.*, 80, 633–642, 2008.
- Marcollì, C. and Krieger, U. K.: Phase changes during hygroscopic cycles of mixed organic/inorganic model systems of tropospheric aerosols, *J. Phys. Chem. A*, 110, 1881–1893, 2006.
- Martin, S. T.: Phase transitions of aqueous atmospheric particles, *Chem. Rev.*, 100, 3403–3453, 2000.
- Murphy, D. M., Thomson, D. S., Middlebrook, A. M., Schein, M. E.: In situ single-particle characterization at Cape Grim, *J. Geophys. Res.*, 103, 16485–16491, 1998.
- Onasch, T. B., Siefert, R. L., Brooks, S. D., Prenni, A. J., Murray, B., Wilson, M. A., and Tolbert, M. A.: Infrared spectroscopic study of the deliquescence and efflorescence of ammonium sulfate aerosol as a function of temperature, *J. Geophys. Res.*, 104, 21317–21326, 1999.
- Parsons, M. T., Knopf, D. A., and Bertram, A. K.: Deliquescence and crystallization of ammonium sulfate particles internally mixed with water-soluble organic compounds, *J. Phys. Chem. A*, 108, 11600–11608, 2004.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Parsons, M. T., Riffell, J. L., and Bertram, A. K.: Crystallization of aqueous inorganic-malonic acid particles: nucleation rates, dependence on size and dependence on the ammonium-to-sulfate ratio, *J. Phys. Chem. A*, 110, 8101–8115, 2006.
- 5 Prenni, A. J., Wise, M. E., Brooks, S. D., and Tolbert, M. A.: Ice nucleation in sulfuric acid and ammonium sulfate particles, *J. Geophys. Res.*, 106, 3037–3044, 2001.
- Robinson, C. B., Schill, G. P., and Tolbert, M. A.: Optical growth of highly viscous organic/sulfate particles, *J. Atmos. Chem.*, 71, 145–156, 2014.
- Russell, L. M., Maria, S. F., and Myneni, S. C. B.: Mapping organic coatings on atmospheric particles, *Geophys. Res. Lett.*, 29, 26-1-26-4, doi:10.1029/2002GL014874, 2002.
- 10 Saxena, P., Hildemann, L. M., McMurry, P. H., and Seinfeld, J. H.: Organics alter hygroscopic behavior of atmospheric particles, *J. Geophys. Res.*, 100, 18755–18770, 1995.
- Schill, G. P., De Haan, D. O., and Tolbert, M. A.: Heterogeneous ice nucleation on simulated secondary organic aerosol, *Environ. Sci. Technol.*, 48, 1675–1682, 2014.
- Smith, M. L., Kuwata, M., and Martin, S. T.: Secondary organic material produced by the dark ozonolysis of α -pinene minimally affects the deliquescence and efflorescence of ammonium sulfate, *Aerosol Sci. Tech.*, 45, 244–261, 2011.
- 15 Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid–liquid phase separation in aerosol particles: dependence on O : C, organic functionalities, and compositional complexity, *Geophys. Res. Lett.*, 39, L19801, doi:10.1029/2012GL052807, 2012.
- 20 Tolocka, M. P., Saul, T. D., and Johnston, M. V.: Reactive uptake of nitric acid into aqueous sodium chloride droplets using real-time single-particle mass spectrometry, *J. Phys. Chem. A*, 108, 2659–2665, 2004.
- Veghte, D. P., Altaf, M. B., and Freedman, M. A.: Size dependence of the structure of organic aerosol, *J. Amer. Chem. Soc.*, 135, 16046–16049, 2013.
- 25 You, Y., Renbaum-Wolff, L., and Bertram, A. K.: Liquid-liquid phase separation in particles containing organics mixed with ammonium sulfate, ammonium bisulfate, ammonium nitrate or sodium chloride, *Atmos. Chem. Phys.*, 13, 11723–11734, doi:10.5194/acp-13-11723-2013, 2013.
- 30 Zappoli, S., Andracchio, A., Fuzzi, S., Facchini, M. C., Gelencsér, A., Kiss, G., Krivácsy, Z., Molnár, Á., Mészáros, E., Hansson, H.-C., Rosman, K., and Zebühr, Y.: Inorganic, organic and macromolecular components of fine aerosol in different areas of Europe in relation to their water solubility, *Atmos. Environ.*, 33, 2733–2743, 1999.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, 2007.
- Zhao, L., Wang, F., Zhang, K., Zeng, Q., and Zhang, Y.: Deliquescence and efflorescence processes of aerosol particles studied by in situ FTIR and Raman spectroscopy, *Chin. J. Chem. Phys.*, 21, 1–11, 2008.
- Zuend, A., Marcolli, C., Peter, T., and Seinfeld, J. H.: Computation of liquid-liquid equilibria and phase stabilities: implications for RH-dependent gas/particle partitioning of organic-inorganic aerosols, *Atmos. Chem. Phys.*, 10, 7795–7820, doi:10.5194/acp-10-7795-2010, 2010.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. List of IR features visible in the FTIR spectra shown in this work and corresponding central wavenumbers.

IR feature	Central wavenumber (cm^{-1})
Water vapor	3750
Condensed water (OH stretch)	3450
NH_4^+ (N-H stretch)	3050
D_2O and DOH vapor	2700
Condensed DOH	2500
Condensed water (HOH bend)	1640
Water vapor	1500
NH_4^+ (deformation mode)	1435
Sulfate	1115
Condensed water (H-bonding)	650
Sulfate	620

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Table 2. SRH (phase separation relative humidity), ERH (efflorescence relative humidity) and DRH (deliquescence relative humidity) points for ternary systems used in this experiment. The organic to ammonium sulfate ratio was 1.0 in all cases.

	O : C	SRH	ERH	DRH
1,4-butanediol	0.5	78.6–80.1 % (Marcolli and Krieger, 2006)	35–40 % (this study)	78–80.1 % (Marcolli and Krieger, 2006, this study)
glycerol	1	not observed (You et al., 2013)	20 % (Parsons et al., 2004) not observed (this study)	72–75 % (Marcolli and Krieger, 2006; Parsons et al., 2004) not observed (this study)
1,2,6-hexanetriol	0.5	76.7 % (You et al., 2013)	32 % (You et al., 2013) not observed (this study)	80 % (You et al., 2013) not observed (this study)
1,2-hexanediol	0.33	79.8–94.0 % (Marcolli and Krieger, 2006)	N/A	78–80.1 % (Marcolli and Krieger, 2006)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 1. Experimental setup. An atomizer was used to produce particles in an aqueous state or crystalline after removal of condensed phase water in a dryer. H₂O or D₂O vapor could then be added to the system from two bubblers coupled to mass flow controllers. The IR spectrum of the aerosol was then determined in a FTIR-coupled flow tube.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 2. A typical FTIR spectrum of ammonium sulfate at high relative humidity obtained in this experiment. Note the condensed and gas-phase features. The slope at higher wave number corresponds to aerosol scattering.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 3. Spectra obtained after subtraction of gas-phase water features. Top panel: ammonium sulfate, middle panel: 1 : 1 $(\text{NH}_4)_2\text{SO}_4$ /1,2,6-hexanetriol (mixture by mass), bottom panel: 1 : 1 $(\text{NH}_4)_2\text{SO}_4$ /glycerol. Note the presence of a condensed-phase water feature at 1640 cm^{-1} (the HOH bend). This was used to quantify liquid water in the aerosol phase.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 4. $(\text{NH}_4)_2\text{SO}_4/1,2,6\text{-hexanetriol}$ solution exposed to D_2O at three different RH values. Note the presence of features related to D_2O : the D_2O vapor features at $2600\text{--}2800\text{ cm}^{-1}$ and the condensed HOD feature at 2500 cm^{-1} . HOD is present in all three spectra, indicating that gas-phase diffusion occurred despite phase separation. Note that the green spectrum, taken at 62% RH, has a higher aerosol concentration than the other cases, resulting in the enhanced scattering slope at higher wavenumbers.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 5. Deliquescence and efflorescence experiments on $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol/water ternary mixtures of variable concentrations. **(a):** pure ammonium sulfate. Note, efflorescence at $\sim 35\%$ RH and deliquescence at $\sim 80\%$ RH. **(b):** 1 : 3 1,4-butanediol/ $(\text{NH}_4)_2\text{SO}_4$. Note, efflorescence at 35–40 % RH and deliquescence at 80 % RH. **(c):** 1 : 2 1,4-butanediol/ $(\text{NH}_4)_2\text{SO}_4$. Note, efflorescence at 35–40 % RH and deliquescence at 80 % RH. **(d):** 1 : 1 1,4-butanediol/ $(\text{NH}_4)_2\text{SO}_4$. Note, efflorescence at 35–40 % RH and deliquescence at 80 % RH.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 6. Deliquescence and efflorescence experiments on $(\text{NH}_4)_2\text{SO}_4$ /glycerol/water ternary mixtures of variable concentrations. **(a):** pure ammonium sulfate. Note, efflorescence at $\sim 35\%$ RH, deliquescence at $\sim 80\%$ RH. **(b):** 1 : 3 glycerol/ $(\text{NH}_4)_2\text{SO}_4$. Note, no clear efflorescence or deliquescence points. **(c):** 1 : 2 glycerol/ $(\text{NH}_4)_2\text{SO}_4$. Note, no clear efflorescence or deliquescence points. **(d):** 1 : 1 glycerol/ $(\text{NH}_4)_2\text{SO}_4$. Note, no clear efflorescence or deliquescence points.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 7. Deliquescence and efflorescence experiments on $(\text{NH}_4)_2\text{SO}_4/1,2,6\text{-hexanetriol/water}$ ternary mixtures of variable concentrations. **(a):** pure ammonium sulfate. Note, efflorescence at $\sim 35\%$ RH, deliquescence at $\sim 80\%$ RH. **(b):** 1 : 3 1,2,6-hexanetriol/ $(\text{NH}_4)_2\text{SO}_4$. Note, efflorescence at 40 % RH, deliquescence at 70–80 % RH. **(c):** 1 : 2 1,2,6-hexanetriol/ $(\text{NH}_4)_2\text{SO}_4$. Note, efflorescence at 40–50 % RH, no clear deliquescence point. **(d):** 1 : 1 1,2,6-hexanetriol/ $(\text{NH}_4)_2\text{SO}_4$. Note, no clear efflorescence or deliquescence points.

Hygroscopic and phase separation properties of ammonium sulfate/organic/water ternary solutions

M. A. Zawadowicz et al.




Figure 8. Gas-phase exchange experiments for organic/ $(\text{NH}_4)_2\text{SO}_4$ mixtures in three RH ranges. Data for pure ammonium sulfate at 30% RH, a crystalline solid, is plotted for reference. No change in the condensed D_2O band area is seen in only the two control cases: ammonium sulfate at 30% RH and a 1 : 1 $(\text{NH}_4)_2\text{SO}_4$ /1,4-butanediol mixture at 20% RH, both of which are below their efflorescence point (i.e., they contain no condensed-phase water). For all particles containing condensed water, an exchange of D_2O vapor is apparent. Liquid–liquid phase separation does not exclude water diffusion into the aqueous core.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion