



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore¹, J. R. Hopkins^{1,2}, R. T. Lidster¹, J. D. Lee^{1,2}, M. J. Evans^{1,2},
A. R. Rickard^{1,2}, A. C. Lewis^{1,2}, and J. F. Hamilton¹

¹Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK

²National Centre for Atmospheric Science, University of York, Heslington, York, YO10 5DD, UK

Received: 11 March 2015 – Accepted: 16 March 2015 – Published: 31 March 2015

Correspondence to: J. F. Hamilton (jacqui.hamilton@york.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Hydrocarbons are key precursors to two priority air pollutants, ozone and particulate matter. Those with two to seven carbons have historically been straightforward to observe and have been successfully reduced in many developed cities through air quality policy interventions. Longer chain hydrocarbons released from diesel vehicles are not considered explicitly as part of air quality strategies and there are few direct measurements of their gaseous abundance in the atmosphere. This study describes the chemically comprehensive and continuous measurements of organic compounds in a developed megacity (London), which demonstrate that on a seasonal median basis, diesel-related hydrocarbons represent only 20–30 % of the total hydrocarbon mixing ratio but comprise more than 50 % of the atmospheric hydrocarbon mass and are a dominant local source of secondary organic aerosols. This study shows for the first time that, 60 % of the winter primary hydrocarbon hydroxyl radical reactivity is from diesel-related hydrocarbons and using the maximum incremental reactivity scale, we predict that they contribute up to 50 % of the ozone production potential in London. Comparing real-world urban composition with regulatory emissions inventories in the UK and US highlights a previously unaccounted for but, very significant under-reporting of diesel related hydrocarbons; an underestimation of a factor ~ 4 for C_9 species rising to a factor of over 70 for C_{12} during winter. These observations show that hydrocarbons from diesel vehicles can dominate gas phase reactive carbon in cities with high diesel fleet fractions. Future control of urban particulate matter and ozone in such locations requires a shift in policy focus onto gas phase hydrocarbons released from diesels as this vehicle type continues to displace gasoline world-wide.

1 Introduction

With an increasing proportion of the world's population living in cities, rising from only 3 % in the 1800's to over 47 % by the end of the 20th Century (Gaffney and Marley,

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2009), the impact of urban air pollution has become a significant factor in global health (Harrison et al., 2012). The costs of air pollution are high even in those locations that have seen considerable improvements in air quality over the past decades (Chameides et al., 1992); in the UK exposure to particulate matter (PM) alone is estimated to reduce life expectancy on average by around 7–8 months, with a cost to society estimated at up to GBP 20 billion per year (House of Commons Environmental Audit Committee, 2012).

Primary urban air pollution emissions are dominated by PM, nitrogen oxides (NO_x), carbon monoxide (CO) and volatile organic compounds (VOCs). Many of these species can react in the atmosphere to create secondary pollutants, such as ozone (O₃), oxygenated VOCs (OVOCs), peroxy acetyl nitrates (PANs) and condensed materials in the form of secondary organic aerosol (SOA), which add to the overall PM load (Atkinson and Arey, 2003; Odum et al., 1997).

Air quality in London has been controlled and monitored for over 60 years, making it in theory one of the better understood atmospheres of the world's megacities. Current measurements in London focus on assessing national compliance with legally prescribed air quality standards, and this includes the hydrocarbons, 1,3-butadiene and benzene. However, compliance measurements in themselves are insufficient to fully describe the chemical and physical processes occurring in the urban atmosphere (McMeeking et al., 2012), and a particular weakness lies in speciating the many different classes of carbon compounds in urban air.

The past two decades have seen declining concentrations of most smaller hydrocarbons in European and US cities, a result of tighter regulation of sources such as vehicle exhaust, evaporation and solvents (von Schneidemesser et al., 2010; Warneke et al., 2012), better control of natural gas leakage and an overall switch from gasoline to diesel powered vehicles. Current national emissions estimates suggest that the bulk of organic emissions to air are associated with smaller hydrocarbons, and this has driven both policy, regulation and observation strategies for compliance. Figure 1 shows Government-estimated emissions for the UK (left) and US (right), categorised

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

into the dominant emission sources. It is clear that based on current emission inventories, gaseous organic emissions from diesel appear to represent a negligible fraction of reactive carbon released into the atmosphere. The demand for diesel fuel is expected to increase by 75 % between 2010 and 2040 and by 2020 it is expected to overtake gasoline as the number one transport fuel used worldwide (Exxon Mobil, 2014). The environmental impacts of this change are evaluated in part based on the national emission inventories that underpin Fig. 1.

The efficiency with which O_3 and SOA can be formed from diesel or gasoline emissions is dependent on the mass of available organic carbon, and the reactivity and volatility of that material (Gros et al., 2007). To quantify this requires individual specification of VOCs in order that each property can be properly estimated. The key urban sources of organic compounds include combustion products, unburnt fuels and evaporative emissions of fuels and solvents, all of which are highly complex, often propagating the original complexity of fossil fuels into the air. Whilst each VOC has a unique set of reaction mechanisms, in general terms, as the carbon number increases, the relative complexity of reactions and yields of SOA and O_3 also increase (Hamilton and Lewis, 2003). The organic mixture in air is complicated further by the presence of secondary oxygenated products, requiring a combined approach to investigate VOC composition.

Several recent field studies have investigated the relative importance of gasoline, diesel and biogenic emissions in generating SOA. These have been carried out predominantly in the US, particularly in California, where current diesel usage is rather low by global standards. The US diesel fleet is dominated by heavy-duty vehicles, leading to a difference in the source strength of diesel and gasoline engines between weekdays and the weekend. In contrast, this trend is not observed in London (see Supplement and Fig. S1), which has a different vehicle fleet composition, with 60 % diesel fuel use, and a large number of diesel buses operating at similar times to domestic vehicles. It is also important to consider that London and Los Angeles (arguably the world's most well studied city for air pollution) have significant differences in terms of urban geography, population density, commuting patterns, amount of green space/trees and upwind

the ground through a condensation finger in an ethylene glycol bath held at -30°C , to remove any moisture from the sample. See Supplement for details about calibrations.

The DC-GC-FID was operated by the National Centre for Atmospheric Science (NCAS) Facility for Ground Atmospheric Measurements (FGAM) with the instrument set up and calibration described in Hopkins et al. (2003). In brief, the system has three GC columns that operate in parallel, where after sampling and desorption the flow is split 50 : 50; one column is an aluminium oxide (Al_2O_3) Porous Layer Open Tubular (PLOT, 50 m, 0.53 mm id) for non-methane hydrocarbons (NMHCs) analysis; and the other column is actually two LOWOX columns in series (10 m, 0.53 mm id) for OVOC analysis.

The GC \times GC-FID comprised of a Markes TT24-7 thermal desorption (TD) unit with an air server attachment (Markes International, Llantrisant, UK) and an Agilent 7890 GC (Agilent Technologies, Wilmington, DE, USA) equipped with an FID operating at 200 Hz. The TD unit sampled at a rate of 100 mL min^{-1} for 55 min, giving a total sample volume of 5.5 L. The trap temperature was set to -10°C , held for 3 min, then on injection heated at $100^{\circ}\text{C min}^{-1}$ to 200°C to ensure all analytes of interest were desorbed.

The GC \times GC-FID system first dimension column was a BPX-5 ($25\text{ m} \times 0.15\text{ mm}$, $0.4\text{ }\mu\text{m df}$), at 50 psi, combined with a second dimension column of a BP-20 ($5\text{ m} \times 0.25\text{ mm}$, $0.25\text{ }\mu\text{m df}$), at 23 psi (SGE, Australia), with column pressures controlled using the Agilent 7890 EPC. The total transfer flow valve modulator (Valco Instruments, Houston, TX, USA) was a 6-port, 2-way diaphragm valve modulator, with actuation achieved using a solenoid valve, controlled by software written "in house". The modulator was held at 120°C throughout the run and had a modulation period of 5 s, with 4.7 s sample and 0.3 s injection times. The chromatographic and modulation configuration of the GC \times GC-FID system is detailed in Lidster et al. (2011). During the injection of sample, liquid carbon dioxide (CO_2) was sprayed onto the first 2 cm portion at the head of the first dimension column for 60 s to re-focus the sample.

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An oven temperature programme was developed which optimised separation and resolution of compounds of interest. The initial oven temperature was 30 °C, held for 1 min, ramped at 2.5 °C min⁻¹ to 130 °C, held for 1 min then ramped at 10 °C min⁻¹ to 200 °C, and held for 1 min; giving a total run time of 50 min. This, combined with the TD run time, gave a total analysis time of 55 min.

During the summer campaign, some parameters had to be changed. Ambient temperatures were higher in comparison to the winter, meaning the oven temperature programme had to be altered to allow the oven to reach its minimum temperature. The initial oven temperature was changed to 35 °C held for 2 min and the final temperature of 200 °C was held for 2 min. All other oven parameters were kept the same. Due to a sensitivity drop, the TD sampling rate was increased to 200 mL min⁻¹ for 55 min, to give a total sample volume of 11 L.

2.3 Supporting measurements

Measurements of NO_x were made using a single channel, chemiluminescence instrument (Air Quality Design Inc., USA), which has a wide linear range (1 ppt to 500 ppb). Lee et al. (2009) Ozone measurements were made using an UV Absorption TEI 49C and 49i (Thermo Scientific) with a limit of detection of 1 ppb.

3 Observations of hydrocarbons in urban air

The two GC instruments individually quantified 78 VOCs (36 aliphatics, 19 monoaromatics, 21 oxygenated and 2 halogenated), as well as many hundreds more included in a lumped carbon number assessment from 2667 samples (1352 winter and 1315 summer). The DC-GC instrument measured volatile VOCs, C₁–C₇ hydrocarbons and a selection of OVOCs, with effective saturation concentrations (Nannoolal et al., 2004, 2008) ranging from 3 × 10⁷ to 1.4 × 10¹² μg m⁻³. The GC × GC-FID instrument measured the less volatile VOC fraction (effective saturation concentration range of 1.8 × 10⁶ to

$2.4 \times 10^9 \mu\text{g m}^{-3}$), with hydrocarbons from C_6 to C_{13} , plus a large group of OVOCs (from C_3 onwards). There was some overlap in species measured by both instruments, with good agreement seen (e.g. benzene R^2 0.92, slope 1.070 ± 0.013 , see Fig. S4).

The number of possible structural isomers increases exponentially with carbon number (Goldstein and Galbally, 2007) and beyond around C_9 it becomes impossible to accurately identify the structure of every hydrocarbon present in air. Using the retention behaviour of each compound on chromatographic columns it is however possible to assign individual species to particular chemical classes and functionalities. Here we group according to carbon number and basic functionality, an example of which can be seen in Fig. S6. The C_2 – C_6 volatility range contains both primary hydrocarbon emissions, with isomers quantified individually, along with several oxygenated compounds. Between C_6 to C_{13} , a wide range of hydrocarbon and OVOC species, including the *n*-alkanes, α -pinene, limonene, monoaromatics with up to 3 substituents and naphthalene were quantified individually (see Fig. S5 and Table S2). Higher aliphatic compounds (C_6 – C_{13} , predominantly alkanes with some alkenes and cycloalkanes), C_4 substituted monoaromatics and C_{10} monoterpenes have been grouped together and the combined class abundance estimated using a response ratio to the corresponding straight-chained alkane, 1,3-diethyl benzene and α -pinene respectively. The number of individual isomer peaks that could be isolated in the aliphatics grouping increased from 9 for the C_6 group to 40 for C_{10} (shown in Fig. 5, black squares). A full table of the observed isomer peaks and group mass concentrations is provided in Table S3.

3.1 Diurnal behaviour

The average diurnal behaviour of a selection of VOCs, O_3 and NO_x , are shown in Fig. 2. VOCs with an anthropogenic source (e.g. ethane, toluene, C_4 substituted monoaromatics and C_{13} aliphatics) have higher mixing ratios in winter, consistent with reduced rates of photochemical removal in the northern European winter and a lower boundary layer height. Toluene and NO_x are strong indicators of traffic related emissions, and

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

both have diurnal profiles with rush hour peaks, commonly observed in urban areas (Borbon et al., 2001; Parrish et al., 2009; Gaffron, 2012). The profiles of the higher carbon number species, C₄ substituted monoaromatics and C₁₃ aliphatics, show similar traffic-related profiles, strongly indicative that this is their major source. Those species with a dominant biogenic source (e.g. isoprene and α -pinene) are higher in summer, due to increased emission. The winter profiles of isoprene and α -pinene show possible anthropogenic sources, traffic and cleaning products respectively.

3.2 Reactivity and mass calculations of grouped compounds

Figure 3 shows the relative emission source contributions of compounds to the total hydrocarbon mixing ratio (top), mass concentration (middle) and primary hydrocarbon hydroxyl radical (OH) reactivity (bottom), calculated by carbon number, and split according to emission source. For each sample, the mass concentration ($\mu\text{g m}^{-3}$) and primary hydrocarbon OH reactivity (s^{-1} , see Supplement) were calculated and a seasonal median calculated using the mixing ratios of the individual components and the summation of all further unidentified species within the ten class groups, not including the OVOCs. The Passant (2002) speciated emissions inventory (discussed previously) was used to determine the main emission sources for each compound. For C₂ to C₅, the main source is either natural gas usage or leakage, followed by road transport use. From C₆ onwards, the main emission source was classified as road transport or other fuel usage categories (i.e. filling of petrol stations). To determine the percentage contributions from diesel or gasoline fuel usage, the detailed fuel characterisation of Gentner et al. (2012) was used with a value of 60 % diesel use in the UK (Department of Energy and Climate Change, 2014).

The winter has generally higher abundances of hydrocarbons, with summer showing a marked increase in the biogenic source compounds. In both winter and summer, in mixing ratio terms, the distribution is dominated by high volatility species (over 85 % of the total) primarily from natural gas and gasoline sources. When viewed in terms of mass concentration, however, the distribution of combined natural gas and gasoline

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vs. diesel is closer to 70 : 30 % in winter and 77 : 23 % in summer, the latter due to the increased loss rates for reactive species in summer disproportionately removing larger hydrocarbons.

In order to calculate primary hydrocarbon OH reactivity of diesel emissions, rate constants have to be estimated as each individual species is not uniquely identified and because the rate constants (and subsequent chemistry) are un-measured in the majority of cases for hydrocarbons larger than C₈. The nearest straight-chain alkane rate constant (Atkinson and Arey, 2003) is applied to all carbon in that aliphatic grouping. This will lead to a conservative estimate of reactivity since cycloalkanes and alkenes would be expected to react faster (i.e. the k_{OH} for the reaction of *n*-dodecane is $1.32 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, compared to that of 1-dodecene which is $5.03 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Aschmann and Atkinson, 2008), for further details see Supplement and Table S4). For the C₄ substituted monoaromatic and C₁₀ monoterpene groups, the rate constants of 1,3 diethyl benzene and α -pinene respectively were used.

3.3 Calculation of unmeasured diesel emissions

The GC measurements stop at C₁₃, however diesel typically has a range of hydrocarbons from C₉ to C₂₂ that peaks with *n*-hexadecane as the most abundant compound. Using the observed distribution of C₁₀–C₁₃, allows for an estimate to be made of the remaining, *unobserved* NMHC fraction of gaseous diesel emissions in the C₁₄–C₂₂ range, using the fuel composition-based emission factors for the gas phase compounds from Gentner et al. (2013). Assuming no atmospheric loss, a reasonable approximation in winter, it is estimated that the GC × GC-FID technique observes around 25–30 % of the total gaseous hydrocarbon emissions from diesel sources. It is then possible to estimate a seasonal average unmeasured gas phase NMHC mass concentration from diesel sources in London as 76.1–97.8 $\mu\text{g m}^{-3}$ in winter and 26.8–34.3 $\mu\text{g m}^{-3}$ in summer. These values can be compared to typical primary organic aerosol measurements of 1 $\mu\text{g m}^{-3}$ (Zhang et al., 2011; Young et al., 2014), indicating that, in ambient air,

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



diesel-related emission of hydrocarbons are overwhelmingly (a factor of 100) to the gas phase, consistent with laboratory and tail pipe studies (Gordon et al., 2013). In contrast, using gasoline liquid fuel speciation, the combined GC approach can observe approximately over 98 % of the mass of gasoline.

The impact of the unobserved diesel emissions on primary hydrocarbon OH reactivity was estimated using the *n*-dodecane rate constant as a proxy. The percentage contributions to mixing ratio (top), mass concentration (middle) and primary hydrocarbon OH reactivity (bottom), divided by emission source, including the unmeasured diesel emissions and OVOCs, are shown in Fig. 4. It is clear that diesel plays an important role in the composition of NMHCs and their subsequent reactivity. The total (measured + calculated) diesel emissions contribute 5.1 and 1.7 s⁻¹ to OH reactivity in winter and summer respectively compared to 1.7 and 0.8 s⁻¹ from gasoline compounds; increasing the contribution of diesel-related hydrocarbons to calculated VOC OH reactivity from 23 % using the measured VOCs to 61 % including the unmeasured I/VOCs in winter and from 8 to 34 % in summer (full details can be found in Table S5). In summer, the primary emissions have undergone a degree of loss due to photochemical ageing and so the values are an underestimate of the fuel sources. These diesel-related hydrocarbons may be partly responsible for the “missing” OH reactivity observed between measurements of OH lifetime vs. the value calculated from observed sinks in many studies (Yoshino et al., 2006, 2012).

3.4 Comparison to emissions inventories

Assuming that; the winter observations are made “at source” (hence atmospheric losses and lifetime differences can be neglected), the measurement location is representative of an urban setting, and the UK inventory correctly estimates the emission for toluene (based on direct flux comparisons made in London by Langford et al., 2010), there appears to be a significant inventory under-reporting for the higher carbon number species. When normalising to toluene, the UK national emissions inventory, which is believed to use best-practice international reporting methodologies, underreports by

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

this study, the uncertainty in the unspiciated fraction has been reduced by grouping ambient observations of VOCs by carbon number and functionality. The $> C_9$ aliphatic groups are dominated by diesel emissions at this location and so the SOA source strength can be more accurately determined. The potential contributions of the higher hydrocarbons to SOA formation has been estimated by multiplying the median measured VOC mass concentration (Fig. 7, top panel, black columns) by the corresponding SOA yield (Fig. 7, top panel, blue circles) (Zhang et al., 2014). The yields applied were measured in high NO_x chamber studies (VOC_0/NO_x (ppbC ppb $^{-1}$) of 0.5) with an organic aerosol mass C_{OA} of $10 \mu\text{g m}^{-3}$, representative of urban areas (Zhang et al., 2014; Presto et al., 2010). Here it is assumed that NMHCs with less than six carbons and aqueous chemistry of water soluble oxidation products, such as glyoxal, do not contribute to SOA mass. In both winter and summer, the observed levels of aliphatic compounds from diesel sources have the potential to form significant quantities of SOA (Fig. 7, bottom panel, red columns). If a diesel SOA yield of 0.15 is applied to the total diesel emissions (as calculated previously), then gas phase emissions from diesel engines represent the dominant traffic related precursor source of urban SOA in a European megacity such as London, where the use of diesel fuel is prevalent, in line with previous studies in the US (Gentner et al., 2012). Each cubic meter of air contains sufficient gas phase hydrocarbons to potentially produce 14.4–17.6 and 4.9–6.1 μg of SOA in winter and summer following atmospheric oxidation.

Recent simulation chamber studies indicate that modern engines fitted with diesel particle filters, such as EURO5 emissions control, have greatly reduced VOC tailpipe emission and form little SOA under chamber conditions (Gordon et al., 2014a, b). However, Carslaw and Rhys-Tyler (2013) have recently shown that when vehicles are driven under real-world urban conditions, the emission of NO_x from diesel engines have not been reduced as expected given the new technologies implemented. It is possible to infer that if NO_x emissions are higher than expected, the VOC emissions are also likely to be higher. In this study we have shown that there is a significant diesel vehicle source emitting sufficient VOCs to impact ozone and SOA formation in the urban atmosphere.

4 Conclusions

From the results presented, it is possible to conclude that current inventories and emissions estimates do not adequately represent emissions of gas phase higher carbon number species from the diesel fleet under real-world conditions and in a developed urban environment. The calculated impact of these species is significant, particularly in terms of OH reactivity, ozone formation potential and SOA production.

In the last decade, there has been a steady shift in fuel use in many locations. For example, in the UK diesel fuel use as a fraction of total fuel has risen from 52 % in 2005 to 62 % in 2011 (see Fig. S8 and Table S7) (Department of Energy and Climate Change, 2014). Although the UK may be considered typical of Europe (where diesel use varies between 45–80 %) (European Commission, 2012), the average US value was around 29 % diesel use (U.S. Energy Information Administration, 2014) in 2013, with the understanding of other geographical regions currently being poor.

This shift to an increasingly diesel-powered fleet in many developed cities, as a response to energy efficiency drivers, has therefore shifted the balance of hydrocarbons in urban air from short to long chain compounds, and these observations provide direct atmospheric evidence of this effect in London. Whilst this has been seen in a small selection of previous studies, as discussed previously, these were carried out in the US where the geographic characteristics are quite different to London. In many instances, this has gone unmeasured in ambient air due to the lack of meaningful measurement infrastructure. Understanding the impact of this change is significantly hindered however by a lack of appropriate physico-chemical data for individual longer chain hydrocarbons. There are already very significant policy challenges for many developed cities relating to the control of NO₂ from modern diesel vehicles, and this study indicates that there may also be a similar, but currently un-recognized, policy challenge to control reactive carbon emissions and their contributions to secondary pollutants.

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Author contributions. R. E. Dunmore and J. R. Hopkins analysed the GC data and J. D. Lee made the O₃ and NO_x measurements. R. T. Lidster and J. F. Hamilton developed the GC × GC-FID instrument. A. R. Rickard provided insight into the kinetics and photochemical ozone creation potentials. R. E. Dunmore, J. F. Hamilton, A. C. Lewis and M. J. Evans wrote the paper. All authors contributed towards the final version of the paper.

Acknowledgements. R. E. Dunmore would like to thank NERC (NE/J500197/1) for PhD funding and to acknowledge David Carslaw for his assistance using Openair and Richard Derwent for assistance with the NAEI. All authors would like to acknowledge NERC for funding the Clearflo project, and the efforts of all participants in the field experiment in supporting these measurements. We would like to acknowledge Stephen Belcher for project co-ordination.

References

- Aschmann, S. and Atkinson, R.: Rate constants for the gas-phase reactions of OH radicals with *E*-7-tetradecene, 2-methyl-1-tridecene and the C₇–C₁₄ 1-alkenes at 295 ± 1 K, *Phys. Chem. Chem. Phys.*, 10, 4159–4164, doi:10.1039/B803527J, 2008. 9550
- Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605–4638, doi:10.1021/cr0206420, 2003. 9543, 9550
- Bigi, A. and Harrison, R.: Analysis of the air pollution climate at a central urban background site, *Atmos. Environ.*, 44, 2004–2012, doi:10.1016/j.atmosenv.2010.02.028, 2010. 9545
- Bohnenstengel, S. I., Belcher, S. E., Aiken, A., Allan, J. D., Allen, G., Bacak, A., Bannan, T. J., Barlow, J., Beddows, D. C. S., Bloss, W. J., Booth, A. M., Chemel, C., Coceal, O., Di Marco, C. F., Dubey, M. K., Faloon, K. H., Fleming, Z. L., Furger, M., Gillett, J. K., Graves, R. R., Green, D. C., Grimmond, C. S. B., Halios, C. H., Hamilton, J. F., Harrison, R. M., Heal, M. R., Heard, D. E., Helfter, C., Herndon, S. C., Holmes, R. E., Hopkins, J. R., Jones, A. M., Kelly, F. J., Kotthaus, S., Langford, B., Lee, J. D., Leigh, R. J., Lewis, A. C., Lidster, R. T., Lopez-Hilfiker, F. D., McQuaid, J. B., Mohr, C., Monks, P. S., Nemitz, E., Ng, N. L., Percival, C. J., Prévôt, A. S. H., Ricketts, H. M. A., Sokhi, R., Stone, D.,

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Thornton, J. A., Tremper, A. H., Valach, A. C., Visser, S., Whalley, L. K., Williams, L. R., Xu, L., Young, D. E. E., and Zotter, P.: Meteorology, air quality, and health in London: the ClearfLo project, *B. Am. Meteorol. Soc.*, online first, doi:10.1175/bams-d-12-00245.1, 2014. 9545

5 Borbon, A., Fontaine, H., Veillerot, M., Locoge, N., Galloo, J., and Guillermo, R.: An investigation into the traffic-related fraction of isoprene at an urban location, *Atmos. Environ.*, 35, 3749–3760, doi:10.1016/S1352-2310(01)00170-4, 2001. 9549

Calvert, J. G., Derwent, R. G., Orlando, J. J., Tyndall, G. S., and Wallington, T. J.: *Mechanisms of Atmospheric Oxidation of the Alkenes*; Oxford University Press: New York, 2000. 9552, 9553

10 Carslaw, D. and Rhys-Tyler, G.: New insights from comprehensive on-road measurements of NO_x, NO₂ and NH₃ from vehicle emission remote sensing in London, UK, *Atmos. Environ.*, 81, 339–347, doi:10.1016/j.atmosenv.2013.09.026, 2013. 9555

Carslaw, D. and Ropkins, K.: openair – an R package for air quality data analysis, *Environ. Modell. Softw.*, 27–28, 52–61, doi:10.1016/j.envsoft.2011.09.008, 2012a. 9566

15 Carslaw, D. and Ropkins, K.: openair – data analysis tools for the Air Quality Community, *The R Journal*, 4, 20–29, 2012b. 9566

Carter, W.: SAPRC-07 Atmospheric Chemistry Mechanisms and VOC Reactivity Scales, available at: <http://www.engr.ucr.edu/~carter/SAPRC/> (last access: 10 February 2015), 2010. 9553

20 Chameides, W., Fehsenfeld, F., Rodgers, M., Cardelino, C., Martinez, J., Parrish, D., Lonnenman, W., Lawson, D., Rasmussen, R., Zimmerman, P., Greenberg, J., Middleton, P., and Wang, T.: Ozone precursor relationships in the ambient atmosphere, *J. Geophys. Res.-Atmos.*, 97, 6037–6055, doi:10.1029/91JD03014, 1992. 9543

25 Department of Energy and Climate Change: Road transport energy consumption at regional and local authority level, available at: www.gov.uk/government/statistical-data-sets/road-transport-energy-consumption-at-regional-and-local-authority-level (last access: 10 February 2015), 2014. 9549, 9556

30 Derwent, R., Jenkin, M., Saunders, S., and Pilling, M.: Photochemical ozone creation potentials for organic compounds in northwest Europe calculated with a master chemical mechanism, *Atmos. Environ.*, 32, 2429–2441, doi:10.1016/S1352-2310(98)00053-3, 1998. 9552

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Derwent, R., Jenkin, M., Pilling, M., Carter, W., and Kaduwela, A.: Reactivity scales as comparative tools for chemical mechanisms, *J. Air Waste Manage.*, 60, 914–924, doi:10.3155/1047-3289.60.8.914, 2010. 9553

European Commission: EU Transport in Figures, available at: <http://ec.europa.eu/transport/facts-fundings/statistics/doc/2012/pocketbook2012.pdf> (last access: 10 February 2015), 2012. 9556

Exxon Mobil: The Outlook for Energy: A view to 2040, Tech. rep., Exxon Mobil Corporation, Texas, 2014. 9544

Gaffney, J. and Marley, N.: The impacts of combustion emissions on air quality and climate – from coal to biofuels and beyond, *Atmos. Environ.*, 43, 23–36, doi:10.1016/j.atmosenv.2008.09.016, 2009. 9542

Gaffron, P.: Urban transport, environmental justice and human daily activity patterns, *Transport Policy*, 20, 116–129, 12th World Conference on Transport Research (WCTR)/14th World Conference of the Air-Transport-Research-Society (ATRS), Lisbon, Portugal, 11–15 July 2010, 2012. 9549

Gentner, D., Isaacman, G., Worton, D., Chan, A., Dallmann, T., Davis, L., Liu, S., Day, D., Russell, L., Wilson, K., Weber, R., Guha, A., Harley, R., and Goldstein, A.: Elucidating secondary organic aerosol from diesel and gasoline vehicles through detailed characterization of organic carbon emissions, *P. Natl. Acad. Sci. USA*, 109, 18318–23, doi:10.1073/pnas.1212272109, 2012. 9549, 9554, 9555

Gentner, D., Worton, D., Isaacman, G., Davis, L., Dallmann, T., Wood, E., Herndon, S., Goldstein, A., and Harley, R.: Chemical composition of gas-phase organic carbon emissions from motor vehicles and implications for ozone production, *Environ. Sci. Technol.*, 47, 11837–11848, doi:10.1021/es401470e, 2013. 9550, 9553, 9554

Goldstein, A. and Galbally, I.: Known and unexplored organic constituents in the Earth's atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, doi:10.1021/es072476p, 2007. 9548

Gordon, T., Tkacik, D., Presto, A., Zhang, M., Jathar, S., Nguyen, N., Massetti, J., Truong, T., Cicero-Fernandez, P., Maddox, C., Rieger, P., Chattopadhyay, S., Maldonado, H., Maricq, M. M., and Robinson, A.: Primary gas- and particle-phase emissions and secondary organic aerosol production from gasoline and diesel off-road engines, *Environ. Sci. Technol.*, 47, 14137–14146, doi:10.1021/es403556e, 2013. 9551

Gordon, T. D., Presto, A. A., May, A. A., Nguyen, N. T., Lipsky, E. M., Donahue, N. M., Gutierrez, A., Zhang, M., Maddox, C., Rieger, P., Chattopadhyay, S., Maldonado, H., Mar-

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

icq, M. M., and Robinson, A. L.: Secondary organic aerosol formation exceeds primary particulate matter emissions for light-duty gasoline vehicles, *Atmos. Chem. Phys.*, 14, 4661–4678, doi:10.5194/acp-14-4661-2014, 2014a. 9554, 9555

5 Gordon, T. D., Presto, A. A., Nguyen, N. T., Robertson, W. H., Na, K., Sahay, K. N., Zhang, M., Maddox, C., Rieger, P., Chattopadhyay, S., Maldonado, H., Maricq, M. M., and Robinson, A. L.: Secondary organic aerosol production from diesel vehicle exhaust: impact of aftertreatment, fuel chemistry and driving cycle, *Atmos. Chem. Phys.*, 14, 4643–4659, doi:10.5194/acp-14-4643-2014, 2014b. 9554, 9555

10 Gros, V., Sciare, J., and Yu, T.: Air-quality measurements in megacities: focus on gaseous organic and particulate pollutants and comparison between two contrasted cities, Paris and Beijing, *CR Geosci.*, 339, 764–774, doi:10.1016/j.crte.2007.08.007, 2007. 9544

Hamilton, J. and Lewis, A.: Monoaromatic complexity in urban air and gasoline assessed using comprehensive GC and fast GC-TOF/MS, *Atmos. Environ.*, 37, 589–602, doi:10.1016/S1352-2310(02)00930-5, 2003. 9544

15 Harrison, R. M., Dall'Osto, M., Beddows, D. C. S., Thorpe, A. J., Bloss, W. J., Allan, J. D., Coe, H., Dorsey, J. R., Gallagher, M., Martin, C., Whitehead, J., Williams, P. I., Jones, R. L., Langridge, J. M., Benton, A. K., Ball, S. M., Langford, B., Hewitt, C. N., Davison, B., Martin, D., Petersson, K. F., Henshaw, S. J., White, I. R., Shallcross, D. E., Barlow, J. F., Dunbar, T., Davies, F., Nemitz, E., Phillips, G. J., Helfter, C., Di Marco, C. F., and Smith, S.: Atmospheric chemistry and physics in the atmosphere of a developed megacity (London): an overview of the REPARTEE experiment and its conclusions, *Atmos. Chem. Phys.*, 12, 3065–3114, doi:10.5194/acp-12-3065-2012, 2012. 9543

20 Hopkins, J., Lewis, A., and Read, K.: A two-column method for long-term monitoring of non-methane hydrocarbons (NMHCs) and oxygenated volatile organic compounds (OVOCs), *J. Environ. Monitor.*, 5, 8–13, doi:10.1039/b202798d, 2003. 9546

25 House of Commons Environmental Audit Committee: Air Quality: A Follow-up Report, Ninth Report, Tech. rep., House of Commons, London, The Stationery Office Limited, 2012. 9543

Jathar, S., Gordon, T., Hennigan, C., Pye, H. T., Pouliot, G., Adams, P., Donahue, N., and Robinson, A.: Unspeciated organic emissions from combustion sources and their influence on the secondary organic aerosol budget in the United States, *P. Natl. Acad. Sci. USA*, 111, 10473–10478, doi:10.1073/pnas.1323740111, 2014. 9554

30 Langford, B., Nemitz, E., House, E., Phillips, G. J., Famulari, D., Davison, B., Hopkins, J. R., Lewis, A. C., and Hewitt, C. N.: Fluxes and concentrations of volatile organic compounds

above central London, UK, *Atmos. Chem. Phys.*, 10, 627–645, doi:10.5194/acp-10-627-2010, 2010. 9551

Lee, J. D., Moller, S. J., Read, K. A., Lewis, A. C., Mendes, L., and Carpenter, L. J.: Year-round measurements of nitrogen oxides and ozone in the tropical North Atlantic marine boundary layer, *J. Geophys. Res.-Atmos.*, 114, D21302, doi:10.1029/2009JD011878, 2009. 9547

Lidster, R., Hamilton, J., and Lewis, A.: The application of two total transfer valve modulators for comprehensive two-dimensional gas chromatography of volatile organic compounds, *J. Sep. Sci.*, 34, 812–821, doi:10.1002/jssc.201000710, 2011. 9546

McMeeking, G. R., Bart, M., Chazette, P., Haywood, J. M., Hopkins, J. R., McQuaid, J. B., Morgan, W. T., Raut, J.-C., Ryder, C. L., Savage, N., Turnbull, K., and Coe, H.: Airborne measurements of trace gases and aerosols over the London metropolitan region, *Atmos. Chem. Phys.*, 12, 5163–5187, doi:10.5194/acp-12-5163-2012, 2012. 9543

Nannoolal, Y., Rarey, J., Ramjugernath, D., and Cordes, W.: Estimation of pure component properties: Part 1. Estimation of the normal boiling point of non-electrolyte organic compounds via group contributions and group interactions, *Fluid Phase Equilibr.*, 226, 45–63, doi:10.1016/j.fluid.2004.09.001, 2004. 9547

Nannoolal, Y., Rarey, J., and Ramjugernath, D.: Estimation of pure component properties: Part 3. Estimation of the vapor pressure of non-electrolyte organic compounds via group contributions and group interactions, *Fluid Phase Equilibr.*, 269, 117–133, doi:10.1016/j.fluid.2008.04.020, 2008. 9547

Odum, J. R., Jungkamp, T. P. W., Griffin, R. J., Forstner, H. J. L., Flagan, R. C., and Seinfeld, J. H.: Aromatics, reformulated gasoline, and atmospheric organic aerosol formation, *Environ. Sci. Technol.*, 31, 1890–1897, doi:10.1021/es960535l, 1997. 9543

Parrish, D., Kuster, W., Shao, M., Yokouchi, Y., Kondo, Y., Goldan, P. D., de Gouw, J., Koike, M., and Shirai, T.: Comparison of air pollutant emissions among mega-cities, *Atmos. Environ.*, 43, 6435–6441, doi:10.1016/j.atmosenv.2009.06.024, 2009. 9549

Passant, N.: Speciation of UK emissions of non-methane volatile organic compounds, Tech. rep., AEA Technology Report ENV-05452002, Culham, Abingdon, UK, 2002. 9552, 9564

Platt, S. M., El Haddad, I., Zardini, A. A., Clairrotte, M., Astorga, C., Wolf, R., Slowik, J. G., Temime-Roussel, B., Marchand, N., Ježek, I., Drinovec, L., Močnik, G., Möhler, O., Richter, R., Barmet, P., Bianchi, F., Baltensperger, U., and Prévôt, A. S. H.: Secondary organic aerosol formation from gasoline vehicle emissions in a new mobile environmental reac-

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Zhang, Q., Jimenez, J., Canagaratna, M., Ulbrich, I., Ng, N., Worsnop, D., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Anal. Bioanal. Chem.*, 401, 3045–3067, doi:10.1007/s00216-011-5355-y, 2011. 9550
- 5 Zhang, X., Cappa, C., Jathar, S., McVay, R., Ensberg, J., Kleeman, M., and Seinfeld, J.: Influence of vapor wall loss in laboratory chambers on yields of secondary organic aerosol, *P. Natl. Acad. Sci. USA*, 111, 5802–5807, doi:10.1073/pnas.1404727111, 2014. 9555

ACPD

15, 9541–9571, 2015

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Figure 1. Total mass by carbon number and functionality from UK 2012 (left) and US 2011 (right) emission inventories. The carbon number and functionality of emissions have been estimated by applying the speciated inventory of emission sources of Passant (2002) to the most recent estimates of non-methane hydrocarbon source apportionment for each country (full details can be found in the Supplement).

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.




Figure 3. Seasonal median values for hydrocarbon mixing ratio, mass concentration and primary hydrocarbon OH reactivity in London air grouped by carbon number and potential emission source.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.




Figure 4. Contributions of emission source to total mixing ratio, mass and OH reactivity for winter and summer. Diesel is the summation of measured and calculated, with error bars indicating the uncertainty of the unobserved diesel NMHC fraction.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.




Figure 5. Winter emissions inventory underestimation (left axis and blue columns) and the number of isomers included in each grouped set of compounds (right axis and black squares). Grey line shows a factor of 1 i.e. inventory emission estimation is consistent with the observations.

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.




Figure 6. Contribution of emission sources to ozone formation potential, where diesel is representative of total diesel emissions and the error bars show the uncertainty of the unobserved diesel fraction calculation.

Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities

R. E. Dunmore et al.




Figure 7. Potential SOA mass estimates. Upper: Mean VOC mass concentration [VOC] shown by black columns, and the corresponding SOA yields (Y) for the VOC precursors in blue circles. Lower: potential SOA mass concentration [SOA], calculated as the product of mean VOC mass and SOA yields. Winter shown on left and summer shown on right hand side panels.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion