

**SCIAMACHY surface  
pressure retrieval**

B. van Dierenhoven et al.

# Surface pressure retrieval from SCIAMACHY measurements in the O<sub>2</sub> A Band: validation of the measurements and sensitivity on aerosols

**B. van Dierenhoven, O. P. Hasekamp, and I. Aben**

SRON National Institute for Space Research, The Netherlands

Received: 7 February 2005 – Accepted: 22 February 2005 – Published: 14 March 2005

Correspondence to: B. van Dierenhoven (b.van.dierenhoven@sron.nl)

© 2005 Author(s). This work is licensed under a Creative Commons License.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

## Abstract

We perform surface pressure retrievals from cloud-free Oxygen A band measurements of SCIAMACHY. These retrievals can be well validated because surface pressure is a quantity that is, in general, accurately known from meteorological models. Therefore, surface pressure retrievals and their validation provide important insight into the quality of the instrument calibration. Furthermore, they can provide insight into retrievals which are affected by similar radiation transport processes, for example the retrieval of total columns of H<sub>2</sub>O, CO, CO<sub>2</sub> and CH<sub>4</sub>. In our retrieval aerosols are neglected. Using synthetic measurements, we show that for low to moderate or high surface albedos this leads to an under- or overestimation of the retrieved surface pressures, respectively. The surface pressures retrieved from the SCIAMACHY measurements indeed show this dependence on surface albedo, when compared to the corresponding pressures from a meteorological database. However, an offset of about 30 hPa was found, which can not be caused by neglecting aerosols in the retrieval. The same offset was found when comparing the retrieved surface pressures to those retrieved from co-located GOME Oxygen A band measurements. This implies a calibration error in the SCIAMACHY measurements. By adding an offset of 1% of the continuum reflectance at 756 nm to the SCIAMACHY reflectance measurements, this systematic bias vanishes.

## 1. Introduction

The satellite instrument SCIAMACHY (Bovensmann et al., 1999), launched March 2002 on ENVISAT, and its precursor GOME (Burrows et al., 1999), launched in 1995 on ERS-2, measure in the Oxygen A absorption band, near 760 nm. These measurements are of great importance for the retrieval of cloud properties (Koelemeijer et al., 2001; Rozanov and Kokhanovsky, 2004) and aerosol properties (Koppers et al., 1997). These retrievals assume the surface pressure is known. Alternatively, surface pressures can be retrieved from the Oxygen A band measurements (Barton and Scott,

## SCIAMACHY surface pressure retrieval

B. van Diedenhoven et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

1986; Bréon and Bouffiés, 1996; Vanbauce et al., 1998; Dubuisson et al., 2001; Ramon et al., 2004). Surface pressure retrievals can be used to scale total column retrievals of other species, e.g. H<sub>2</sub>O, CO, CO<sub>2</sub>, CH<sub>4</sub>, in order obtain volume mixing ratios (Buchwitz et al., 2004) or for light path corrections (Kuang et al., 2002). Furthermore, surface pressures retrievals can be well validated because surface pressure is a quantity that is, in general, accurately known from meteorological models. Therefore, surface pressure retrievals and their validation provide important insight into the quality of the instrument calibration (Dubuisson et al., 2001). Furthermore, they can provide insight into retrievals which are affected by similar radiation transport processes, for example the retrieval of total columns of H<sub>2</sub>O, CO, CO<sub>2</sub> and CH<sub>4</sub> from SCIAMACHY measurements. The most important potential error source for surface pressure retrievals from cloud-free measurements is scattering by aerosols (Bréon and Bouffiés, 1996; Dubuisson et al., 2001). As the relevant aerosol information is usually not available, surface pressures retrieved from the Oxygen A band neglecting aerosol scattering are normally referred to as apparent surface pressures.

In this paper we will retrieve apparent surface pressures from SCIAMACHY Oxygen A band measurements. To interpret the retrieved apparent surface pressures, we first study the effect of aerosols on the retrieval of surface pressure using simulated measurements. The surface pressures retrieved from SCIAMACHY measurements are validated with reference surface pressures from a meteorological model. Additionally, the retrieved surface pressures are compared with surface pressures retrieved from coinciding GOME Oxygen A band measurements. Measurements of GOME overlap very well with those of SCIAMACHY with only about a 30 min time lag, allowing a good comparison. Although SCIAMACHY has been calibrated on ground, this calibration needs to be validated in flight. Therefore, the results and validation of our retrievals are useful to identify calibration inaccuracies.

The paper is constructed as follows. First the retrieval method is presented in Sect. 2. The effect of aerosols on the surface pressure retrieval and on the Oxygen A band measurements is studied using simulated measurements in Sect. 3. Then, in Sect. 4,

---

**SCIAMACHY surface  
pressure retrieval**

 B. van Diedenhoven et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

EGU

the surface pressures retrieved from SCIAMACHY measurements are presented and compared with the findings from the simulated measurements and with surface pressures retrieved from GOME measurements. The discussion and conclusions follow in Sect. 5.

## 2. The retrieval method

For the retrieval of a state vector  $\mathbf{x}$  from a measurement vector  $\mathbf{y}$ , a forward model  $\mathbf{F}$  is needed that describes how  $\mathbf{y}$  depends on  $\mathbf{x}$ , i.e.

$$\mathbf{y} = \mathbf{F}(\mathbf{x}) + \mathbf{e}, \quad (1)$$

with error term  $\mathbf{e}$ . The measurement vector  $\mathbf{y}$  contains the reflectances measured at different wavelengths and the state vector  $\mathbf{x}$  contains the unknown parameters to be retrieved. For the surface pressure retrievals in this paper the state vector  $\mathbf{x}$  contains at least the following parameters: (1) the surface pressure, (2) the surface albedo at 756 nm and (3) the linear spectral dependence of the surface albedo. Additionally,  $\mathbf{x}$  may contain instrument parameters such as a wavelength shift and response function parameters.

Since SCIAMACHY and GOME are polarisation sensitive instruments, the intensity  $I_{pol}$  measured by a certain detector pixel at wavelength  $\lambda_i$  is not only affected by the intensity of the light that enters the instrument but also by its state of polarisation, viz.

$$I_{pol}(\lambda_i) = \bar{I}_{TOA}(\lambda_i) + m_{12}(\lambda_i)\bar{Q}_{TOA}(\lambda_i) + m_{13}(\lambda_i)\bar{U}_{TOA}(\lambda_i), \quad (2)$$

where  $m_{12}$  and  $m_{13}$  are elements of the instrument's Müller matrix normalised to its element (1,1). Furthermore,  $\bar{I}_{TOA}$ ,  $\bar{Q}_{TOA}$ , and  $\bar{U}_{TOA}$  are the elements of the intensity

vector  $\bar{I}_{TOA}$  defined by

$$\bar{I}_{TOA}(\lambda_i) = \int_0^{\infty} d\lambda S(\lambda_i, \lambda) I_{TOA}(\lambda), \quad (3)$$

where  $I_{TOA}(\lambda)$  is the intensity vector of the light at the entrance of the instrument. The spectral response function  $S$  is described by

$$S(\lambda_i, \lambda) = \frac{a_1^2}{((\lambda_i - \lambda)/\Delta_p)^4 + a_0^2}, \quad (4)$$

where  $\Delta_p$  is the width of the detector pixels, assumed to be 0.217 nm. The width of the response function is determined by  $a_0$ . For SCIAMACHY and GOME  $a_0$  is approximately 1.1772 and 0.7377, respectively. The normalisation factor  $a_1$  is determined by the requirement

$$\int_0^{\infty} S(\lambda_i, \lambda) d\lambda = 1. \quad (5)$$

Similarly to the earth radiances, SCIAMACHY and GOME also measure the solar irradiance  $F_o$  to obtain the polarisation sensitive reflectances  $R_{pol}$  defined by

$$R_{pol} = \frac{I_{pol}}{F_o}. \quad (6)$$

The common approach to account for the polarisation sensitivity of SCIAMACHY and GOME is to apply a polarisation correction to the radiances  $I_{pol}$  using the Polarisation Measurement Devices (PMDs). However, the broadband PMD measurements are not sufficient to correct for the polarisation sensitivity in spectral regions where the state of polarisation is varying rapidly with wavelength, as is the case for the Oxygen A band (Stam et al., 2000; Schutgens and Stammes, 2003). Moreover, the SCIAMACHY

---

[Title Page](#)
[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

PMDs are not yet well calibrated. In our retrieval approach these errors due to polarisation sensitivity are avoided, by using the polarisation sensitive reflectances  $R_{pol}$  as the elements of the measurement vector  $\mathbf{y}$ . This means that the forward model  $\mathbf{F}$  directly models the polarisation sensitive reflectances  $R_{pol}$ . This approach was introduced by Hasekamp et al. (2002) for the retrieval of Ozone profiles from GOME.

The main part of the forward model  $\mathbf{F}$  is an atmospheric radiative transfer model. In this study the SRON radiative transfer model (Hasekamp and Landgraf, 2002) is used, which includes polarisation and multiple scattering. A Rayleigh scattering atmosphere is assumed with 60 1 km thick layers. The absorption line parameters are taken from the HITRAN2000 spectroscopic database (Rothman et al., 2003) and a Voigt lineshape is assumed. The calculations are performed on a 0.02 nm spectral resolution. Figure 1 shows an example of the Oxygen A band at 0.02 nm resolution and at SCIAMACHY resolution. The Oxygen A band consists of 2 branches which are resolved in the SCIAMACHY resolution, i.e. the deep R branch around 761 nm and the broader P branch around 765 nm.

All retrievals are performed using iterative non-linear least squares fitting. For these fits, we use  $\ln(R)$  instead of  $R$ .

### 3. Retrieval from simulated measurements

Dubuisson et al. (2001), among others, have shown that aerosols significantly affect surface pressure retrievals using the Oxygen A band. Furthermore, they showed that the effects of aerosols depend on surface albedo. To study the effects of aerosols on the retrieval of surface pressure from SCIAMACHY, SCIAMACHY measurements are simulated for atmospheres with different aerosol loads and above several surface albedos. Subsequently, the apparent surface pressure from these simulated measurements are retrieved using a forward model with only Rayleigh scattering included. For these retrievals, the state vector  $\mathbf{x}$  (see Eq. 1) contains only the surface pressure and surface albedo and the linear spectral dependence of the surface albedo. The SCIA-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

MACHY measurements are simulated for a nadir viewing geometry with a solar zenith angle of  $40^\circ$  and the US standard atmosphere with a surface pressure of 1000 hPa. Desert dust aerosols are included of which the characteristics are given in Table 1. The optical aerosol properties are calculated using Mie theory.

Figures 2a and b show the difference between the true surface pressure and the retrieved apparent surface pressure ( $\Delta P = P_{true} - P_{apparent}$ ) as a function of retrieved apparent surface albedo, for different aerosol optical thicknesses and height distributions, respectively. Although depending on aerosol optical thickness, height distribution and type and geometry, two general effects are observed: (1) For low and moderate surface albedos the surface pressures are underestimated in the retrieval. This is due to the relatively high contribution to the measured radiances by photons that are scattered back by the aerosol layer, reducing the photon path. Furthermore, most of the light penetrating through the aerosol layer is absorbed by the surface. This effect was also identified by Dubuisson et al. (2001). (2) At large surface albedos, the surface pressures are, in general, overestimated in the retrieval. Here, most photons penetrating through the aerosol layer are reflected by the surface and travel through the aerosol layer once more. Due to (multiple) scattering in the aerosol layer the photon path is enhanced, resulting in an overestimation of the surface pressure. This effect was not previously recognised by Dubuisson et al. (2001).

Furthermore, the results show that the effects of aerosols on the surface pressure retrieval increase with increasing aerosol optical thickness, while the aerosol height distribution only significantly impacts the surface pressure retrieval above low albedos.

The spectral fitting residuals between the simulated measurements for an atmosphere including aerosols and the fitted reflectances ignoring aerosols are shown in Fig. 3a and b, for different aerosol optical thicknesses and height distributions, respectively. The residuals show a large negative peak in the centre of the deep R branch and positive values in the P branch. The amplitude of the residuals increases with increasing aerosol optical thickness (Fig. 3a) and increasing height of the bulk of the aerosols (Fig. 3b). Furthermore, it depends on type of the aerosols and slightly on the surface

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

albedo (not shown). The shape of the residuals, however, is rather characteristic.

Figures 2 and 3 show a clear dependence of both the retrieved  $\Delta P$  and the spectral fitting residual on the aerosol height distribution and optical thickness. This implies that, when the true surface pressure and the aerosol type are known, information about the aerosol optical thickness and height distribution can be retrieved from the Oxygen A band measurements. This was previously demonstrated by Koppers et al. (1997).

In addition to aerosols, Rayleigh scattering also has a large effect on the surface pressure retrievals, although the Rayleigh scattering optical thickness at these wavelengths is relatively low ( $\sim 0.025$ ). This effect is illustrated in Fig. 4, which shows  $\Delta P$  retrieved while neglecting all scattering from simulated SCIAMACHY measurements for atmospheres including aerosols and, additionally, for a clear-sky atmosphere including only Rayleigh scattering. In the clear-sky case, neglecting Rayleigh scattering leads to an underestimation of the surface pressure above all surface albedos. For aerosol loaded atmospheres, a comparison of Fig. 4 with Fig. 2 shows that, for low albedos, the effect of neglecting Rayleigh scattering dominates the effect of neglecting aerosols for a low aerosol optical thickness. Furthermore, for large albedos the effect of neglecting Rayleigh scattering can partly compensate for the effect of neglecting aerosols. The large effect of Rayleigh scattering can be explained by the fact that the Oxygen A band contains a large number of optically thick absorption lines (see Fig. 1) for which few photons penetrate through the atmosphere to high pressure levels and thus the reflectance at the wavelengths of these absorption lines is mainly determined by Rayleigh scattering occurring at low pressure levels. Therefore, neglecting Rayleigh scattering in the forward model leads to an increase in the depth of these absorption lines which leads to an underestimation of the surface pressure.

The simulations in this section show aerosols significantly affect the Oxygen A band measurements. When retrieving surface pressure neglecting aerosol scattering, characteristic residuals as shown in Fig. 3 are expected. Furthermore, surface pressures retrieved using the Oxygen A band are expected to underestimate the actual surface pressure at low and moderate albedos and overestimate them at high albedos.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

## 4. Retrieval from SCIAMACHY measurements

### 4.1. Data

For this study we analyse the reflectances measured by SCIAMACHY and GOME above southern Europe and northern Africa on 23 January 2003. False colour images of the data used are shown in Fig. 5. In SCIAMACHY Nadir and Limb measurements are alternated. Since for this study only Nadir measurements are used, the SCIAMACHY orbits are broken up into several blocks, generally referred to as states. The characteristics of the Oxygen A band measurements for both instruments are summarised in Table 2.

The SCIAMACHY reflectances (level 1b data) used in this study are improved by replacing calibration data using SCIA\_PATCH\_NL1<sup>1</sup>. For both instruments all calibration options are applied, with the exception of the polarisation sensitivity correction, which is not needed because we directly model the polarisation sensitive measurement (see Sect. 2). Furthermore, on the reflectances measured by SCIAMACHY a multiplicative correction factor of 1.2 is applied. This correction factor was found by Acarreta et al. (2004) and Acarreta and Stammes (2005) by comparing the measured reflectances with those measured by MERIS, also on ENVISAT.

Cloudy pixels are excluded from the data. To identify clouds in SCIAMACHY measurements, we use the PRICE method developed by Krijger et al. (2005) using the PMD measurements. For GOME, an adapted version is used.

As reference surface pressures for validation, the sea-level pressure data from the UKMO Stratospheric Assimilated dataset are combined with the TerrainBase surface elevation database. The UKMO dataset has a spatial resolution of  $2.5 \times 3.75^\circ$  and the grid-box containing the centre of the instrument footprint is taken. The TerrainBase data has a resolution of  $5 \times 5$  arc-minutes and is averaged over the instrument footprint. To calculate the absorption cross-sections, the temperature profiles are taken from the

<sup>1</sup>Available at <http://www.sron.nl/~hees/SciaDC>.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## 4.2. Retrieval results

To retrieve apparent surface pressures from SCIAMACHY measurements, the state vector  $x$  (Eq. 1) contains (1) the surface pressure, (2) the surface albedo at 756 nm, (3) the linear spectral dependence of the surface albedo, (4) a wavelength shift  $\Delta\lambda$  and (5) the response function parameter  $a_0$  (see Eq. 4). The instrument parameters (4 and 5) did not significantly interfere with the other retrieved parameters and are very similar for each retrieval. The mean and standard deviation of the retrieved instrument parameters are given in Table 3.

Figure 6 shows the absolute differences between retrieved apparent surface pressures and the reference surface pressures for the SCIAMACHY data ( $\Delta P = P_{reference} - P_{apparent}$ ). Figure 7 shows  $\Delta P$  as a function of the retrieved surface albedo. Much variation is seen, ranging from  $\Delta P = 300$  to  $-60$ . Clearly, systematic differences are seen between retrievals above land and sea, i.e. high and low surface albedos. In general,  $\Delta P$  is positive for albedos lower than 0.1 and negative for higher surface albedos. The few low surface pressures retrieved above high surface albedos are probably due to clouds not flagged by our cloud filter.

Also shown in Fig. 7 are results from the simulated measurements as presented in Fig. 2, for an aerosol optical thickness of 0.1 and 0.3. The retrieved values match the results from the simulated measurements well for low albedos ( $<0.1$ ). However, for higher albedos, the retrieved values for  $\Delta P$  are systematically about 30 hPa lower. Several other aerosol types and height distributions were tried, none of which match these retrievals.

For comparison, surface pressures retrieved from GOME measurements are shown in Fig. 8. Similar variations as in the SCIAMACHY retrievals of Figs. 6 and 7 are seen. However, the values of  $\Delta P$  are systematically about 30 hPa higher than those retrieved from SCIAMACHY and are more in line with the simulations presented in Sect. 3. Thus, it seems likely that the surface pressures retrieved from SCIAMACHY suffer from a

## SCIAMACHY surface pressure retrieval

B. van Diedenhoven et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

systematic error(s), probably due to a calibration error(s) in the measurements.

We investigated the effect of several SCIAMACHY calibration errors on the retrieved surface pressures. First, it was found that changing the multiplicative factor applied on the measured reflectances (Acarreta et al., 2004, see Sect. 4.1) only minimally affects the retrieved surface pressure. The reason for this is that the “relative” depth of the Oxygen A band contains the surface pressure information. On the other hand, the retrieved surface pressures are affected when an offset is added to the measured reflectances. Several offsets, defined as a percentage of the continuum reflectance around 756 nm (after multiplication by 1.2), were applied and the effect on the retrievals was evaluated. A linear dependence of  $\Delta P$  with this offset is found. Adding an offset of 1% of the continuum reflectance around 756 nm to the reflectances, increases  $\Delta P$  by approximately 30 hPa. Figure 9 shows  $\Delta P$  as a function of surface albedo when this correction is applied. With this correction, the retrievals behave as predicted by theory (see Sect. 3) and compare well with the GOME retrievals shown in Fig. 8b. The same correction was found by evaluating SCIAMACHY data from other geolocations and dates. This offset is probably related to the inaccuracies in the reflectance calibration (Acarreta et al., 2004; Noël, 2004).

The different colours in Fig. 9 correspond to different SCIAMACHY states, i.e. different geolocations (see Figs. 6 and 7). Significant differences in the retrieved values of  $\Delta P$  between states are seen. As concluded from the retrievals using simulated measurements shown in Fig. 2a, it is likely that these differences are due to the varying aerosol optical thickness. In particular, the retrievals from data from the most northern state of orbit 4700 (in black) match well with results from simulated measurements corresponding to an aerosol optical thickness of 0.05–0.2, while the most southern state of orbit 4700 (in red) matches better with results from simulated measurements for an aerosol optical thickness of 0.2–0.6. This increase in aerosol optical thickness at the location of the most southern state is confirmed by the aerosol optical thickness retrieved by the MISR instrument on EOS Terra (Diner et al., 2001) on the same day, as shown in Fig. 10. However, the absolute values of the MISR optical thicknesses do

---

**SCIAMACHY surface  
pressure retrieval**B. van Diedenhoven et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

not match those derived from Fig. 9. As is illustrated in Fig. 2b, this can be due to an incorrect assumption about the aerosol height distribution in our forward model. Also, the true aerosol type can be different than the one assumed. Furthermore, the MISR optical thicknesses are given at a wavelength of 558 nm, while we use a wavelength of 765 nm. However, our results are in qualitative agreement with the aerosol optical thickness retrievals from MISR.

### 4.3. Spectral fitting residuals

A closer inspection of the spectral fitting residuals yields valuable information. Figure 11 shows residuals from a SCIAMACHY and a GOME measurement above approximately the same location with an apparent surface albedo of  $\sim 0.16$ . The residuals are very much alike. Also shown is a residual from a simulated measurement similar to those shown in Fig. 3. The residual from the simulated measurement is very similar to those from the real SCIAMACHY and GOME measurements. This confirms that the spectral fitting residuals are for a large part caused by neglecting aerosols in our retrieval.

However, also some high frequency scatter of about 2–5% is seen in the residuals from the SCIAMACHY and GOME measurements, which are not observed in the simulations and can thus not be caused by neglecting aerosols in the retrieval. Again these features are very common in all our spectral fitting residuals. They might be due to errors in the spectroscopy data (Chance, 1997; Rothman et al., 2003) or the instrument response functions. Furthermore, solar Fraunhofer lines (e.g. at 766.5 nm) could appear in the reflectances due to calibration errors.

## 5. Conclusions and discussion

In this paper we presented surface pressure retrievals from cloud-free SCIAMACHY Oxygen A band measurements. The retrievals were performed using a radiative trans-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

fer model that fully includes multiple Rayleigh scattering and polarisation but does not include the effect of aerosols. Using synthetic measurements, the effects of neglecting aerosols on the retrieved surface pressures were investigated. It was found that for low and moderate surface albedo aerosols lead to an underestimation of the retrieved surface pressures due to a reduction of the light path. For high surface albedos, scattering by aerosols leads to an enhancement of the light path and thus to an overestimation of the surface pressure. The magnitude of these effects depends on aerosol optical thickness and on the aerosol height distribution. It was also observed that aerosols cause a characteristic spectral feature in the fitting residuals, of which the amplitude also depends on the aerosol optical thickness and height distribution.

The apparent surface pressures retrieved from SCIAMACHY were validated with reference surface pressures from a meteorological model. The difference between the SCIAMACHY apparent surface pressures and the reference surface pressure showed a dependence on surface albedo that can be explained by the fact that aerosols are neglected in the retrieval. However, also a systematic offset of about 30 hPa was found that cannot be explained by the effect of aerosols. The same offset was found by comparing the surface pressures retrieved from SCIAMACHY with surface pressures retrieved from co-located GOME measurements. This indicates the presence of a calibration error in the SCIAMACHY measurements. Adding an offset of 1% of the continuum reflectance at 756 nm to the SCIAMACHY reflectance measurements removes this systematic bias. The surface pressures retrieved then also agree well with the corresponding GOME retrievals. The fitting residuals found for the GOME and SCIAMACHY measurements are very similar and both show the characteristic spectral features due to neglecting aerosols in the retrieval, as was also found in the study using synthetic measurements.

The clear dependence of both the retrieved surface pressure and the spectral fitting residuals on the aerosol optical thickness and height distribution, as shown in this paper, indicates that both parameters may be retrieved from Oxygen A band measurements, when one assumes the surface pressure as known. This was previously

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

demonstrated by [Koppers et al. \(1997\)](#).

For similar retrievals of total columns of other species than O<sub>2</sub>, e.g. H<sub>2</sub>O, CO, CO<sub>2</sub> and CH<sub>4</sub>, similar effects due to aerosols as discussed in this paper, i.e. under- or over-estimation of the total column, can be expected ([Dubuisson et al., 2004](#), [Houweling et al., 2005](#)<sup>2</sup>). The retrieved total column of these species are often scaled to oxygen to correct for the light path ([Noël et al., 1999](#); [O'Brien and Rayner, 2002](#); [Kuang et al., 2002](#); [Dufour et al., 2004](#)) or to obtain volume mixing ratios of the species ([Buchwitz et al., 2004](#)). As pointed out by [O'Brien and Rayner \(2002\)](#), the light paths are affected by the height distribution and optical thickness of the absorption lines of individual species, which needs to be considered when applying such a scaling. Furthermore, results shown in this paper imply that care must be taken when using the Oxygen A band for scaling of retrievals at other wavelengths, because the effects of aerosols are shown to depend significantly on surface albedo, which is spectral dependent. Additionally, the optical thickness of Rayleigh scattering and therewith its relative contribution to the total scattering varies with wavelength as well. These problems can be avoided by retrieving aerosol properties simultaneously with the total column of the trace gas, the surface albedos and the surface pressure ([Kuang et al., 2002](#)). To conclude, we showed that surface pressure retrievals can be used to validate the calibration of the Oxygen A band measurements. This approach could also be used to validate the calibration of the SCIAMACHY measurements of other oxygen bands, i.e. O<sub>2</sub> B and γ and O<sub>4</sub> bands.

*Acknowledgements.* SCIAMACHY and GOME data were provided by ESA. The SCIAMACHY data were processed using SCIA\_PATCH\_NL1, available at <http://www.sron.nl/~hees/SciaDC>. The MISR data were obtained from the NASA Langley Research Center Atmospheric Sciences Data Center. Part of this work was supported by the Dutch User Support Programme 2001–2005 (USP) under project number EO-069.

<sup>2</sup>Houweling, S., Hartmann, W., Aben, I., Schrijver, J., Skidmore, J., and Bréon, F.-M.: Evidence of systematic errors in Sciamachy observed CO<sub>2</sub> due to aerosols, Atmos. Chem. Phys. Discuss., in preparation, 2005.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## References

- Acarreta, J. R. and Stammes, P.: Calibration comparison between SCIAMACHY and MERIS on board ENVISAT, *IEEE Geosci.*, 2, 31–35, 2005. [1477](#)
- Acarreta, J. R., Stammes, P., and Tilstra, L. G.: Reflectance Comparison Between SCIAMACHY and MERIS, in: *ESA SP-562: Atmospheric Chemistry Validation of ENVISAT (ACVE-2)*, 2004. [1477](#), [1479](#)
- Barton, I. J. and Scott, J. C.: Remote measurements of surface pressure using absorption in the Oxygen A band, *Appl. Optics*, 25, 3502–3507, 1986. [1470](#)
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noel, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, 1999. [1470](#)
- Bréon, F. and Bouffies, S.: Land Surface Pressure Estimate from Measurements in the Oxygen A Absorption Band, *J. Appl. Met.*, 35, 69–77, 1996. [1471](#)
- Buchwitz, M., de Beek, R., Burrows, J. P., Bovensmann, H., Warneke, T., Notholt, J., Meirink, J. F., Goede, A. P. H., Bergamaschi, P., Körner, S., Heimann, M., Muller, J. F., and Schulz, A.: Atmospheric methane and carbon dioxide from SCIAMACHY satellite data: initial comparison with chemistry and transport models, *Atmos. Chem. Phys. Discuss.*, 4, 7217–7279, 2004,  
[SRef-ID: 1680-7375/acpd/2004-4-7217](#). [1471](#), [1482](#)
- Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstätter-Weissenmayer, A., Richter, A., Debeek, R., Hoogen, R., Bramstedt, K., Eichmann, K., Eisinger, M., and Perner, D.: The Global Ozone Monitoring Experiment (GOME): Mission Concept and First Scientific Results, *J. Atmos. Sci.*, 56, 151–175, 1999. [1470](#)
- Chance, K.: Improvement of the O<sub>2</sub> A band spectroscopic database for satellite-based cloud detection, *J. Quant. Spectrosc. Radiat. Transfer*, 58, 375–378, 1997. [1480](#)
- Diner, D., Abdou, W., Ackerman, T., Crean, K., Gordon, H., Kahn, R., Martonchik, J., McMurdock, S., Paradise, S., Pinty, B., Verstraete, M., Wang, M., and West, R.: Level 2 Aerosol Retrieval, *ATBD-MISR-09*, 2001. [1479](#)
- Dubuisson, P., Borde, R., Schmechtig, C., and Santer, R.: Surface pressure estimates from satellite data in the Oxygen A-band: Applications to the MOS sensor over land, *J. Geophys. Res.*, 106(D21), 27 277–27 286, 2001. [1471](#), [1474](#), [1475](#)
- Dubuisson, P., Dessailly, D., Vesperini, M., and Frouin, R.: Water vapor retrieval over ocean us-

---

### SCIAMACHY surface pressure retrieval

B. van Diedenhoven et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

ing near-infrared radiometry, *J. Geophys. Res.*, 109, D19106, doi:10.1029/2004JD004516, 2004. [1482](#)

Dufour, E., Bréon, F., and Peylin, P.: CO<sub>2</sub> column averaged mixing ratio from inversion of ground-based solar spectra, *J. Geophys. Res.*, 109, D09304, doi:10.1029/2003JD004469, 2004. [1482](#)

Hasekamp, O. P. and Landgraf, J.: A linearized vector radiative transfer model for atmospheric trace gas retrieval, *J. Quant. Spectrosc. Radiat. Transfer*, 75, 221–238, 2002. [1474](#)

Hasekamp, O. P., Landgraf, J., and van Oss, R.: The need of polarization modeling for ozone profile retrieval from backscattered sunlight, *J. Geophys. Res.*, 107(D23), doi:10.1029/2002JD002387, 2002. [1474](#)

Koelemeijer, R. B. A., Stammes, P., Hovenier, J. W., and de Haan, J. F.: A fast method for retrieval of cloud parameters using Oxygen A band measurements from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, 106(D4), 3475–3490, doi:10.1029/2000JD900657, 2001. [1470](#)

Koppers, G. A. A., Jansson, J., and Murtagh, D. P.: Aerosol optical thickness retrieval from GOME satellite data in the Oxygen A band, in: *ESA SP-414: Third ERS Symposium on Space at the service of our Environment*, 693–696, 1997. [1470](#), [1476](#), [1482](#)

Krijger, J., Aben, I. and Schrijver, H.: Distinction between clouds and ice/snow covered surfaces in the identification of cloud-free observations using SCIAMACHY PMDs, *Atmos. Chem. Phys. Discuss.*, 5, 815–845, 2005, [SRef-ID: 1680-7375/acpd/2005-5-815](#). [1477](#)

Kuang, Z., Margolis, J., Toon, G., Crisp, D., and Yung, Y.: Spaceborne measurements of atmospheric CO<sub>2</sub> by high-resolution NIR spectrometry of reflected sunlight: An introductory study, *Geophys. Res. Lett.*, 29(15), doi:10.1029/2001GL014298, 2002. [1471](#), [1482](#)

Noël, S.: Determination of correction factors for SCIAMACHY radiances and irradiances, Technical Report IFE-SCIA-SN-20040514, University of Bremen, Germany, 2004. [1479](#)

Noël, S., Buchwitz, M., Bovensmann, H., Hoogen, R., and Burrows, J. P.: Atmospheric water vapor amounts retrieved from GOME satellite data, *Geophys. Res. Lett.*, 26, 1841–1844, 1999. [1482](#)

O'Brien, D. M. and Rayner, P. J.: Global observations of the carbon budget, 2, CO<sub>2</sub> column from differential absorption of reflected sunlight in the 1.61 μm band of CO<sub>2</sub>, *J. Geophys. Res.*, 107(D18), doi:10.1029/2001JD000617, 2002. [1482](#)

Ramon, D., Santer, R., Dilligeard, E., Jolivet, D., and Vidot, J.: MERIS Land Products Validation,

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

in: ESA SP-549: Proceedings of MERIS User Workshop, 10.1, 2004. [1471](#)

Rothman, L. S., Barbe, A., Benner, D. C., Brown, L. R., Camy-Peyret, C., Carleer, M. R., Chance, K., Clerbaux, C., Dana, V., Devi, V. M., Fayt, A., Flaud, J.-M., Gamache, R. R., Goldman, A., Jacquemart, D., Jucks, K. W., Lafferty, W. J., Mandin, J.-Y., Massie, S. T.,  
5 Nemtchinov, V., Newnham, D. A., Perrin, A., Rinsland, C. P., Schroeder, J., Smith, K. M.,  
Smith, M. A. H., Tang, K., Toth, R. A., Vander Auwera, J., Varanasi, P., and Yoshino, K.: The  
HITRAN molecular spectroscopic database: edition of 2000 including updates through 2001,  
J. Quant. Spectrosc. Radiat. Transfer, 82, 5–44, 2003. [1474](#), [1480](#)

10 Rozanov, V. V. and Kokhanovsky, A. A.: Semianalytical cloud retrieval algorithm as applied  
to the cloud top altitude and the cloud geometrical thickness determination from top-of-  
atmosphere reflectance measurements in the Oxygen A band, J. Geophys. Res., 109(D5),  
doi:10.1029/2003JD004104, 2004. [1470](#)

Schutgens, N. A. J. and Stammes, P.: A novel approach to the polarization correction of space-  
borne spectrometers, J. Geophys. Res., 108(D7), doi:10.1029/2002JD002736, 2003. [1473](#)

15 Stam, D. M., De Haan, J. F., Hovenier, J. W., and Aben, I.: Detecting radiances in the O<sub>2</sub> A  
band using polarization-sensitive satellite instruments with application to the Global Ozone  
Monitoring Experiment, J. Geophys. Res., 105(D17), 22 379–22 392, 2000. [1473](#)

Torres, O., Decea, R., Veefkind, P., and de Leeuw, G.: OMI aerosol Retrieval Algorithm, ATBD-  
OMI-03, 2001. [1486](#)

20 Vanbauce, C., Buriez, J. C., Parol, F., Bonnel, B., Sèze, G., and Couvert, P.: Apparent pressure  
derived from ADEOS-POLDER observations in the Oxygen A-band over ocean, Geophys.  
Res. Lett., 25, 3159–3162, 1998. [1471](#)

---

**SCIAMACHY surface  
pressure retrieval**

 B. van Dierenhoven et al.
 

---

**Table 1.** Characteristics at 765 nm of the two modes of the desert dust aerosols used in this study, based on aerosol model DL from Torres et al. (2001). A log-normal size distribution is assumed. The relative contribution of the modes to the number concentration is represented by the fraction. The aerosols are evenly distributed over the first 2 km and decrease in number as  $P^3$  from 2 to 12 km, unless indicated otherwise.

mode	Effective radius	Size variance	Refr. index		fraction
			real	imag.	
1	0.105 $\mu\text{m}$	0.323 $\mu\text{m}^2$	1.53	0.001	0.99565
2	1.605 $\mu\text{m}$	0.418 $\mu\text{m}^2$	1.53	0.001	0.00435

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

**Table 2.** Characteristics of the SCIAMACHY and GOME measurements in the Oxygen A band from 755–775 nm.

	SCIAMACHY	GOME
Channel	4	4
Spectral resolution	~0.4 nm	~0.25 nm
Spectral sampling	~0.22 nm	~0.22 nm
Footprint size	30 × 60 km <sup>2</sup>	320 × 40 km <sup>2</sup>

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)


EGU

**Table 3.** Mean and standard deviation of the retrieved wavelength shifts  $\Delta\lambda$  and response function parameters  $a_0$ .

	mean	standard deviation
$\Delta\lambda$	-0.0772 nm	0.00115 nm
$a_0$	1.09573	0.0356539

**SCIAMACHY surface  
pressure retrieval**

B. van Diedenhoven et al.




**Fig. 1.** Example of an Oxygen A band reflectance spectrum at 0.02 nm spectral resolution (black) and at SCIAMACHY resolution (red).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

**SCIAMACHY surface  
pressure retrieval**

B. van Dierenhoven et al.



**Fig. 2.** The difference between the true surface pressure used in the simulated measurements including aerosols and the apparent surface pressure retrieved ignoring aerosols,  $\Delta P = P_{true} - P_{apparent}$ , as a function of retrieved apparent surface albedo. The true surface albedos are 0, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5 and 0.6. The retrieved values are indicated by pluses. Panel (a) shows the retrieved values for different values of the total aerosol optical thickness  $\tau$  at 765 nm. To obtain  $\tau$  at the more common wavelength 550 nm the values have to be multiplied by 1.11. Panel (b) shows the retrieved values for different height distributions, when an aerosol optical thickness of 0.2 is taken. Here, our standard aerosol distribution (see Table 1) is modified such that the lower layer with evenly distributed aerosols is extended to higher levels, indicated by the layer top.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU




**Fig. 3.** Residuals between the simulated reflectance measurements  $R_{meas}$  for an atmosphere including aerosols and the fitted reflectances  $R_{mod}$  ignoring aerosols. The residuals are defined as  $(R_{mod} - R_{meas}) / R_{meas} \times 100\%$ . A surface albedo of 0.1 is used. The residuals are shown for different values of the aerosol optical thickness **(a)** and different height distributions **(b)**, similar to Fig. 2

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.




**Fig. 4.** Similar to Fig. 2a, but when  $\Delta P$  and the apparent surface albedo are retrieved while neglecting all scattering. The solid black line shows results of retrievals from simulated measurements for a clear-sky case including only Rayleigh scattering.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.




**Fig. 5.** False colour images from SCIAMACHY (left) and GOME (right) measurements. For this study data from SCIAMACHY orbits 4700 (West) and 4699 (East) and GOME orbits 30 123 106 (West) and 30 123 092 (East) are used. The PMDs of SCIAMACHY and GOME used to produce these images measure in different wavelength ranges, resulting in different colouring.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.




**Fig. 6.** The absolute differences between the reference surface pressures and the retrieved apparent surface pressures ( $\Delta P = P_{reference} - P_{apparent}$ ) for the SCIAMACHY measurements from orbit 4700 (west) and 4699 (east).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval


B. van Diedenhoven et al.



**Fig. 7.**  $\Delta P$  as a function of apparent surface albedo for SCIAMACHY orbits 4700 (left) 4699 (right). Results from the 5 individual SCIAMACHY states per orbit as seen in Fig. 6 are colour-coded in black, blue, green, yellow and red going from north to south. Two results from simulated measurements for atmospheres with aerosol optical thicknesses of 0.1 and 0.3, as presented in Fig. 2, are plotted with dashed lines.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU




**Fig. 8.**  $\Delta P$  retrieved by the GOME instrument, as a map-projection **(a)** and as function of apparent surface albedo **(b)**, similar to Figs. 6 and 7, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.




**Fig. 9.** Same as Fig. 7, but when the SCIAMACHY reflectances are corrected by adding an offset of 1% of the continuum reflectance to the reflectances.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.




**Fig. 10.** Aerosol optical thicknesses at 558 nm retrieved using the MISR instrument on EOS Terra on 23 January 2003. Data from MISR orbits 16 486 (west) and 16 485 (east) are shown. These data were obtained from the NASA Langley Research Centre Atmospheric Sciences Data Centre. The contours of the SCIAMACHY states are indicated in red.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

SCIAMACHY surface  
pressure retrieval

B. van Diedenhoven et al.



**Fig. 11.** Residuals of spectral fits of a SCIAMACHY (black) and a GOME (red) Oxygen A band measurement with a nadir viewing geometry and a solar zenith angle of approximately  $40^\circ$ . Both measurements are taken at approximately  $13^\circ$  latitude and  $-10^\circ$  longitude and have an apparent surface albedo of  $\sim 0.16$ . Note that on the SCIAMACHY measurements the correction suggested in Sect. 4.2 is applied. In blue, a residual from a simulated SCIAMACHY measurement, similar to those in Fig. 3, is shown. For this, an atmosphere including aerosols with an optical thickness of 0.45 and a layer top of 9 km (see Fig. 2) is taken. The true surface pressure and apparent surface albedo are similar to those in the SCIAMACHY and GOME measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU