

Temperature
climatology and
trend estimates over
Durban, South Africa

H. Bencherif et al.

Temperature climatology and trend estimates in the UTLS region as observed over a southern subtropical site, Durban, South Africa

H. Bencherif¹, R. D. Diab², T. Portafaix¹, B. Morel¹, P. Keckhut³, and A. Moorgawa²

¹Laboratoire de l'Atmosphère et des Cyclones, UMR-CNRS 8105, Université de La Réunion, B.P 7151, Saint-Denis, Reunion Island, France

²University of KwaZulu-Natal, Durban 4041, South Africa

³Service d'Aéronomie, UMR-CNRS 7620, B.P. 3, Verrières-le-Buisson, France

Received: 20 September 2005 – Accepted: 29 November 2005 – Published: 20 February 2006

Correspondence to: H. Bencherif (hassan.bencherif@univ-reunion.fr)

© 2006 Author(s). This work is licensed under a Creative Commons License.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Abstract

Temperature trends in the UTLS region are under-reported, particularly in the Southern Hemisphere, and yet temperature is one of the most important indicators of changes in dynamical and radiative processes in the atmosphere. Here radiosonde data from Durban, South Africa (30.0° S, 30.9° E) over the period 1980 to 2001 (22 years) between 250 and 20 hPa are used to derive a mean temperature climatology and to determine trends. The seasonal cycle at the 250-hPa level is anti-correlated with the seasonal cycles at the 150-hPa and 100-hPa heights. The 100-hPa level (local tropopause) exhibits a minimum temperature in late summer and a maximum in winter, and closely corresponds to previous results for tropical regions. Based on a Fourier analysis, both the annual cycle (AO) and the semi-annual cycle (SAO) are dominant, although the former is about 4 times stronger. The AO is strongest at the 100-hPa height. A trend analysis reveals a cooling trend at almost all heights in the UTLS region, with a maximum cooling rate of 1.09 ± 0.27 K per decade, at 70-hPa. Cooling rates are in good agreement with other studies and are slightly higher in summer than in winter.

1. Introduction

Temperature controls the rates of chemical reactions and thus ozone abundance, and is one of the most important parameters in terms of its influence on dynamical and radiative processes in the terrestrial atmosphere, particularly in the upper troposphere – lower stratosphere (UTLS) region. The UTLS is a region where significant changes are taking place that have the potential to perturb the Earth's climate in the coming decades.

Since dynamical activity is closely linked to temperature, temperature trends are a useful indicator of changes in dynamical processes and can contribute to a better understanding thereof. Indeed, any change in activity of gravity waves, equatorial Kelvin waves and Rossby planetary waves may have a consequent effect on mass and energy

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

fluxes, including stratosphere-troposphere exchanges as well as tropical-midlatitude exchanges.

5 Relatively few studies of trends in temperature in the UTLS region have been undertaken. One of the most complete studies is that of Oort and Liu (1993). Their study is based on the global rawinsonde network of more than 700 stations, of which 70% are in the NH and about 50% in the 30–60° N latitude band. Only slightly more than 15% of the stations are situated in the Southern Hemisphere (SH).

10 Most of the temperature trend studies that have been performed for the 25–70 km altitude range, have used Indian, American, Japanese and Russian rocket-sondes (Mohanakumar, 1994; Keckhut et al., 1999a, b; Kokin and Lysenko, 1994), and French lidar measurements (Hauchecorne et al., 1991; Keckhut et al., 1995). Some authors have performed trend analyses from the large ozonesonde databases (Logan, 1994; Harris et al., 1997), but again, these are predominantly available for the NH.

15 Within this context, this study focuses on a climatological and trend analysis of temperature, in a subtropical region where data are sparse. Regular upper-air radiosonde data recorded by the South African Weather Service (SAWS) over Durban (30.0° S, 30.9° E), situated on the east coast of South Africa, are used as the basis of the temperature climatology and trend analysis.

20 Theoretical studies, transport models and in situ observations have shown that at subtropical latitudes, vertical and horizontal exchange processes are expected and have been observed (Grant et al., 1994; Holon et al., 1995; Vaughan et Timmis, 1998; O'Connor et al., 1999). Furthermore, recent observational studies have shown that in the southern subtropics, within the UTLS region, many dynamical processes, such as subtropical tropopause breaks (Baray et al., 1998), planetary-scale tropopause fold events (Baray et al., 2000) and isentropic horizontal exchanges in the low stratosphere
25 through the southern subtropical barrier (Bencherif et al., 2003; Portafaix et al., 2003) take place. These processes play an important role in the transfer of energy, mass and tracers between the tropics and the mid-latitudes.

Temperature trend analyses in tropical regions based on rocket data (Keckhut et al.,

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

1999) have revealed a significant long-term cooling (1 to 3 K per decade), greater by a factor of 2, than model predictions. In general, Stratospheric Temperature Trend Assessments performed as part of the SPARC/WCRP project (Ramaswamy et al., 1999) and the WMO (1999) assessment have noted that numerical models predict a smaller cooling in the stratosphere than that detected in observed data series.

The outline of this paper is as follows. Details on the data and data analysis are presented in the next section. Section 3 presents the variability and trends of temperature and its climatology over Durban. In the last section our results are discussed and compared with previous observational analyses.

2. Data set

The data used in this study are daily radiosonde temperature measurements recorded at the Durban international airport (30.0° S, 30.9° E) by the SAWS for the 22-year period from January 1980 to December 2001. The study focuses on temperature time series recorded at 8 pressure levels, i.e., 250, 200, 150, 100, 70, 50, 30 and 20 hPa, corresponding approximately to the heights of 10, 12, 14, 16, 18, 20, 23 and 26 km, respectively. Hereafter we will refer to levels 250, 200 and 150 hPa as tropospheric, and levels with pressure less than 100 hPa as stratospheric. According to Randel et al. (2000), based on ECMWF, UKMO and NCEP data, the 100-hPa height tracks nearly identically the tropopause height. Hence the 100-hPa level will be referred to as the tropopause level.

The SAWS undertakes regular radiosonde launches twice a day, at approximately the same time (within ± 15 min), in the early morning and afternoon, i.e., $\cong 01:00$ and $\cong 12:35$ LT, respectively. In order to limit tidal biases, only afternoon data were used. Moreover, data quality checks were performed and all unrealistic readings were removed. As expected, the percentage of reliable afternoon data decreases with increasing height, and is 96.6%, 96.3%, 95.9%, 94.7%, 91.3%, 86.1%, 73.1%, 54.8%, at 250, 200, 150, 100, 70, 50, 30 and 20 hPa levels, respectively. No adjustments or

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

interpolations were made.

Many authors have emphasised that changes in instruments may lead to apparent discontinuities in temperature records with magnitudes that can be as large as the observed upper-air trends, notably in the tropics and SH (Angell, 1988; Gaffen, 1994).

5 The introduction of new types of radiosondes is one of the potential sources of data heterogeneity, and should be taken into account. For the studied location of Durban, no instrumental change was reported during the 1980–2001 period according to the SAWS archives. The Väisälä RS80-15 model was used throughout the period.

3. Results

10 3.1. Climatological temperatures

For both climatology and trend assessments data have been reduced to monthly averages at each of the pressure levels (see Fig. 1). Mean monthly climatological temperature values, as depicted in Figs. 2a and 2b, were calculated from more than 520 daily values. All the mean monthly climatological temperatures and corresponding standard deviations are given in Table 1.

15 Figure 2 shows that the seasonal variations of temperature at the 150-hPa and local tropopause (100-hPa) heights are quite similar, but are anti-correlated with the 250-hPa temperature variations. The minimum temperature at the tropopause is about 199 K, and appears during austral summer in February; while the maximum temperature (about 208 K) at the tropopause is obtained during austral winter (June–August).
20 Even though Durban is a subtropical site, the observed annual cycle at the tropopause level is very similar to results obtained by Randel et al. (2000) for tropical regions. Based on radiosonde and NCEP data they found that the tropical tropopause exhibited a minimum temperature during austral summer and a maximum during austral winter.

25 As suggested visually by the temperature time-height cross-section shown in Fig. 1, Fourier analysis, performed at different heights, highlights two dominant seasonal os-

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

5 cillations: an annual oscillation (AO) and a semi-annual oscillation (SAO). The corresponding vertical profiles of the AO and SAO in the subtropical UTLS region under consideration are depicted in Fig. 3. In general, the subtropical UTLS is dominated by the AO, with the AO amplitudes about 4 times stronger than the SAO amplitudes. It is noted that in the lower stratosphere (70-hPa), the SAO shows weaker amplitudes than in the upper troposphere (200-hPa).

10 The SAO has its maximum amplitude at 200-hPa where the AO shows its minimum (Fig. 3). The seasonal variation of temperature at 200-hPa appears to be driven by the SAO cycle. Thus, it is evident in Fig. 2a that the coldest temperatures at 200-hPa appear in May and November, during the autumn and spring equinoxes, respectively, while the highest temperatures are observed in February and August.

The AO shows a maximum-amplitude at the tropopause height, and is dominant in the lower stratosphere, at the 70- and 50-hPa levels (Fig. 3). At the 30- and 20-hPa heights both the SAO and AO show weak amplitudes.

15 3.2. Trend estimation

In the present study, temperature trends from 22 years of upper-air data over Durban, have been investigated using a statistical trend model adapted from the AMOUNT and AMOUNT-O₃ models for ozone and temperature trend assessment (Hauchecorne et al., 1991; Mohanakumar, 1994; Keckhut et al., 1995; Guirlet et al. 2000). It is based on a multivariate least-squares method and takes into account seasonal variations, i.e., annual and semi-annual cycles, and uses parametric functions in order to consider the QBO, the ENSO, and the 11-years solar cycles. The solar cycle is defined as a linear function and is correlated with the solar flux at 10.7 cm.

25 In the trend model, temperature T at time t and height z is represented by the equation:

$$T(t, z) = \sum_{k=1}^p a(z, k) \times c(t, k) + T'(t, z)$$

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

where $c(t,k)$ describes the temporal evolution of the forcing k ; $a(z,k)$ is a coefficient computed by the model for the forcing k ; and $T'(t,z)$ is the residual term. A least-squares method computes the coefficients $a(z,k)$ in order to minimise the quantity:

$$\chi^2(z) = \left[\sum_t T(t,z) - \sum_k a(z,k) \times c(t,z) \right]^2$$

5 The variance of the residual term is defined by:

$$\sigma_r^2(z) = \frac{\chi^2(z)}{n-p}$$

where n and p represent measurement and forcing numbers, respectively.

One of the most significant factors that influence temperature trends, notably in the lower stratosphere, is the large aerosol enhancement following a volcanic eruption (WMO, 1995). Due to solar radiation and terrestrial infrared radiation increases caused by a major eruptions, aerosol loading results in a warming in the stratosphere, particularly in the tropical and subtropical regions (Labitzke and McCormick, 1992).

Taking into account the temporal coverage of the dataset used in the present study, trend analyses may be biased if post-Pinatubo data are not removed or adjusted for volcanic influence. The Pinatubo erupted in June 1991 and has injected a large quantity of gases and aerosols into the stratosphere up to a height of 30 km, with most of the volcanic clouds in the 20–25 km altitude range (McCormick and Viega, 1992; Nardi et al., 1993). Bencherif et al. (2003) have shown from SAGE II data that the aerosol amounts in the southern subtropical UTLS decreased gradually to return to normal levels by late 1995 and early 1996.

Once post-Pinatubo data (June 1991–December 1995) were removed, linear trends in temperature at all UTLS pressure levels were estimated. Figure 4a shows the vertical distribution of the linear trend in temperature per decade. A cooling trend is observed at almost all heights in the UTLS region (250–20 hPa). The maximum cooling rate is

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

1.09±0.27 K per decade, and is observed in the lower stratosphere, at 70-hPa. Figure 4b depicts the time evolution of mean monthly temperatures at that level, with the cooling trend line superimposed.

The mean tropospheric (250–100 hPa) and stratospheric (70–20 hPa) temperature trends are 0.10±0.14 and 0.76±0.23 K/decade, respectively. The cooling rate in the lower stratosphere is indeed significant and higher than in the upper troposphere. It is noted that for tropospheric heights below the 200-hPa height, temperature trend rates are found to be positive but not significant.

Using a global rawinsonde network Oort and Liu (1993) considered the latitudinal structure of long-term trends for the 1968–1988 period. They obtained cooling trends in the SH UTLS region that increased with latitude from the tropics to the South Pole. More recently, Parker et al. (1997) have combined the rawinsonde data from the Australian region with the satellite-based Microwave Sounding Unit (MSU) retrievals, to calculate temperature trends over the period 1979–1996. They also found cooling trends in the southern UTLS, notably at subtropical latitudes.

Stratospheric and tropospheric temperature trends as derived by Angell (1988), Oort and Liu (1993) and Parker et al. (1997) are compared with values obtained in this study in Table 2. The cooling rate obtained in the lower stratosphere appears to be the most important. It is in good agreement with the other results, taking into account that the periods of analysed datasets are not identical.

A study on the variance and standard deviation (STD) of the mean temperature over a region is considered important for climatological use. Variability is defined as the STD of all temperature observations relevant to a given month. Monthly average temperature variability (STD) obtained for Durban is reported in Table 1. From March to October STD values are found to be relatively higher in the troposphere, including at the tropopause level (100 hPa), in comparison with stratospheric levels. Yet, at all pressure levels, STD shows a maximum during austral winter, i.e., during July and August. The highest temperature STD value over Durban is obtained at 200 hPa in July; while the highest STD in the lower stratosphere is recorded during the June–July period, at

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

70 hPa. This is most likely related to temperature variations caused by the development of synoptic disturbances in the troposphere in winter. Indeed, such wave disturbances propagate vertically through to the stratospheric layers in the winter hemisphere when the zonal winds are westerly (Charney and Drazin, 1961). Hauchecorne et al. (1987) have also drawn attention to the role of wintertime propagations of planetary and gravity waves as the causes of increases in temperature variability.

In order to understand the seasonal characteristics of temperature trends, knowledge of trends in dynamical activity is needed. Dynamical trends in the UTLS may be induced by changes in concentrations of radiatively active gases or by any tropospheric changes.

In the present study, having separated the initial dataset into monthly time-series, the least-squares model (introduced above) was applied separately to winter (June) and summer (December) datasets in the UTLS, in order to estimate seasonal temperature trends. In the upper troposphere, there were no seasonal differences in temperature (trends were nearly the same as depicted in Fig. 4a). Table 3 gives the winter and summer temperature trends estimated in the lower stratosphere (70, 50 and 30 hPa) over Durban. Taking into account the reduction in size of the datasets, the derived seasonal trends may be used qualitatively, but should not be used as a precise quantitative result.

It is observed that the cooling trends are slightly higher in summer than in winter. Maximum cooling is obtained at the 70 hPa height for both summer and winter time series. As suggested above, the seasonal dependence of temperature trends may be related to trends in wave activity in the southern UTLS. In fact, Bodeker and Scourfield (1995), using TOMS data recorded over the period 1979–1992 and an EOF analysis on the vertical component of Eliassen-Palm flux, have reported an increasing trend in wave activity during winter in the southern UTLS. This is consistent with our seasonal analysis and shows that in the lower stratosphere, trends in temperature should be examined in conjunction with trends in dynamics.

4. Conclusion

Daily afternoon radiosonde data over Durban (30.0° S, 30.9° E), which is situated on the east coast of South Africa, over the period January 1980 to December 2001, i.e. 22 years, were used to derive a mean temperature climatology and to determine trends in temperature in the upper-troposphere and lower-stratosphere (UTLS) region. Previous studies have highlighted the importance of this region because of the many dynamical processes that occur here and their role in the vertical and horizontal transfer of energy, mass and atmospheric constituents.

The mean seasonal temperature climatology revealed differences between the upper troposphere and the lower stratosphere. The seasonal cycle of temperature at the 250-hPa height is anti-correlated with that at 150-hPa and 100-hPa. The 100-hPa level (tropopause) exhibits a minimum temperature in late summer (February) and a maximum in winter (June to August), and closely corresponds to the results obtained by Randel et al. (2000) for tropical regions.

A Fourier analysis performed at different heights highlighted two dominant seasonal oscillations: an annual (AO) and a semi-annual oscillation (SAO). The former is dominant (about 4 times stronger) in the UTLS region and maximises at the 100-hPa height, whereas the SAO has its maximum amplitude at 200-hPa, where the AO is at its minimum.

A trend analysis based on a multivariate least-squares regression technique was undertaken. Seasonal variations (both AO and SAO), the QBO, the ENSO and the 11-year solar cycle were taken into account. Furthermore, the data were adjusted for the influence of the Pinatubo eruption. Cooling trends were observed at almost all heights in the UTLS region (250–20 hPa), although cooling rates in the lower stratosphere were higher than in the upper troposphere. The maximum cooling rate of 1.09 ± 0.27 K per decade was observed at the 70-hPa height. Cooling trends were slightly higher in summer than in winter and were consistent with cooling rates derived by other authors.

Temperature variability as indicated by the magnitude of the standard deviation was

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

found to be higher in the troposphere than in the stratosphere and also to be greater in winter. The higher temperature variations in winter were attributed to the greater frequency of tropospheric synoptic disturbances in winter and to the vertical propagation of such wave disturbances through to the stratosphere under westerly zonal winds.

5 *Acknowledgements.* We are grateful to the South African Weather Service (SAWS) for providing upper-air data. We thank F. Sokolic from the School of Environmental Sciences, University of KwaZulu-Natal. This work was undertaken under the French-South African co-operative programme and supported by the French Centre National de la Recherche Scientifique (CNRS),
10 the South African National Research Foundation (NRF) and the Regional Council, Conseil Régional de La Réunion.

References

- Angell, J. K.: Variations and trends in tropospheric and stratospheric global temperatures, 1958–87, *J. Climate*, 1(12), 1296–1313, 1988.
- Baray, J. L., Ancellet, G., Taupin, F. G., Bessafi, M., Baldy, S., and Keckhut, P.: Subtropical tropopause break as a possible stratospheric source of ozone in the tropical troposphere, *J. Atmos. Sol.-Terr. Phys.*, 60, 27–36, 1998.
- 15 Baray, J. L., Daniel, V., Ancellet, G., and Legras, B.: Planetary-scale tropopause folds in the southern subtropics, *Geophys. Res. Lett.*, 27, 353–356, 2000.
- Bencherif, H., Portafaix, T., Baray, J. L., Morel, B., Baldy, S., Leveau, J., Moorgawa, A., Michaelis, M. M., Hauchecorne, A., Keckhut, P., and Diab, R.: LIDAR observations of lower stratospheric aerosols over South Africa linked to large scale transport across the southern subtropical barrier, *J. Atmos. Sol.-Terr. Phys.*, 65, 707–715, 2003.
- 20 Bodeker, G. E. and Scourfield, M. W. J.: Planetary waves in total ozone and their relation to Antarctic ozone depletion, *Geophys. Res. Lett.*, 22, 21, 2949–2952, 1995.
- Charney, J. G. and Drazin, P. G.: Propagation of planetary-scale disturbances from the lower into the upper atmosphere, *J. Geophys. Res.*, 66, 83–109, 1961.
- Gaffen, D. J.: Temporal inhomogeneities in radiosonde temperature records, *J. Geophys. Res.*, 99(D2), 3667–3676, 1994.
- Grant, W. B., Browell, E. V., Fishman, J., Brakett, V. G., Veiga, R. E., Nganga, D., Minga, A.,

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Cros, B., Butler, C. F., Fenn, M. A., Long, C. S., and Sowe, L. L.: Aerosol associated changes in tropical stratospheric ozone following the eruption of Mount Pinatubo, *J. Geophys. Res.*, 99(D4), 8197–8211, 1994.

5 Guirlet M., Keckhut, P., Godin, S., and Megie, G.: Description of the long-term ozone data series obtained from different instrumental techniques at a single location: the Observatoire de Haute-Provence (43.9° N, 5.7° E), *Ann. Geophys.*, 18, 1325–1339, 2000,
[SRef-ID: 1432-0576/ag/2000-18-1325](#).

Harris, N. R. P., Ancellet, G., Bishop, L., Hofmann, D. J., Kerr, J. B., McPeters, R. D., Prenz-
10 dez, M., Randel, W. J., Staehelin, J., Subbaraya, B. H., Volz-Thomas, A., Zawodny, J.,
and Zerefos, C. S.: Trends in stratospheric and free tropospheric ozone, *J. Geophys. Res.*,
102(D1), 1571–1590, 1997.

Hauchecorne, A. and Wilson, R.: Mesospheric temperature inversion and gravity wave break-
ing, *Geophys. Res. Lett.*, 14, 933–936, 1987.

Hauchecorne, A., Chanin, M. L., and Keckhut, P.: Climatology and trends of the middle at-
15 mospheric temperature (33–87 km) as seen by Rayleigh lidar over the south of France, *J.*
Geophys. Res., 96(15), 15 297–15 309, 1991.

Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglas, A. R., Rood, R. B., and Pfister, L.:
Stratosphere-Troposphere Exchange, *Rev. Geophys.*, 1995.

Keckhut, P., Hauchecorne, A., and Chanin, M. L.: Mid-latitude long-term variability of the middle
20 atmosphere : trends and cyclic and episodic changes, *J. Geophys. Res.*, 1995.

Keckhut, P., Schmidlin, F. J., Hauchecorne, A., and Chanin, M.-L.: Trend estimates from rock-
etsondes at low latitude station (8S–34N), taking into account instrumental changes and
natural variability, *J. Atmos. Sol. Terr. Phys.*, 61, 447–459, 1999.

Kokin, G., and Lysenko, E.: On temperature trends of the atmosphere from rocket and ra-
25 diosonde data, *J. Atmos. Terr. Phys.*, 56, 1035–1044, 1994.

Labitzke, K. and McCormick M. P.: Stratospheric temperature increases due to Pinatubo
aerosols, *Geophys. Res. Lett.*, 19(2), 207–210, 1992.

Logan, J. A.: Trends in the vertical distribution of ozone: An analysis of ozonesonde data, *J.*
Geophys. Res., 99(D12), 25 553–25 586, 1994.

30 McCormick, M. P. and Veiga, R. E.: SAGE II measurements of early Pinatubo aerosols, *Geo-*
phys. Res. Lett., 19, 155–158, 1992.

Mohanakumar, K.: Temperature variability over the tropical middle atmosphere, *Ann. Geophys.*,
12, 448–456, 1994,

- Nardi, B, Chanin, M.-L., Hauchecorne, A., Avdyushin, S. I., Tulinov, G. F., Ivanov, M. S., Kuzmenko, B. N., and Mezhuev, I. R.: Morphology and dynamics of the Pinatubo aerosol layer in the Northern Hemisphere as detected from a ship-borne lidar, *Geophys. Res. Lett.*, 20(18), 1967–1970, 1993.
- O'Connor, F. M., Vaughan, G., and De Backer, H.: Observation of subtropical air in the European mid-latitude lower stratosphere, *Q. J. R. Meteorol. Soc.*, 125, 2965–2986, 1999.
- Oort, A. H. and Liu, H.: Upper-air temperature trends over the globe, 1958–1989, *J. Clim.*, 6, 292–307, 1993.
- Parker, D. E., Gordon, M., Cullum, D. P. N., Sexton, D. M. H., Folland, C. K., and Rayner, N.: A new global gridded radiosonde temperature database and recalculated temperature trends, *Geophys. Res. Lett.*, 24, 1499–1502, 1997.
- Portafaix, T., Morel, B., Bencherif, H., Godin-Beekmann, S., Baldy, S., and Hauchecorne, A.: Fine scale study of a thick stratospheric ozone lamina at the edge of the southern subtropical barrier, *J. Geophys. Res.*, 108(D6), 4196, doi:10.1029/2002002741, 2003.
- Ramaswamy, V., Chanin, M.-L., Angell, J., Barnett, J., Gaffen, D., Gelman, M., Keckhut, P., Koshelkov, Y., Labitzke, K., Lin, J.-J. R., O'Neill, A., Nash, J., Randel, W., Rood, R., Shine, K., Shiotani, M., and Swinbank, R.: Stratospheric temperature trends: observations and model simulations, *Rev. Geophys.*, 39(1), 71–122, 2001.
- Randel, W. J., Wu, F., and Gaffen, D. J.: Interannual variability of the tropical tropopause derived from radiosonde data and NCEP reanalyses, *J. Geophys. Res.*, 105(D12), 15 509–15 524, 2000.
- Vaughan, G. and Timmis, C.: Transport of near-tropopause air into the lower midlatitude stratosphere, *Q. J. R. Meteorol. Soc.*, 124(549), 1559–1578, 1998.
- WMO: Scientific assessment of ozone depletion: 1994, *Global Ozone Res. and Monit. Proj.*, Rep. 37, 1995.
- WMO: Scientific assessment of ozone depletion: 1998, *Global Ozone Res. and Monit. Proj.*, Rep. 44, 1999.

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Table 1. Mean monthly temperature (in Kelvin) and corresponding standard deviations derived from Durban-SAWS data covering the period from January 1980 to December 2001.

height (hPa)	Jan	Feb	March	April	May	June	July	Aug	Sep	Oct	Nov	Dec
250	234.5	235.3	233.2	230.1	228.6	228.3	228.2	228.8	228.9	229.7	231.4	232.6
	2.6	2.8	2.4	2.3	2.6	2.7	2.8	3.3	2.6	2.2	2.1	2.2
200	219.4	220.5	218.8	217.3	217.1	218.2	219.0	219.1	218.1	217.6	217.3	218.0
	2.6	2.7	2.6	2.9	3.4	3.8	3.9	3.7	3.7	3.2	2.8	2.4
150	208.9	209.2	209.7	211.1	212.6	213.8	214.1	213.9	213.1	212.6	210.9	209.8
	2.3	2.2	2.5	3.0	3.5	3.4	3.3	3.4	3.6	3.1	2.7	2.8
100	200.1	199.7	201.6	204.5	206.8	207.6	207.9	207.9	207.4	206.7	204.1	202.4
	2.8	2.7	3.2	3.3	3.2	3.1	3.3	3.7	3.6	3.2	2.8	2.7
70	202.1	201.9	203.8	205.8	207.6	208.3	208.6	209.2	209.7	208.3	205.9	203.4
	3.1	2.7	2.9	2.8	2.9	3.4	3.4	3.2	3.0	2.8	2.6	2.9
50	209.9	209.5	210.5	211.0	212.1	212.6	213.0	213.8	214.3	213.0	211.2	210.2
	2.8	2.5	2.4	2.3	2.5	2.7	3.0	2.8	2.9	2.4	2.3	2.8
30	218.5	218.8	219.5	219.1	219.0	219.0	219.2	219.4	219.4	218.5	217.4	218.1
	2.2	2.2	2.1	2.1	2.2	2.5	3.2	2.7	2.6	2.6	2.2	2.6
20	224.4	224.8	225.3	224.3	223.5	223.2	222.9	222.7	222.6	222.1	221.5	223.1
	2.3	2.2	2.0	2.1	2.5	2.8	3.2	2.9	2.6	2.8	2.3	2.7

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.

Table 2. Upper troposphere and lower stratosphere temperature trends (in Kelvin per decade) from published studies and from the current study.

	Angel (1988) 1959–1988	Oort and Liu (1993) 1964–1988	Parker et al. (1997) 1979–1996	Current study 1980–2001
Upper Troposphere	-0.16 ± 0.11	-0.11 ± 0.11	-0.11	-0.10 ± 0.14
Lower Stratosphere	-0.64 ± 0.29	-0.43 ± 0.16	-0.73	-0.76 ± 0.23

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

**Temperature
climatology and
trend estimates over
Durban, South Africa**H. Bencherif et al.

Table 3. Winter and summer temperature trends (in Kelvin per decade) as estimated in the lower stratosphere (70, 50 and 30 hPa) over Durban.

Pressure Level	Winter Trend (K/year)	Summer Trend (K/year)
30 hPa	−0.051	−0.124
50 hPa	−0.075	−0.080
70 hPa	−0.110	−0.125

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.




Fig. 1. Time-height section of temperature over Durban, South Africa, obtained with data from January 1980 to December 2001.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Temperature climatology and trend estimates over Durban, South Africa

H. Bencherif et al.




Fig. 2. Time evolution of monthly averaged temperatures per heights: **(a)** in the upper troposphere and **(b)** in the lower stratosphere over Durban (see legends).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.




Fig. 3. Vertical profiles of the annual and the semi-annual oscillations as derived from SAWS data recorded over Durban (see legend).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**Temperature
climatology and
trend estimates over
Durban, South Africa**

H. Bencherif et al.




Fig. 4. (a) Linear temperature trend (in Kelvin per decade) height profile framed with the 95% confidence profiles. (b) Time evolution of the monthly averaged temperature values in the lower stratosphere (at 70 hPa), over Durban. The superimposed red line indicates the obtained linear trend, i.e., -1.09 ± 0.27 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)