

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters in atmospheric conditions using quantum chemical methods

T. Kurtén¹, L. Torpo¹, M. R. Sundberg², V.-M. Kerminen³, H. Vehkamäki¹, and M. Kulmala¹

¹Division of Atmospheric Sciences, Department of Physical Sciences, P.O.Box 64, FI-00014 University of Helsinki, Finland

²Laboratory of Inorganic Chemistry, Department of Chemistry, P.O. Box 55, FI-00014 University of Helsinki, Finland

³Finnish Meteorological Institute, Air Quality Research, Sahaajankatu 20 E, FI-00880 Helsinki, Finland

Received: 31 January 2007 – Accepted: 16 February 2007 – Published: 23 February 2007

Correspondence to: T. Kurtén (theo.kurten@helsinki.fi)

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

We study the ammonia addition reactions of $\text{H}_2\text{SO}_4\cdot\text{NH}_3$ molecular clusters containing up to four ammonia and two sulfuric acid molecules using the ab initio method RI-MP2 (Resolution of Identity 2nd order Møller-Plesset perturbation theory). Together with results from previous studies, we use the computed values to estimate an upper limit for the ammonia content of small atmospheric clusters, without having to explicitly include water molecules in the quantum chemical simulations. Our results indicate that the $\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratio of small molecular clusters in typical atmospheric conditions is probably around 1:2. High ammonia concentrations or low temperatures may lead to the formation of ammonium bisulfate (1:1) clusters, but our results rule out the formation of ammonium sulfate clusters (2:1) anywhere in the atmosphere. A sensitivity analysis indicates that the qualitative conclusions of this study are not affected even by relatively large errors in the calculation of electronic energies or vibrational frequencies.

1 Introduction

In most atmospheric conditions, particle formation by nucleation is thought to involve sulfuric acid and water, with possible contributions from ions, ammonia or some organic species. Experimental studies have indicated that the presence of ammonia has a clear nucleation-enhancing effect (Ball et al., 1999) in the sulfuric acid - water system. However, previous estimates of $\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratios in atmospheric nucleating clusters have varied widely. Predictions of theoretical studies have ranged from zero (corresponding to no nucleation enhancement by ammonia) to 1:1 or even 2:1, corresponding to ammonium bisulfate and ammonium sulfate formation, respectively. Quantum chemical studies (Ianni and Bandy, 1999) have predicted low values for the $\text{NH}_3:(\text{H}_2\text{SO}_4)$ -ratio, while state of the art thermodynamics (Clegg et al., 1998) predicts extensive ammonium bisulfate formation (Vehkamäki et al., 2004). Older versions of the classical nucleation theory indicated that ammonia enhances $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nucleation very strongly (Napari et al., 2002), while the updated models result in more modest, but still significant enhancement (Anttila et al., 2005).

Our PW91/DNP study (Kurtén et al., 2007) on two-acid clusters containing up to one ammonia and up to seven water molecules predicted the fraction of ammonia-containing two-acid clusters in ambient conditions¹ to be quite low. However, ammonia was observed to significantly enhance the addition of sulfuric acid molecules to the clusters, and preliminary unpublished MPW1B95 calculations² on three-acid clusters demonstrate that the enhancement increases with the number of sulfuric acid molecules. This might tentatively indicate a lower limit of around 1:3 or 1:4 for the NH₃:(H₂SO₄) mole ratio in atmospheric conditions. In this study, we have extended the analysis along the ammonia co-ordinate in order to complete the picture of the role of ammonia in ternary nucleation.

We apply an accurate ab initio method to study the thermochemistry of (H₂SO₄)₂·(NH₃)_y (y=0–4) molecular clusters and investigate reactions potentially relevant to atmospheric nucleation by calculating their Gibbs free energies. Using the derived cluster distributions, we further estimate an upper limit for the NH₃:(H₂SO₄) - ratio in atmospheric conditions.

2 Computational details

Previous quantum chemistry studies (Ianni and Bandy, 1999; Kurtén et al., 2007; Larson et al., 1999; Ianni and Bandy, 2000; Re et al., 1999; Bandy and Ianni, 1998; Ding et al., 2003) on sulfuric acid - ammonia - water clusters have mainly used various density functionals (e.g. B3LYP or PW91) to treat the exchange-correlation of electrons. Currently, all generally available density functionals are, by construction,

¹Cluster distributions were derived at a temperature of 265 K, a RH of 50%, an ammonia partial pressure of 14 ppt and a sulfuric acid partial pressure of 0.7 ppt.

²Torpo, L.; Kurtén, T.; Vehkamäki, H.; Laasonen K., and Kulmala, M.: The significance of ammonia in growth of atmospheric nanoclusters, in preparation, 2007.

Estimating the NH₃:H₂SO₄ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

unable to accurately treat electron dispersion (see e.g. Jensen, 2007), which plays a moderately important role in hydrogen bonding. This is a major reason for the large differences observed between the experimental formation free energies for sulfuric acid-water clusters (Hanson and Eisele, 2000) and the computed values (see e.g. Kurtén et al., 2006). There are two main pathways to obtain accurate thermochemistry for hydrogen-bonded clusters. One approach is to use density functionals that are sufficiently well parametrized to replicate binding energies of various benchmarking sets of weakly bound clusters, despite the physically inaccurate treatment of dispersion. Another alternative is to use correlated wavefunction-based methods such as Møller-Plesset (MP) perturbation theory (Møller and Plesset, 1934) or coupled-cluster methods. Unfortunately, the computational cost of even the most affordable correlated wavefunction-based method, MP2, is normally orders of magnitude greater than that of density functional methods. However, the recently developed and implemented (Weigend and Häser, 1997) resolution of identity (RI) approximation MP2 (RI-MP2) produces binding energies that are essentially identical to MP2, at a small fraction of the computational effort (Weigend et al., 1998). In this study we have used the RI-MP2 method implemented in the Turbomole program suite (Ahlrichs et al., 1989; Häser and Ahlrichs, 1989; Turbomole version 5.8) for all calculations.

For structure optimization and numerical frequency calculations, we have used the augmented correlation-consistent basis set of valence double-zeta quality (Dunning Jr. et al., 2001) aug-cc-pV(D+d)Z. The electronic energies are then calculated at these geometries using the larger triple-zeta aug-cc-pV(T+d)Z basis set. The corresponding auxiliary basis sets needed for the RI expansion are given by Weigend et al. (2002). From the aug-cc-pV(D+d)Z and aug-cc-pV(T+d)Z energies, basis-set limit (denoted as aug-cc-pV(∞ +d)Z) energies can be extrapolated using the formula of Helgaker et al. (1997):

$$E(X) = E(\infty) + AX^{-3} \quad (1)$$

where $E(\infty)$ is the basis-set limit energy, A is a constant, $X=2$ for the aug-cc-pV(D+d)Z set and $X=3$ for the aug-cc-pV(T+d)Z set. The purpose of the extrapolation is to

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

remove both basis-set superposition errors and basis-set incompleteness errors, and thus provide a more reliable estimate of the basis-set limit than that given e.g. by the counterpoise correction of Boys and Bernadi (1970). Test calculations up to the RI-MP2/aug-cc-pV(5+d)Z level carried out in connection with our recent high-level study on small neutral and ionic sulfuric acid - water clusters³ indicate that for the RI-MP2 method, basis-set effects beyond the aug-cc-pV(T+d)Z level are relatively small (well under 0.5 kcal/mol in terms of the binding energy per molecule). Neglect of higher-order correlation may cause some errors but these are also likely to be relatively small.

The convergence with respect to the electronic energy in the self-consistent field (SCF) step was 10^{-7} a.u. (atomic units), and the convergence with respect to the gradient was 10^{-4} a.u. For the numerical frequency calculations, a stepsize of 0.01 a.u. and a SCF convergence limit of 10^{-8} a.u. were used. Thermal contributions to the enthalpies and entropies are computed using the rigid rotor and harmonic oscillator approximations.

To obtain the initial geometries for the moieties under study, a Monte Carlo-based conformational search installed in Spartan '02 (Wavefunction Inc., 2002) was first applied. The search was carried out by semiempirical methods using the AM1 basis. Since the chosen level of theory was low, several structures for each stoichiometry were subsequently optimized at the B3LYP/6-31G* level using the Gaussian 03 program suite (Frisch et al., 2004). The most stable structures from this optimization were then selected for the accurate RI-MP2 computations. For the (H₂SO₄)₂·(NH₃)₄ stoichiometry, several structures were optimized at the RI-MP2 level in order to ensure that the final energies (and corresponding NH₃:H₂SO₄ ratios) are representative.

³Kurtén, T., Noppel, M., Vehkamäki, H., Salonen, M., and Kulmala, M.: Quantum chemical studies of hydrate formation of H₂SO₄ and HSO₄⁻, submitted, Boreal Env. Res, 2007.

3 Results and discussion

Modeling the hydration of sulfuric acid - ammonia clusters is problematic. In typical atmospheric conditions, two-acid clusters can be bound to as many as seven or eight water molecules according to our recent PW91/DNP study (Kurtén et al., 2007). The addition of water molecules to the clusters increases the computational effort in two ways. First, the CPU time and memory required for one energy, or gradient, evaluation increases as an n :th power of the system size. For the RI-MP2 method used in this study, this so-called scaling coefficient n is around 4. For example, with the aug-cc-pV(D+d)Z basis set, a $(\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_8$ cluster contains about 1.6 times as many basis functions as a $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_4$ cluster, implying approximately a 6-fold increase in the computing time for each energy evaluation. For numerical frequency calculations (required for the computing of free energies), the scaling is even worse, as the hydrated $(\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_8$ cluster possesses more than 1.8 times the number of normal modes as the unhydrated $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_4$ cluster, and each normal mode requires two energy evaluations. Second, the number of possible bonding patterns (cluster conformers) increases combinatorially with the number of molecules. Finding a decent representative minimum energy configuration (the so-called configurational sampling problem) thus becomes prohibitively difficult as the cluster size increases. Also, as the degree of water-water bonding grows, the clusters become increasingly floppy (i.e. the number of very low frequency vibrational modes grows), and the harmonic oscillator and rigid rotor approximations become less reliable. Fortunately, an analysis of previous quantum chemical results (Ianni and Bandy, 1999; Kurtén et al., 2007) indicates that though the energetics of individual water addition reactions vary significantly, the average free energy for the addition of several water molecules to a sulfuric-acid-ammonia cluster is only weakly dependent on the ammonia content of the cluster. Similarly, the binding of ammonia to the clusters is weakly dependent on the water content. The free energies for the addition of ammonia to various clusters calculated in previous studies are presented in Table 1.

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

It can be seen from Table 1 that the effect of water molecules on the addition free energies of ammonia is relatively weak at least for the first few water molecules. The effect is also to some extent systematic: the addition of water molecules tends to somewhat decrease the binding of ammonia to the cluster. (The deviations from this pattern may be related to configurational sampling issues rather than true chemical effects.) This is reasonable from a chemical point of view. The binding between sulfuric acid and ammonia is much stronger than that between sulfuric acid and water or ammonia and water. Thus, the binding of a few water molecules to the cluster is unlikely to significantly disrupt the bonding pattern of the sulfuric acid – ammonia core. However, as the number of water molecules increases, the core will eventually start to dissolve, and both the acid and ammonia molecules will be bonded mainly to water molecules.

The addition of ammonia to a unhydrated (H₂SO₄)_x·(NH₃)_y core thus represents the minimum case with respect to the free energy (i.e. the most negative values for ΔG are obtained for the unhydrated or almost unhydrated cluster), and hydration will probably only serve to decrease the H₂SO₄–H₃ binding. Therefore, cluster distributions derived for the (H₂SO₄)_x·(NH₃)_y system with no water present represent the maximum case with respect to the amount of ammonia molecules bound to the clusters. While the competition of water molecules with ammonia will lead to lower NH₃:(H₂SO₄) mole ratios in the real atmosphere, there exist, to our knowledge, no mechanisms which would tend to lead to greater ratios than that predicted for this reference case. NH₃:(H₂SO₄) ratios calculated from these distributions at some temperatures and partial pressures will thus yield useful upper limits for atmospheric cluster distributions.

3.1 Structural and thermochemical data

The cluster structures studied here are shown in Fig. 1. The clusters have been drawn using the MOLEKEL 4.3 visualization package (Portmann, 2002). The corresponding electronic energies, enthalpies and entropies are presented in Table 2. The cartesian coordinates for the most stable cluster structures are given in the auxiliary information. <http://www.atmos-chem-phys-discuss.net/7/2937/2007/>

acpd-7-2937-2007-supplement.pdf The electronic and standard Gibbs free energies for the ammonia addition reactions (at 298 K and reactant pressures of 1 atm) are given in Table 3.

It can be seen from Table 3 that the free energy of the ammonia addition reactions increases steadily with the number of ammonia molecules. Thus, while the addition of the first ammonia molecule is strongly exothermic, the third and fourth addition reactions are close to endothermic.

The mode of intermolecular interaction in the sulfuric acid dimers varies depending on the number of attached ammonia molecules (see Fig. 1). In the pure dimer [structure a)] each sulfuric acid has two “outer” and two “inner” hydrogen atoms. The latter atoms form a typical hydrogen bonding system found e.g. in organic acids. In structure b), an attached ammonia molecule has taken one of the two outer hydrogens. However, when the number of the ammonia molecule increases to two, one of the inner hydrogen atoms is also taken. In structure d), the remaining inner hydrogen still participates in the intermolecular hydrogen bonding between the sulfuric acid moieties. There is also an intermolecular interaction between an ammonia molecule and the vicinal ammonium cation. In structure e), the inner hydrogen still remains, and there are two similar ammonia – ammonium dimers approximately on the opposite sides of the sulfuric acid dimer.

3.2 Cluster distributions

The computed free energies can be used to derive cluster size distributions given some ambient monomer pressures. Specifically, the relative fraction of two-acid clusters bound to n ammonia molecules is:

$$\frac{[(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_n]}{\sum_{k=0}^4 [(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_k]} = \frac{[\text{NH}_3]^n e^{-\frac{\Delta G_n}{RT}}}{1 + \sum_{k=1}^4 [\text{NH}_3]^k e^{-\frac{\Delta G_k}{RT}}} \quad (2)$$

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**T. Kurtén et al.

where the ammonia vapor pressure [NH₃] is given in atmospheres and ΔG_n is the free energy change for the reaction $(\text{H}_2\text{SO}_4)_2 + n \text{NH}_3 \Rightarrow (\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_n$. (Note that as we are comparing the relative concentrations of two-acid clusters, the sulfuric acid concentration does not enter the final expression.) Table 4 shows the relative concentrations of the five studied cluster types for three temperatures and four ammonia concentrations; 1 ppt–1 ppm. The concentrations were chosen to span the range of measured atmospheric values. According to Seinfeld and Pandis (1998), typical atmospheric ammonia concentrations in continental air are around 0.1–10 ppb. In a recent (spring 2005) campaign at the Hyytiälä measurement station (Riipinen et al., 2006), ammonia concentrations in the range 10^9 – 10^{11} cm⁻³ were measured, corresponding to mixing ratios of about 0.04–4 ppb. Similarly, results from the BIOFOR campaign (Kulmala et al., 2001) in 2001 yielded average ammonia mixing ratios of 49 ppt and 52 ppt in spring and summer, respectively.

The results are shown in Table 4. It can be seen that the two-acid clusters of our hypothetical unhydrated cluster distribution are bound to at least one ammonia molecule in almost all conditions. The ammonium bisulfate dimer $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_2$ is formed in high ammonia concentrations or low temperatures, but the ammonium sulfate dimer $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_4$ is not formed in any atmospherically relevant conditions. It should be noted that these values relate to the very first steps of gas-to-particle nucleation only: ammonium sulfate clusters may be produced in the atmosphere by other mechanisms (e.g. growth processes), which are beyond the scope of this study.

3.3 Sensitivity analysis

Error analysis of previous studies (Kurtén et al., 2006) indicates that far-reaching atmospheric conclusions drawn from relatively small differences in computed thermochemical data may be unreliable. Though the RI-MP2/aug-cc-pV(T+d)Z data should be reliable (more so than previous density functional data), our thermochemical parameters may be contaminated by several error sources. Basis-set superposition errors not accounted for by the extrapolation procedure of Eq. (1) would tend to decrease the

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

computed Gibbs free formation energies (i.e. exaggerate the stability of the clusters) compared to the real values, while the neglect of vibrational anharmonicity would tend to increase them. Conformational sampling may cause problems in two ways. First, we can not be certain that our studied cluster structures correspond to the global minimum geometries. Second, especially the largest (H₂SO₄)₂·(NH₃)₄ cluster possesses some very low vibrational frequencies, indicating that treating the cluster as a single structure vibrating harmonically around some minimum energy geometry may cause large errors in the thermochemistry. Moreover, our assumption that the binding of water molecules will tend to decrease the H₂SO₄ – NH₃ binding might at least in some conditions be incorrect, and the atmospheric NH₃:H₂SO₄ mole ratios could therefore be greater than those predicted here.

The effects of these error sources on our atmospheric conclusions were estimated by recomputing the cluster size distributions using formation energetics that have been modified in two ways. First, we have scaled the vibrational frequencies by a factor of 0.75, which is somewhat lower than the factors computed in our previous study (see the footnote on page 4) on small ionic clusters using accurate anharmonic thermochemical expressions, and significantly lower than the scaling factors usually used at the MP2 level (which are obtained by using the anharmonic fundamental frequencies in the expressions derived for harmonic oscillators, and are usually around 0.94–0.96, see e.g. Foresman and Frisch, 1996). This should be more than enough to account for vibrational anharmonicity. Second, we have assumed that the calculated electronic energy change for each ammonia addition reaction is too positive by 2 kcal/mol. (Thus, the electronic energy change for the (H₂SO₄)₂ + 4 NH₃ => (H₂SO₄)₂·(NH₃)₄ reaction used in the sensitivity analysis is 8 kcal/more more negative than that computed from Table 3.) This is intended to give an upper limit to the correction for all possible errors relating, for example, to the configurational sampling, the omission of higher-order correlation, and the binding of water molecules to the cluster. (It should be noted that any basis-set superposition error not corrected by the extrapolation procedure would tend to act in the opposite direction). The energetics used for the sensitivity analy-

sis are given in the auxiliary information. <http://www.atmos-chem-phys-discuss.net/7/2937/2007/acpd-7-2937-2007-supplement.pdf>

The results of the sensitivity analysis are presented graphically in Fig. 2. The solid lines correspond to the average $\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratio (computed as a weighted sum of the relative concentrations) for two-acid clusters computed from the original data in Tables 2 and 3. The dashed lines are the corresponding mole ratios computed with the data from the sensitivity analysis. We expect that the upper limits to the real atmospheric mole ratios will lie somewhere between the two lines, as the systematic errors assumed in the sensitivity analysis are rather extreme. The step-function - like behavior of the curves is explained by the fact that the mole ratio is a sum of polynomial terms, with e.g. the relative fraction of $(\text{H}_2\text{SO}_4)_2\cdot\text{NH}_3$ depending on $[\text{NH}_3]$, the relative fraction of $(\text{H}_2\text{SO}_4)_2\cdot(\text{NH}_3)_2$ on $[\text{NH}_3]^2$ and so on.

The dashed curves predict somewhat greater mole ratios (with e.g. significant amounts of ammonium bisulfate forming at around 2 orders of magnitude lower ammonia concentrations for each temperature), but mole ratios of 3:2 or higher are still not reached in any atmospherically realistic conditions. We thus conclude that ammonium sulfate dimer clusters are very unlikely to form by gas-to-particle nucleation in the atmosphere. While this does not directly prove anything about the composition of larger cluster structures (containing more than two sulfuric acid molecules), it is a strong indication that in nucleating molecular clusters, $\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratios significantly greater than 1:1 are unlikely.

3.4 Comparison to experimental and classical simulation results

Extensive experimental information on the NH_3 to H_2SO_4 mole ratio in the atmosphere is only available for bulk samples consisting of a large number of particles with diameter below $1\ \mu\text{m}$. In these samples, the NH_3 to H_2SO_4 mole ratio (in experimental studies usually expressed as the NH_4^+ to SO_4^{2-} mole ratio) depends in a complicated manner on the relative magnitudes of regional ammonia and sulfur emissions, on the

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

conversion of gaseous sulfur compounds to particulate sulfate, and on the removal rate of gaseous and particulate compounds from the atmosphere. In continental areas, the most important ammonia source is agriculture, whereas in marine areas gaseous ammonia originates mainly from the oceans (Bouwman et al., 1997). The principal sulfate precursor over the continents is sulfur dioxide (SO_2) with its largest emissions taking place in industrial regions (see e.g. Berglen et al., 2004). Over marine areas, aerosol sulfate is being produced from gaseous dimethylsulfide (DMS) originating from natural biological activities in the oceans (Kettle and Andreae, 2000). The conversion of SO_2 and DMS to aerosol sulfate takes place via photochemical reactions in the gas phase followed by H_2SO_4 condensation, and by aqueous-phase reactions taking place in clouds or wet aerosol particles (Berglen et al., 2004). Acidic sulfate aerosols rapidly take up gaseous ammonia if available. If there is ammonia in excess of that needed to neutralize all particulate sulfate, which corresponds to a $\text{NH}_4^+:\text{SO}_4^{2-}$ mole ratio of 2:1, also ammonium nitrate (and other ammonium salts) may be formed, leading to $\text{NH}_4^+:\text{SO}_4^{2-}$ mole ratios larger than 2:1 (Adams et al., 1999; Feng and Penner, 2007).

Observed NH_4^+ to SO_4^{2-} mole ratios display a large variability in fine ($<2.5 \mu\text{m}$ diameter) lower-tropospheric aerosols. Over continental mid-latitudes, these ratios are commonly in the range 1:1–2:1, even though larger ratios can be observed close to agricultural areas (Malm et al., 2004; Schaap et al., 2004). Ratios close to or even below 1:1 are frequently observed over Arctic or sub-Arctic regions (Ricard et al., 2002). In the marine boundary layer, NH_4^+ to SO_4^{2-} mole ratios are practically always below 2:1, with typical values being in the range 1:2–3:2 (Bates et al., 2001; Virkkula et al., 2006). Ratios smaller than unity are typical for the Southern Ocean, and ratios approaching zero, indicative of almost pure $\text{H}_2\text{SO}_4\text{--H}_2\text{O}$ aerosols, have been measured in Antarctic air masses (O'Dowd et al., 2006).

Macroscopic model simulations confirm the above patterns and show a few additional features (Adams et al., 1999). For example, NH_4^+ to SO_4^{2-} mole ratios are predicted to be close to or above 2:1 over most low-latitude continental regions. Due to the lack of significant ammonia sources, substantially lower ratios are expected to be

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

found in remote continental areas as well as in the free troposphere.

The experimental values described above are completely dominated by particles very much larger than the molecular clusters studied here. Neutral particles below 2.5 nm in diameter (corresponding to ca. 100–200 molecules) or so can not be detected at all with present instrumentation (McMurry, 2000), though there have been some improvements recently (Kulmala et al., 2005). Single-particle composition measurements are only possible for particles in the 10 nm size range (corresponding to ca 10^5 molecules) or larger (Nash et al., 2006). In contrast, the clusters studied here contain only up to 6 molecules (perhaps 10–20 when hydration is accounted for) and are below 1 nm in diameter. Also, the computational methods used to calculate thermochemical parameters in this study are based on the assumption that the translational, rotational and vibrational contributions to the entropy and enthalpy of the molecular clusters can be calculated as if they were large gas-phase molecules. In the larger particles, ammonium sulfate may exist in the solid phase, and the thermochemistry determining the $\text{NH}_3:(\text{H}_2\text{SO}_4)$ mole ratios may be quite different.

Our computational results indicate that the $\text{NH}_3:(\text{H}_2\text{SO}_4)$ mole ratio for molecular clusters formed in the very first steps of atmospheric nucleation is not likely to be greater than 1:1. This does not contradict the measurement results (which in turn do not cast doubt on the validity of the computations), but simply indicates that the chemical composition of nucleating particles is different from that of larger particles. Thus, using e.g. ammonium sulfate as a model for the chemical or physical properties of nucleating molecular clusters may not be justified, even though experimental evidence indicates that it is in many conditions a good model species for predicting the properties of larger particles.

4 Conclusions

We have studied $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_y$ (with $y=0-4$) clusters at the RI-MP2/aug-cc-pV(T+d)//RI-MP2/aug-cc-pV(D+d)Z-level in order to determine an upper limit to the

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratio of small molecular clusters in atmospheric conditions. Our calculations indicate that depending on the temperature, ammonia concentrations of around 10–10 000 ppb are required for the ratio to reach 1:1 (corresponding to ammonium bisulfate). Ratios of 2:1 (corresponding to ammonium sulfate) are unlikely to be found anywhere in the atmosphere. A sensitivity analysis indicates that errors in the computational method might lead to an overestimation of the ammonia concentration required for ammonium bisulfate to form by at most 2 orders of magnitude. However, the finding that ammonium sulfate molecular clusters do not form in the atmosphere is insensitive even to relatively large systematic errors.

Acknowledgements. The authors thank the Scientific Computing Center (CSC) in Espoo, Finland for computing time, and the Academy of Finland for financial support.

References

Adams, P. J., Seinfeld, J. H., and Koch, D. M.: Global concentrations of tropospheric sulfate, nitrate, and ammonium aerosol simulated in a general circulation model, *J. Geophys. Res.*, 104, D13791, 1999.

Ahlich, R., Bär, M., Häser, M., Horn, H., and Kölmel, C.: Electronic structure calculations on workstation computers: The program system Turbomole, *Chem. Phys. Lett.*, 162, 165–169, 1989.

Anttila, T., Vehkamäki, H., Napari, I., and Kulmala, M.: Effect of ammonium bisulphate formation on atmospheric water-sulphuric acid-ammonia nucleation, *Boreal Env. Res.*, 10, 511–523, 2005.

Ball, S. M., Hanson, D. R., Eisele, F. L., and McMurry, P. H.: Laboratory studies of particle nucleation: Initial results for H_2SO_4 , H_2O , and NH_3 vapors, *J. Geophys. Res. D*, 104, 23 709–23 718, 1999.

Bandy, A. R. and Ianni, J. C.: Study of Hydrates of H_2SO_4 Using Density Functional Theory, *J. Phys. Chem. A*, 102, 6533–6539, 1998.

Bates, T. S., Quinn, P. K., Coffman, D. J., Johnson, J. E., Miller, T. L., Covert, D. S., Wiedensohler, A., Leinert, S., Nowak, A., and Neususs, C.: Regional physical and chemical proper-

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ties of the marine boundary layer aerosol across the Atlantic during Aerosols99: An overview, *J. Geophys. Res.* 106, 20767–20782, 2001.

Berglen, T. F., Berntsen, T. K., Isaksen, I. S. A., and Sundet, J. K.: A global model of the coupled sulfur/oxidant chemistry in the troposphere: The sulfur cycle. *J. Geophys. Res.*, 109, D19310, doi:10.1029/2003JD003948, 2004.

Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochem. Cycles*, 11, 561–587, 1997.

Boys, S. F. and Bernardi, F.: The Calculation of Small Molecular Interactions by the Differences of Separate Total Energies, *Mol. Phys.*, 19, 553, 1970.

Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: A thermodynamic model of the system H-NH₄-SO₄-NO₃-H₂O at tropospheric temperatures, *J. Phys. Chem. A*, 102, 2137–2154, 1998.

Ding, C.-G., Laasonen, K., and Laaksonen, A.: Two Sulphuric Acids in Small Water Clusters, *J. Phys. Chem A*, 107, 8648–8658, 2003.

Dunning Jr., T. H., Peterson, K. A., and Wilson, A. K.: Gaussian basis sets for use in correlated molecular calculations. X. The atoms aluminum through argon revisited, *J. Chem. Phys.*, 114, 9244–9253, 2001.

Feng, Y. and Penner, J. E.: Global modeling of nitrate and ammonium: Interaction of aerosols and tropospheric chemistry, *J. Geophys. Res.*, 112, D01304, doi:10.1029/2005JD006404, 2007.

Foresman, J. B. and Frisch, Æ: *Exploring Chemistry with Electronic Structure Methods*, 2nd edition, Gaussian, Inc., Wallingford CT, U.S.A., 1996.

Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Montgomery, Jr., J. A., Vreven, T., Kudin, K. N., Burant, J. C., Millam, J. M., Iyengar, S. S., Tomasi, J., Barone, V., Mennucci, B., Cossi, M., Scalmani, G., Rega, N., Petersson, G. A., Nakatsuji, H., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Klene, M., Li, X., Knox, J. E., Hratchian, H. P., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Ayala, P. Y., Morokuma, K., Voth, G. A., Salvador, P., Dannenberg, J. J., Zakrzewski, V. G., Dapprich, S., Daniels, A. D., Strain, M. C., Farkas, O., Malick, D. K., Rabuck, A. D., Raghavachari, K., Foresman, J. B., Ortiz, J. V., Cui, Q., Baboul, A. G., Clifford, S., Cioslowski, J., Stefanov, B. B., Liu, G., Liashenko,

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**T. Kurtén et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A., Piskorz, P., Komaromi, I., Martin, R. L., Fox, D. J., Keith, T., Al-Laham, M. A., Peng, C. Y., Nanayakkara, A., Challacombe, M., Gill, P. M. W., Johnson, B., Chen, W., Wong, M. W., Gonzalez, C., and Pople, J. A.: Gaussian 03, Revision C.02, Gaussian, Inc., Wallingford CT, 2004.

5 Hanson, D. R. and Eisele, F. L.: Diffusion of H₂SO₄ in Humidified Nitrogen: Hydrated H₂SO₄, J. Phys. Chem. A, 104, 1715–1719, 2000.

Häser, M. and Ahlrichs, R: Improvements on the direct SCF method, J. Comput. Chem., 10, 104–111, 1989.

Helgaker, T., Klopper, W., Koch, H. and Noga. J.: Basis-set convergence of correlated calculations on water, Chem. Phys., 106, 9639–9646, 1997.

10 Ianni, J. C. and Bandy, A. R.: A theoretical study of the hydrates of (H₂SO₄)₂ and its implications for the formation of new atmospheric particles, J. Mol. Struct. (THEOCHEM), 497, 19–37, 2000.

Ianni, J. C. and Bandy, A. R.: A Density Functional Theory Study of the Hydrates of NH₃*H₂O*H₂SO₄ and Its Implications for the Formation of New Atmospheric Particles, J. Phys. Chem. A, 103, 2801–2811, 1999.

Jensen, F.: Introduction to Computational Chemistry, 2nd edition, John Wiley & Sons, Ltd, West Sussex, UK, 2007.

20 Kettle, A. J. and Andreae, M. O.: Flux of dimethylsulfide from the oceans: A comparison of updated data sets and flux models, J. Geophys. Res., 105, 26 793–26 808, 2000.

Kulmala, M., Hämeri, K., Aalto, P. P., Mäkelä, J. M., Pirjola, L., Nilsson, E. D., Buzorius, G., Rannik, Ü., dal Maso, M., Seidl, W., Hoffman, T., Janson, R., Hansson, H.-C., Viisanen, Y., Laaksonen, A. and O'Dowd, C. D.: Overview of the international project on biogenic aerosol formation in the boreal forest (BIOFOR), Tellus, 53B, 324–343, 2001.

25 Kulmala, M., Lehtinen, K. E. J., Laakso, L., Mordas, G. and Hämeri, K.: On the existence of neutral atmospheric clusters, Boreal Env. Res., 10, 79–87, 2005.

Kurtén, T., Sundberg, M. R., Vehkamäki, H., Noppel, M., Blomqvist, J., and Kulmala, M.: An Ab Initio and Density Functional Theory Reinvestigation of Gas-Phase Sulfuric Acid Monohydrate and Ammonium Hydrogensulfate, J. Phys. Chem. A 110, 7178–7188, 2006.

30 Kurtén, T., Torpo, L., Ding, C.-G., Vehkamäki, H., Sundberg, M. R., Laasonen, K. and Kulmala, M.: A density functional study on water-sulfuric acid-ammonia clusters and implications for atmospheric cluster formation, J. Geophys. Res., in press, 2007.

Larson, L. J., Largent, A., and Tao, F.-M.: Structure of the sulfuric acid-ammonia system and

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**T. Kurtén et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the effect of water molecules in the gas phase, *J. Phys. Chem. A.*, 103, 6786–6792, 1999.

Malm, W. C., Schichter, B. A., Pitchford, M. L., Ashbaugh, L. L., and Eldred, R. A.: Spatial and monthly trends in speciated fine particle concentration in the United States, *J. Geophys. Res.*, 109, D03306, doi:10.1029/2003JD003739, 2004.

5 McMurry, P. H.: A review of atmospheric aerosol measurements, *Atmos. Environ.*, 34, 1959–1999, 2000.

Møller, C. and Plesset, M. S.: Note on an Approximation Treatment for Many-Electron Systems, *J. Phy. Rev.*, 46, 618–622, 1934.

Napari, I., Noppel, M., Vehkamäki, H. and Kulmala, M.: An improved model for ternary nucleation of sulfuric acid - ammonia - water, *J. Chem. Phys.*, 116, 4221–4227, 2002.

10 Nash, D. G., Baer, T. and Johnson, M. V.: Aerosol mass spectrometry: An introductory review, *Int. J. of Mass Spectrometry*, 258, 2–12, 2006.

O’Dowd, C. D., Lowe, J. A., Smith, M. H., Davison, B., Hewitt, N., and Harrison, R. M.: (Biogenic sulphur emissions and inferred non-sea-salt-sulphate cloud condensation nuclei in and around Antrartica, *J. Geophys. Res.*, 102, 12 839–12 854, 1997.

15 Portmann, S.: MOLEKEL, Version 4.3.win32. Swiss Center for Scientific Computing (CSCS)/ETHZ, Switzerland, 2002.

Re, S., Osamura, Y., and Morokuma, K.: Coexistence of Neutral and Ion-Pair Clusters of Hydrated Sulfuric Acid (H₂SO₄)(H₂O)_n (n= 1-5) – A Molecular Orbital Study, *J. Phys. Chem. A.*, 103, 3535–3547, 1999.

20 Ricard, V., Jaffrezo, J.-L., Kerminen, V.-M., Hillamo, R. E., Sillanpää, M., Ruellan, S., Liousse, C., and Cachier, H.: Two years of continuous aerosol measurements in northern Finland, *J. Geophys. Res.*, 107(D11), doi:10.1029/2001JD000952, 2002.

Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III–IV campaigns in Heidelberg and Hyytiälä, *Atmos. Chem. Phys. Discuss.*, 6, 10 837–10 882, 2006.

25 Schaap, M., Denier Van Der Gon, H. A. C., Dentener, F. J., Visschedijk, A. J. H., Van Loon, M., ten Brink, H. M., Putaud, J.-P., Guillaume, B., Liousse, C., and Builtjes, P. J. H.: Anthropogenic black carbon and fine aerosol distribution over Europe, *J. Geophys. Res.*, 109, D18207, doi:10.1029/2003JD004330, 2004.

30 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley&Sons: New York, U.S.A, 1998.

Vehkamäki, H., Napari, I., Kulmala M., and Noppel, M: Stable ammonium bisulphate clusters in the atmosphere, *Phys. Rev. Lett.*, 93, 148501-1–148501-4, 2004.

Virkkula, A., Teinilä, K., Kerminen, V.-M., Saarikoski, S., Aurela, M., Viidanoja, J., Paatero, J., Koponen, I. K., and Kulmala, M.: Chemical composition of boundary layer aerosol over Atlantic Ocean and at an Antarctic site, *Atmos. Chem. Phys.* 6, 3407–3421, 2006.

Wavefunction, Inc: Spartan '02 Windows, Wavefunction, Inc. Irvine, CA, U.S.A., 2002. See also: <http://www.wavefun.com>.

Weigend, F. and Häser, M: RI-MP2: first derivatives and global consistency, *Theor. Chem. Acc.*, 97, 331–340, 1997.

Weigend, F., Häser, M., Patzelt, H., and Ahlrichs, R.: RI-MP2: Optimized auxiliary basis sets and demonstration of efficiency, *Chem. Phys. Lett.*, 294, 143–152, 1998.

Weigend, F., Köhn, A., and Hättig, C.: Efficient use of the correlation consistent basis sets in resolution of the identity MP2 calculations, *J. Chem. Phys.*, 116, 3175–3183, 2002.

ACPD

7, 2937–2960, 2007

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. The computed reaction Gibbs free energies, in units of kcal mol⁻¹, for the addition of one ammonia molecule to various clusters, with monomer pressures of 1 atm.

reaction	ΔG , kcal/mol
$\text{H}_2\text{SO}_4 + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3$	-8.7 ^a -5.3 ^b
$\text{H}_2\text{SO}_4 \cdot \text{H}_2\text{O} + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3 \cdot \text{H}_2\text{O}$	-7.6 ^a -4.4 ^b
$\text{H}_2\text{SO}_4 \cdot (\text{H}_2\text{O})_2 + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_2$	-9.1 ^a -4.3 ^b
$\text{H}_2\text{SO}_4 \cdot (\text{H}_2\text{O})_3 + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_3$	-7.8 ^a -3.6 ^b
$\text{H}_2\text{SO}_4 \cdot (\text{H}_2\text{O})_4 + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_4$	-7.5 ^a -4.2 ^b
$\text{H}_2\text{SO}_4 \cdot (\text{H}_2\text{O})_5 + \text{NH}_3 \leftrightarrow \text{H}_2\text{SO}_4 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_5$	-6.8 ^a -5.7 ^b
$(\text{H}_2\text{SO}_4)_2 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3$	-15.1 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot \text{H}_2\text{O} + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot \text{H}_2\text{O}$	-15.9 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_2 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_2$	-16.7 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_3 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_3$	-15.0 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_4 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_4$	-9.8 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_5 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_5$	-12.1 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_6 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_6$	-8.3 ^a
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{H}_2\text{O})_7 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot \text{NH}_3 \cdot (\text{H}_2\text{O})_7$	-6.4 ^a

a)At the PW91/DNP level and 265 K, Kurtén et al. (2007).

b)At the B3LYP/6-311++G(2d,2p) level and 273 K, Ianni and Bandy (1999).

**Estimating the
NH₃:H₂SO₄ ratio of
nucleating clusters**

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating the NH₃:H₂SO₄ ratio of nucleating clusters

T. Kurtén et al.

Table 2. The electronic energy E_0 (in Hartree), thermal contribution to the enthalpy H_{term} (in kcal mol⁻¹) and entropy S_{term} (in cal (Kmol)⁻¹) from calculations at T=298 K. E_{ODZ} and E_{OTZ} (a.u.) correspond to the aug-cc-pV(D+d)Z and aug-cc-pV(T+d)Z basis sets, respectively. All values are computed using the RI-MP2 method.

species	E_{ODZ} hartree	E_{OTZ} hartree	H_{term} kcal/mol	S_{term} cal/Kmol
NH ₃	-56.40765	-56.47724	23.99	49.80
(H ₂ SO ₄) ₂	-1398.38942	-1399.25446	57.58	113.04
(H ₂ SO ₄) ₂ ·NH ₃	-1454.84352	-1455.78118	84.13	119.73
(H ₂ SO ₄) ₂ ·(NH ₃) ₂	-1511.28801	-1512.29423	110.75	131.42
(H ₂ SO ₄) ₂ ·(NH ₃) ₃	-1567.71964	-1568.79619	136.35	147.29
(H ₂ SO ₄) ₂ ·(NH ₃) ₄	-1624.15452	-1625.30224	162.14	155.60

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating the NH₃:H₂SO₄ ratio of nucleating clusters

T. Kurtén et al.

Table 3. The electronic and free energies of ammonia addition reactions, in kcal mol⁻¹, with electronic energies extrapolated to the RI-MP2/aug-cc-pV(∞+d)Z level and thermal contributions computed at the RI-MP2/aug-cc-pV(D+d)Z level. All values correspond to a temperature of 298 K and monomer pressures of 1 atm.

Reaction	ΔE_0	ΔG (298 K)
$(\text{H}_2\text{SO}_4)_2 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)$	-31.85	-16.39
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3) + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_2$	-22.20	-8.18
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_2 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_3$	-15.71	-3.94
$(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_3 + \text{NH}_3 \leftrightarrow (\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_4$	-18.50	-4.28

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Table 4. Relative concentrations of different two-acid cluster types at different temperatures and ammonia concentrations.

	T=298 K	T=298 K	T=298 K	T=298 K
	[NH ₃]=1 ppt	[NH ₃]=100 ppt	[NH ₃]=10 ppb	[NH ₃]=1 ppm
(H ₂ SO ₄) ₂	48.792%	0.944%	0.009%	0
(H ₂ SO ₄) ₂ ·NH ₃	51.208%	99.046%	99.009%	50.198%
(H ₂ SO ₄) ₂ ·(NH ₃) ₂	0	0.010%	0.982%	49.763%
(H ₂ SO ₄) ₂ ·(NH ₃) ₃	0	0	0	0.039%
(H ₂ SO ₄) ₂ ·(NH ₃) ₄	0	0	0	0
	T=273 K	T=273 K	T=273 K	T=273 K
	[NH ₃]=1 ppt	[NH ₃]=100 ppt	[NH ₃]=10 ppb	[NH ₃]=1 ppm
(H ₂ SO ₄) ₂	0.937%	0.009%	0	0
(H ₂ SO ₄) ₂ ·NH ₃	99.060%	99.767%	81.706%	4.244%
(H ₂ SO ₄) ₂ ·(NH ₃) ₂	0.002%	0.223%	18.292%	95.025%
(H ₂ SO ₄) ₂ ·(NH ₃) ₃	0	0	0.001%	0.717%
(H ₂ SO ₄) ₂ ·(NH ₃) ₄	0	0	0	0.014%
	T=248 K	T=248 K	T=248 K	T=248 K
	[NH ₃]=1 ppt	[NH ₃]=100 ppt	[NH ₃]=10 ppb	[NH ₃]=1 ppm
(H ₂ SO ₄) ₂	0.004%	0	0	0
(H ₂ SO ₄) ₂ ·NH ₃	99.910%	92.095%	10.425%	0.101%
(H ₂ SO ₄) ₂ ·(NH ₃) ₂	0.086%	7.905%	89.481%	86.814%
(H ₂ SO ₄) ₂ ·(NH ₃) ₃	0	0	0.094%	9.076%
(H ₂ SO ₄) ₂ ·(NH ₃) ₄	0	0	0	4.009%

Estimating the NH₃:H₂SO₄ ratio of nucleating clusters

T. Kurtén et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.




Fig. 1. The optimized structures of the clusters in this study.

(a) $(\text{H}_2\text{SO}_4)_2$

(b) $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)$

(c) $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_2$

(d) $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_3$

(e) $(\text{H}_2\text{SO}_4)_2 \cdot (\text{NH}_3)_4$.

The hydrogen bonds are displayed with dotted lines. Some relevant bond lengths and other interatomic distances (in Ångström) are given in green. Color coding: white=hydrogen, red=oxygen, yellow=sulfur and blue=nitrogen.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Estimating the $\text{NH}_3:\text{H}_2\text{SO}_4$ ratio of nucleating clusters

T. Kurtén et al.




Fig. 2. Average $\text{NH}_3:\text{H}_2\text{SO}_4$ mole ratios as a function of the ammonia concentration, at three different temperatures. Solid lines correspond to the RI-MP2/aug-cc-pV(∞ +d)Z//RI-MP2/aug-cc-pV(D+d)Z data with harmonic frequencies. Dashed lines correspond to the sensitivity analysis data, in which vibrational frequencies have been scaled by 0.75 and a further term of -2 kcal/mol has been added to each ammonia addition reaction energy.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)