

# **Electronic supplement for “Simple measures of ozone depletion in the polar stratosphere”**

Rolf Müller<sup>1</sup>, Jens-Uwe Grooß<sup>1</sup>, Carsten Lemmen<sup>1,2</sup>, Daniel Heinze<sup>1</sup>, Martin Dameris<sup>3</sup>, and Greg Bodeker<sup>4</sup>

<sup>1</sup>ICG-1, Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

<sup>2</sup>now at Copernicus Instituut voor Duurzame Ontwikkeling en Innovatie, Universiteit Utrecht, 3584CS Utrecht, The Netherlands  
and Institut für Küstenforschung, GKSS-Forschungszentrum Geesthacht GmbH, 21502 Geesthacht, Germany

<sup>3</sup>DLR, IPA, Oberpfaffenhofen, Germany

<sup>4</sup>NIWA, Private Bag 50061, Omakau Central Otago, New Zealand

Manuscript submitted to

**Atmospheric Chemistry and Physics Discussions**

May 15, 2007

The purpose of this electronic supplement is to provide additional information on the mean ozone over the polar cap (see Figs. 1 and 2 of the main paper) for the Arctic in April (Fig. 1) and for the Antarctic in October (Fig. 2).

Further, we provide tabulated numbers for the values shown in several plots in  
5 the main paper for easier use. The ozone column and vortex data for individual  
years of the time slice experiments (shown in Fig. 6 of the main paper) are given  
in the file timeslice\_column.tsv. The values for mean column ozone over the polar  
cap (see Figs. 1 and 2 of the main paper) are given in avg\_o3column\_march.dat  
(for the Arctic) and avg\_o3column\_october.dat (for the Antarctic). The minimum  
10 of the daily average ozone in spring poleward of a threshold value (see Figs. 8 and  
9 of the main paper) are given in min\_o3column\_march.dat (for the Arctic) and  
min\_o3column\_october.dat (for the Antarctic). The Arctic data files also contain  
the values for  $V_{PSC}$  that are used in the paper. Finally, the file pfp.tsv provide the  
PFP values employed here (which constitute a slightly updated version of the values  
15 presented earlier, Tilmes et al. 2006).

## References

Tilmes, S., Müller, R., Engel, A., Rex, M., and Russell III, J.: Chemical ozone loss in the Arctic and Antarctic stratosphere between 1992 and 2005, *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL026925, 2006.


## Figure Captions

**Fig. 1.** latitude boundary at  $60^{\circ}\text{N}$ ,  $63^{\circ}\text{N}$ ,  $65^{\circ}\text{N}$ , and  $70^{\circ}\text{N}$ . Bottom panel: the April mean of Arctic ozone for a latitude boundary at  $63^{\circ}\text{N}$  is compared with calculations using the equivalent latitude of  $63^{\circ}\text{N}$  and the maximum gradient in potential vorticity (applied on the 475 K potential temperature surface) as vortex edge definitions. All averages are area weighted averages.


**Fig. 2.** Top panel: the September mean of Antarctic ozone for a latitude boundary at  $60^{\circ}\text{S}$ ,  $63^{\circ}\text{S}$ ,  $65^{\circ}\text{S}$ , and  $70^{\circ}\text{S}$ . Bottom panel: the September mean of Antarctic ozone for a latitude boundary at  $63^{\circ}\text{S}$  is compared with calculations using the equivalent latitude of  $63^{\circ}\text{S}$  and the Nash-criterion (applied on the 475 K potential temperature surface) as vortex edge definitions.



## Figures



**Fig. 1.** latitude boundary at 60°N, 63°N, 65°N, and 70°N. Bottom panel: the April mean of Arctic ozone for a latitude boundary at 63°N is compared with calculations using the equivalent latitude of 63°N and the maximum gradient in potential vorticity (applied on the 475 K potential temperature surface) as vortex edge definitions. All averages are area weighted averages.



**Fig. 2.** Top panel: the September mean of Antarctic ozone for a latitude boundary at 60°S, 63°S, 65°S, and 70°S. Bottom panel: the September mean of Antarctic ozone for a latitude boundary at 63°S is compared with calculations using the equivalent latitude of 63°S and the Nash-criterion (applied on the 475 K potential temperature surface) as vortex edge definitions.