

Seasonal changes in gravity wave activity measured by lidars at mid-latitudes

M. Rauthe^{1,*}, M. Gerding¹, and F.-J. Lübken¹

¹Leibniz-Institut für Atmosphärenphysik an der Universität Rostock, Kühlungsborn, Germany

* now at: Institut für Meteorologie und Klimaforschung, Universität Karlsruhe, Karlsruhe, Germany

Received: 18 June 2008 – Accepted: 23 June 2008 – Published: 17 July 2008

Correspondence to: M. Rauthe (monika.rauthe@imk.uka.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

M. Rauthe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

More than 230 nights of temperature measurements between 1 and 105 km have been performed at the Leibniz-Institute of Atmospheric Physics in Kühlungsborn with a combination of two different lidars, i.e. a Rayleigh-Mie-Raman lidar and a potassium lidar.

- 5 About 1700 h of measurements have been collected between 2002 and 2006. Apart from some gaps due to the adverse weather conditions the measurements are well distributed throughout the year. Comprehensive information about the activity of medium- and low-frequency gravity waves was extracted from this data set. The dominating vertical wavelengths found are between 10 and 20 km and do not show any seasonal
10 variation. In contrast the temperature fluctuations due to gravity waves experience a clear annual cycle with a maximum in winter. The most significant differences exist around 60 km where the fluctuations in winter are more than two times larger than they are in summer. Only small seasonal differences are observed above 90 km and below 35 km. Generally, the fluctuations grow from about 0.5 K up to 8 K between 20 and
15 100 km. Damping of waves is observed at nearly all altitudes and in all seasons. The planetary wave activity shows a similar structure in altitude and season as the gravity wave activity which indicates a strong coupling between the processes of the different scales. Combining the monthly mean temperatures and the fluctuations we show that the transition between winter and summer season and vice versa seems to start in the
20 mesopause region and to penetrate downward.

1 Introduction

In recent years gravity waves in the atmosphere have been the subject of intense research because they play a decisive role for atmospheric circulation, structure, variability and composition (see review by [Fritts and Alexander, 2003](#), and references therein).

- 25 Generated in the troposphere/lower stratosphere gravity waves propagate upward and deposit their energy and momentum when breaking or being filtered. They greatly

ACPD

8, 13741–13773, 2008

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



influence the winds and thermal structure of the middle and upper atmosphere (e.g. Lindzen, 1981; Holton, 1982; McLandress, 1998; Holton and Alexander, 2000).

A number of different techniques have been used to observe and analyse gravity waves in the lower, middle and upper atmosphere (e.g. Wilson et al., 1991b; Eckermann et al., 1995; Fritts and Hoppe, 1995; Hertzog et al., 2002). However, there have been only few measurements which resolve gravity waves with small vertical and horizontal scales and short periods. These include measurements by satellites (e.g. Wu and Waters, 1996a; Krebsbach and Preusse, 2007; Sofieva et al., 2007; Alexander et al., 2008), radars (e.g. Nakamura et al., 1996; Serafimovich et al., 2005), OH-
10 Imagers (e.g. Stockwell and Lowe, 2001; Suzuki et al., 2004), radiosonde soundings (e.g. Allen and Vincent, 1995; Vincent and Alexander, 2000) and rocket soundings (e.g. Eckermann et al., 1995; Rapp et al., 2004). The lidar is the only technique which resolves low- and medium-frequency gravity waves, which are an important part of the wave spectrum (see review by Fritts and Alexander, 2003), with an appropriate
15 time and height resolution over the entire altitude range from the troposphere up to the lower thermosphere. But most gravity wave analyses of lidar measurements cover only a small altitude range because only one single lidar technique is used (e.g. Gardner and Voelz, 1987; Wilson et al., 1991a,b; Schöch et al., 2004). Just a few case studies include the whole altitude range relevant for gravity waves from their source
20 (troposphere/lower stratosphere) to their main dissipation region (mesosphere) (e.g. Dao et al., 1995; Sica et al., 1995; Williams et al., 2004). These case studies do not allow to analyse the seasonal variation of gravity wave activity.

The purpose of this paper is to present mean features of gravity wave activity at mid-latitudes as function of season and altitude using the combined data set of two lidars.
25 The main focus is the analysis of vertical wavelengths, amplitudes and potential energies in the stratosphere and the mesosphere. This kind of ground-based observation is needed to understand the spectrum of gravity waves present in the atmosphere. Here, temperatures between 1 and 105 km are measured. They are calculated from data of two different lidars, namely a Rayleigh-Mie-Raman lidar (RMR lidar) and a potas-

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



sium lidar. Both are operated at the Leibniz-Institute of Atmospheric Physics (IAP) in Kühlungsborn, Germany (54° N, 12° E). For the calculation of temperatures temporal and vertical resolutions of 15 min and 1 km are used, respectively. This range of scales is unique and can only be covered by the combination of different lidar techniques
5 (Alpers et al., 2004). Altogether, more than 230 nights with more than 3 h measurement duration are examined and the presented results extend the winter-summer study of Rauthe et al. (2006).

The paper is organised as follows. In Sect. 2 short descriptions of the observational methods and the calculation of relevant gravity wave parameters are presented.
10 An example including three consecutive nights of measurements finishes this section. The seasonal variability of vertical wavelengths, amplitudes and potential energies are shown in Sect. 3. Section 4 includes a comparison between the monthly mean temperature fluctuations and temperatures to study the transition between the seasons and the different scales of variability. In Sect. 5 the discussion of the results is presented.
15 The conclusions are summarised in Sect. 6.

2 Lidar systems and data analysis

This section is divided in two parts which contain the technical base of the paper. First a brief description of our lidar systems is given. More details can be found in Alpers et al. (2004). Afterwards the analyses of gravity wave parameters are presented.

20 2.1 Measurement techniques

Temperature soundings from 1 to 105 km are preformed at IAP by the combination of two different lidar systems. As shown in Fig. 1 three different methods are applied in four altitude regions from the troposphere up to the lower thermosphere. A potassium resonance lidar examines the Doppler broadening of the potassium D1 resonance line by tunable narrow band laser. This technique provides absolute temperatures be-
25

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



tween ~85–105 km depending on the existing potassium layer. With the RMR lidar the Rayleigh backscatter at 532 nm wavelength is detected which yields atmospheric density profiles (20–90 km). These density profiles can be converted to temperature profiles through integration assuming hydrostatic equilibrium and using an initial temperature value from the potassium lidar. The altitude range is divided in two parts to manage the wide range of signal dynamics. Since February 2004 the effect of stratospheric aerosols are corrected via vibrational N₂ Raman signal (608 nm), before the correction was empirical (for details see Alpers et al., 2004; Gerding et al., 2004). Below 20 km the rotational Raman backscatter in two narrow wavelength ranges are used to measure the temperature.

One example of a temperature profile integrated over 1 h is shown in Fig. 1. The statistical uncertainty is typically 1.5–2.5 K and does not exceed 10 K. Only in the troposphere the typical temperature fluctuations are larger than the uncertainties of the measurement. Using the start values for Rayleigh temperature retrieval from potassium lidar avoid errors in the upper part of the Rayleigh temperature profile. Comparisons between climatological mean and observed temperature show even differences up to 20 K. Case studies have been shown that taking the start temperature from a climatology mean instead of the potassium lidar would give temperature fluctuation errors of up to 8 K.

To ensure continuous temperature profiles with a good trade-off between statistical uncertainties and temperature fluctuations the following procedure is necessary. The raw data profiles are smoothed applying an altitude-dependent filter (0.6 and 3 km at 40 and 80 km, respectively) and a running mean over 1 h with 15 min shift in time. At the end, the temperature data are binned into 1 km altitude ranges.

2.2 Gravity wave analysis

The calculation of parameters which characterise the gravity wave activity is based on the temperature deviations from the nightly mean (ΔT). These values are used to calculate the mean temperature fluctuations ($|\overline{\Delta T}|$) as an altitude-dependent parameter.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ter and are also called temperature fluctuations. With the temperature deviations the gravity wave potential energy density (GWPED) is derived. Two different values are distinguished: the potential energy per mass $E_{\text{pot,M}}$ and the potential energy per volume $E_{\text{pot,V}}$. The potential energy per mass can be calculated directly from our measurements

$$E_{\text{pot,M}} = \frac{1}{2} \frac{g^2}{N^2} \left(\frac{\tilde{\rho}}{\bar{\rho}} \right)^2 \approx \frac{1}{2} \frac{g^2}{N^2} \left(\frac{\tilde{T}}{\bar{T}} \right)^2. \quad (1)$$

This value depends on gravitational acceleration (g), Brunt-Väisälä frequency (N) and density (ρ) or temperature (T). Since the amplitudes (\tilde{T}) of the single waves are usually not known in observational studies which measure the net amplitude of many superimposed waves, means of amplitudes are used ($\tilde{T} = \Delta T$). Multiplying $E_{\text{pot,M}}$ with density gives $E_{\text{pot,V}}$. For lidar measurements the density has to be taken from a reference atmosphere like CIRA-86 (Fleming et al., 1990). In contrast to $E_{\text{pot,M}}$, which for an undamped propagating wave increases exponentially due the decreasing air density, $E_{\text{pot,V}}$ is approximately constant with altitude in the absence of wave dissipation.

Apart from amplitudes the vertical wavelength (λ_z) is also an important but not well-known property of gravity waves. Vertical wavelengths are calculated by wavelet analyses of single altitude profiles. A Morlet wavelet of fifth order, which is a good approach for many geophysical applications, is used (Torrence and Compo, 1998). Due to our enormous altitude range it is possible to derive vertical wavelengths from 5 up to 50 km. The nightly mean spectrum provides the mean characteristics of the single night. To determine the dominating wavelength at each altitude up to three maxima are calculated from this nightly mean spectrum. The periods are also determined by the wavelet analyses in a similar way as for λ_z . Periods are calculated from the time series at every altitude. Note that the periods (P) are obtained relative to the ground-based station (intrinsic values may be different because of Doppler shift).

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.3 Example of three consecutive nights

ACPD

8, 13741–13773, 2008

To demonstrate the night-to-night variability of our observations, three consecutive nights in October 2005 are presented in this section. Besides measurements at night from both lidars between 1 and 105 km there are measurements between 85 and 95 km during daylight conditions from the potassium lidar. Due to the weak signal the time resolution has to be reduced to a running mean over 2 h with 30 min shift for the continuous time series from the potassium lidar. Otherwise a calculation of temperature is not possible around noon because of high background noise level.

In Fig. 2 the temperature deviation from the nightly mean with the regular time resolution of a running mean over 1 h with 15 min shift are presented. In all three nights the observed wave pattern is dominated by only few waves. Comparing the wave structures of the different nights it is obvious that the temperature variations of the last night seem to be more unstructured than those of the two nights before. Furthermore, in this night the amplitudes are about 25% smaller than in the first night.

For extracting the details of these differences vertical wavelengths and periods are calculated by wavelet analyses. The mean wavelet spectra of the vertical wavelengths are shown in Fig. 3. The small amplitudes of the last night are noticeable. In the first night the dominating modes exist at wavelengths of about 13 km and 30 km in the whole possible altitude range. In contrast there is only one dominating mode of about 18 km in the second night. Below 60 km two other dominating vertical wavelengths of about 35 km and smaller than 10 km are also present. In the last night two dominating vertical wavelengths of about 12 km and 25 km are found.

The continuous temperature measurements of the potassium lidar allow to determine the periods longer than 6 h. In Fig. 4a the time series of temperature deviations at 90 km altitude are shown. It has to be noted, that for this analysis the time resolution is reduced to a running mean of 2 h with 30 min shift to calculate temperatures also at noon. Variations of about 12 h are already visible in the time series (Fig. 4a). In the results of the wavelet analyses (Fig. 4b) the two first days are dominated by a 12 h

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



variation with 10–14 K amplitude, which vanishes in the last night. These variations are indicative of tides. Furthermore, the semidiurnal tide generates about a third or a half of the total fluctuations, which is in agreement with other measurements in mid-latitudes (cp. [Shepherd and Fricke-Begemann, 2004](#)). Shorter periodic variations (2–6 h), which are often gravity waves, occur intermittently but do not persist more than three hours. The amplitudes of this shorter variations are lower than 6 K.

Combining the different spectral analyses the semidiurnal tide seems to appear with a vertical wavelength of 30–35 km, because this dominating mode of vertical wavelength is the only one which is not present in the third night. In this night the semidiurnal tide is not visible in the period analysis. Simultaneously, the absence of the semidiurnal tide produces more chaotic wave structures, which were also seen by [Fricke-Begemann and Höffner \(2005\)](#).

As shown before the tides influence largely the temperature fluctuations. Furthermore, the maximum duration for nighttime observations in summer are restricted due to the shorter nighttime conditions. In [Rauthe et al. \(2006\)](#) examples for winter and summer are described and the influences of the measurement duration are studied in detail. For these reasons the lidar measurements are restricted to a duration of 3–5 h in the following seasonal examination. Due to the restriction we can assume that the observed variations are mainly generated by gravity waves.

20 3 Seasonal gravity wave activity

The seasonal variation of the gravity wave parameters as observed by the combination of two lidars above Kühlungsborn are shown in the following paragraphs. As presented in the previous section the observed wave patterns are dominated by only few waves. Their mean characteristics regarding amplitudes, potential energies and vertical wavelengths are presented here. This analysis includes 236 measurements between June 2002 and October 2006 with a minimum duration of 3 h. The different seasons winter (Nov–Jan), spring (Feb–May), summer (June–July) and autumn (Aug–

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Oct) are represented by 27, 66, 56, and 87 measurements, respectively. The distribution throughout the different years and month is listed in Table 1. Apart from some gaps in winter due to adverse weather conditions, the observations are well distributed throughout the year and allow to study the seasonal behaviour.

5 3.1 Vertical wavelengths

Due to the large altitude range covered by the observation the analysis of the vertical wavelengths is very effective. As described in the section before one dominating vertical wavelength at each altitude is derived from the nightly mean wavelet spectrum. In Fig. 5a this dominating vertical wavelength is shown at the altitude of 50 km, in which the largest range of vertical wavelengths can be resolved. The crosses represent the results of the single nights. Independent of seasons the dominating modes found are between 6 and 48 km, i.e. they cover nearly the entire possible range. The mean vertical wavelength is about 20 km (cf. monthly means in red).

15 Due to the large day-to-day variability the dependence of vertical wavelengths on altitude is studied on the basis of monthly mean values. No seasonal variation is found at any altitude (cf. Fig. 5b), even though a seasonal dependence would be expected from a simplified theory. Assuming that the wave period does not change the following relation can be derived from a simplified dispersion relation for medium-frequency gravity waves (e.g. Fritts and Alexander, 2003):

$$20 \lambda_z \propto 1/N. \quad (2)$$

Since N is up to 25% smaller values in summer than in winter, it would be expected that λ_z is larger in summer than in winter. Possible reasons for the differences between theory and observation are the small number of measurements in winter, the large variability of λ_z and the missing resolution of vertical wavelengths smaller than 5 km. 25 However, in Fig. 5 the smaller vertical wavelengths ($\lambda_z < 25$ km) are more frequent than larger vertical wavelengths (for winter and summer cp. Fig. 8 of Rauthe et al., 2006). The distributions of dominating vertical wavelengths for spring and autumn are similar

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(not shown). It is likely that the small vertical wavelengths are typical for internal gravity waves, whereas large vertical wavelengths can also be caused by tides.

The interpretation of the altitude dependence in Fig. 5b is difficult because at the upper and lower edge of the observed altitude range the wavelet analysis does not resolve the largest vertical wavelengths due to the edge effects (e.g. Fig. 3). Only altitudes which have the same distance to the edge of the observed altitude range are comparable, i.e. 20 with 80 km, 30 with 70 km or 40 with 60 km. It seems that the mean vertical wavelengths increase with altitude. This result agrees well with the theoretical consideration in Eq. (2). In the stratosphere N is up to 30% larger compared to the mesosphere which results in smaller vertical wavelengths in the stratosphere.

The amplitudes of dominating vertical wavelengths can also be derived from wavelet spectra (not shown). In contrast to the vertical wavelengths the amplitudes display a clear annual cycle with maximum in winter in all altitudes between 30 and 80 km. Furthermore, the amplitudes of the dominating modes increase with altitude from about 1 K at 25 km up to 2.5 K (4.5 K) at 80 km in summer (winter). These results agree with the temperature fluctuations of all waves, which will be described in the next paragraph.

3.2 Temperature fluctuations

One of the important characteristics of gravity waves is the amplitude. The nightly mean temperature fluctuations of the whole year and in the altitude range between 20 and 100 km are shown in Fig. 6. For the first time this gravity wave activity is analysed in such a large altitude range. Due to the different lengths of the measurements the data evaluation is limited to durations of 3–5 h to get a comparable data set in all seasons. A more detailed analysis of the influence of the measurement duration on the amplitude can be found in Rauthe et al. (2006). To underline the general gravity wave variation the data in Fig. 6 are smoothed applying a Hanning filter of ± 15 d and ± 2 km. Generally, the temperature fluctuations increase with height from about 0.5 K at 25 km up to about 8 K at 95 km. This is expected due to the decreasing air density. But the growth rate of the fluctuations with altitude varies throughout the year. Between 35 and 70 km

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



altitude the increase is weaker in summer than in winter. But between 70 and 90 km the fluctuations grow faster in summer than in winter. This difference results in an annual variation of temperature fluctuations between 35 and 90 km altitude than are clearly seen in Fig. 6. The fluctuations in winter are up to 2.5 times larger than in summer.

5

3.3 Potential energy

As described before the largest seasonal differences of the gravity activity appear between winter and summer. To extract these seasonal differences the seasonal values of the potential energy per volume are calculated and shown in Fig. 7. The largest values of potential energy arise in winter and the smallest in summer. The values in autumn and spring are very similar and inbetween the values of summer and winter. As is well known the potential energy per volume is constant with height in case of undamped wave propagation. Nearly undamped propagation is only found below 40 km in summer and autumn in our observations. Furthermore, the weakest damping occurs in the summer mesosphere (60–85 km). A more detailed analysis of damping layers in winter and summer can be found in Rauthe et al. (2006). The large variations of the potential energy with height in winter may be a result of the small number of measurements in this season. However, the frequent occurrence of inversion layers in winter blocking gravity waves may produce the observed altitudinal variation of potential energy.

10

15

15

20

20

4 Planetary and gravity wave activity

For more detailed analyses of the different types of wave activity the standard deviations of the monthly mean temperatures and the monthly mean temperature fluctuations are used. In this connection the variability of the mean temperatures is used as measure for planetary wave activity, which is then compared with the gravity wave activity. Moreover, the transition between the strong gravity wave activity in winter and

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the weaker activity in summer is studied.

The standard deviations of the monthly mean of temperatures are displayed in Fig. 8. These values show the long-periodic variations of the temperature field due to the large scale atmospheric change, i.e. the planetary wave activity. Throughout the year 5 the planetary wave activity in the troposphere is large due to the weather activity with high and low pressure systems. Furthermore, the variability in the stratosphere and lower mesosphere between November and February is relatively high, especially in December and January, which is due to the displacement of the polar vortex position and the resulting favourable propagation conditions for planetary waves. In April and 10 September the transition between the high winter mesopause and the lower summer mesopause is marked by a large temperature variability. Between May and August the variability of the temperatures is low, which means that the planetary wave activity is weak. To characterise the different seasons we shortly describe the absolute temperatures, which are published in Gerding et al. (2008¹, Figs. 3 and 4). In the winter 15 stratosphere region high temperature of more than 300 K are found in our observations due to so called stratospheric warming events. The so called two level mesopause structure (She and von Zahn, 1998) is identifiable. But because of the variable extension of the metal layer from night to night, the higher winter-level (~103 km) of the mesopause is not always detectable. However, for the month between May and August 20 the lower and extremely cold mesopause at about 86 km is clearly visible.

The monthly mean temperature fluctuations are shown in Fig. 9. As described in Sect. 3.2 the largest seasonal differences between the monthly mean temperature fluctuations can be found between 40 and 70 km. Furthermore, the variability of the monthly means below 80 km is much smaller in summer than in winter. In further studies, which will be published in a separate paper, we will show that during so called 25 stratospheric warmings and connected mesospheric coolings the gravity wave activity

¹Gerding, M., Höffner, J., Lautenbach, J., Rauthe, M., and Lübken, F.-J.: Seasonal variation of temperatures between 1 and 105 km altitude at 54° N by lidar, *Atmos. Chem. Phys. Discuss.*, submitted, 2008.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



is smaller than for mean winter conditions. So the total variability of the gravity wave activity increases in the winter stratopause region and also in the lower mesosphere. Above 80 km the variability of the monthly mean fluctuations is too high to identify seasonal differences. In contrast to the temperatures and the planetary wave activity the temperature fluctuations (i.e. the gravity wave activity) are very similar and show typical summer characteristics between March and September and not only between May and August, i.e. the fluctuations increase only slowly up to 70 km.

Combining the variability of temperatures and fluctuations allows a comparison between the large scale wave activity, which is mainly caused by planetary waves, and the small scale wave activity caused by gravity waves. In our observations the planetary wave activity and the gravity wave activity show similar structures with season and altitude. Only below 35 km the gravity wave activity is very low throughout the year and the planetary wave activity is large in winter due to the changes in the polar vortex. In the winter stratosphere and mesosphere stratospheric warmings modify the planetary and gravity wave activity. The observations show a larger planetary wave activity and an increased variability of gravity wave activity, especially in the lower mesosphere. The wave activity is large in the upper mesosphere throughout the year due to the general growth of the wave amplitude with altitude. Furthermore, the total variability increases due to the inversion layers (Gerding et al., 2008¹). In summary, the observations show that the change of planetary wave activity leads to a change in the gravity wave activity and vice versa. Impressive examples of this coupling are the stratospheric warmings, in which the planetary wave activity increases and the gravity wave activity decreases because of stronger filtering. We suggest that this coupling between the different scales is due to the background fields, i.e. the temperature changes due to waves which imply also changes in the wind field and vice versa. The filtering and damping of the waves are also modified.

Combining the results of the temperatures (Fig. 4 of Gerding et al., 2008¹) and fluctuations (Fig. 9) some conclusions about the direction of the transition from summer to winter state are possible. In September and October the temperatures of the

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



mesopause region are clearly similar to the winter state, but the fluctuations in the lower mesosphere and upper stratosphere is clearly similar to the summer state. This result indicates that the temperature and its fluctuations change first in the mesopause region and later in the stratopause region. There are some indications for a similar mechanism in spring. This top-down motion in spring and autumn was also observed in the winds (e.g. Hoffmann et al., 2002) and confirms theoretical conceptions as described e.g. by Fritts (2003).

5 Discussion

Comparing our results with other measurements it has to be noted that depending on the measurement method different parts of the gravity wave spectrum are resolved. Furthermore, there are no long-term lidar observations which include the whole altitude range from the lower stratosphere up to the lower thermosphere. In the following discussion only lidar measurements in mid-latitudes are used from the RMR lidar in southern France (Haute-Provence, 44° N, 6° E and Biscarrosse, 44° N, 1° W), in Wales (Aberystwyth, 52° N, 4° E) and in Canada (Toronto, 44° N, 88° E), and for the mesopause region from the Na lidar in the USA (Urbana, 40° N, 88° W). In Urbana there are also RMR lidar measurements. In addition results of satellite and rocket measurements are discussed.

Due to vertical wavelengths all comparable measurements include only a part of the wave spectrum which is covered by our measurements. For this reason the comparison of absolute values is less useful. No indication of a seasonal dependence of vertical wavelengths are found in the lidar measurements of Gardner and Voelz (1987) and Gardner et al. (1989), which is in agreement with our results from the whole altitude range from the troposphere up to the thermosphere. However the data of the CRISTA satellite instrument from August 1997 show a dominating vertical wavelength of 14 km in 25 km altitude in winter on the Southern Hemisphere and a 2 km larger vertical wavelength in the summer of the Northern Hemisphere (Ern et al., 2004). Similar

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



results have also been found for HIRDLS in the altitude range from 20 up to 55 km (Alexander et al., 2008). But these differences are probably not only caused by seasonal effects but also by hemispheric effects, e.g. different sources of gravity waves.

In contrast to the paper of Offermann et al. (2006) there is no indication of a dramatic change in the temperature fluctuations in the upper mesosphere in our data, which also vary with the season. Only between March and September the fluctuations increase up to 70 km slowly and above strongly. But in the other months the increase of the amplitudes is more or less continuous in our measurements.

Because of the seasonal similarity of the temperature fluctuations and the potential energy both parameters will be discussed together as gravity wave activity in the following. As shown before by Rauthe et al. (2006) there is a clear seasonal variation of the gravity wave activity in mid-latitudes with maximum in winter, especially between 35 and 90 km altitude. This result is in agreement with other lidar measurements and also with data from radiosondes, satellites and rockets. Between 30 and 60 km all lidar measurements vary annually with maximum values in winter (Gardner et al., 1989; Wilson et al., 1991b; Mitchell et al., 1991; Whiteway and Carswell, 1995). In the data of Wilson et al. (1991b) from France an additional semiannual component of the gravity wave activity is observed above 60 km. Hence a second maximum in summer occurs, whose reason is not completely understood. They assume that the minima occur during weak mean wind intensity. The two times larger values of gravity wave activity in winter compared to the summer is also measured by the satellite instruments (e.g. Wu and Waters, 1996b; Ern et al., 2004; Alexander et al., 2008) and also by rockets (e.g. Hirota, 1984; Eckermann et al., 1995). An analysis of the SABER data shows also a semiannual variation (Krebsbach and Preusse, 2007) but in our latitude of 54° N the interpretation of the data is problematic because every two months the line of sight of the instrument changes. Only if the instrument is looking northward the latitude of 54° N is observed. In the mesopause region and also in the upper troposphere and lower stratosphere the differences between the seasons vanish more and more in all types of measurements (cp. Gardner and Voelz, 1987; Allen and Vincent, 1995; Tsuda

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The cause of the observed annual variation of the gravity wave activity was discussed by different authors. On the one hand the background temperature and density play an important role to explain the winter-summer-differences as shown by Eckermann et al. (1995), Alexander (1998). In our data the winter-summer-ratio of the temperature fluctuation is larger than one like the ratio of these background parameters. Moreover the altitudinal structure agrees roughly in our data (Rauthe et al., 2006). On the other hand the influence of the background wind via filtering is often discussed (e.g. Wilson et al., 1991b; Whiteway and Carswell, 1995). Lindzen (1981) demonstrated this connection of wind and waves, i.e. that the wind filtering of waves is different in winter and summer due to the eastward wind and westward wind in the middle atmosphere, respectively. But with our lidar data no statement is possible yet. First the gravity waves observed by a lidar station are not necessarily influenced by the local wind field below because of the tilted propagation of the waves. Second the sources of the gravity waves may be seasonally different. At the moment direct correlation between the strength of gravity wave activity and the wind direction and/or wind strength is not found in our data (not shown). Advanced analyses with ray tracing to determine the sources are necessary and also the combination of the lidar results with wind measurements of radars.

6 Conclusions and outlook

In this paper the measurements with two lidars (RMR lidar and potassium lidar) include the whole altitude range from the troposphere up to the lower thermosphere. The presented data set contains more than 230 nights with more than 1700 measurement hours. The data were analysed in terms of waves. As shown by a case study the night-to-night variability is high and depends also on tides. It seems that the presence of the semidiurnal tide produces clearer wave structures. In the whole data set monthly mean values of dominating vertical wavelengths between 10 and 20 km are observed. Even though the dependence of vertical wavelength and static stability would suggest

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



smaller vertical wavelengths in summer, no seasonal differences for the dominating vertical wavelengths are found in our data. The missing seasonal dependence may be due to not resolved waves with vertical wavelengths smaller than 5 km. Unlike the vertical wavelengths the temperature fluctuations and also the potential energy show a strong seasonal variation with maximum values in winter, which are up to 2.5 times larger than in summer. The causes of the annual variations are discussed in a previous paper (Rauthe et al., 2006). It is clear that the background temperature and density explain a main part of the winter-summer-differences of the fluctuations. The differences with season are small below 35 km and above 90 km. The nightly mean fluctuations increase with altitude from less than 1 K up to 8 K. Nevertheless, damping of waves occurs nearly in all seasons and altitudes. But the weakest damping in the mesosphere is found in the summer between 60 and 85 km, which confirms the results of the damping layer analysis of Rauthe et al. (2006). Comparing the planetary wave activity with the gravity wave activity the altitudinal structure and also the seasonal structure of the variations are similar. This similarity indicates that both types of activity are coupled, i.e. the different scales interact due to the background fields, e.g. due to the wind. In combination with the temperature results (cp. Gerding et al., 2008¹) we can conclude that the transition between the winter and summer season and vice versa in the temperature and their fluctuations starts in the mesopause region and then continues in the mesosphere and stratosphere, which confirms the theoretical concept of downward-control.

The lidar technique includes the whole altitude range from the troposphere/lower stratosphere up to the lower thermosphere. The main focus of our observations is the middle atmosphere, where propagation, filtering and dissipation occur. We have shown that the large altitude range and the high temporal and spatial resolution allow to study an important part of the wave spectrum and the coupling between the scales and also between the different layers of the atmosphere. Limitations of lidar measurements are that they are only done at a single location and do not bring any information about the wind. A combination with other measurements e.g. from radars is planned to eliminate

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the deficiencies and to extend and affirm the analyses presented here.

5 Acknowledgements. We thank Matthias Alpers (now at DLR Bonn) for developing the IAP RMR lidar until autumn 2002 and Josef Höffner for making the potassium lidar data available. Furthermore, we appreciate the support in lidar operation and maintenance of Torsten Köpnick and Michael Priester. In addition, we would like to thank our colleagues at the IAP and numerous students for their help in nighttime lidar soundings. Parts of this work were supported by the Deutsche Forschungsgemeinschaft under grant GE 1625/1-1.

References

- Alexander, M. J.: Interpretations of observed climatological patterns in stratospheric gravity wave variance, *J. Geophys. Res.*, 103, 8627–8640, doi:10.1029/97JD03325, 1998. [13756](#)
- 10 Alexander, M. J., Gillw, J., Cavanaugh, C. M. C., Eden, T., Francis, G., Halvorson, C., Hannigan, J., Khosravi, R., Kinnison, D., Lee, H., Massie, S., Nardi, B., Barnett, J., Hepplewhite, C., Lambert, A., and Dean, V.: Global estimates of gravity wave momentum flux from High Resolution Dynamics Limb Sounder observations, *J. Geophys. Res.*, 113, D15S18, doi:10.1029/2007JD008807, 2008. [13743](#), [13755](#)
- 15 Allen, S. J. and Vincent, R. A.: Gravity wave activity in the lower atmosphere: Seasonal and latitudinal variations, *J. Geophys. Res.*, 100, 1327–1350, doi:10.1029/94JD02688, 1995. [13743](#), [13755](#)
- 20 Alpers, M., Eixmann, R., Fricke-Begemann, C., Gerding, M., and Höffner, J.: Temperature lidar measurements from 1 to 105 km altitude using resonance, Rayleigh, and rotational Raman scattering, *Atmos. Chem. Phys.*, 4, 793–800, 2004, <http://www.atmos-chem-phys.net/4/793/2004/>. [13744](#), [13745](#)
- 25 Dao, P. D., Farely, R., Tao, X., and Gardner, C. S.: Lidar observations of the temperature profile between 25 and 105 km: Evidence of strong tidal perturbations, *Geophys. Res. Lett.*, 22, 2825–2828, doi:10.1029/95GL02950, 1995. [13743](#)
- Eckermann, S. D., Hirota, I., and Hocking, W. K.: Gravity wave and equatorial wave morphology of the stratosphere derived from long-term rocket soundings, *Quart. J. Roy. Meteorol. Soc.*, 121, 149–186, doi:10.1256/smsqj.52107, 1995. [13743](#), [13755](#), [13756](#)
- Ern, M., Preusse, P., Alexander, M. J., and Warner, C. D.: Absolute values of gravity wave

M. Rauthe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- momentum flux derived from satellite data, *J. Geophys. Res.*, 109, D20103, doi:10.1029/2004JD004752, 2004. [13754](#), [13755](#)
- Fleming, E. L., Chandra, S., Barnett, J. J., and Corney, M.: Zonal mean temperature, pressure, zonal wind, and geopotential height as functions of latitude, COSPAR international reference atmosphere: 1986, Part II: Middle atmosphere models, *Adv. Space Res.*, 10, 11–59, doi:10.1016/0273-1177(90)90386-E, 1990. [13746](#)
- Fricke-Begemann, C. and Höffner, J.: Temperature tides and waves near the mesopause from lidar observations at two latitudes, *J. Geophys. Res.*, 110, D19103, doi:10.1029/2005JD00770, 2005. [13748](#)
- Fritts, D. C.: Gravity waves, in: *Encyclopedia of Atmospheric Science*, edited by: Holton, J. R., Curry, J. A., and Pyle, J. A., 1308–1314, Academic Press, San Diego, USA, 2003. [13754](#)
- Fritts, D. C. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, *Rev. Geophys.*, 41, 3/1–64, doi:10.1029/2001RG000106, 2003. [13742](#), [13743](#), [13749](#)
- Fritts, D. C. and Hoppe, U.-P.: High-resolution measurements of vertical velocity with the European incoherent scatter VHF radar 2. Spectral observations and model comparisons, *J. Geophys. Res.*, 100, 16 827–16 838, doi:10.1029/95JD01467, 1995. [13743](#)
- Gardner, C. S. and Voelz, D. G.: Lidar studies of the nighttime sodium layer over Urbana, Illinois, 2. Gravity waves, *J. Geophys. Res.*, 92, 4673–4694, 1987. [13743](#), [13754](#), [13755](#)
- Gardner, C. S., Miller, M. S., and Liu, C. H.: Rayleigh lidar observations of gravity wave activity in the upper stratosphere at Urbana, Illinois, *J. Atmos. Sci.*, 46, 1838–1854, doi:10.1175/1520-0469(1989)046<1838:RLOOGW>2.0.CO;2, 1989. [13754](#), [13755](#)
- Gerding, M., Rauthe, M., and Höffner, J.: Temperature soundings from 1 to 105 km altitude by combination of co-located lidars, and its application for gravity wave examination, in: Reviewed and revised papers presented at the 22nd International Laser Radar Conference, edited by: Pappalardo, G. and Amodeo, A., vol. ESA SP-561, pp. 567–570, Nordwijk, The Netherlands, 2004. [13745](#)
- Hertzog, A., Vial, F., Mechoso, C. R., Basdevant, C., Cocquerez, P., Dubourg, V., and Nouel, F.: Planetary and gravity wave activity in the equatorial lower stratosphere as seen by ultra-long duration balloons, *Adv. Space Res.*, 30, 1381–1386, doi:10.1016/S0273-1177(02)00555-0, 2002. [13743](#)
- Hirota, I.: Climatology of gravity waves in the middle atmosphere, *J. Atmos. Terr. Phys.*, 46, 767–773, doi:10.1016/0021-9169(84)90057-6, 1984. [13755](#)
- Hoffmann, P., Singer, W., and Keuer, D.: Variability of the mesospheric wind field at middle and

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic latitudes in winter and its relation to stratospheric circulation disturbances, *J. Atmos. Solar-Terr. Phys.*, 64, 1229–1240, doi:10.1016/S1364-6826(02)00071-8, 2002. [13754](#)

Holton, J. R.: The role of gravity wave induced drag and diffusion in the momentum budget of the mesosphere, *J. Atmos. Sci.*, 39, 791–799, doi:10.1175/1520-0469(1982)039(0791:TROGWI)2.0.CO;2, 1982. [13743](#)

Holton, J. R. and Alexander, M. J.: The role of waves in the transport circulation of the middle atmosphere, in: *Atmospheric science across the stratopause*, edited by: Siskind, D. E., Eckermann, S. D., and Summers, M. E., vol. 123 of AGU Monograph Series, 21–35, American Geophysical Union, Washington D.C., USA, 2000. [13743](#)

Krebsbach, M. and Preusse, P.: Spectral analysis of gravity wave activity in SABER temperature data, *Geophys. Res. Lett.*, 34, L03814, doi:10.1029/2006GL028040, 2007. [13743](#), [13755](#)

Lindzen, R. S.: Turbulence and stress owing to gravity wave and tidal breakdown, *J. Geophys. Res.*, 86, 9707–9714, 1981. [13743](#), [13756](#)

McLandress, C.: On the importance of gravity waves in the middle atmosphere and their parameterization in general circulation models, *J. Atmos. Solar-Terr. Phys.*, 60, 1357–1383, doi:10.1016/S1364-6826(98)00061-3, 1998. [13743](#)

Mitchell, N. J., Thomas, L., and Marsh, A. K. P.: Lidar observations of long-period gravity waves in the stratosphere, *Ann. Geophys.*, 9, 588–596, 1991, <http://www.ann-geophys.net/9/588/1991/>. [13755](#)

Nakamura, T., Tsuda, T., Fukao, S., Manson, A. H., Meek, C. E., Vincent, R. A., and Reid, I. M.: Mesospheric gravity waves at Saskatoon (52° N), Kyoto (35° N), and Adelaide (35° S), *J. Geophys. Res.*, 101, 7005–7012, doi:10.1029/95JD03826, 1996. [13743](#)

Offermann, D., Jarisch, M., Oberheide, J., Gusev, O., Wohltmann, I., Russell III, J. M., and Mlynczak, M. G.: Global wave activity from upper stratosphere to lower thermosphere: A new turbopause concept, *J. Atmos. Solar-Terr. Phys.*, 68, 1709–1729, doi:10.1016/j.jastp.2006.01.013, 2006. [13755](#)

Rapp, M., Strelnikov, B., Mülleman, A., Lübken, F.-J., and Fritts, D. C.: Turbulence measurements and implications for gravity wave dissipation during the MaCWave/MIDAS rocket program, *Geophys. Res. Lett.*, 31, L24S07, doi:10.1029/2003GL019325, 2004. [13743](#)

Rauthe, M., Gerding, M., Höffner, J., and Lübken, F.-J.: Lidar temperature measurements of gravity waves over Kühlungsborn (54° N) from 1 to 105 km: A winter-summer comparison, *J. Geophys. Res.*, 111, D24108, doi:10.1029/2006JD007354, 2006. [13744](#), [13748](#), [13749](#),

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Schöch, A., Baumgarten, G., Fritts, D. C., Hoffmann, P., Serafimovich, A., Wang, L., Dalin, P., Mülleman, A., and Schmidlin, F. J.: Gravity waves in the troposphere and stratosphere during the MaCWAVE/MIDAS summer rocket program, *Geophys. Res. Lett.*, 31, L24S04, doi:10.1029/2004GL019837, 2004. [13743](#)
- Serafimovich, A., Hoffmann, P., Peters, D., and Lehmann, V.: Investigation of inertia-gravity waves in the upper troposphere/lower stratosphere over Northern Germany observed with collocated VHF/UHF radars, *Atmos. Chem. Phys.*, 5, 295–310, 2005, <http://www.atmos-chem-phys.net/5/295/2005/>. [13743](#)
- She, C. Y. and von Zahn, U.: Concept of a two-level mesopause: Support through new lidar observations, *J. Geophys. Res.*, 103, 5855–5863, doi:10.1029/97JD03450, 1998. [13752](#)
- Shepherd, M. and Fricke-Begemann, C.: Study of tidal variations in mesospheric temperature at low and mid latitudes from WINDII and potassium lidar observations, *Ann. Geophys.*, 22, 1513–1528, 2004, <http://www.ann-geophys.net/22/1513/2004/>. [13748](#)
- Sica, R. J., Sargoytchev, S., Argall, P. S., Borra, E. F., and Girard, L.: Lidar measurements taken with a large-aperture liquid mirror: 1. Rayleigh-scatter system, *Appl. Optics*, 34, 6925–6936, 1995. [13743](#)
- Sofieva, V. F., Kyrölä, E., Hassinen, S., Backman, L., Tamminen, J., Seppälä, A., Thölix, L., Gurvich, A. S., Kan, V., Dalaudier, F., Hauchecorne, A., Bertaux, J.-L., Fussen, D., Van hellemont, F., Fanton d'Andon, O., Barrot, G., Mangin, A., Guirlet, M., Fehr, T., Snoeij, P., Saavedra, L., Koopman, R., and Fraisse, R.: Global analysis of scintillation variance: Indication of gravity wave breaking in the polar winter upper stratosphere, *Geophys. Res. Lett.*, 34, L03812, doi:10.1029/2006GL028132, 2007. [13743](#)
- Stockwell, R. G. and Lowe, R. P.: Airglow imaging of gravity waves – 1. Results from a small network of OH nightglow scanning imagers, *J. Geophys. Res.*, 106, 17 185–17 203, doi:10.1029/2001JD900035, 2001. [13743](#)
- Suzuki, S., Shiokawa, K., Otsuka, Y., Ogawa, T., and Wilkinson, P.: Statistical characteristics of gravity waves observed by an all-sky imager at Darwin, Australia, *J. Geophys. Res.*, 109, D20S07, doi:10.1029/2003JD004336, 2004. [13743](#)
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Am. Meteorol. Soc.*, 79, 61–78, doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2, 1998. [13746](#)
- Tsuda, T., Nishida, M., Rocken, C., and Ware, R. H.: A global morphology of gravity wave activity in the stratosphere revealed by the GPS occultation data (GPS/MET), *J. Geophys.*

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Res., 105, 7257–7273, doi:10.1029/1999JD901005, 2000. [13755](#)

Vincent, R. A. and Alexander, M. J.: Gravity waves in the tropical lower stratosphere: An observational study of seasonal and annual variability, *J. Geophys. Res.*, 105, 17 971–17 982, doi:10.1029/2000JD900196, 2000. [13743](#)

5 Wang, L. and Geller, M. A.: Morphology of gravity-wave energy as observed from 4 years (1998–2001) of high vertical resolution U.S. radiosonde data, *J. Geophys. Res.*, 108, 4489, doi:10.1029/2002JD002786, 2003. [13756](#)

Whiteway, J. A. and Carswell, A. I.: Lidar observations of gravity wave activity in the upper stratosphere over Toronto, *J. Geophys. Res.*, 100, 14 113–14 124, doi:10.1029/95JD00511, 10 1995. [13755](#), [13756](#)

Williams, B. P., Fritts, D. C., Wang, L., She, C.-Y., Vance, J. D., Schmidlin, F. J., Goldberg, R. A., Müllemann, A., and Lübken, F.-J.: Gravity waves in the arctic mesosphere during the MaCWAVE/MIDAS summer rocket program, *Geophys. Res. Lett.*, 31, L24S05, doi:10.1029/2004GL020049, 2004. [13743](#)

15 Wilson, R., Chanin, M. L., and Hauchecorne, A.: Gravity waves in the middle atmosphere observed by Rayleigh lidar – 1. Case studies, *J. Geophys. Res.*, 96, 5153–5167, 1991a. [13743](#)

Wilson, R., Chanin, M. L., and Hauchecorne, A.: Gravity waves in the middle atmosphere observed by Rayleigh lidar – 2. Climatology, *J. Geophys. Res.*, 96, 5169–5183, 1991b. [13743](#), 20 20 [13755](#), [13756](#)

Wu, D. L. and Waters, J. W.: Gravity-wave-scale temperature fluctuations seen by UARS MLS, *Geophys. Res. Lett.*, 23, 3289–3292, doi:10.1029/96GL02924, 1996a. [13743](#)

Wu, D. L. and Waters, J. W.: Satellite observations of atmospheric variances: A possible indication of gravity waves, *Geophys. Res. Lett.*, 23, 3631–3634, doi:10.1029/96GL02907, 1996b. 25 [13755](#)

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

Table 1. Number of measurements longer than 3 h per year and month. Data before June 2002 and after October 2006 are not included in this paper.

Jahr	J	F	M	A	M	J	J	A	S	O	N	D	Σ
2002	—	—	—	—	—	1	0	1	2	0	2	2	8
2003	0	5	0	4	1	5	1	7	7	8	2	3	43
2004	1	6	3	8	7	4	6	10	7	9	2	1	64
2005	3	3	7	11	2	10	5	7	7	10	2	1	68
2006	8	1	2	0	6	8	16	1	10	1	—	—	53
Σ	12	15	12	23	16	28	28	26	33	28	8	7	236

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		



Table 2. Monthly mean temperatures fluctuations (K) between 20 and 100 km from all soundings longer than 3 h as shown in Fig. 6

altitude (km)	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
20	0.7	0.8	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.5	0.6	0.5
22	0.7	0.7	0.6	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.6
24	0.7	0.7	0.6	0.5	0.5	0.4	0.4	0.4	0.5	0.5	0.6	0.6
26	0.8	0.7	0.6	0.6	0.5	0.5	0.4	0.4	0.6	0.5	0.6	0.6
28	0.8	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.6	0.6	0.7	0.7
30	0.8	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.7	0.7	0.7	0.8
32	0.9	0.9	0.9	0.8	0.7	0.7	0.7	0.7	0.8	0.8	0.8	0.9
34	1.1	1.1	1.0	1.0	0.8	0.8	0.8	0.8	0.9	1.0	1.0	1.2
36	1.3	1.3	1.2	1.1	1.0	1.0	1.0	1.0	1.0	1.2	1.1	1.5
38	1.4	1.4	1.3	1.2	1.2	1.1	1.1	1.1	1.0	1.4	1.3	1.8
40	1.5	1.6	1.4	1.3	1.3	1.2	1.1	1.2	1.1	1.5	1.5	2.1
42	1.6	1.7	1.5	1.3	1.3	1.1	1.2	1.2	1.0	1.5	1.6	2.2
44	1.6	1.8	1.5	1.3	1.3	1.1	1.2	1.2	1.0	1.4	1.7	2.3
46	1.7	1.9	1.5	1.3	1.3	1.1	1.3	1.3	1.0	1.5	1.6	2.4
48	1.8	1.9	1.5	1.3	1.3	1.1	1.3	1.3	1.1	1.6	1.6	2.6
50	1.8	1.9	1.4	1.3	1.3	1.1	1.2	1.4	1.2	1.7	1.8	2.6
52	1.9	1.9	1.5	1.4	1.3	1.2	1.2	1.4	1.3	1.7	2.3	2.5
54	2.1	2.1	1.6	1.5	1.3	1.3	1.2	1.5	1.5	1.8	2.6	2.6
56	2.2	2.4	1.8	1.4	1.3	1.3	1.2	1.5	1.5	2.0	2.6	3.2
58	2.2	2.5	1.9	1.4	1.3	1.3	1.2	1.6	1.5	2.2	2.5	3.7
60	2.2	2.6	2.1	1.4	1.4	1.4	1.3	1.6	1.5	2.4	2.7	3.9
62	2.4	2.9	2.1	1.6	1.4	1.5	1.3	1.7	1.5	2.7	3.0	3.7
64	2.9	3.2	2.2	1.8	1.4	1.6	1.4	1.8	1.6	2.9	3.3	3.4
66	3.1	3.4	2.4	1.9	1.5	1.8	1.5	1.8	1.7	2.9	3.8	3.2
68	3.3	3.7	2.7	1.9	1.7	1.8	1.6	1.9	1.8	2.9	4.1	3.0
70	3.7	4.0	3.0	2.0	1.8	1.9	1.7	2.0	2.0	2.8	4.0	2.9
72	4.1	3.9	3.2	2.3	1.9	2.1	1.9	2.4	2.3	2.9	3.8	3.0
74	4.2	4.0	3.4	2.6	2.1	2.3	2.3	2.8	2.7	3.0	4.1	3.5
76	4.2	4.3	3.4	2.8	2.3	2.5	2.7	3.1	3.0	3.3	4.5	3.9
78	4.4	4.7	3.7	3.2	2.6	2.8	3.0	3.3	3.4	3.6	4.9	4.2
80	4.7	5.0	4.1	3.7	3.0	3.2	3.3	3.5	3.8	3.9	5.3	4.7
82	5.3	5.0	4.6	4.3	3.3	3.5	3.6	3.8	4.2	4.4	5.6	5.4
84	6.0	5.2	5.1	4.8	3.7	4.0	4.2	4.2	4.7	4.9	6.2	5.7
86	6.3	5.5	5.5	5.0	4.2	4.4	4.9	4.6	4.9	5.3	6.8	5.3
88	6.1	5.9	5.8	5.1	4.8	4.5	5.3	4.9	5.0	5.4	7.0	5.1
90	6.0	6.2	6.1	5.0	5.3	4.5	5.2	5.1	4.9	5.3	6.5	5.3
92	6.1	6.5	6.2	5.0	5.6	4.5	5.2	5.2	4.9	5.3	6.2	5.6
94	6.3	7.2	6.2	5.3	5.9	4.8	5.7	5.5	5.3	5.5	6.4	5.9
96	6.5	7.9	6.4	5.7	6.5	5.6	6.3	6.1	5.9	6.1	7.2	6.1
98	7.4	8.1	6.9	6.4	7.0	6.7	6.4	6.9	6.4	6.9	7.8	6.1
100	8.2	7.9	7.2	7.0	7.0	7.4	6.5	7.6	6.6	7.6	8.1	6.5



Seasonal changes in gravity wave activity measured by lidars
M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.




Fig. 1. Temperature profile obtained from a combination of different lidar techniques measured on 5 October 2005, 19:45–20:45 UT, above Kühlungsborn. The colored dots at 88 km and 48 km indicate the start value for the Rayleigh temperature retrieval.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 2. Temperature deviations from the nightly mean for three consecutive nights (5–8 Oct 2005). The dotted lines indicate the transition heights between the different measurement methods.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 3. Mean wavelet spectrum of vertical wavelengths for the temperature profiles shown in Fig. 2 (5–8 Oct 2005). Pluses indicate the local maxima. The hatched area indicates the “cone of influence”.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.




Fig. 4. Measurements of the potassium lidar (5–8 Oct 2005) at 90 km altitude using a running mean over 2 h with 30 min shift in time: **(a)** Time series of temperature deviations from the nightly mean. **(b)** Wavelet analyses of periods. The hatched area indicates the “cone of influence”. Note the non linear y-axis .

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)




Fig. 5. Dominating vertical wavelengths: **(a)** Results for a single altitude at 50 km. The black crosses indicate the results of single measurements. The red line and dots show the monthly mean. **(b)** Results for different altitudes as monthly means. Note the different y-axes.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 6. Seasonal variation of nightly mean temperature fluctuations between 20 and 100 km, i.e. the gravity wave activity. 236 measurements are used and the duration of the single measurements is restricted to intervals of 3–5 h. The data are smoothed with a Hanning filter of ± 15 d and ± 2 km.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.




Fig. 7. Profiles of potential energy per volume for different altitudes. A Hanning filter of ± 5 km is used to smooth the data.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 8. Standard deviation of the monthly mean temperatures between 0 and 105 km characterising the planetary wave activity. 236 measurements are used.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 9. Monthly mean temperature fluctuations for (a) October–March and (b) April–September characterising the gravity wave activity. The numbers indicate the number of measurements per month. The bars show the standard deviation. A Hanning filter of ± 5 km is used to smooth the data.

Seasonal changes in gravity wave activity measured by lidars

M. Rauthe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

