

**Interpretation of
radiative signature in
transition zones**

J. C. Chiu et al.

Physical interpretation of the spectral radiative signature in the transition zone between cloud-free and cloudy regions

J. C. Chiu¹, A. Marshak², Y. Knyazikhin³, P. Pilewskie⁴, and W. J. Wiscombe^{2,5}

¹University of Maryland Baltimore County, Baltimore, MD, USA

²NASA/Goddard Space Flight Center, Greenbelt, MD, USA

³Boston University, Boston, MA, USA

⁴University of Colorado at Boulder, Boulder, CO, USA

⁵Brookhaven National Laboratory, New York, NY, USA

Received: 29 July 2008 – Accepted: 22 August 2008 – Published: 26 September 2008

Correspondence to: J. C. Chiu (christine.chiu@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

One-second-resolution zenith radiance measurements from the Atmospheric Radiation Measurement program's new shortwave spectrometer (SWS) provide a unique opportunity to analyze the transition zone between cloudy and cloud-free air, which has considerable bearing on the aerosol indirect effect. In the transition zone, we find a remarkable linear relationship between the sum and difference of radiances at 870 and 1640 nm wavelengths. The intercept of the relationship is determined primarily by aerosol properties, and the slope by cloud properties. We then show that this linearity can be predicted from simple theoretical considerations and furthermore that it supports the hypothesis of inhomogeneous mixing, whereby optical depth increases as a cloud is approached but the effective drop size remains unchanged. In addition, the width of transition zones from SWS data is in the range of 50–150 m, which differs from the width in satellite observations (a few kilometers) and in airborne lidar data (1–2 km).

1 Introduction

The aerosol indirect effect is the largest source of uncertainty in the radiative forcing of climate (The Intergovernmental Panel on Climate Change, IPCC, Fourth Assessment Report, 2007). Using 11 GCM models, Stephens (2002) also showed the importance of cloud feedbacks in modeling responses of climate to a doubling of carbon dioxide. We cannot evaluate performance of climate models without accurate knowledge of aerosol forcing and cloud feedbacks (Diner et al., 2004).

Studies on aerosol direct and indirect effects demand a precise separation of cloud-free and cloudy areas (Charlson et al., 2007; Koren et al., 2007). However, separation between cloud-free and cloudy areas from remotely-sensed measurements is ambiguous. From the ground, separations have been made using broadband pyranometer, microwave radiometer, total sky imager, radar, lidar, and ceilometer data (Long and

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ackerman, 2000; Berendes et al., 2004; Long et al., 2006a, b; Taylor et al., 2008). Each instrument has a different field of view, sensitivity, and sampling resolution; in addition, each method uses different thresholds for the separation. From satellites, the separation depends on spatial resolution, illumination and observation geometry, surface types, and screening algorithms (Ackerman et al., 1998; Martins et al., 2002; Brennan et al., 2005; Gomez-Chova et al., 2007). While a separation is not free of ambiguity at any scale (Koren et al., 2008), it is important to understand the transition zone between cloud-free and cloudy areas.

Yet it has been difficult to study the transition zone using conventional data. Both satellite and in situ aircraft data are inadequate. Satellite data are hampered both by lack of high enough spatial resolution and by ambiguity in interpreting radiances due to 3-D radiative transfer effects. As a result, aerosol retrievals in the vicinity of clouds may be contaminated by undetected clouds (Zhang et al., 2005) as well as by radiation reflected from clouds (Marshak et al., 2008). In addition, cloud properties in the collection 5 Moderate Resolution Imaging Spectroradiometer (MODIS) product are not reported for cloudy pixels that border clear-sky pixels, because those cloudy pixels may include both clear and cloudy areas, and the retrievals of their microphysical properties are not reliable (Coakley et al., 2005).

On the other hand, in situ data are hampered by lack of fast enough time resolution. For measuring cloud microphysical properties, most probes have no better than 1 to 10 Hz sampling rate, though some probes can sample faster (e.g., 2000 Hz from Gerber probes, Davis et al., 1999; and 1000 Hz from Fast Forward Scattering Spectrometer probe, Brenguier et al., 1998). However, due to tiny sample volumes, averaging over longer time periods is usually necessary to achieve statistical significance. If measurements are averaged to 1–2 Hz with an aircraft speed of 100 m/s, the spatial resolution will be 50–100 m, which is not fine enough to study physical processes around cloud edges.

In situ data are also hampered by noise in measurements. For example, standard hot-wire liquid water probes are noisy at the 0.2 g/m^3 level. Uncertainty of cloud micro-

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



physical measurements is in an order of 20% (Allan et al., 2008). The variations in the transition zone would be masked by such large noise values.

This paper aims to study changes of aerosol and cloud properties in the transition zone from radiative signatures measured by the new Atmospheric Radiation Measurement (ARM) Program shortwave spectrometer (SWS). The SWS is the first ground-based instrument that measures zenith radiances with high temporal (1-s) and spectral resolution in the visible and near-infrared region. Spectra from SWS contain rich information on radiative properties of aerosols and clouds to advance our understanding of physical processes in the transition zone, such as activation and evaporation of cloud droplets and humidification of aerosols.

2 Shortwave spectrometer and ancillary ARM data

The SWS, a ground-based instrument based upon the design of the airborne Solar Spectral Flux Radiometer (Pilewskie et al., 2003), was first deployed in March 2006 at the ARM Oklahoma site. The SWS measures zenith radiance at 418 wavelengths between 350 and 2170 nm. The spectral resolution for visible and near infrared regions is 8 and 12 nm, respectively. The field of view is 1.4° . The integration time of each 1-s measurement is about 300 ms. The SWS is calibrated biweekly using the on-site ARM 12'' integrating sphere that is in turn calibrated by the 30'' sphere at the NASA Ames Research Center. Therefore, the absolute accuracy of measurements depends on the accuracy of the transfer standard from the 30'' sphere to the 12'' sphere. The 30'' sphere has an accuracy of 1–2%.

We used two discrete SWS wavelengths of 870 and 1640 nm to explore their spectral changes in more details. We selected these wavelengths to minimize Rayleigh scattering and maximize the sensitivity of zenith radiance to cloud optical depth and cloud drop size. Neither liquid water nor vapor absorb sunlight at 870 nm wavelength. On the other hand, liquid water absorbs weakly at 1640 nm (and is amplified by multiple scattering within cloud), where there is negligible absorption by water vapor and car-

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bon dioxide. We further normalized zenith radiance measurement $I_{m,\lambda}$ at wavelength λ using:

$$I_{\lambda} = \frac{\pi \cdot I_{m,\lambda}}{\mu_0 \cdot F_{\text{TOA},\lambda}}, \quad (1)$$

where I_{λ} is the normalized zenith radiance; $F_{\text{TOA},\lambda}$ is the solar irradiance at the top of the atmosphere; and μ_0 is the cosine of solar zenith angle (SZA). We denote normalized zenith radiances at 870 and 1640 nm hereafter as I_{870} and I_{1640} , respectively, as well as

$$\text{SUM} = I_{870} + I_{1640} \quad (2a)$$

$$\text{DIF} = I_{870} - I_{1640} \quad (2b)$$

which are in many cases more informative and illustrative than radiances themselves.

We also used ancillary instruments and products to better understand atmospheric state and cloud field for case study. First, the ARM Total Sky Imager (TSI) captures cloud field images at a 30-s sampling interval with a half-hemispheric field of view (Long et al., 2001). A shadowband on the mirror blocks the intense direct-normal light from the sun. Although clouds of interest are those at the center of TSI images and are often blocked by the shadowband, one can see cloud evolution and movement from time series of cloud images. Second, cloud boundary heights were obtained at a 10-s resolution from the ARM Active Remotely Sensed Clouds Locations (ARSCL) product, based on measurements of cloud radar, micropulse lidar, and ceilometer (Clothiaux et al., 2000). Third, liquid water path was retrieved from ARM microwave radiometer measurements at a 20-s resolution (Turner et al., 2007). Finally, wind speed was estimated from the merged sounding product (Miller et al., 2003). Note that the temporal resolutions of all ancillary data (10, 20, 30-s, and 1-min) are much lower than that of the SWS (1-s).

In addition to clouds, aerosols and the underlying surface also affect zenith radiance. Aerosol properties were obtained from the NASA Aerosol Robotic Network standard

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



products (Holben et al., 1998). The underlying surface around the site is vegetated. For a single leaf, the reflectance at 870 nm is generally 1.2–1.5 times greater than that at 1640 nm (Walter-Shea and Norman, 1991). However, surface albedo depends not only on properties of single leaf, but also on canopy structures (Knyazikhin et al., 1997).

5 In this paper, values of surface albedo were based on MODIS collection 5 retrievals at a 500 m resolution (Schaaf et al., 2002).

3 Observed spectral signatures from SWS in the transition zone

We chose three cases (Table 1) to show how the spectral signature of the transition zone changes between cloud-free and cloudy areas for different solar zenith angles.

10 We made our choices using the following criteria. First, we separated cloudy from cloud-free times in the SWS data using radiances at wavelengths of 673 and 870 nm. Marshak et al. (2004) suggested that over a vegetated surface, a larger radiance at 673 nm than at 870 nm indicates a cloud-free situation. For cloudy situations, the situation reverses; the radiance at 870 nm becomes larger. Second, we limited ourselves
15 to cases in which the cloud-free and cloudy periods both lasted at least one minute, to avoid very small clouds and gaps. This criterion leads to 300–500 m sizes of cloud and gap, typical for fair weather cumulus (Joseph and Cahalan, 1990; Lane et al., 2002). Third, to avoid messy situations in which the cloud itself is very fragmented, we excluded cases in which the ratio of radiances at 673 nm and 870 nm changed by more
20 than 10% during cloudy periods.

3.1 Case 1

Case 1 is a single small cumulus cloud passing over; we look at the transitions at both the beginning and end of the passage. Images of the sky dome, taken by the TSI, are shown in Fig. 1a. The cumulus cloud of interest was blocked by the shadowband at
25 21:35 UTC, but could be seen in TSI images after 21:36 UTC. The cloud moved toward

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the northeast (upper-right corner of the sky images) with a 3 m s^{-1} speed at its cloud-base height of 2 km. Liquid water path of the cloud was $\sim 16 \text{ g m}^{-2}$. The sun was in the west with a solar zenith angle of 45° . Because the sun was shining behind the cloud, the radiometer was on the cloud's shadowed side before the cloud passed over, and the illuminated side after.

Figure 1b shows the time series of Case 1 SWS spectral radiances as a color contour plot. The structure of this plot will be similar for the other two cases, so we describe it here briefly. The times ($T1$, etc.) indicated correspond to the images in Fig. 1a. The brighter colors indicate the cloudy period, the darker colors the cloud-free period, and the intermediate colors the transition period, which obviously has a finite temporal (and hence horizontal) extent. The vertical black bands during the cloudy period are absorption bands of water vapor.

Performing cuts of Fig. 1b at wavelengths of 870 and 1640 nm (marked by arrows on the x-axis) leads to the two time series in Fig. 1c. During the cloud-free period, both radiances I_{870} and I_{1640} are small, and I_{870} is greater than I_{1640} due to stronger molecular (Rayleigh) and aerosol scattering; neither radiance changes significantly during this period. In the transition period from cloud-free to cloudy, I_{870} and I_{1640} both increase sharply and switch order.

Figure 1d re-plots the data from Fig. 1c on the DIF vs SUM plane (cf. Eqs. 2a and b). Here, one can follow the evolution from cloud-free (upper left corner) to cloudy (lower right corner) and back again to near the starting point; the arrows indicate the flow of time. The main message of this figure is that, in both transition periods, there is a linear relationship between SUM and DIF. In the cloudy period, Fig. 1c shows three peaks of I_{870} and I_{1640} due to internal cloud variability; this variation causes the wandering behavior displayed by the black cloudy points.

The linear relationships shown in the two transition periods have a similar slope. Radiances in the second transition period are slightly higher because the spectrometer is viewing the illuminated side of the cloud. At the illuminated cloud edge, more photons are scattered into the radiometer than for the shadowed cloud edge in the first transition

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



period. Note that this type of radiance enhancement in cloud-free areas is similar to what Wen et al. (2007) and Marshak et al. (2008) found in satellite reflectance.

3.2 Case 2

Case 2 involves the edge of a large cumulus cloud that just strikes a kind of grazing blow, and then is followed by a small puff of cloud; we look at the transition out of the large cloud and into the small puff. The large cloud (around the shadowband in TSI images of Fig. 2a) moved toward the south with a 6 m s^{-1} speed at its cloud-base height of 3.5 km. Liquid water path of the cloud was $\sim 16 \text{ g m}^{-2}$. The small puffy cloud approached the FOV of SWS (the center of TSI images) at 18:54:00 and 18:54:30 UTC. The sun was near overhead with a SZA of 15° . For this case, the sun illuminated the cloud edge in the first transition period.

The contour plot of SWS radiances (Fig. 2b) clearly shows two transition periods, even before the small puff cloud was about to enter the FOV. Radiances I_{870} and I_{1640} in this case behave similar to what we have found in Case 1. First, Fig. 2c shows that during the clear sky period, I_{870} is greater than I_{1640} , and neither fluctuates substantially. When clouds approach, I_{870} and I_{1640} increase sharply. Second, for both transition periods there is a linear relationship between SUM and DIF (Fig. 2d). The SUM was slightly greater in the first transition period than that in the second one because of cloud edge illuminations. The slopes of the linear relationships are close.

3.3 Case 3

Case 3 is a low large cumulus cloud passing over; we look at the transitions at the end of the passage. Figure 3a shows that in the beginning of this case, the low cumulus cloud was imaged at the center of the TSI images, and high cirrus and alto-cumulus clouds were to the west. The low cumulus cloud moved toward the north with a 7 m s^{-1} speed at its cloud-based height of 1 km. The liquid water path retrieval is not available for the cloud. The sun was in the west with a SZA of 65° , illuminating the cumulus

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cloud during the whole period.

Figure 3b shows that SWS spectral radiances are higher (brighter colors) during the cloud's passage, due to enhanced cloud scattering. In the beginning of the cloudy period, radiance I_{870} is much larger than I_{1640} , and the difference between two gradually becomes smaller (Fig. 3c). In the transition to the cloud-free period, radiances I_{870} and I_{1640} both decrease sharply and switch order back and forth. Similar to the previous two cases, Fig. 3d shows a linear relationship between SUM and DIF in the transition period.

3.4 Radiative signature regimes

Based on the above cases, we have found the following:

- During cloud-free periods, radiances at 870 and 1640 nm are small. The radiance at 870 nm is higher than that at 1640 nm because of stronger molecular (Rayleigh) and aerosol scattering at shorter wavelengths.
- During transition periods from cloud-free to cloudy, radiances at both wavelengths increase sharply in the vicinity of cloud edges. A remarkable linear relationship is found between $I_{870} - I_{1640}$ (DIF) and $I_{870} + I_{1640}$ (SUM) at various solar zenith angles. The slopes of linear relationships for different transition periods are close, but their intercepts differ and depend on sun-cloud-radiometer illumination.
- During cloudy periods, I_{870} and I_{1640} are much higher than those in cloud-free periods. Whether the difference $I_{870} - I_{1640}$ is positive or negative depends on a number of factors, such as aerosol and cloud optical depth, particle and droplet size, 3-D cloud structure, surface reflectance, and solar zenith angle.

Figure 4 is a schematic plot to show the above distinct spectral signatures found in those three cases. This plot is drawn based on 1-D plane-parallel radiative transfer calculations. In this plot, we define 5 regimes on the DIF vs SUM plane. These regimes are:

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of radiative signature in transition zones

J. C. Chiu et al.

- Regime 0 corresponds to cloud-free areas. It is a single point on the DIF vs SUM plane.
- Regime 1 corresponds to transition zones in which cloud optical depth τ up to ~ 0.2 . In this regime, SUM increases, but DIF decreases. The relationship between DIF and SUM is linear. (In cases of very large SZAs and small cloud droplets, the slope could be also positive. See Sect. 3.)
- Regime 2 corresponds to areas with very thin clouds (τ up to ~ 1). This regime is same as Regime 1, but the relationship between SUM and DIF is no longer linear.
- Regime 3 corresponds to areas with thin clouds (τ up to ~ 5). In this regime, both SUM and DIF increase, and the relationship between two is nonlinear.
- Regime 4 corresponds to areas with thicker clouds ($\tau > 5$). In this regime, SUM decreases while DIF increases and the relationship between two is strongly nonlinear.

Physical interpretations of these radiative signature regimes are discussed next.

4 Physics of radiative transfer behind the spectral signatures

For plane-parallel clouds over a Lambertian surface, any ground-based measurement of radiance I can be expressed as the sum of the downward radiation calculated over a non-reflecting (black) surface and the radiation introduced by interactions between clouds and the underlying surface (Box et al., 1988). The downward radiance over a black surface is determined by scattering from atmospheric molecules, aerosols, and clouds. The cloud-surface interactions are determined by surface albedo and cloud reflective and transmissive properties. In short, the spectral signatures in zenith radiance are primarily determined by four factors:

1. Molecular (Rayleigh) and aerosol scattering,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2. in-cloud single scattering,
3. in-cloud multiple scattering, and
4. cloud-surface interactions.

4.1 Regime 0

5 Rayleigh and aerosol scattering dominate Regime 0. Due to stronger Rayleigh and aerosol scatterings at 870 than those at 1640 nm, I_{870} is larger than I_{1640} , and thus $I_{870}-I_{1640}$ is positive.

This regime strongly depends on aerosol loading and aerosol particle size. We used three different aerosol loadings: no aerosols, low aerosol optical depth (AOD, $\tau_{870}^a=0.05$, $\tau_{1640}^a=0.02$, Ångström exponent $\approx 2/3$), and high AOD ($\tau_{870}^a=0.15$, $\tau_{1640}^a=0.08$, Ångström exponent ≈ 1). With increasing aerosol loading (Fig. 5a), Regime 0 moves toward the upper-right direction, i.e., both $I_{870}+I_{1640}$ and $I_{870}-I_{1640}$ increase. The sum $I_{870}+I_{1640}$ increases because of the increase of aerosol scattering at both two wavelengths, while $I_{870}-I_{1640}$ increases because of greater impact of stronger scattering on the shorter wavelength.

Regime 0 also strongly depends on aerosol phase function. We show an example here by increasing aerosol particle size from effective radius of $0.25\ \mu\text{m}$ to $4\ \mu\text{m}$ for a given SZA of 45° (Fig. 5b). From aerosol phase functions (Fig. 6), a larger aerosol particle size results in stronger forward scatterings and weaker scattering at scattering angles greater than 20° . It leads to decreases in both I_{870} and I_{1640} at larger scattering angles. However, the rate of decreases in I_{870} is different from that in I_{1640} . In the scattering angle range between 20° and $\sim 50^\circ$, the rate of decrease in I_{870} is faster than that in I_{1640} , and vice versa for scattering angles greater than 50° . Therefore, at a given SZA of 45° (i.e., scattering angle of 45°), both I_{870} and I_{1640} decrease, and I_{870} decreases faster than I_{1640} with increasing aerosol particle size. As a result, Regime 0 moves toward the lower-left direction with increasing aerosol particle size.

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 5 also shows that the locations of Regime 1 are sensitive to changes in aerosol properties, but the slope of the linear relationship is only weakly sensitive to those changes. We use radiative transfer calculations to understand this behavior in next section.

5 4.2 Regime 1

Aerosol scattering and in-cloud single scattering dominate Regime 1. In this regime, photons are scattered not only by aerosols, but also by cloud droplets. Therefore, approaching clouds leads to an increase in I_{870} and I_{1640} . However, the behavior of DIF vs SUM is complex because it depends on a number of variables: the cosine of SZA μ_0 , the single scattering albedo, phase function and optical depth of aerosols and clouds. We denote them as ϖ_λ^a , P_λ^a , and τ_λ^a , respectively for aerosols, and ϖ_λ^c , P_λ^c , τ^c for clouds. The subscript λ shows the wavelength dependency of ϖ and P . Note that aerosol optical depth (AOD) is wavelength dependent, but cloud optical depth (COD) is wavelength independent at the wavelengths we used.

15 Ignoring molecular (Rayleigh) scattering, the total optical depth is given as:

$$\tau_\lambda = \tau_\lambda^a + \tau^c. \quad (3)$$

Using the single-scattering approximation and assuming a unit incident flux at the top of the atmosphere, the downward zenith radiance is derived as (Thomas and Stamnes, 2002, p. 219):

$$20 I_\lambda \propto \varpi_\lambda \cdot P_\lambda \cdot \frac{\mu_0}{1 - \mu_0} [\exp(-\tau_\lambda) - \exp(-\tau_\lambda / \mu_0)], \quad (4a)$$

where ϖ_λ and P_λ are the total single scattering albedo and phase function. For a very small optical depth, Eq. (4a) can be simplified as

$$I_\lambda \propto \varpi_\lambda \cdot P_\lambda \cdot \tau_\lambda. \quad (4b)$$

Figure 7 illustrates the right side of Eqs. (4a) and (4b) with $\varpi_\lambda=1$ and $P_\lambda=1$ for three SZAs: 15° , 45° and 60° . Comparing curves of Eq. (4a) with the straight line of Eq. (4b),

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



we see that the linear approximation is good for small optical depths up to ~ 0.1 – 0.2 , depending on μ_0 .

Let us first assume that cloud droplets are the only scatterers in Regime 1. Then, because $\varpi_{870}^c = 1$, Eq. (4b) determines the ratio of DIF to SUM as:

$$\frac{I_{870} - I_{1640}}{I_{870} + I_{1640}} = \frac{P_{870}^c - \varpi_{1640}^c P_{1640}^c}{P_{870}^c + \varpi_{1640}^c P_{1640}^c} = \frac{1 - \varpi_{1640}^c \chi}{1 + \varpi_{1640}^c \chi}, \quad (5)$$

where χ is the ratio between P_{1640}^c and P_{870}^c , i.e.,

$$\chi = \chi(\mu_0; r_{c,\text{eff}}) = \frac{P_{1640}^c}{P_{870}^c}, \quad (6)$$

where $r_{c,\text{eff}}$ is the cloud droplet effective radius. Equation (5) shows that the ratio is independent of cloud optical depth. It follows from here that at a given μ_0 the slope of the DIF vs. SUM relationship plane is fully determined by cloud drop size in the case of $\tau_\lambda^a = 0$. Figure 8 shows the dependence of the slope on cloud droplet effective radius for three μ_0 values. We see that for small droplets less than $4 \mu\text{m}$, the slope derived from Eq. (5) is very sensitive to droplet size while it asymptotes for larger droplets. We also notice that at smaller μ_0 , the slope can be positive when cloud effective radius is very small ($< 2.5 \mu\text{m}$).

Now we assume a more general case of $\tau_\lambda^a > 0$. Then in the frame of linear approximation Eq. (4b), Eq. (5) can be rewritten as

$$\frac{I_{870} - I_{1640}}{I_{870} + I_{1640}} = \frac{a_- + (P_{870}^c - \varpi_{1640}^c P_{1640}^c) \cdot \tau^c}{a_+ + (P_{870}^c + \varpi_{1640}^c P_{1640}^c) \cdot \tau^c}, \quad (7)$$

where

$$a_\pm = \varpi_{870}^a P_{870}^a \tau_{870}^a \pm \varpi_{1640}^a P_{1640}^a \tau_{1640}^a. \quad (8)$$

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



From Eq. (7), we see that DIF vs. SUM is a linear function and the corresponding slope is:

$$\text{Slope} = \frac{P_{870}^c - \omega_{1640}^c P_{1640}^c}{P_{870}^c + \omega_{1640}^c P_{1640}^c} = \frac{1 - \omega_{1640}^c \chi}{1 + \omega_{1640}^c \chi}, \quad (9)$$

where χ is defined in Eq. (6). This slope is the same as that in the special case of $\tau_\lambda^a = 0$ (Eq. 5). It shows that the slope is determined by cloud drop size only and independent of aerosol properties, which has been observed in Fig. 5.

Thus, assuming constant aerosol optical depth and constant effective sizes of cloud droplets and aerosol particles, Regime 1 shows a linear relationship between DIF and SUM as:

$$\text{DIF} = a + b \cdot \text{SUM}, \quad (10)$$

where as a first approximation, intercept a is determined primarily by aerosol properties, and slope b by cloud properties, defined in Eq. (9). It shows that the linear relationship nicely separates aerosol and cloud effects into a and b , respectively. However, aerosol properties are not the sole factor that would affect the intercept a . For the same aerosol properties, the Sun's relative location with respect to clouds could change the intercept a through 3-D radiative effects. Case 1 (Fig. 1) is the perfect example, showing that the Sun's location introduces a difference in the intercept for the two transition periods.

Note that this linear relationship becomes less pronounced with increasing cloud optical depth, in which first the linear approximation (Eq. 4b) and then the single scattering approximation (Eq. 4a) is no longer held. Generally, the relationship remains linear at COD smaller than ~ 0.2 depending on μ_0 .

4.3 Regime 2–4

Because our main purpose here is to study the transition zone, the detailed description and simulations of Regime 2–4 are beyond the scope of the paper. Hence we

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



briefly highlight main features for each regime. For the difference and the sum between 670 nm and 870 nm we can reference to Marshak et al. (2004) and Chiu et al. (2006).

Regime 2 is a continuation of Regime 1 in which in-cloud single scattering dominates. However, unlike Regime 1, the linear approximation Eq. (4b) is no longer valid in Regime 2 and radiances show a non-linear relationship with increasing optical depth (as shown in Fig. 7).

In-cloud multiple scatterings and cloud-surface interactions dominate Regime 3 and 4. Equations (4a) and (4b) are no longer valid in these two regimes. In Regime 3, cloud optical depths are typically less than 4–5. This regime is in a situation in which in-cloud multiple scattering dominates and surface-cloud interaction starts playing an important role. Recall that the surface-cloud interactions increase zenith radiance. Because the surface is brighter at 870 nm than at 1640 nm, I_{870} increases faster than I_{1640} , and thus the DIF increases with increasing COD.

Finally, Regime 4 associates with larger cloud optical depths (>5). Because less transmission gets through thicker clouds, zenith radiances at 870 and 1640 nm decrease. Similar to Regime 3, I_{870} increases faster than I_{1640} due to cloud-surface interactions. As a result, SUM gradually decreases while DIF increases.

5 Discussion

From radiative transfer calculations in Eqs. (4)–(7), we showed that linearity between SUM and DIF in the transition zone can be explained using the single scattering approximation and the following assumptions: constant cloud droplet and aerosol particle sizes, and constant AOD. To further discuss whether these assumptions are realistic, we review cloud microphysical measurements and some theoretical studies in more details.

A number of studies showed that liquid water content increased when approaching clouds (Paluch and Baumgardner, 1989; Stith, 1992; Blyth et al., 2005). The increase of liquid water content is the result of either/both increase in droplet number concen-

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Interpretation of
radiative signature in
transition zones**J. C. Chiu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tration or/and effective radius. Which process dominates? Recall that in all cases we observed, zenith radiances sharply increased from cloud-free to cloudy areas (Figs. 1c, 2c, 3c). An increase of number concentration with constant effective radius leads directly to the increase of cloud optical depth and thus of zenith radiance. The assumption of a constant effective radius is also consistent with the theory of inhomogeneous mixing that results in fast evaporation of droplets of all sizes leaving effective radius unchanged (e.g., Baker et al., 1980; Freud et al., 2008). Therefore, as a very first approximation, our assumption of constant effective radius seems realistic.

Let us now assume that effective radius increases when clouds approach. According to Eq. (4b) zenith radiance is proportional to a product of cloud optical depth and cloud scattering phase function. Cloud optical depth is also proportional to the droplet concentration and the square of droplet size. The phase function, in general, has a non-monotonic dependency on effective radius (Fig. 9). Because it is more likely that droplets are small in the transition zone, we focus on drop size from 2 to 4 μm . For these droplet sizes, at SZA of 15° , the scattering phase function increases slightly at 870 nm (Fig. 9c), but decreases sharply at 1640 nm (Fig. 9d). Therefore, no matter whether the number of droplet concentration increases or remains unchanged, the decrease in phase function at 1640 nm leads to a slower increase in zenith radiance at 1640 nm than that at 870 nm. This contradicts SWS observations of Case 2, because zenith radiance increased faster at 1640 nm (Fig. 2c).

We have assumed that aerosol particle size and optical depth remain unchanged in our theoretical derivations. However, depending on aerosol compositions, aerosol extinction coefficients and sizes could increase dramatically around cloud edges because of increasing uptake of water vapor (Kotchenruther et al., 1999; Clarke et al., 2002; Loeb and Manalo-Smith, 2005; Zhang et al., 2005). Recent observations from an airborne HSRL (High Spectral Resolution Lidar) have demonstrated that when approaching clouds, aerosol optical depth and extinctions could increase by 10% and 20%, respectively (R. Ferrare, personal communication). More detailed model simulations are required to understand how these changes in aerosol optical properties

modulate spectral signatures of SWS in the transition zone.

Radiative signatures in the transition zone are characterized not only by changes in aerosol and cloud properties, but also by the 3-D cloud effects. As shown in our cases, these effects could enhance zenith radiance, depending on sun-cloud-radiometer locations. In addition, the field of view (FOV) of the SWS is 1.4° . When the FOV is not fully covered by clouds or by clear-sky, radiative signatures could be affected as well. For more details see Chiu et al. (2006).

6 Summary

Although to the casual observer clouds appear to have distinct boundaries, to define them from remote-sensed measurements is difficult (Koren et al., 2008). This difficulty arises from the sensitivity of instruments, the temporal and spatial resolutions, and the thresholds used to distinguish cloud-free and cloudy areas. This problem has major climatic consequences, in particular on aerosol indirect effect studies and on calculations of aerosol forcing.

The ARM program developed a new shortwave spectrometer to measure zenith radiance in the visible and near-infrared region with high spectral and temporal resolutions. We have taken advantage of the SWS rich spectra to study the transition zone around clouds where strong aerosol-cloud interactions are taking place. We demonstrate the general spectral radiative characteristics, and provide physics interpretations to better understand optical properties of aerosols and clouds in the transition zone.

From cloud-free to cloudy areas, we define 5 regimes based on spectral signatures of zenith radiance and the factors that dominate the signatures. While Regimes 2–4 correspond to a typical well-defined cloud, in this paper we focus on Regime 0 and 1 that represent the cloud-free area and the transition zone around clouds, respectively.

In Regime 0, because of the absence of clouds, spectral signature is mainly determined by Rayleigh and aerosol scattering, and thus strongly depends on aerosol loading and particle size in the environment. This indicates that one can qualitatively

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Interpretation of
radiative signature in
transition zones**

J. C. Chiu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

study aerosol optical properties using SWS zenith radiance in the cloud-free areas. In Regime 1, spectral signature is mainly determined by aerosol and in-cloud single scatterings. Due to in-cloud single scatterings, we observed and theoretically proved that there would be a linear relationship between the sum and difference of zenith radiances at wavelengths of 870 and 1640 nm. The linear behavior allowed us to separate effects of aerosols and clouds: the intercept is more affected by aerosol properties while the slope by clouds. This linear relationship holds for small optical depths only.

To quantitatively interpret the slope of the linear relationship observed in Regime 1, we have assumed that droplet sizes remain unchanged. Analyzing SWS data, we have found that this assumption is reasonable; it is also consistent with the predictions of the inhomogeneous mixing process. Thus, based on zenith radiance measurements at 870 and 1640 nm, the decrease in droplet number concentrations with the distance away from clouds likely dominates over the decrease in droplet sizes.

Using SWS radiance measurements to study transition zones has a few defects, though. We need to wait for clouds to pass overhead, and the SWS only works in daytime. In addition, other ARM instruments do not measure on the 1-sec timescale of SWS and thus cannot help resolve ambiguities.

Acknowledgements. This research was supported by the Office of Science (BER, US Department of Energy, Interagency Agreement No. DE-AI02-95ER61961) as part of the ARM program.

References

- Ackerman, S. A., Strabala, K. I., Menzel, W. P., Frey, R. A., Moeller, C. C., and Gumley, L. E.: Discriminating clear sky from clouds with MODIS, *J. Geophys. Res.*, 103, 32 141–32 158, 1998.
- Allan, J. D., Baumgardner, D., Raga, G. B., Mayol-Bracero, O. L., Morales-García, F., García-García, F., Montero-Martínez, G., Borrmann, S., Schneider, J., Mertes, S., Walter, S., Gysel, M., Dusek, U., Frank, G. P., and Krämer, M.: Clouds and aerosols in Puerto Rico – a new evaluation, *Atmos. Chem. Phys.*, 8, 1293–1309, 2008, <http://www.atmos-chem-phys.net/8/1293/2008/>.

- Baker, M. B., Corbin, R. G., and Latham, J.: The influence of entrainment on the evolution of cloud droplet spectra: I. A model of inhomogeneous mixing. *Q. J. Roy. Meteor. Soc.*, 106, 581–598, 1980.
- Berendes, T. A., Berendes, D. A., Welch, R. M., Dutton, E. G., Uttal, T., and Clothiaux, E. E.: Cloud cover comparisons of the MODIS daytime cloud mask with surface instruments at the North Slope of Alaska ARM site, *IEEE T. Geosci. Remote*, 42, 2584–2593, 2004.
- Blyth, A. M., Lasher-Trapp, S. G., and Cooper, W. A.: A study of thermals in cumulus clouds, *Q. J. Roy. Meteor. Soc.*, 131, 1171–1190, 2005.
- Box, M. A., Gerstl, S. A. W., and Simmer, C.: Application of the adjoint formulation to the calculation of atmospheric radiative effects, *Beitr. Phys. Atmos.*, 61, 303–311, 1988.
- Brenguier, J.-L., Bourriane, T., Coelho, A. D., Isbert, J., Peytavi, R., Trevarin, D., and Weschler, P.: Improvements of droplet size distribution measurements with the Fast-FSSP (Forward Scattering Spectrometer Probe), *J. Atmos. Ocean. Tech.*, 15, 1077–1090, 1998.
- Brennan, J. I., Kaufman, Y. J., Koren, I., and Li, R. R.: Aerosol-cloud interaction – Misclassification of MODIS clouds in heavy aerosol, *IEEE T. Geosci. Remote*, 43, 911–915, 2005.
- Charlson, R. J., Ackerman, A. S., Bender, F. A.-M., Anderson, T. L., and Liu, Z.: On the climate forcing consequences of the albedo continuum between cloudy and clear air, *Tellus B*, 59, 715–727, 2007.
- Chiu, J. C., Marshak, A., Knyazikhin, Y., Wiscombe, W. J., Barker, H. W., Barnard, J. C., and Luo, Y.: Remote sensing of cloud properties using ground-based measurements of zenith radiance, *J. Geophys. Res.*, 111, D16201, doi:10.1029/2005JD006843, 2006.
- Clarke, A. D., Howell, S., Quinn, P. K., Bates, T. S., Ogren, J. A., Andrews, E., Jefferson, A., Massling, A., Mayol-Bracero, O., Maring, H., Savoie, D., and Cass, G.: INDOEX aerosol: A comparison and summary of chemical, microphysical, and optical properties observed from land, ship, and aircraft, *J. Geophys. Res.*, 107(D19), 8033, doi:10.1029/2001JD00572, 2002.
- Clothiaux, E. E., Ackerman, T. P., Mace, G. G., Moran, K. P., Marchand, R. T., Miller, M. A., and Martner, B. E.: Objective determination of cloud heights and radar reflectivities using a combination of active remote sensors at the ARM CART sites, *J. Appl. Meteorol.*, 39, 645–665, 2000.
- Coakley, J. A., Friedman, M. A., and Tahnk, W. R.: Retrieval of cloud properties for partly cloudy imager pixels, *J. Atmos. Ocean. Tech.*, 22, 3–17, 2005
- Davis, A., Marshak, A., Gerber, H., and Wiscombe, W. J.: Horizontal structure of marine bound-

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ary layer clouds from centimeter to kilometer scales, *J. Geophys. Res.*, 104, 6123–6144, 1999.

Diner, D. J., Ackerman, T. P., Anderson, T. L., Bosenberg, J., Braverman, A. J., Charlson, R. J., Collins, W. D., Davies, R., Holben, B. N., Hostetler, C. A., Kahn, R. A., Martonchik, J. V., Menzies, R. T., Miller, M. A., Ogren, J. A., Penner, J. E., Rasch, P. J., Schwartz, S. E., Seinfeld, J. H., Stephens, G. L., Torres, O., Travis, L. D., Wielicki, B. A., and Yu, B.: PARAGON – an integrated approach for characterizing aerosol climate impacts and environmental interactions, *B. Am. Meteorol. Soc.*, 85, 1491–1501, 2004.

Freud, E., Rosenfeld, D., Andreae, M. O., Costa, A. A., and Artaxo, P.: Robust relations between CCN and the vertical evolution of cloud drop size distribution in deep convective clouds, *Atmos. Chem. Phys.*, 8, 1661–1675, 2008, <http://www.atmos-chem-phys.net/8/1661/2008/>.

Gomez-Chova, L., Camps-Valls, G., Calpe-Maravilla, J., Guanter, L., and Moreno, J.: Cloud-screening algorithm for ENVISAT/MERIS multispectral images, *IEEE T. Geosci. Remote*, 45, 4105–4118, 2007.

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET – a federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.* 66, 1–16, 1998.

Joseph, J. H. and Cahalan, R. F.: Nearest neighbor spacing of fair weather cumulus, *J. Appl. Meteor.*, 29, 793–805, 1990.

Knyazikhin, Y., Miessen, G., Panfyorov, O., and Gravenhorst, G.: Small-scale study of three-dimensional distribution of photosynthetically active radiation in a forest, *Agr. Forest Meteorol.*, 88, 215–239, 1997.

Koren, I., Oreopoulos, L., Feingold, G., Remer, L. A., and Altartatz, O.: How small is a small cloud?, *Atmos. Chem. Phys.*, 8, 3855–3864, 2008, <http://www.atmos-chem-phys.net/8/3855/2008/>.

Koren, I., Remer, L. A., Kaufman, Y. J., Rudich, Y., and Martins, J. V.: On the twilight zone between clouds and aerosols, *Geophys. Res. Lett.*, 34, L08805, doi:10.1029/2007GL029253, 2007.

Kotchenruther, R. A., Hobbs, P. V., and Hegg, D. A.: Humidification factors for atmospheric aerosols off the mid-Atlantic coast of the United States, *J. Geophys. Res.*, 104(D2), 2239–2251, 1999.

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Interpretation of
radiative signature in
transition zones**J. C. Chiu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Lane, D. E., Goris, K., and Somerville, R. C. J.: Radiative transfer through broken clouds: observations and model validation, *J. Climate*, 15, 2921–2933, 2002.
- Loeb, N. G. and Manalo-Smith, N.: Top-of-atmosphere direct radiative effect of aerosols over global oceans from merged CERES and MODIS observations, *J. Climate*, 18, 3506–3526, 2005.
- 5 Long, C. N. and Ackerman, T. P.: Identification of clear skies from broadband pyranometer measurements and calculation of downwelling shortwave cloud effects, *J. Geophys. Res.*, 105(D12), 15 609–15 626, 2000.
- Long, C. N., Slater, D. W., and Tooman, T.: Total Sky Imager Model 880 status and testing results, ARM Technical Report ARM TR-006, U.S. Department of Energy, Washington, D.C., 2001.
- 10 Long, C. N., Ackerman, T. P., Gaustad, K. L., and Cole, J. N. S.: Estimation of fractional sky cover from broadband shortwave radiometer measurements, *J. Geophys. Res.*, 111, D11204, doi:10.1029/2005JD006475, 2006a.
- 15 Long, C. N., Sabburg, J. M., Calbo, J., and Pages, D.: Retrieving cloud characteristics from ground-based daytime color all-sky images, *J. Atmos. Ocean. Tech.*, 23, 633–652, 2006b.
- Marshak, A., Knyazikhin, Y., Evans, K. D., and Wiscombe, W. J.: The “RED versus NIR” plane to retrieve broken-cloud optical depth from ground-based measurements, Cloud-vegetation interaction: Use of normalized difference cloud index for estimation of cloud optical thickness, *J. Atmos. Sci.*, 61, 1911–1925, 2004.
- 20 Marshak, A., Wen, G., Coakley, J., Remer, L., Loeb, N. G., and Cahalan, R. F.: A simple model for the cloud adjacency effect and the apparent bluing of aerosols near clouds, *J. Geophys. Res.*, 113, D14S17, doi:10.1029/2007JD009196, 2008.
- Martins, J. V., Tanré, D., Remer, L. A., Kaufman, Y. J., Mattoo, S., and Levy, R.: MODIS cloud screening for remote sensing of aerosol over oceans using spatial variability, *Geophys. Res. Lett.*, 29(12), 8009, doi:10.1029/2001GL013252, 2002.
- 25 Miller, M. A., Johnson, K. L., Troyan, D. T., Clothiaux, E. E., Mlawer, E. J., and Mace, G. G.: ARM value-added cloud products: description and status, Proceedings of the 13th ARM Science Team Meeting, 31 March to 4 April 2003, Broomfield, Colorado, available at: http://www.arm.gov/publications/proceedings/conf13/extended_abs/miller-ma.pdf, 2003.
- 30 Paluch, I. R. and Baumgardner, D. G.: Entrainment and fine-scale mixing in a continental convective cloud, *J. Atmos. Sci.*, 46, 261–278, 1989.
- Pilewskie, P., Pommier, J., Bergstrom, R., Gore, W., Howard, S., Rabbette, M., Schmid, B.,

**Interpretation of
radiative signature in
transition zones**

J. C. Chiu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hobbs, P. V., and Tsay, S. C.: Solar spectral radiative forcing during the Southern African Regional Science Initiative, *J. Geophys. Res.*, 108(D13), 8486, doi:10.1029/2002JD002411, 2003.

5 Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X. W., Tsang, T., Strugnell, N. C., Zhang, X. Y., Jin, Y. F., Muller, J. P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., d'Entremont, R. P., Hu, B. X., Liang, S. L., Privette, J. L., and Roy, D.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote Sens. Environ.*, 83, 135–148, 2002.

10 Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z. E., Illingworth, A. J., O'Connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., and Mitrescu, C.: The CloudSat Mission and the A-Train – A new dimension of space-based observations of clouds and precipitation, *B. Am. Meteorol. Soc.*, 83, 1771–1790, 2002.

Stith, J. L.: Observations of cloud-top entrainment in cumuli, *J. Atmos. Sci.*, 49, 1334–1347, 1992.

15 Taylor, T. E., L'Ecuyer, T. S., Slusser, J. R., Stephens, G. L., and Goering, C. D.: An operational retrieval algorithm for determining aerosol optical properties in the ultraviolet, *J. Geophys. Res.*, 113, D03201, doi:10.1029/2007JD008661, 2008.

Thomas, G. E. and Stamnes, K.: Radiative transfer in the atmosphere and ocean, Cambridge University Press, 517 pp., 2002.

20 Turner, D. D., Clough, S. A., Lijegren, J. C., Clothiaux, E. E., Cady-Pereira, K. E., and Gaustad, K. L.: Retrieving liquid water path and precipitable water vapor from the atmospheric radiation measurement (ARM) microwave radiometers, *IEEE T. Geosci. Remote*, 45, 3680–3690, 2007.

25 Walte-Shea, E. A. and Norman, J. M.: Leaf optical properties, in: Photon-vegetation interactions: applications in plant physiology and optical remote sensing, edited by: Myneni, R. B. and Ross, J., Springer Verlag, 229–251, 1991.

Wen, G., Marshak, A., Cahalan, R. F., Remer, L. A., and Kleidman, R. G.: 3-D aerosol-cloud radiative interaction observed in collocated MODIS and ASTER images of cumulus cloud fields, *J. Geophys. Res.*, 112, D13204, doi:10.1029/2006JD008267, 2007.

30 Zhang, J. L., Reid, J. S., and Holben, B. N.: An analysis of potential cloud artifacts in MODIS over ocean aerosol optical thickness products, *Geophys. Res. Lett.*, 32, L15803, doi:10.1029/2005GL023254, 2005.

Table 1. Case summary.

Case ID	1	2	3
Date	2007/05/18 ^a	2006/06/27 ^b	2007/09/05 ^b
Starting time (UTC)	21:35:24	18:52:48	22:46:01
Solar zenith angle	45	15	65
Aerosol optical depth ^c at 870 and 1640 nm	(0.14, 0.09)	(0.07, –)	(0.09, 0.07)
Aerosol effective radius (μm) ^c	0.4	0.25	0.45
Peaks of bimodal size distribution (μm) ^c	(0.15, 1.7)	(0.15, 2.7)	(0.15, 5.0)
Cloud-base height (km) ^d	2.0	3.5	1.2 for Cu
Cloud top height (km) ^d	2.1	3.8	1.7 for Cu
Liquid water path (gm^{-2}) ^e	16	16	–
Wind speed at cloud-base height (ms^{-1}) ^f	3	6	7
Wind direction (deg) ^f	197	343	180
Surface albedo at 870 and 1640 nm ^g	(0.30, 0.24)	(0.32, 0.35)	(0.36, 0.35)

^a Data length is 300 s

^b Data length is 120 s

^c Estimated from NASA's Aerosol Robotic Network (AERONET)

^d Estimated from ARM's Active Remotely Sensed Clouds Locations (ARSCL) product

^e Estimated from an ARM PI product (MWRRET)

^f Estimated from ARM's Merged Sounding product

^g Estimated from NASA's Moderate Resolution Imaging Spectroradiometer (MODIS)

Note that aerosol properties at 1640 nm are not available at the Oklahoma site until May 2007.

Interpretation of radiative signature in transition zones

J. C. Chiu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of radiative signature in transition zones

J. C. Chiu et al.




Fig. 1. Case 1, 18 May 2007 with a SZA of 45° : **(a)** total sky images (from TSI instrument), and **(b)** time-wavelength contour plot of SWS spectra measured from 21:35:24 to 21:40:24 UTC (300 s). SWS-observed zenith radiances have been normalized by the extraterrestrial solar spectrum and by $\cos(\text{SZA})$. In (b), arrows pointed at the time axis correspond to the times of the sky images shown in (a), while arrows pointed at the wavelength axis correspond to 870 and 1640 nm. We also see strong water vapor absorption bands at wavelengths of 930, 1120, 1400, and 1900 nm. **(c)** is time series of radiances at 870 and 1640 nm corresponding to two slices of (b). **(d)** is the plot of normalized radiance difference versus sum at wavelengths of 870 and 1640 nm. Letters S and E indicate the start and end of the time series, while two thick arrows indicate the flow of time evolution.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of radiative signature in transition zones

J. C. Chiu et al.




Fig. 2. Same as Fig. 1, but for Case 2, 27 June 2006 during 18:52:50–18:54:50 UTC (120 s). Solar zenith angle is around 15° . Note that some points of the cloud are not shown in **(d)** to emphasize points in the transition periods.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of radiative signature in transition zones

J. C. Chiu et al.




Fig. 3. Same as Fig. 1, but for Case 3, 5 September 2007 during 22:46:01–22:48:00 UTC (120 s). Solar zenith angle for this case is around 65° .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of
radiative signature in
transition zones

J. C. Chiu et al.




Fig. 4. The modeled track of the sum and difference of intensities (I_{870} and I_{1640}) as the cloud optical depth is varied from zero (point labeled 0) to 15. Model used is DISORT (one-dimensional plane-parallel radiative transfer). Red dots, corresponding to different cloud optical depths, separate the regimes discussed in the text. Regime 0 (a single point) represents a cloud-free condition. Regime 1 is the transition between clear and cloudy (Regimes 2–4). The dashed line is drawn to emphasize the linearity in Regime 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Interpretation of radiative signature in transition zones

J. C. Chiu et al.




Fig. 5. Modeled difference vs. sum of intensities, as in previous figures, for clouds with various aerosol situations below the clouds. **(a)** Cloud-free Regime 0 (indicated by diamonds) is affected by three assumed aerosol loadings: no aerosols, low aerosol optical depth ($\tau_{870}^a=0.05$, $\tau_{1640}^a=0.02$, Ångström exponent $\approx 2/3$), and high aerosol optical depth ($\tau_{870}^a=0.15$, $\tau_{1640}^a=0.08$, Ångström exponent ≈ 1). Assumed aerosol effective radius is $0.25 \mu\text{m}$. Data points correspond to cloud optical depth from 0 to 0.5 in steps of 0.1, one set for each cloud drop effective radius ($4 \mu\text{m}$ and $8 \mu\text{m}$). **(b)** Repeats the high aerosol loading case in (a), then adds a second case with a 16 times larger aerosol particle size ($4 \mu\text{m}$) leading to the lower two curves. Plots are based on 1-D plane-parallel simulations at a SZA of 45° .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interpretation of
radiative signature in
transition zones

J. C. Chiu et al.




Fig. 6. Aerosol scattering phase functions at wavelengths of 870 and 1640 nm for two effective particle sizes (0.25 or 4 μm). The dash-dot line corresponds to a 45° scattering angle.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Interpretation of
radiative signature in
transition zones

J. C. Chiu et al.




Fig. 7. The right-hand side of Eqs. (4a) and (4b) with $\varpi_\lambda=1$ and $P_\lambda=1$ vs. cloud optical depth in the single scattering approximation at solar zenith angles of 15° , 45° and 65° . The straight line is the linear approximation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Interpretation of
radiative signature in
transition zones

J. C. Chiu et al.




Fig. 8. Model calculations of the slope of the $I_{870} - I_{1640}$ vs. $I_{870} + I_{1640}$ relationship as a function of cloud drop effective radius.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Interpretation of
radiative signature in
transition zones

J. C. Chiu et al.




Fig. 9. Cloud phase functions for wavelength 870 nm (left column) and 1640 nm (right column). Top row shows phase functions versus scattering angle for effective cloud drop radii 2, 4, 8 μm . Bottom row shows phase functions at specific angles versus effective drop radius; the angles chosen are the same as the solar zenith angles in Cases 1–3 for the purpose of understanding single-scattering behavior.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)