Technical notes: Kinetic data for MISTRA

M. Piot and R. von Glasow

The Chemistry Influencing ODEs in the Polar Boundary Layer in Spring: A Model Study

Atmos. Chem. Phys. Discuss., 2008

1 Tables of reaction rates

This collection comprises a complete listing of all gas and aqueous phase species (Table 1), gas phase (Table 2) and aqueous phase (Table 3) reaction rates, as well as rates for the heterogeneous (particle surface) reactions (Table 4), aqueous phase equilibrium constants (Table 5), Henry constants and accommodations coefficients (Table 6).

Table 1: Species in MISTRA

Gas phase
O^1D, O_2, O_3, OH, HO_2, H_2O_2, H_2O
NO, NO_2, NO_3, N_2O_5, HONO, HNO_3, HNO_4, PAN, NH_3, $RONO_2$
CO, CO_2, CH_4, C_2H_6, C_2H_4, HCHO, HCOOH, ALD (i.e., CH_3CHO), CH_2O_2, $HOCH_2O_2$, CH_3CO_3, CH_3O_2, $C_2H_5O_2$, H_3CO_2, EO_2 (i.e., $H_2C(OH)CH_2OO$), CH_2O_2, ROOH (i.e., alkylhydroperoxides), DOM
SO_2, SO_3, $HOSO_2$, H_2SO_4, DMS, CH_3SCH_2OO, DMSO, $DMSO_2$, CH_3S, CH_3SO, CH_3SO_2, CH_3SO_3, CH_3SO_2H, CH_3SO_3H
Cl, ClO, OClO, HCl, HOCl, Cl_2, Cl_2O_2 $ClNO_2$, $ClNO_3$
Br, BrO, HBr, HOBr, Br_2, $BrNO_2$, $BrNO_3$, BrCl, $CHBr_3$, CH_3Br
Liquid phase (neutral)
O_2, O_3, OH, HO_2, H_2O_2, H_2O
NO, NO_2, NO_3, HONO, HNO_3, HNO_4, NH_3
CO_2, HCHO, HCOOH, CH_3OH, CH_3OO, CH_3OOH
SO_2, H_2SO_4, DMSO, $DMSO_2$, CH_3SO_2H, CH_3SO_3H
Cl, HCl, HOCl, Cl_2
Br, HBr, HOBr, Br_2, BrCl
Liquid phase (ions)
H^+, OH^-, O_2^-
NO_2^-, NO_3^-, NO_4^-, NH_4^+
HCO_3^-, CO_3^-, $HCOO^-$
HSO_3^-, $SO_3{}^2-$, HSO_4^-, SO_4^{2-}, HSO_5^-, SO_3^-, SO_4^-, SO_5^-, $CH_3SO_3^-$, $CH_2OHSO_2^-$, $CH_2OHSO_3^-$
Cl^-, Cl_2^-, ClO^-, $ClOH^-$
Br^-, Br_2^-, BrO^-, $BrCl_2^-$, Br_2Cl^-, $BrOH^-$

Table 2: Gas phase reactions.

no	reaction	n	A $[(cm^{-3})^{1-n}s^{-1}]$	$-E_a$ / R [K]	reference
O 1	$O^1D + O_2 \longrightarrow O_3$	2	3.2×10^{-11}	70	Atkinson et al. (2006)
O 2	$O^1D + N_2 \longrightarrow O_3$	2	1.8×10^{-11}	110	Atkinson et al. (2006)
O 3	$O^1D + H_2O \longrightarrow 2\ OH$	2	2.2×10^{-10}		Atkinson et al. (2006)
O 4	$OH + O_3 \longrightarrow HO_2 + O_2$	2	1.7×10^{-12}	-940	Atkinson et al. (2006)
O 5	$OH + HO_2 \longrightarrow H_2O + O_2$	2	4.8×10^{-11}	250	Atkinson et al. (2006)
O 6	$OH + H_2O_2 \longrightarrow HO_2 + H_2O$	2	2.9×10^{-12}	-160	Atkinson et al. (2006)
O 7	$HO_2 + O_3 \longrightarrow OH + 2O_2$	2	1.0×10^{-14}	-490	Atkinson et al. (2004)
O 8	$HO_2 + HO_2 \longrightarrow H_2O_2 + O_2$	2	2.3×10^{-13}	600	Atkinson et al. (2006)
O 9	$O_3 + h\nu \longrightarrow O_2 + O^1D$	1	[1]		DeMore et al. (1997)
O 10	$H_2O_2 + h\nu \longrightarrow 2OH$	1	[1]		DeMore et al. (1997)
N 1	$NO + OH \xrightarrow{M} HONO$	3	[2]		Sander et al. (2003)
N 2	$NO + HO_2 \longrightarrow NO_2 + OH$	2	3.5×10^{-12}	250	Atkinson et al. (2004)
N 3	$NO + O_3 \longrightarrow NO_2 + O_2$	2	3.0×10^{-12}	-1500	Sander et al. (2003)
N 4	$NO + NO_3 \longrightarrow 2NO_2$	2	1.5×10^{-11}	170	Sander et al. (2003)
N 5	$NO_2 + OH \xrightarrow{M} HNO_3$	3	[2]		Sander et al. (2003)
N 6	$NO_2 + HO_2 \xrightarrow{M} HNO_4$	3	[2]		Atkinson et al. (2006)
N 7	$NO_2 + O_3 \longrightarrow NO_3 + O_2$	2	1.2×10^{-13}	-2450	Sander et al. (2003)
N 8	$NO_2 + h\nu \longrightarrow NO + O_3$	1	[1]		DeMore et al. (1997)
N 9	$NO_2 + NO_3 \xrightarrow{M} N_2O_5$	3	[2]		Sander et al. (2003)
N 10	$NO_3 + h\nu \longrightarrow NO + O_2$	1	[1]		Wayne et al. (1991)
N 11	$NO_3 + HO_2 \longrightarrow 0.3\ HNO_3 + 0.7\ OH + 0.7\ NO_2 + O_2$	2	4.0×10^{-12}		Atkinson et al. (2006)
N 12	$NO_3 + NO_3 \longrightarrow NO_2 + NO_2 + O_2$	2	8.5×10^{-13}	-2450	Sander et al. (2003)
N 13	$NO_3 + h\nu \longrightarrow NO_2 + O_3$	1	[1]		Wayne et al. (1991)
N 14	$N_2O_5 \xrightarrow{M} NO_2 + NO_3$	2	[2]		Sander et al. (2003)
N 15	$N_2O_5 + H_2O \longrightarrow 2HNO_3$	2	2.6×10^{-22}		Atkinson et al. (2006)
N 16	$N_2O_5 + h\nu \longrightarrow NO_2 + NO_3$	1	[1]		DeMore et al. (1997)
N 17	$HONO + OH \longrightarrow NO_2$	2	1.8×10^{-11}	-390	Sander et al. (2003)
N 18	$HONO + h\nu \longrightarrow NO + OH$	1	[1]		DeMore et al. (1997)
N 19	$HNO_3 + h\nu \longrightarrow NO_2 + OH$	1	[1]		DeMore et al. (1997)
N 20	$HNO_3 + OH \longrightarrow NO_3 + H_2O$	2	[2]		Atkinson et al. (2006)
N 21	$HNO_4 \xrightarrow{M} NO_2 + HO_2$	2	[2]		Sander et al. (2003)
N 22	$HNO_4 + OH \longrightarrow NO_2 + H_2O + O_2$	2	1.3×10^{-12}	380	Haggerstone et al. (2005)
N 23	$HNO_4 + h\nu \longrightarrow NO_2 + HO_2$	1	[1]		DeMore et al. (1997)
N 24	$HNO_4 + h\nu \longrightarrow OH + NO_3$	1	[1]		DeMore et al. (1997)
N 25	$RONO_2 + OH \longrightarrow R + H_2O + NO_2$	2	1.3×10^{-12}		Sander et al. (1997)
N 26	$RONO_2 + h\nu \longrightarrow RO + NO_2$	1	assumed similar as N 19		

Table 2: Continued.

no	reaction	n	A [(cm^{-3})$^{1-n}$s^{-1}]	$-E_a$ / R [K]	reference
C 1	$CO + OH \xrightarrow{O_2} HO_2 + CO_2$	2	[2]		Sander et al. (2003)
C 2	$CH_4 + OH \xrightarrow{O_2} CH_3OO + H_2O$	2	2.4×10^{-12}	-1775	Sander et al. (2003)
C 3	$C_2H_6 + OH \longrightarrow C_2H_5O_2 + H_2O$	2	1.7×10^{-11}	-1232	Lurmann et al. (1986)
C 4	$C_2H_4 + OH \longrightarrow EO_2$	2	1.66×10^{-12}	474	Sander et al. (1997), see note
C 5	$C_2H_4 + O_3 \longrightarrow HCHO + 0.4CH_2O_2 + 0.12HO_2 + 0.42CO + 0.06CH_4$	2	1.2×10^{-14}	-2633	Lurmann et al. (1986), see note
C 6	$HO_2 + CH_3OO \longrightarrow ROOH + O_2$	2	4.1×10^{-13}	750	Sander et al. (2003)
C 7	$HO_2 + C_2H_5O_2 \longrightarrow ROOH + O_2$	2	7.5×10^{-13}	700	Sander et al. (2003)
C 8	$HO_2 + CH_3CO_3 \longrightarrow ROOH + O_2$	2	4.5×10^{-13}	1000	DeMore et al. (1997)
C 9	$CH_3OO + CH_3OO \longrightarrow 1.4HCHO + 0.8HO_2 + O_2$	2	1.5×10^{-13}	220	Lurmann et al. (1986)
C 10	$C_2H_5O_2 + NO \longrightarrow ALD + HO_2 + NO_2$	2	4.2×10^{-12}	180	Lurmann et al. (1986)
C 11	$2C_2H_5O_2 \longrightarrow 1.6ALD + 1.2HO_2$	2	5.00×10^{-14}		Lurmann et al. (1986)
C 12	$EO_2 + NO \longrightarrow NO_2 + 2.0HCHO + HO_2$	2	4.2×10^{-12}	180	Lurmann et al. (1986)
C 13	$EO_2 + EO_2 \longrightarrow 2.4HCHO + 1.2HO_2 + 0.4ALD$	2	5.00×10^{-14}		Lurmann et al. (1986)
C 14	$HO_2 + EO_2 \longrightarrow ROOH + O_2$	2	3.00×10^{-12}		Lurmann et al. (1986)
C 15	$HCHO + h\nu \longrightarrow 2HO_2 + CO$	1	[1]		DeMore et al. (1997)
C 16	$HCHO + h\nu \longrightarrow CO + H_2$	1	[1]		DeMore et al. (1997)
C 17	$HCHO + OH \xrightarrow{O_2} HO_2 + CO + H_2O$	2	1.00×10^{-11}		DeMore et al. (1997)
C 18	$HCHO + HO_2 \longrightarrow HOCH_2O_2$	2	6.7×10^{-15}	600	Sander et al. (2003)
C 19	$HCHO + NO_3 \xrightarrow{O_2} HNO_3 + HO_2 + CO$	2	5.8×10^{-16}		DeMore et al. (1997)
C 20	$ALD + OH \longrightarrow CH_3CO_3 + H_2O$	2	6.9×10^{-12}	250	Lurmann et al. (1986)
C 21	$ALD + NO_3 \longrightarrow HNO_3 + CH_3CO_3$	2	1.40×10^{-15}		DeMore et al. (1997)
C 22	$ALD + h\nu \longrightarrow CH_3OO + HO_2 + CO$	1	[1]		Lurmann et al. (1986)
C 23	$ALD + h\nu \longrightarrow CH_4 + CO$	1	[1]		Lurmann et al. (1986)
C 24	$HOCH_2O_2 + NO \longrightarrow HCOOH + HO_2 + NO_2$	2	4.2×10^{-12}	180	Lurmann et al. (1986)
C 25	$HOCH_2O_2 + HO_2 \longrightarrow HCOOH + H_2O + O_2$	2	2.00×10^{-12}		Lurmann et al. (1986)
C 26	$2\ HOCH_2O_2 \longrightarrow 2HCOOH + 2HO_2 + 2O_2$	2	1.00×10^{-13}		Lurmann et al. (1986)
C 27	$HCOOH + OH \xrightarrow{O_2} HO_2 + H_2O + CO_2$	2	4.0×10^{-13}		DeMore et al. (1997)
C 28	$CH_3CO_3 + NO_2 \longrightarrow PAN$	2	4.70×10^{-12}		Lurmann et al. (1986)
C 29	$PAN \longrightarrow CH_3CO_3 + NO_2$	1	1.9×10^{16}	-13543	DeMore et al. (1997)
C 30	$CH_3CO_3 + NO \longrightarrow CH_3OO + NO_2 + CO_2$	2	4.2×10^{-12}	180	Lurmann et al. (1986)
C 31	$CH_3OO + NO \xrightarrow{O_2} HCHO + NO_2 + HO_2$	2	3.0×10^{-12}	280	DeMore et al. (1997)
C 32	$ROOH + OH \longrightarrow 0.7\ CH_3OO + 0.3\ HCHO + 0.3\ OH$	2	3.8×10^{-12}	200	DeMore et al. (1997), see note
C 33	$ROOH + h\nu \longrightarrow HCHO + OH + HO_2$	1	[1]		DeMore et al. (1997), see note

Table 2: Continued.

no	reaction	n	A [$(cm^{-3})^{1-n}s^{-1}$]	$-E_a$ / R [K]	reference
S 1	$SO_2 + OH \xrightarrow{M} HOSO_2$	3	^{2}		Atkinson et al. (2006)
S 2	$HOSO_2 + O_2 \longrightarrow HO_2 + SO_3$	2	1.3×10^{-12}	330	Atkinson et al. (2006)
S 3	$SO_3 \xrightarrow{H_2O} H_2SO_4$	1	^{2}		Jayne et al. (1997)
S 4	$CH_3SCH_3 + OH \longrightarrow CH_3SCH_2OO + H_2O$	2	^{2}		Atkinson et al. (1997)
S 5	$CH_3SCH_3 + OH \xrightarrow{O_2} CH_3SOCH_3 + HO_2$	2	^{2}		Atkinson et al. (1997)
S 6	$CH_3SCH_3 + NO_3 \xrightarrow{O_2} CH_3SCH_2OO + HNO_3$	2	1.9×10^{-13}	520	Atkinson et al. (1999)
S 7	$CH_3SCH_3 + Cl \xrightarrow{O_2} CH_3SCH_2OO + HCl$	2	3.3×10^{-10}		Atkinson et al. (1999)
S 8	$CH_3SCH_3 + Br \xrightarrow{O_2} CH_3SCH_2OO + HBr$	2	9.0×10^{-11}	-2386	Jefferson et al. (1994)
S 9	$CH_3SCH_3 + BrO \longrightarrow CH_3SOCH_3 + Br$	2	2.54×10^{-14}	850	Ingham et al. (1999)
S 10	$CH_3SCH_3 + ClO \longrightarrow CH_3SOCH_3 + Cl$	2	9.5×10^{-15}		Barnes et al. (1991)
S 11	$CH_3SCH_3 + IO \longrightarrow CH_3SOCH_3 + I$	2	1.4×10^{-14}		THALOZ (2005)
S 12	$CH_3SCH_2OO + NO \longrightarrow HCHO + CH_3S + NO_2$	2	4.9×10^{-12}	263	Urbanski et al. (1997)
S 13	$CH_3SCH_2OO + CH_3SCH_2OO \xrightarrow{O_2} 2\ HCHO + 2\ CH_3S$	2	1.0×10^{-11}		Urbanski et al. (1997); Atkinson et al. (2006)
S 14	$CH_3S + O_3 \longrightarrow CH_3SO + O_2$	2	1.15×10^{-12}	432	Atkinson et al. (2006)
S 15	$CH_3S + NO_2 \longrightarrow CH_3SO + NO$	2	3.0×10^{-11}	210	Atkinson et al. (2006)
S 16	$CH_3SO + NO_2 \xrightarrow{O_2} 0.82\ CH_3SO_2 + 0.18\ SO_2 + 0.18\ H_3CO_2 + NO$	2	1.2×10^{-11}		Atkinson et al. (2006); Kukui et al. (2000), product ratios from van Dingenen et al. (1994)
S 17	$CH_3SO + O_3 \xrightarrow{O_2} CH_3SO_2$	2	6.0×10^{-13}		Atkinson et al. (2006)
S 18	$CH_3SO_2 \longrightarrow SO_2 + CH_3OO$	1	1.9×10^{13}	-8661	Barone et al. (1995)
S 19	$CH_3SO_2 + NO_2 \longrightarrow CH_3SO_3 + NO$	2	2.2×10^{-12}		Ray et al. (1996)
S 20	$CH_3SO_2 + O_3 \longrightarrow CH_3SO_3$	2	$3. \times 10^{-13}$		Barone et al. (1995)
S 21	$CH_3SO_3 + HO_2 \longrightarrow CH_3SO_3H$	2	$5. \times 10^{-11}$		Barone et al. (1995)
S 22	$CH_3SO_3 \xrightarrow{H_2O,O_2} CH_3OO + H_2SO_4$	1	1.36×10^{14}	-11071	Barone et al. (1995)
S 23	$CH_3SOCH_3 + OH \longrightarrow 0.95\ CH_3SO_2H + 0.95\ CH_3OO + 0.05\ DMSO_2$	2	8.7×10^{-11}		Urbanski et al. (1998)
S 24	$CH_3SO_2H + OH \longrightarrow 0.95\ CH_3SO_2 + 0.05\ CH_3SO_3H + 0.05\ HO_2 + H_2O$	2	$9.\times 10^{-11}$		Kukui et al. (2003)
S 25	$CH_3SO_2H + NO_3 \longrightarrow CH_3SO_2 + HNO_3$	2	1.0×10^{-13}		Yin et al. (1990)

Table 2: Continued.

no	reaction	n	A [$(cm^{-3})^{1-n}s^{-1}$]	$-E_a$ / R [K]	reference
Cl 1	$Cl + O_3 \longrightarrow ClO + O_2$	2	2.8×10^{-11}	-250	Atkinson et al. (2006)
Cl 2	$Cl + HO_2 \longrightarrow HCl + O_2$	2	1.8×10^{-11}	170	Sander et al. (2003)
Cl 3	$Cl + HO_2 \longrightarrow ClO + OH$	2	4.1×10^{-11}	-450	Sander et al. (2003)
Cl 4	$Cl + H_2O_2 \longrightarrow HCl + HO_2$	2	1.1×10^{-11}	-980	Atkinson et al. (2006)
Cl 5	$Cl + CH_3OO \longrightarrow 0.5\ ClO + 0.5\ HCHO + 0.5\ HO_2 + 0.5\ HCl + 0.5\ CO + 0.5\ H_2O$	2	1.6×10^{-10}		Sander et al. (2003)
Cl 6	$Cl + NO_3 \longrightarrow ClO + NO_2$	2	2.4×10^{-11}		Mellouki et al. (1987)
Cl 7	$Cl + CH_4 \xrightarrow{O_2} HCl + CH_3OO$	2	9.6×10^{-12}	-1360	Sander et al. (2003)
Cl 8	$Cl + C_2H_6 \xrightarrow{O_2} HCl + C_2H_5O_2$	2	7.7×10^{-11}	-90	Sander et al. (2003)
Cl 9	$Cl + C_2H_4 \xrightarrow{O_2} HCl + C_2H_5O_2$	2	$1. \times 10^{-10}$		Sander et al. (1997), see note
Cl 10	$Cl + HCHO \xrightarrow{O_2} HCl + HO_2 + CO$	2	8.1×10^{-11}	-30	Sander et al. (2003)
Cl 11	$Cl + ROOH \longrightarrow CH_3OO + HCl$	2	5.7×10^{-11}		Wallington et al. (1990), see note
Cl 12	$Cl + OClO \longrightarrow ClO + ClO$	2	3.2×10^{-11}	170	Atkinson et al. (2006)
Cl 13	$Cl + ClNO_3 \longrightarrow Cl_2 + NO_3$	2	6.5×10^{-12}	135	Sander et al. (2003)
Cl 14	$Cl + PAN \longrightarrow HCl + HCHO + NO_3$	2	1.0×10^{-14}		Tsalkani et al. (1988)
Cl 15	$Cl + HNO_3 \longrightarrow HCl + NO_2$	2	1.0×10^{-16}		Wine et al. (1988)
Cl 16	$Cl + RONO_2 \longrightarrow HCl + NO_2$	2	7.7×10^{-11}		estimated from Muthuramu et al. (1994)
Cl 17	$ClO + OH \longrightarrow Cl + HO_2$	2	7.4×10^{-12}	-270	Sander et al. (2003)
Cl 18	$ClO + OH \longrightarrow HCl + O_2$	2	6.0×10^{-13}	-230	Sander et al. (2003)
Cl 19	$ClO + HO_2 \longrightarrow HOCl + O_2$	2	2.2×10^{-12}	340	Atkinson et al. (2006)
Cl 20	$ClO + CH_3OO \longrightarrow Cl + HCHO + HO_2$	2	3.3×10^{-12}	-115	Sander et al. (2003)
Cl 21	$ClO + NO \longrightarrow Cl + NO_2$	2	6.2×10^{-12}	295	Atkinson et al. (2006)
Cl 22	$ClO + NO_2 \xrightarrow{M} ClNO_3$	3	[2]		Atkinson et al. (2006)
Cl 23	$ClO + ClO \longrightarrow Cl_2O_2$	2	[2]		Atkinson et al. (2006)
Cl 24	$ClO + ClO \longrightarrow Cl_2 + O_2$	2	1.0×10^{-12}	-1590	Atkinson et al. (2006)
Cl 25	$ClO + ClO \longrightarrow Cl_2O_2$	2	3.0×10^{-11}	-2450	Atkinson et al. (2006)
Cl 26	$ClO + ClO \longrightarrow Cl + OClO$	2	3.5×10^{-13}	-1370	Atkinson et al. (2006)
Cl 27	$OClO + OH \longrightarrow HOCl + O_2$	2	4.5×10^{-13}	800	Atkinson et al. (2006)
Cl 28	$OClO + NO \longrightarrow ClO + NO_2$	2	1.1×10^{-13}	350	Atkinson et al. (2006)
Cl 29	$Cl_2O_2 \longrightarrow ClO + ClO$	1	[2]		Atkinson et al. (2006)
Cl 30	$HOCL + OH \longrightarrow ClO + H_2O$	2	3.0×10^{-12}	-500	Sander et al. (2003)
Cl 31	$HCl + OH \longrightarrow H_2O + Cl$	2	1.8×10^{-12}	-240	Atkinson et al. (2006)
Cl 32	$ClNO_2 + OH \longrightarrow HOCl + NO_2$	2	2.4×10^{-12}	-1250	Atkinson et al. (2006)

Table 2: Continued.

no	reaction	n	A [(cm^{-3})$^{1-n}$s^{-1}]	$-E_a$ / R [K]	reference
Cl 33	$ClNO_3 + OH \longrightarrow 0.5\ ClO + 0.5\ HNO_3 + 0.5\ HOCl + 0.5\ NO_3$	2	1.2×10^{-12}	-330	Atkinson et al. (2006)
Cl 34	$ClNO_3 \longrightarrow ClO + NO_2$	1	[2]		Anderson and Fahey (1990)
Cl 35	$OClO + h\nu \xrightarrow{O_2,O_3} O_3 + ClO$	1	[1]		DeMore et al. (1997)
Cl 36	$Cl_2O_2 + h\nu \longrightarrow Cl + Cl + O_2$	1	[1]		DeMore et al. (1997)
Cl 37	$Cl_2 + h\nu \longrightarrow 2\ Cl$	1	[1]		DeMore et al. (1997)
Cl 38	$HOCl + h\nu \longrightarrow Cl + OH$	1	[1]		DeMore et al. (1997)
Cl 39	$ClNO_2 + h\nu \longrightarrow Cl + NO_2$	1	[1]		DeMore et al. (1997)
Cl 40	$ClNO_3 + h\nu \longrightarrow Cl + NO_3$	1	[1]		DeMore et al. (1997)
Br 1	$Br + O_3 \longrightarrow BrO + O_2$	2	1.7×10^{-11}	-800	Atkinson et al. (2006)
Br 2	$Br + HO_2 \longrightarrow HBr + O_2$	2	7.7×10^{-12}	-450	Atkinson et al. (2006)
Br 3	$Br + C_2H_4 \xrightarrow{O_2} HBr + C_2H_5O_2$	2	$5. \times 10^{-14}$		Sander et al. (1997), see note
Br 4	$Br + HCHO \xrightarrow{O_2} HBr + CO + HO_2$	2	1.7×10^{-11}	-800	Sander et al. (2003)
Br 5	$Br + ROOH \longrightarrow CH_3OO + HBr$	2	2.66×10^{-12}	-1610	Mallard et al. (1993), see note
Br 6	$Br + NO_2 \longrightarrow BrNO_2$	2	[2]		Sander et al. (2003)
Br 7	$Br + BrNO_3 \longrightarrow Br_2 + NO_3$	2	4.9×10^{-11}		Orlando and Tyndall (1996)
Br 8	$BrO + OH \longrightarrow Br + HO_2$	2	1.8×10^{-11}	250	Atkinson et al. (2006)
Br 9	$BrO + HO_2 \longrightarrow HOBr + O_2$	2	4.5×10^{-12}	500	Atkinson et al. (2006)
Br 10	$BrO + CH_3OO \longrightarrow HOBr + HCHO$	2	4.1×10^{-12}		Aranda et al. (1997)
Br 11	$BrO + CH_3OO \longrightarrow Br + HCHO + HO_2$	2	1.6×10^{-12}		Aranda et al. (1997)
Br 12	$BrO + HCHO \xrightarrow{O_2} HOBr + CO + HO_2$	2	1.5×10^{-14}		Hansen et al. (1999)
Br 13	$BrO + NO \longrightarrow Br + NO_2$	2	8.7×10^{-12}	260	Atkinson et al. (2006)
Br 14	$BrO + NO_2 \xrightarrow{M} BrNO_3$	3	[2]		Atkinson et al. (2006)
Br 15	$BrO + BrO \longrightarrow 2\ Br + O_2$	2	2.4×10^{-12}	40	Sander et al. (2003)
Br 16	$BrO + BrO \longrightarrow Br_2 + O_2$	2	2.9×10^{-14}	860	Sander et al. (2003)
Br 17	$HBr + OH \longrightarrow Br + H_2O$	2	5.5×10^{-12}	205	Atkinson et al. (2006)
Br 18	$BrNO_3 \longrightarrow BrO + NO_2$	1	[2]		Orlando and Tyndall (1996)
Br 19	$BrO + h\nu \xrightarrow{O_2} Br + O_3$	1	[1]		DeMore et al. (1997)
Br 20	$Br_2 + h\nu \longrightarrow 2\ Br$	1	[1]		Hubinger and Nee (1995)
Br 21	$HOBr + h\nu \longrightarrow Br + OH$	1	[1]		Ingham et al. (1999)
Br 22	$BrNO_2 + h\nu \longrightarrow Br + NO_2$	1	[1]		Scheffler et al. (1997)
Br 23	$BrNO_3 + h\nu \longrightarrow Br + NO_3$	1	[1]		DeMore et al. (1997)
Br 24	$Br_2 + OH \longrightarrow HOBr + Br$	2	2.0×10^{-11}	240	Atkinson et al. (2006)
Br 25	$CH_3Br + OH \longrightarrow H_2O + Br$	2	1.7×10^{-12}	-1215	Atkinson et al. (2006)
Br 26	$CHBr_3 + OH \longrightarrow H_2O + Br$	2	1.35×10^{-12}	-600	Atkinson et al. (2004)

Table 2: Continued.

no	reaction	n	A [$(cm^{-3})^{1-n}s^{-1}$]	$-E_a$ / R [K]	reference
Hx 1	$Cl + BrCl \longrightarrow Br + Cl_2$	2	1.5×10^{-11}		Mallard et al. (1993)
Hx 2	$Cl + Br_2 \longrightarrow BrCl + Br$	2	1.2×10^{-10}		Mallard et al. (1993)
Hx 3	$Br + OClO \longrightarrow BrO + ClO$	2	2.6×10^{-11}	-1300	Atkinson et al. (2006)
Hx 4	$Br + Cl_2 \longrightarrow BrCl + Cl$	2	1.1×10^{-15}		Mallard et al. (1993)
Hx 5	$Br + BrCl \longrightarrow Br_2 + Cl$	2	3.3×10^{-15}		Mallard et al. (1993)
Hx 6	$BrO + ClO \longrightarrow Br + OClO$	2	1.6×10^{-12}	430	Atkinson et al. (2006)
Hx 7	$BrO + ClO \longrightarrow Br + Cl + O_2$	2	2.9×10^{-12}	220	Atkinson et al. (2006)
Hx 8	$BrO + ClO \longrightarrow BrCl + O_2$	2	5.8×10^{-13}	170	Atkinson et al. (2006)
Hx 9	$BrCl + h\nu \longrightarrow Br + Cl$	1	[1]		DeMore et al. (1997)

n is the order of the reaction. [1] photolysis rates calculated online, [2] special rate functions (pressure dependent and/or humidity dependent). Notes: The rates for ROOH were assumed as that of CH_3OOH; C_2H_4 is used as generic alkene as in the Lurmann et al. (1986) mechanism. The rate coefficients are calculated with $k = A \times \exp(\frac{-E_a}{RT})$.

Table 3: Aqueous phase reactions.

no	reaction	n	k_0 [$(M^{1-n})s^{-1}$]	$-E_a$ / R [K]	reference
O 1	$O_3 + OH \longrightarrow HO_2$	2	1.1 x 10^8		Sehested et al. (1984)
O 2	$O_3 + O_2^- \longrightarrow OH + OH^-$	2	1.5 x 10^9		Sehested et al. (1983)
O 3	$OH + OH \longrightarrow H_2O_2$	2	5.5 x 10^9		Buxton et al. (1988)
O 4	$OH + HO_2 \longrightarrow H_2O$	2	7.1 x 10^9		Sehested et al. (1968)
O 5	$OH + O_2^- \longrightarrow OH^-$	2	1.0 x 10^{10}		Sehested et al. (1968)
O 6	$OH + H_2O_2 \longrightarrow HO_2$	2	2.7 x 10^7	-1684	Christensen et al. (1982)
O 7	$HO_2 + HO_2 \longrightarrow H_2O_2$	2	9.7 x 10^5	-2500	Christensen and Sehested (1988)
O 8	$HO_2 + O_2^- \xrightarrow{H^+} H_2O_2$	2	1.0 x 10^8	-900	Christensen and Sehested (1988)
N 1	$HONO + OH \longrightarrow NO_2$	2	1.0 x 10^{10}		assumed =N7 Barker et al. (1970)
N 2	$HONO + H_2O_2 \xrightarrow{H^+} HNO_3$	3	4.6 x 10^3	-6800	Damschen and Martin (1983)
N 3	$NO_3 + OH^- \longrightarrow NO_3^- + OH$	2	8.2 x 10^7	-2700	Exner et al. (1992)
N 4	$NO_2 + NO_2 \longrightarrow HNO_3 + HONO$	2	1.0 x 10^8		Lee and Schwartz (1981)
N 5	$NO_2 + HO_2 \longrightarrow HNO_4$	2	1.8 x 10^9		Warneck (1999)
N 6	$NO_2^- + O_3 \longrightarrow NO_3^- + O_2$	2	5.0 x 10^5	-6950	Damschen and Martin (1983)
N 7	$NO_2^- + OH \longrightarrow NO_2 + OH^-$	2	1.0 x 10^{10}		Barker et al. (1970)
N 8	$NO_4^- \longrightarrow NO_2^- + O_2$	1	8.0 x 10^{-1}		Warneck (1999)
C 1	$HCHO + OH \longrightarrow HCOOH + HO_2$	2	7.7 x 10^8	-1020	Chin and Wine (1994)
C 2	$HCOOH + OH \longrightarrow HO_2 + CO_2$	2	1.1 x 10^8	-991	Chin and Wine (1994)
C 3	$HCOO^- + OH \longrightarrow OH^- + HO_2 + CO_2$	2	3.1 x 10^9	-1240	Chin and Wine (1994)
C 4	$CH_3OO + HO_2 \longrightarrow CH_3OOH$	2	4.3 x 10^5		estimated by Jacob (1986)
C 5	$CH_3OO + O_2^- \longrightarrow CH_3OOH + OH^-$	2	5.0 x 10^7		estimated by Jacob (1986)
C 6	$CH_3OH + OH \longrightarrow HCHO + HO_2$	2	9.7 x 10^8		Buxton et al. (1988)
C 7	$CH_3OOH + OH \longrightarrow CH_3OO$	2	2.7 x 10^7	-1715	estimated by Jacob (1986)
C 8	$CH_3OOH + OH \longrightarrow HCHO + OH$	2	1.1 x 10^7	-1715	estimated by Jacob (1986)
C 9	$CO_3^- + O_2^- \longrightarrow HCO_3^- + OH^-$	2	6.5 x 10^8		Ross et al. (1992)
C 10	$CO_3^- + H_2O_2 \longrightarrow HCO_3^- + HO_2$	2	4.3 x 10^5		Ross et al. (1992)
C 11	$CO_3^- + HCOO^- \longrightarrow HCO_3^- + HCO_3^- + HO_2$	2	1.5 x 10^5		Ross et al. (1992)
C 12	$HCO_3^- + OH \longrightarrow CO_3^-$	2	8.5 x 10^6		Ross et al. (1992)
C 13	$DOM + OH \longrightarrow HO_2$	2	5.0 x 10^9		estimated by (C. Anastasio, pers. comm.) from Ross et al. (1998)

Table 3: Continued.

no	reaction	n	k_0 [$(M^{1-n})s^{-1}$]	$-E_a$ / R [K]	reference
S 1	$SO_3^- + O_2 \longrightarrow SO_5^-$	2	1.5 x 10^9		Huie and Neta (1987)
S 2	$HSO_3^- + O_3 \longrightarrow SO_4^{2-} + H^+ + O_2$	2	3.7 x 10^5	-5500	Hoffmann (1986)
S 3	$SO_3^{2-} + O_3 \longrightarrow SO_4^{2-} + O_2$	2	1.5 x 10^9	-5300	Hoffmann (1986)
S 4	$HSO_3^- + OH \longrightarrow SO_3^-$	2	4.5 x 10^9		Buxton et al. (1988)
S 5	$SO_3^{2-} + OH \longrightarrow SO_3^- + OH^-$	2	5.5 x 10^9		Buxton et al. (1988)
S 6	$HSO_3^- + HO_2 \longrightarrow SO_4^{2-} + OH + H^+$	2	3.0 x 10^3		upper limit D. Sedlak pers. comm. with R. Sander
S 7	$HSO_3^- + O_2^- \longrightarrow SO_4^{2-} + OH$	2	3.0 x 10^3		upper limit D. Sedlak pers. comm. with R. Sander
S 8	$HSO_3^- + H_2O_2 \longrightarrow SO_4^{2-} + H^+$	2	$5.2 \times 10^6 \times \frac{[H^+]}{[H^+]+0.1M}$	-3650	Damschen and Martin (1983)
S 9	$HSO_3^- + NO_2 \xrightarrow{NO_2} HSO_4^- + HONO + HONO$	2	2.0 x 10^7		Clifton et al. (1988)
S 10	$SO_3^{2-} + NO_2 \xrightarrow{NO_2} SO_4^{2-} + HONO + HONO$	2	2.0 x 10^7		Clifton et al. (1988)
S 11	$HSO_3^- + NO_3 \longrightarrow SO_3^- + NO_3^- + H^+$	2	1.4 x 10^9	-2000	Exner et al. (1992)
S 12	$HSO_3^- + HNO_4 \longrightarrow HSO_4^- + NO_3^- + H^+$	2	3.1 x 10^5		Warneck (1999)
S 13	$HSO_3^- + CH_3OOH \xrightarrow{H^+} SO_4^{2-} + H^+ + CH_3OH$	3	1.6 x 10^7	-3800	Lind et al. (1987)
S 14	$SO_3^{2-} + CH_3OOH \xrightarrow{H^+} SO_4^{2-} + CH_3OH$	3	1.6 x 10^7	-3800	Lind et al. (1987)
S 15	$HSO_3^- + HCHO \longrightarrow CH_2OHSO_3^-$	2	4.3 x 10^{-1}		Boyce and Hoffmann (1984)
S 16	$SO_3^{2-} + HCHO \xrightarrow{H^+} CH_2OHSO_3^-$	2	1.4 x 10^4		Boyce and Hoffmann (1984)
S 17	$CH_2OHSO_3^- + OH^- \longrightarrow SO_3^{2-} + HCHO$	2	3.6 x 10^3		Seinfeld and Pandis (1998)
S 18	$HSO_3^- + HSO_5^- \xrightarrow{H^+} SO_4^{2-} + SO_4^{2-} + H^+ + H^+$	3	7.1 x 10^6		Betterton and Hoffmann (1988)
S 19	$SO_4^- + OH \longrightarrow HSO_5^-$	2	1.0 x 10^9		Jiang et al. (1992)

Table 3: Continued.

no	reaction	n	k_0 [$(M^{1-n})s^{-1}$]	$-E_a$ / R [K]	reference
S 20	$SO_4^- + HO_2 \longrightarrow SO_4^{2-} + H^+$	2	3.5 x 10^9		Jiang et al. (1992)
S 21	$SO_4^- + O_2^- \longrightarrow SO_4^{2-}$	2	3.5 x 10^9		assumed =S20
S 22	$SO_4^- + H_2O \longrightarrow SO_4^{2-} + H^+ + OH$	2	1.1 x 10^1	-1110	Herrmann et al. (1995)
S 23	$SO_4^- + H_2O_2 \longrightarrow SO_4^{2-} + H^+ + HO_2$	2	1.2 x 10^7		Wine et al. (1989)
S 24	$SO_4^- + NO_3^- \longrightarrow SO_4^{2-} + NO_3$	2	5.0 x 10^4		Exner et al. (1992)
S 25	$SO_4^- + HSO_3^- \longrightarrow SO_3^- + SO_4^{2-} + H^+$	2	8.0 x 10^8		Huie and Neta (1987)
S 26	$SO_4^- + SO_3^{2-} \longrightarrow SO_3^- + SO_4^{2-}$	2	4.6 x 10^8		Huie and Neta (1987)
S 27	$SO_4^{2-} + NO_3 \longrightarrow NO_3^- + SO_4^-$	2	1.0 x 10^5		Logager et al. (1993)
S 28	$SO_5^- + HSO_3^- \longrightarrow SO_4^- + SO_4^{2-} + H^+$	2	7.5 x 10^4		Huie and Neta (1987)
S 29	$SO_5^- + SO_3^{2-} \longrightarrow SO_4^- + SO_4^{2-}$	2	9.4 x 10^6		Huie and Neta (1987)
S 30	$SO_5^- + HSO_3^- \longrightarrow SO_3^- + HSO_5^-$	2	2.5 x 10^4		Huie and Neta (1987); Deister and Warneck (1990)
S 31	$SO_5^- + SO_3^{2-} \xrightarrow{H^+} SO_3^- + HSO_5^-$	2	3.6 x 10^6		Huie and Neta (1987); Deister and Warneck (1990)
S 32	$SO_5^- + O_2^- \xrightarrow{H^+} HSO_5^- + O_2$	2	2.3 x 10^8		Buxton et al. (1996)
S 33	$SO_5^- + SO_5^- \longrightarrow H_2O$	2	1.0 x 10^8		Ross et al. (1992)
S 34	$DMS + O_3 \longrightarrow O_2 + DMSO$	2	8.6 x 10^8	-2600	Gershenzon et al. (2001)
S 35	$DMS + OH \longrightarrow 0.5\ CH_3SO_3^- + 0.5\ CH_3OO + 0.5\ HSO_4^- + HCHO + H^+$	2	1.9 x 10^{10}		Ross et al. (1998)
S 36	$DMSO + OH \longrightarrow CH_3SO_2^- + CH_3OO + H^+$	2	4.5 x 10^9		Bardouki et al. (2002)
S 37	$CH_3SO_2^- + OH \longrightarrow CH_3SO_3^-$ +H2O - O2	2	1.2 x 10^{10}		Bardouki et al. (2002)
S 38	$CH_3SO_3^- + OH \longrightarrow SO_4^{2-} + H^+ + CH_3OO$	2	1.2 x 10^7		Bonsang et al. (1991)

Table 3: Continued.

no	reaction	n	k_0 [$(M^{1-n})s^{-1}$]	$-E_a$ / R [K]	reference
Cl 1	$Cl + H_2O_2 \longrightarrow HO_2 + Cl^- + H^+$	2	2.0 x 10^9		Yu (2001)
Cl 2	$Cl + H_2O \longrightarrow H^+ + ClOH^-$	2	1.8 x 10^5		Yu (2001)
Cl 3	$Cl + NO_3^- \longrightarrow NO_3 + Cl^-$	2	1.0 x 10^8		Buxton et al. (1999b)
Cl 4	$Cl + DOM \longrightarrow Cl^- + HO_2$	2	5.0 x 10^9		estimated (C. Anastasio, pers. comm.) from Ross et al. (1998)
Cl 5	$Cl + SO_4^{2-} \longrightarrow SO_4^- + Cl^-$	2	2.1 x 10^8		Buxton et al. (1999a)
Cl 6	$Cl + Cl \longrightarrow Cl_2$	2	8.8 x 10^7		Wu et al. (1980)
Cl 7	$Cl^- + OH \longrightarrow ClOH^-$	2	4.2 x 10^9		Yu (2001)
Cl 8	$Cl^- + O_3 \longrightarrow ClO^- + O_2$	2	3.0 x 10^{-3}		Hoigné et al. (1985)
Cl 9	$Cl^- + NO_3 \longrightarrow NO_3^- + Cl$	2	9.3 x 10^6	-4330	Exner et al. (1992)
Cl 10	$Cl^- + SO_4^- \longrightarrow SO_4^{2-} + Cl$	2	2.5 x 10^8		Buxton et al. (1999a)
Cl 11	$Cl^- + HSO_5^- \longrightarrow HOCl + SO_4^{2-}$	2	1.8 x 10^{-3}	-7352	Fortnum et al. (1960)
Cl 12	$Cl^- + HOCl + H^+ \longrightarrow Cl_2$	3	2.2 x 10^4	-3508	Ayers et al. (1996)
Cl 13	$Cl_2 \longrightarrow Cl^- + HOCl + H^+$	1	2.2 x 10^1	-8012	Ayers et al. (1996)
Cl 14	$Cl_2^- + OH \longrightarrow HOCl + Cl^-$	2	1.0 x 10^9		Ross et al. (1998)
Cl 15	$Cl_2^- + OH^- \longrightarrow Cl^- + Cl^- + OH$	2	4.0 x 10^6		Jacobi (1996)
Cl 16	$Cl_2^- + HO_2 \longrightarrow Cl^- + Cl^- + H^+ + O_2$	2	3.1 x 10^9		Yu (2001)
Cl 17	$Cl_2^- + O_2^- \longrightarrow Cl^- + Cl^- + O_2$	2	6.0 x 10^9		Jacobi (1996)
Cl 18	$Cl_2^- + H_2O_2 \longrightarrow Cl^- + Cl^- + H^+ + HO_2$	2	7.0 x 10^5	-3340	Jacobi (1996)
Cl 19	$Cl_2^- + NO_2^- \longrightarrow Cl^- + Cl^- + NO_2$	2	6.0 x 10^7		Jacobi (1996)
Cl 20	$Cl_2^- + CH_3OOH \longrightarrow Cl^- + Cl^- + H^+ + CH_3OO$	2	7.0 x 10^5	-3340	assumed by Jacobi (1996)
Cl 21	$Cl_2^- + DOM \longrightarrow Cl^- + Cl^- + HO_2$	2	1.0 x 10^6		estimated (C. Anastasio, pers. comm.) from Ross et al. (1998)
Cl 22	$Cl_2^- + HSO_3^- \longrightarrow SO_3^- + Cl^- + Cl^- + H^+$	2	4.7 x 10^8	-1082	Shoute et al. (1991)
Cl 23	$Cl_2^- + SO_3^{2-} \longrightarrow SO_3^- + Cl^- + Cl^-$	2	6.2 x 10^7		Jacobi et al. (1996)
Cl 24	$Cl_2^- + Cl_2^- \longrightarrow Cl_2 + 2Cl^-$	2	6.2 x 10^9		Yu (2001)
Cl 25	$Cl_2^- + Cl \longrightarrow Cl^- + Cl_2$	2	2.7 x 10^9		Yu (2001)
Cl 26	$Cl_2^- + DMS \longrightarrow 0.5\ CH_3SO_3^- + 0.5\ CH_3OO + 0.5\ HSO_4^- + HCHO + 2\ Cl^- + 2\ H^+$	2	3.0 x 10^9		rate from Ross et al. (1998)
Cl 27	$ClOH^- \longrightarrow Cl^- + OH$	1	6.0 x 10^9		Yu (2001)
Cl 28	$ClOH^- + H^+ \longrightarrow Cl$	2	4.0 x 10^{10}		Yu (2001)
Cl 29	$HOCl + HO_2 \longrightarrow Cl + O_2$	2	7.5 x 10^6		assumed = Cl30 Long and Bielski (1980)
Cl 30	$HOCl + O_2^- \longrightarrow Cl + OH^- + O_2$	2	7.5 x 10^6		Long and Bielski (1980)
Cl 31	$HOCl + SO_3^{2-} \longrightarrow Cl^- + HSO_4^-$	2	7.6 x 10^8		Fogelman et al. (1989)
Cl 32	$HOCl + HSO_3^- \longrightarrow Cl^- + HSO_4^- + H^+$	2	7.6 x 10^8		assumed = Cl31 Fogelman et al. (1989)
Cl 33	$Cl_2 + HO_2 \longrightarrow Cl_2^- + H^+ + O_2$	2	1.0 x 10^9		Bjergbakke et al. (1981)
Cl 34	$Cl_2 + O_2^- \longrightarrow Cl_2^- + O_2$	2	1.0 x 10^9		assumed = Cl33 Bjergbakke et al. (1981)
Cl 35	$Cl^- + HNO_4 \longrightarrow HOCl + NO_3^-$	2	1.4 x 10^{-2}		Régimbal and Mozurkewich (1997)

Table 3: Continued.

no	reaction	n	k_0 $[(M^{1-n})s^{-1}]$	$-E_a$ / R [K]	reference
Br 1	$Br + OH^- \longrightarrow BrOH^-$	2	1.3 x 10^{10}		Zehavi and Rabani (1972)
Br 2	$Br + DOM \longrightarrow Br^- + HO_2$	2	2.0 x 10^{8}		estimated (C. Anastasio, pers. comm.) from Ross et al. (1998)
Br 3	$Br^- + OH \longrightarrow BrOH^-$	2	1.1 x 10^{10}		Zehavi and Rabani (1972)
Br 4	$Br^- + O_3 \longrightarrow BrO^-$	2	2.1 x 10^{2}	-4450	Haag and Hoigné (1983)
Br 5	$Br^- + NO_3 \longrightarrow Br + NO_3^-$	2	3.8 x 10^{9}		Zellner et al. 1996 in Herrmann et al. (2000)
Br 6	$Br^- + SO_4^- \longrightarrow Br + SO_4^{2-}$	2	2.1 x 10^{9}		Jacobi (1996)
Br 7	$Br^- + HSO_5^- \longrightarrow HOBr + SO_4^{2-}$	2	1.0	-5338	Fortnum et al. (1960)
Br 8	$Br^- + HOBr + H^+ \longrightarrow Br_2$	3	1.6 x 10^{10}		Liu and Margerum (2001)
Br 9	$Br_2 \longrightarrow Br^- + HOBr + H^+$	1	9.7 x 10^{1}	7457	Liu and Margerum (2001)
Br 10	$Br_2^- + O_2^- \longrightarrow Br^- + Br^-$	2	1.7 x 10^{8}		Wagner and Strehlow (1987)
Br 11	$Br_2^- + HO_2 \longrightarrow Br_2 + H_2O_2$ - H^+	2	4.4 x 10^{9}		Matthew et al. (2003)
Br 12	$Br_2^- + H_2O_2 \longrightarrow Br^- + Br^- + H^+ + HO_2$	2	5.0 x 10^{2}		Chameides and Stelson (1992)
Br 13	$Br_2^- + Br_2^- \longrightarrow Br^- + Br^- + Br_2$	2	1.9 x 10^{9}		Ross et al. (1992)
Br 14	$Br_2^- + CH_3OOH \longrightarrow Br^- + Br^- + H^+ + CH_3OO$	2	1.0 x 10^{5}		assumed by Jacobi (1996)
Br 15	$Br_2^- + DOM \longrightarrow Br^- + Br^- + HO_2$	2	1.0 x 10^{5}		estimated (C. Anastasio, pers. comm.) from Ross et al. (1998)
Br 16	$Br_2^- + NO_2^- \longrightarrow Br^- + Br^- + NO_2$	2	1.7 x 10^{7}	-1720	Shoute et al. (1991)
Br 17	$Br_2^- + HSO_3^- \longrightarrow Br^- + Br^- + H^+ + SO_3^-$	2	6.3 x 10^{7}	-782	Shoute et al. (1991)
Br 18	$Br_2^- + SO_3^{2-} \longrightarrow Br^- + Br^- + SO_3^-$	2	2.2 x 10^{8}	-650	Shoute et al. (1991)
Br 19	$Br_2^- + DMS \longrightarrow 0.5\ CH_3SO_3^- + 0.5\ CH_3OO + 0.5\ HSO_4^- + HCHO + 2\ Br^- + 2\ H^+$	2	3.2 x 10^{9}		rate from Ross et al. (1998)
Br 20	$BrOH^- \longrightarrow Br^- + OH$	1	3.3 x 10^{7}		Zehavi and Rabani (1972)
Br 21	$BrOH^- \longrightarrow Br + OH^-$	1	4.2 x 10^{6}		Zehavi and Rabani (1972)
Br 22	$BrOH^- + H^+ \longrightarrow Br$	2	4.4 x 10^{10}		Zehavi and Rabani (1972)
Br 23	$BrOH^- + Br^- \longrightarrow Br_2^- + OH^-$	2	1.9 x 10^{8}		Zehavi and Rabani (1972)
Br 24	$BrO^- + SO_3^{2-} \longrightarrow Br^- + SO_4^{2-}$	2	1.0 x 10^{8}		Troy and Margerum (1991)
Br 25	$HOBr + HO_2 \longrightarrow Br + O_2$	2	1.0 x 10^{9}		Herrmann et al. (1999)
Br 26	$HOBr + O_2^- \longrightarrow Br + OH^- + O_2$	2	3.5 x 10^{9}		Schwarz and Bielski (1986)
Br 27	$HOBr + H_2O_2 \longrightarrow Br^- + H^+ + O_2$	2	1.2 x 10^{6}		von Gunten and Oliveras (1998)
Br 28	$HOBr + SO_3^{2-} \longrightarrow Br^- + HSO_4^-$	2	5.0 x 10^{9}		Troy and Margerum (1991)
Br 29	$HOBr + HSO_3^- \longrightarrow Br^- + HSO_4^- + H^+$	2	5.0 x 10^{9}		assumed = Br28
Br 30	$Br_2 + HO_2 \longrightarrow Br_2^- + H^+ + O_2$	2	1.1 x 10^{8}		Ross et al. (1998)
Br 31	$Br_2 + O_2^- \longrightarrow Br_2^- + O_2$	2	5.6 x 10^{9}		Ross et al. (1998)
Br 32	$Br^- + HNO_4 \longrightarrow HOBr + NO_3^-$	2	5.4 x 10^{-1}		Régimbal and Mozurkewich (1997)
Br 33	$Br^- + O_3 + H^+ \longrightarrow HOBr + O_2$	3	11.7		Haag and Hoigné (1983)

Table 3: Continued.

no	reaction	n	k_0 [$(M^{1-n})s^{-1}$]	$-E_a$ / R [K]	reference
Hx 1	$Br^- + HOCl + H^+ \longrightarrow BrCl$	3	1.3 x 10^6		Liu and Margerum (2001)
Hx 2	$Cl^- + HOBr + H^+ \longrightarrow BrCl$	3	2.3 x 10^{10}		Liu and Margerum (2001)
Hx 3	$BrCl \longrightarrow Cl^- + HOBr + H^+$	1	3.0 x 10^6		Liu and Margerum (2001)
Hx 4	$Br^- + ClO^- + H^+ \longrightarrow BrCl + OH^-$	3	3.7 x 10^{10}		Kumar and Margerum (1987)
Hx 5	$Cl_2 + Br^- \longrightarrow BrCl_2^-$	2	7.7 x 10^9		Liu and Margerum (2001)
Hx 6	$BrCl_2^- \longrightarrow Cl_2 + Br^-$	1	1.83 x 10^3		Liu and Margerum (2001)
hv 1	$O_3 + hv \longrightarrow OH + OH + O_2$	[1]	1		assumed 2x gas phase
hv 2	$H_2O_2 + hv \longrightarrow OH + OH$	[1]	1		assumed 2x gas phase
hv 3	$NO_3^- + hv \xrightarrow{H^+} NO_2 + OH$	[1]	1		Zellner et al. (1990)
hv 4	$NO_2^- + hv \xrightarrow{H^+} NO + OH$	[1]	1		Zellner et al. (1990); Burley and Johnston (1992)
hv 5	$HOCl + hv \longrightarrow OH + Cl$	[1]	1		assumed 2x gas phase
hv 6	$Cl_2 + hv \longrightarrow Cl + Cl$	[1]	1		assumed 2x gas phase
hv 7	$HOBr + hv \longrightarrow OH + Br$	[1]	1		assumed 2x gas phase
hv 8	$Br_2 + hv \longrightarrow Br + Br$	[1]	1		assumed 2x gas phase
hv 9	$BrCl + hv \longrightarrow Cl + Br$	[1]	1		assumed 2x gas phase

n is the order of the reaction. [1] photolysis rates calculated online. The temperature dependence is $k = k_0 \times \exp(\frac{-E_a}{R}(\frac{1}{T} - \frac{1}{T_0}))$, $T_0 = 298$ K.

Table 4: Heterogeneous reactions.

no	reaction	k	reference
H 1	$N_2O_5 \xrightarrow{H_2O} HNO_{3aq} + HNO_{3aq}$	$\overline{k_t}(N_2O_5) w_{l,i} [H_2O]/Het_T$	Behnke et al. (1994), Behnke et al. (1997)
H 2	$N_2O_5 \xrightarrow{Cl^-} ClNO_2 + NO_3^-$	$\overline{k_t}(N_2O_5) w_{l,i} f(Cl^-)[Cl^-]/Het_T$	Behnke et al. (1994), Behnke et al. (1997)
H 3	$N_2O_5 \xrightarrow{Br^-} BrNO_2 + NO_3^-$	$\overline{k_t}(N_2O_5) w_{l,i} f(Br^-)[Br^-]/Het_T$	Behnke et al. (1994), Behnke et al. (1997)
H 4	$ClNO_3 \xrightarrow{H_2O} HOCl_{aq} + HNO_{3aq}$	$\overline{k_t}(ClNO_3) w_{l,i} [H_2O]/Het_T$	see note
H 5	$ClNO_3 \xrightarrow{Cl^-} Cl_{2aq} + NO_3^-$	$\overline{k_t}(ClNO_3) w_{l,i} f(Cl^-)[Cl^-]/Het_T$	see note
H 6	$ClNO_3 \xrightarrow{Br^-} BrCl_{aq} + NO_3^-$	$\overline{k_t}(ClNO_3) w_{l,i} f(Br^-)[Br^-]/Het_T$	see note
H 7	$BrNO_3 \xrightarrow{H_2O} HOBr_{aq} + HNO_{3aq}$	$\overline{k_t}(BrNO_3) w_{l,i} [H_2O]/Het_T$	see note
H 8	$BrNO_3 \xrightarrow{Cl^-} BrCl_{aq} + NO_3^-$	$\overline{k_t}(BrNO_3) w_{l,i} f(Cl^-)[Cl^-]/Het_T$	see note
H 9	$BrNO_3 \xrightarrow{Br^-} Br_{2aq} + NO_3^-$	$\overline{k_t}(BrNO_3) w_{l,i} f(Br^-)[Br^-]/Het_T$	see note

For a definition of $\overline{k_t}$ and $w_{l,i}$ see von Glasow et al. (2002) or von Glasow (2000). $Het_T = [H_2O + f(Cl^-)[Cl^-] + f(Br^-)[Br^-]$, with $f(Cl^-) = 5.0 \times 10^2$ and $f(Br^-) = 3.0 \times 10^5$. H4 - H9: the total rate is determined by $\overline{k_t}$, the distribution among the different reaction paths was assumed to be the same as for reactions H1 - H3.

Table 5: Aqueous phase equilibrium constants.

no	reaction	m	n	K_0 [M^{n-m}]	-$\Delta H/R$ [K]	reference
EQ 1	$CO_{2aq} \longleftrightarrow H^+ + HCO_3^-$	1	2	4.3×10^{-7}	-913	Chameides (1984)
EQ 2	$NH_{3aq} \longleftrightarrow OH^- + NH_4^+$	1	2	1.7×10^{-5}	-4325	Chameides (1984)
EQ 3	$H_2O_{aq} \longleftrightarrow H^+ + OH^-$	1	2	1.0×10^{-14}	-6716	Chameides (1984)
EQ 4	$HCOOH_{aq} \longleftrightarrow H^+ + HCOO^-$	1	2	1.8×10^{-4}		Weast (1980)
EQ 5	$HSO_3^- \longleftrightarrow H^+ + SO_3^{2-}$	1	2	6.0×10^{-8}	1120	Chameides (1984)
EQ 6	$H_2SO_{4aq} \longleftrightarrow H^+ + HSO_4^-$	1	2	1.0×10^{3}		Seinfeld and Pandis (1998)
EQ 7	$HSO_4^- \longleftrightarrow H^+ + SO_4^{2-}$	1	2	1.2×10^{-2}	1120	Weast (1980)
EQ 8	$HO_{2aq} \longleftrightarrow O_2^- + H^+$	1	2	1.6×10^{-5}		Weinstein-Lloyd and Schwartz (1991)
EQ 9	$SO_{2aq} \longleftrightarrow H^+ + HSO_3^-$	1	2	1.7×10^{-2}	2090	Chameides (1984)
EQ 10	$Cl_2^- \longleftrightarrow Cl_{aq} + Cl^-$	1	2	5.2×10^{-6}		Jayson et al. (1973)
EQ 11	$HOCl_{aq} \longleftrightarrow H^+ + ClO^-$	1	2	3.2×10^{-8}		Lax (1969)
EQ 12	$HBr_{aq} \longleftrightarrow H^+ + Br^-$	1	2	1.0×10^{9}		Lax (1969)
EQ 13	$Br_2^- \longleftrightarrow Br_{aq} + Br^-$	1	2	9.1×10^{-6}		Mamou et al. (1977)
EQ 14	$HOBr_{aq} \longleftrightarrow H^+ + BrO^-$	1	2	2.3×10^{-9}	-3091	Kelley and Tartar (1956)
EQ 15	$BrCl_{aq} + Cl^- \longleftrightarrow BrCl_2^-$	2	1	3.8	1143	Wang et al. (1994)
EQ 16	$BrCl_{aq} + Br^- \longleftrightarrow Br_2Cl^-$	2	1	1.8×10^{4}		Wang et al. (1994)
EQ 17	$Br_{2aq} + Cl^- \longleftrightarrow Br_2Cl^-$	2	1	1.3		Wang et al. (1994)
EQ 18	$HNO_{3aq} \longleftrightarrow H^+ + NO_3^-$	1	2	1.5×10^{1}		Davis and de Bruin (1964)
EQ 19	$HCl_{aq} \longleftrightarrow H^+ + Cl^-$	1	2	1.7×10^{6}		Marsh and McElroy (1985)
EQ 20	$HONO_{aq} \longleftrightarrow H^+ + NO_2^-$	1	2	5.1×10^{-4}	-1260	Schwartz and White (1981)
EQ 21	$HNO_{4aq} \longleftrightarrow NO_4^- + H^+$	1	2	1.0×10^{-5}	8700	Warneck (1999)

The temperature dependence is $K = K_0 \times \exp(\frac{-\Delta H}{R}(\frac{1}{T} - \frac{1}{T_0}))$, $T_0 = 298$ K.

Table 6: Henry constants and accommodation coefficients.

specie	K_H^0 [M/atm]	$-\Delta_{soln}H/R$ [K]	reference	α^0	$-\Delta_{obs}H/R$ [K]	reference
O_3	1.2×10^{-2}	2560	Chameides (1984)	0.002	(at 292 K)	DeMore et al. (1997)
O_2	1.3×10^{-3}	1500	Wilhelm et al. (1977)	0.01	2000	estimated
OH	3.0×10^{1}	4300	Hanson et al. (1992)	0.01	(at 293 K)	Takami et al. (1998)
HO_2	3.9×10^{3}	5900	Hanson et al. (1992)	0.2	(at 293 K)	DeMore et al. (1997)
H_2O_2	1.0×10^{5}	6338	Lind and Kok (1994)	0.077	2769	Worsnop et al. (1989)
NO_2	6.4×10^{-3}	2500	Lelieveld and Crutzen (1991)	0.0015	(at 298 K)	Ponche et al. (1993)
NO_3	2.0	2000	Thomas et al. (1993)	0.04	(at 273? K)	Rudich et al. (1996)
HONO	4.9×10^{1}	4780	Schwartz and White (1981)	0.04	(at 247-297 K)	DeMore et al. (1997)
HNO_3	1.7×10^{5}	8694	Lelieveld and Crutzen (1991)	0.5	(at RT)	Abbatt and Waschewsky (1998)
HNO_4	1.2×10^{4}	6900	Régimbal and Mozurkewich (1997)	0.1	(at 200 K)	DeMore et al. (1997)
NH_3	5.8×10^{1}	4085	Chameides (1984)	0.06	(at 295 K)	DeMore et al. (1997)
CH_3OO	6.0	$=HO_2$	Pandis and Seinfeld (1989)	0.01	2000	estimated
ROOH	3.0×10^{2}	5322	Lind and Kok (1994)	0.0046	3273	Magi et al. (1997)
HCHO	7.0×10^{3}	6425	Chameides (1984)	0.04	(at 260-270 K)	DeMore et al. (1997)
HCOOH	3.7×10^{3}	5700	Chameides (1984)	0.014	3978	DeMore et al. (1997)
CO_2	3.1×10^{-2}	2423	Chameides (1984)	0.01	2000	estimated
HCl	1.2	9001	Brimblecombe and Clegg (1989)	0.074	3072	Schweitzer et al. (2000)
HOCl	6.7×10^{2}	5862	Huthwelker et al. (1995)	=HOBr	=HOBr	estimated
$ClNO_3$	∞	—		0.1	(at RT)	Koch and Rossi (1998)
Cl_2	9.1×10^{-2}	2500	Wilhelm et al. (1977)	0.038	6546	Hu et al. (1995)
HBr	1.3	10239	Brimblecombe and Clegg (1989)	0.031	3940	Schweitzer et al. (2000)
HOBr	9.3×10^{1}	=HOCl	Vogt et al. (1996)	0.5	(at RT)	Abbatt and Waschewsky (1998)
$BrNO_3$	∞	—		0.8	0	Hanson et al. (1996)
Br_2	7.6×10^{-1}	4094	Dean (1992)	0.038	6546	Hu et al. (1995)
BrCl	9.4×10^{-1}	5600	Bartlett and Margerum (1999)	$=Cl_2$	$=Cl_2$	estimated
DMSO	5.0×10^{4}	=HCHO	De Bruyn et al. (1994)	0.048	2578	De Bruyn et al. (1994)
$DMSO_2$	∞	—	assumed	0.03	5388	De Bruyn et al. (1994)
SO_2	1.2	3120	Chameides (1984)	0.11	0	DeMore et al. (1997)
H_2SO_4	∞	—		0.65	(at 303 K)	Pöschl et al. (1998)
CH_3SO_2H	∞	—	assumed	0.0002	0	Lucas and Prinn (2002)
CH_3SO_3H	∞	—	assumed	0.076	1762	De Bruyn et al. (1994)

For ROOH the values of CH_3OOH have been assumed. The temperature dependence is for the Henry constants is $K_H = K_H^0 \times \exp(\frac{-\Delta_{soln}H}{R}(\frac{1}{T} - \frac{1}{T_0}))$, $T_0 = 298$ K and for the accommodation coefficients $dln(\frac{\alpha}{1-\alpha})/d(\frac{1}{T}) = \frac{-\Delta_{obs}H}{R}$. RT stands for “room temperature”.

References

Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO_3, O_3 and NO_2 with deliquescent NaCl aerosols at room temperature, J. Phys. Chem. A, 102, 3719 – 3725, 1998.

Anderson, L. C. and Fahey, D. W.: Studies with $ClONO_2$: Thermal dissociation rate and catalytic conversion to NO using an NO/O_3 chemiluminescence detector, J. Phys. Chem., 94, 644 – 652, 1990.

Aranda, A., Le Bras, G., La Verdet, G., and Poulet, G.: The BrO + CH_3O_2 reaction: Kinetics and role in the atmospheric ozone budget, Geophys. Res. Lett., 24, 2745 – 2748, 1997.

Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry: Supplement VI, J. Phys. Chem. Ref. Data, 26, 1329 – 1499, 1997.

Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, http://www.iupac-kinetic.ch.cam.ac.uk, 1999.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Jenkin, R. G. H. M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, Jul. 2004, http://www.iupac-kinetic.ch.cam.ac.uk, 2004.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, February 2006, http://www.iupac-kinetic.ch.cam.ac.uk, 2006.

Ayers, G. P., Penkett, S. A., Gillett, R. W., Bandy, B., Galbally, I. E., Meyer, C. P., Elsworth, C. M., Bentle, S. T., and Forgan, B. W.: The Annual Cycle of Peroxides and Ozone in Marine Air at Cape Grim, Tasmania, J. Atmos. Chem., 23, 221 – 252, 1996.

Bardouki, H., da Rosa, M. B., Mihalopoulos, N., Palm, W.-U., and Zetzsch, C.: Kinetics and mechanism of the oxidation of dimethylsulfoxide (DMSO) and methanesulfinate (MSI^-) in aqueous medium, Atmos. Environ., 36, 4627 – 4634, 2002.

Barker, G. C., Fowles, P., and Stringer, B.: Pulse radiolytic induced transient electrical conductance in liquid solutions, Trans. Faraday Soc., 66, 1509–1519, 1970.

Barnes, I., Bastian, V., Becker, K. H., and Overath, R. D.: Kinetic studies of the reactions of IO, BrO and ClO with DMS, Int. J. Chem. Kinet., 23, 579 – 591, 1991.

Barone, S. B., Turnipseed, A. A., and Ravishankara, A. R.: Role of adducts in the atmospheric oxidation of dimethyl sulfide, Faraday Discuss., 100, 39 – 54, 1995.

Bartlett, W. P. and Margerum, D. W.: Temperature dependencies of the Henry's law constant and the aqueous phase dissociation constant of bromine chloride, Environ. Sci. Technol., 33, 3410–3414, 1999.

Behnke, W., Scheer, V., and Zetzsch, C.: Production of $BrNO_2$, Br_2 and $ClNO_2$ from the Reaction between Sea Spray Aerosol and N_2O_5, J. Aerosol Sci., 25, Suppl.1, S277 – S278, 1994.

Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of $ClNO_2$ from the reaction of gaseous N_2O_5 with NaCl solution: Bulk and aerosol experiments, J. Geophys. Res., 102, 3795 – 3804, 1997.

Betterton, E. A. and Hoffmann, M. R.: Oxidation of aqueous SO_2 by peroxymonosulfate, J. Phys. Chem., 92, 5962–5965, 1988.

Bjergbakke, E., Navartnam, S., Parsons, B. J., and Swallow, A. J.: Reaction between $HO_2\cdot$ and chlorine in aqueous solution, J. Am. Chem. Soc., 103, 5926–5928, 1981.

Bonsang, B., Martin, D., Lambert, G., Kanakidou, M., de Roulley, J., and Sennequir, G.: Vertical distribution of nonmethane hydrocarbons in the remote marine boundary layer, J. Geophys. Res., 96, 7313 – 7324, 1991.

Boyce, S. D. and Hoffmann, M. R.: Kinetics and mechanism of the formation of hydroxymethanesulfonic acid at low pH, J. Phys. Chem., 88, 4740–4746, 1984.

Brimblecombe, P. and Clegg, S. L.: Erratum, J. Atmos. Chem., 8, 95, 1989.

Burley, J. D. and Johnston, H. S.: Ionic mechanisms for heterogeneous stratospheric reactions and ultraviolet photoabsorption cross-sections for NO_2^+, HNO_3, and NO_3^- in sulfuric-acid, Geophys. Res. Lett., 19, 1359 – 1362, 1992.

Buxton, G. V., Greenstock, C. L., Helman, W. P., and Ross, A. B.: Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot OH/\cdot O^-$) in aqueous solution, J. Phys. Chem. Ref. Data, 17, 513–886, 1988.

Buxton, G. V., McGowan, S., Salmon, G. S., Williams, J. E., and Wood, N. D.: A study of the spectra and reactivity of oxysulphur-radical anions involved in the chain oxidation of S(IV): A pulse and γ-radiolysis study, Atmos. Environ., 30, 2483–2493, 1996.

Buxton, G. V., Bydder, M., and Salmon, G. A.: The reactivity of chlorine atoms in aqueous solution Part II: The equilibrium $SO_4^- + Cl^- \longleftrightarrow Cl + SO_4^{2-}$, Phys. Chem. Chem. Phys., 1, 269 – 273, 1999a.

Buxton, G. V., Salmon, G. A., and Wang, J.: The equilibrium $NO_3 + Cl^- \longleftrightarrow NO_3^- + Cl$: A laser flash photolysis and pulse radiolysis study of the reactivity of NO_3 with chloride ion in aqueous solution, Phys. Chem. Chem. Phys., 1, 3589 – 3593, 1999b.

Chameides, W. L.: The Photochemistry of a Remote Marine Stratiform Cloud, J. Geophys. Res., 89, 4739 – 4755, 1984.

Chameides, W. L. and Stelson, A. W.: Aqueous-Phase Chemical Processes in Deliquescent Sea-Salt Aerosols: A Mechanism That Couples the Atmospheric Cycles of S and Sea Salt, J. Geophys. Res., 97, 20 565 – 20 580, 1992.

Chin, M. and Wine, P. H.: A temperature-dependent competitive kinetics study of the aqueous-phase reactions of OH radicals with formate, formic acid, acetate, acetic acid, and hydrated formaldehyde, in Aquatic and Surface Photochemistry, edited by G. R. Helz, R. G. Zepp, and D. G. Crosby, pp. 85–96, A. F. Lewis, NY, 1994.

Christensen, H. and Sehested, K.: HO_2 and O_2^- radicals at elevated temperatures, J. Phys. Chem., 92, 3007–3011, 1988.

Christensen, H., Sehested, K., and Corfitzen, H.: Reactions of hydroxyl radicals with hydrogen peroxide at ambient and elevated temperatures, J. Phys. Chem., 86, 1588–1590, 1982.

Clifton, C. L., Altstein, N., and Huie, R. E.: Rate constant for the reaction of NO_2 with sulfur(IV) over the pH range 5.3–13, Environ. Sci. Technol., 22, 586–589, 1988.

Damschen, D. E. and Martin, L. R.: Aqueous aerosol oxidation of nitrous acid by O_2, O_3 and H_2O_2, Atmos. Environ., 17, 2005–2011, 1983.

Davis, Jr., W. and de Bruin, H. J.: New activity coefficients of 0-100 per cent aqueous nitric acid, J. Inorg. Nucl. Chem., 26, 1069 – 1083, 1964.

De Bruyn, W. J., Shorter, J. A., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Uptake of gas phase sulfur species methanesulfonic acid, dimethylsulfoxide, and dimethyl sulfone by aqueous surface, J. Geophys. Res., 99, 16 927 – 16 932, 1994.

Dean, J. A.: Lange's Handbook of Chemistry, McGraw-Hill, Inc., 1992.

Deister, U. and Warneck, P.: Photooxidation of SO_3^{2-} in aqueous solution, J. Phys. Chem., 94, 2191–2198, 1990.

DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling, Tech. Rep. JPL Publication 97-4, Jet Propulsion Laboratory, Pasadena, CA, 1997.

Exner, M., Herrmann, H., and Zellner, R.: Laser-based studies of reactions of the nitrate radical in aqueous solution, Ber. Bunsenges. Phys. Chem., 96, 470 – 477, 1992.

Fogelman, K. D., Walker, D. M., and Margerum, D. W.: Non-metal redox kinetics: Hypochlorite and hypochlorous acid reactions with sulfite, Inorg. Chem., 28, 986 – 993, 1989.

Fortnum, D. H., Battaglia, C. J., Cohen, S. R., and Edwards, J. O.: The kinetics of the oxidation of halide ions by monosubstituted peroxides, J. Am. Chem. Soc., 82, 778–782, 1960.

Gershenzon, M., Davidovits, P., Jayne, J. T., Kolb, C. E., and Worsnop, D. R.: Simultaneous Uptake of DMS and Ozone on Water, J. Phys. Chem. A, 105, 7031 – 7036, 2001.

Haag, W. R. and Hoigné, J.: Ozonation of bromide-containing waters: Kinetics of formation of hypobromous acid and bromate, Environ. Sci. Technol., 17, 261 – 267, 1983.

Haggerstone, A.-L., Carpenter, L. J., Carslaw, N., and McFiggans, G.: Improved model predictions of HO_2 with gas to particle mass transfer rates calculated using aerosol number size distributions, J. Geophys. Res., 110, D04 303, doi: 10.1029/2004JD005 282, 2005.

Hansen, J. C., Li, Y., Francisco, J. S., and Li, Z.: On the Mechanism of the BrO + CH_2O Reaction, J. Phys. Chem. A, 103, 8543 – 8546, 1999.

Hanson, D. R., Burkholder, J. B., Howard, C. J., and Ravishankara, A. R.: Measurement of OH and HO_2 radical uptake coefficients on water and sulfuric acid surfaces, J. Phys. Chem., 96, 4979–4985, 1992.

Hanson, D. R., Ravishankara, A. R., and Lovejoy, E. R.: Reaction of $BrONO_2$ with H_2O on submicron sulfuric acid aerosol and the implications for the lower stratosphere, J. Geophys. Res., 101D, 9063–9069, 1996.

Herrmann, H., Reese, A., and Zellner, R.: Time resolved UV/VIS diode array absorption spectroscopy of SO_x^- (x=3, 4, 5) radical anions in aqueous solution, J. Mol. Struct., 348, 183–186, 1995.

Herrmann, H., Ervens, B., Nowacki, P., Wolke, R., and Zellner, R.: A chemical aqueous phase radical mechanism for tropospheric chemistry, Chemosphere, 38, 1223 – 1232, 1999.

Herrmann, H., Ervens, B., Jacobi, H.-W., Wolke, R., Nowacki, P., and Zellner, R.: CAPRAM2.3: A Chemical Aqueous Phase Radical Mechanism for Tropospheric Chemistry, J. Atmos. Chem., 36, 231 – 284, 2000.

Hoffmann, M. R.: On the kinetics and mechanism of oxidation of aquated sulfur dioxide by ozone, Atmos. Environ., 20, 1145 – 1154, 1986.

Hoigné, J., Bader, H., Haag, W. R., and Staehelin, J.: Rate constants of reactions of ozone with organic and inorganic compounds in water — III Inorganic compounds and radicals, Wat. Res., 19, 993–1004, 1985.

Hu, J. H., Shi, Q., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Reactive uptake of Cl_2 and Br_2 by aqueous surfaces as a function of Br^- and I^- ion concentration: The effect of chemical reaction at the interface, J. Phys. Chem., 99, 8768 – 8776, 1995.

Hubinger, S. and Nee, J. B.: Absorption spectra of Cl_2, Br_2 and BrCl between 190 and 600 nm, J. Photochem. Photobiol. A: Chem., 86, 1–7, 1995.

Huie, R. E. and Neta, P.: Rate constants for some oxidations of S(IV) by radicals in aqueous solutions, Atmos. Environ., 21, 1743–1747, 1987.

Huthwelker, T., Clegg, S. L., Peter, T., Carslaw, K., Luo, B. P., and Brimblecombe, P.: Solubility of HOCl in water and aqueous H_2SO_4 to stratospheric temperatures, J. Atmos. Chem., 21, 81–95, 1995.

Ingham, T., Bauer, D., Sander, R., Crutzen, P. J., and Crowley, J. N.: Kinetics and Products of the Reactions BrO + DMS and Br + DMS at 298 K, J. Phys. Chem. A, 103, 7199 – 7209, 1999.

Jacob, D. J.: Chemistry of OH in Remote Clouds and Its Role in the Production of Formic Acid and Peroxymonosulfate, J. Geophys. Res., 91, 9807 – 9826, 1986.

Jacobi, H.-W.: Kinetische Untersuchungen und Modellrechnungen zur troposphärischen Chemie von Radikalanionen und Ozon in wässriger Phase, Ph.D. thesis, Universität Essen, Germany, 1996.

Jacobi, H.-W., Herrmann, H., and Zellner, R.: Kinetic investigation of the Cl_2^- radical in the aqueous phase, in Air Pollution Research Report 57: Homogenous and hetrogenous chemical Processes in the Troposphere, edited by P. Mirabel, pp. 172–176, Office for official Publications of the European Communities, Luxembourg, 1996.

Jayne, J. T., Pöschl, U., Chen, Y., Dai, D., Molina, L. T., Worsnop, D. R., Kolb, C. E., and Molina, M. J.: Pressure and Temperature Dependence of the Gas-Phase Reaction of SO_3 with H_2O and the Heterogeneous Reaction of SO_3 with H_2O/H_2SO_4 Surfaces, J. Phys. Chem. A, 101, 10 000 – 10 011, 1997.

Jayson, G. G., Parsons, B. J., and Swallow, A. J.: Some simple, highly reactive, inorganic chlorine derivatives in aqueous solution, J. Chem. Soc. Faraday Trans., 69, 1597–1607, 1973.

Jefferson, A., Nicovich, J. M., and Wine, P. H.: Temperature-dependent kinetics studies of the reactions $Br(^2P_{3/2})$ + CH_3SCH_3 ↔ CH_3SCH_2 + HBr. Heat of formation of the CH_3SCH_2 radical, J. Phys. Chem., 98, 7128 – 7135, 1994.

Jiang, P.-Y., Katsumura, Y., Nagaishi, R., Domae, M., Ishikawa, K., Ishigure, K., and Yoshida, Y.: Pulse radiolysis study of concentrated sulfuric acid solutions. Formation mechanism, yield and reactivity of sulfate radicals, J. Chem. Soc. Faraday Trans., 88, 1653–1658, 1992.

Kelley, C. M. and Tartar, H. V.: On the system: bromine-water, J. Am. Chem. Soc., 78, 5752–5756, 1956.

Koch, T. G. and Rossi, M. J.: Direct measurement of surface residence times: Nitryl chloride and chlorine nitrate on alkali halides at room temperature, J. Phys. Chem. A, 102, 9193–9201, 1998.

Kukui, A., Bossoutrot, V., Laverdet, G., and Bras, G. L.: Mechanism of the Reaction of CH_3SO with NO_2 in Relation to Atmospheric Oxidation of Dimethyl Sulfide: Experimental and Theoretical Study, J. Phys. Chem. A, 104, 935 – 946, 2000.

Kukui, A., Borissenko, D., Laverdet, G., and Bras, G. L.: Gas phase reactions of OH radicals with dimethyl sulfoxide and methane sulfonic acid using turbulent flow reactor and chemical ionization mass spectrometry, J. Phys. Chem. A, 107, 5732 – 5742, 2003.

Kumar, K. and Margerum, D. W.: Kinetics and mechanism of general-acid-assisted oxidation of bromide by hypochlorite and hypochlorous acid, Inorg. Chem., 26, 2706 – 2711, 1987.

Lax, E.: Taschenbuch für Chemiker und Physiker, Springer Verlag, Berlin, 1969.

Lee, Y.-N. and Schwartz, S. E.: Reaction kinetics of nitrogen dioxide with liquid water at low partial pressure, J. Phys. Chem., 85, 840–848, 1981.

Lelieveld, J. and Crutzen, P. J.: The Role of Clouds in Tropospheric Photochemistry, J. Atmos. Chem., 12, 229 – 267, 1991.

Lind, J. A. and Kok, G. L.: Correction to “Henry’s law determinations for aqueous solutions of hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid” by John A. Lind and Gregory L. Kok, J. Geophys. Res., 99D, 21 119, 1994.

Lind, J. A., Lazrus, A. L., and Kok, G. L.: Aqueous phase oxidation of sulfur(IV) by hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid, J. Geophys. Res., 92D, 4171–4177, 1987.

Liu, Q. and Margerum, D. W.: Equilibrium and Kinetics of Bromine Chloride Hydrolysis, est, 35, 1127 – 1133, 2001.

Logager, T., Sehested, K., and Holcman, J.: Rate constants of the equilibrium reactions SO_4 + HNO_3 $\longleftrightarrow$ HSO_4^- + NO_3 and SO_4 + NO_3 $\longleftrightarrow$ SO_4^{2-} + NO_3., Radiat. Phys. Chem., 41, 539 – 543, 1993.

Long, C. A. and Bielski, B. H. J.: Rate of reaction of superoxide radical with chloride-containing species, J. Phys. Chem., 84, 555–557, 1980.

Lucas, D. D. and Prinn, R. G.: Mechanistic studies of dimethylsulfide oxidation products using an observationally constrained model, J. Geophys. Res., 107, doi: 10.1029/2001JD000 843, 2002.

Lurmann, F. W., Lloyd, A. C., and Atkinson, R.: A Chemical Mechanism for Use in Long-Range Transport/Acid Deposition Computer Modeling, J. Geophys. Res., 91, 10 905 – 10 936, 1986.

Magi, L., Schweitzer, F., Pallares, C., Cherif, S., , Mirabel, P., and George, C.: Investigation of the uptake rate of ozone and methyl hydroperoxide by water surfaces, J. Phys. Chem. A, 101, 4943 – 4949, 1997.

Mallard, W. G., Westley, F., Herron, J. T., Hampson, R. F., and Frizzel, D. H.: NIST Chemical Kinetics Database: Version 5.0, National Institute of Standards and Technology, Gaithersburg, MD, 1993.

Mamou, A., Rabani, J., and Behar, D.: On the oxidation of aqueous Br^- by OH radicals, studied by pulse radiolysis, J. Phys. Chem., 81, 1447–1448, 1977.

Marsh, A. R. W. and McElroy, W. J.: The dissociation constant and Henry's law constant of HCl in aqueous solution, Atmos. Environ., 19, 1075 – 1080, 1985.

Matthew, B. M., George, I., and Anastasio, C.: Hydroperoxyl radical (HO_2) oxidizes dibromide radical anion (Br_2^-) to bromine (Br_2) in aqueous solution: Implications for the formation of Br_2 in the marine boundary layer, Geophys. Res. Lett., 30 (24), 2297, doi: 10.1029/2003GL018 572, 2003.

Mellouki, A., Bras, G. L., and Poulet, G.: Discharge Flow Kinetic Study of NO_3 Reactions with Free Radicals: The Reaction of NO_3 with Cl, J. Phys. Chem., 91, 5760–5764, 1987.

Muthuramu, K., Shepson, P. B., Bottenheim, J. W., Jobson, T., Niki, H., and Anlauf, K. G.: Relationships Between Organic Nitrates and Surface Ozone Destruction During Polar Sunrise Experiment 1992, J. Geophys. Res., 99 (D12), 25 369–25 378, doi:10.1029/94JD01 309, 1994.

Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of $BrONO_2$ and the heat of formation of $BrONO_2$, J. Phys. Chem., 100, 19 398 – 19 405, 1996.

Pandis, S. N. and Seinfeld, J. H.: Sensitivity Analysis of a Chemical Mechanism for Aqueous-Phase Atmospheric Chemistry, J. Geophys. Res., 94, 1105 – 1126, 1989.

Ponche, J. L., George, C., and Mirabel, P.: Mass transfer at the air/water interface: Mass accommodation coefficients of SO_2, HNO_3, NO_2 and NH_3, J. Atmos. Chem., 16, 1–21, 1993.

Pöschl, U., Canagaratna, M., Jayne, J. T., Molina, L. T., Worsnop, D. R., Kolb, C. E., and Molina, M. J.: Mass accommodation coefficient of H_2SO_4 vapor on aqueous sulfuric acid surfaces and gaseous diffusion coefficient of H_2SO_4 in N_2/H_2O, J. Phys. Chem. A, 102, 10 082–10 089, 1998.

Ray, A., Vassalli, I., G.Laverdet, and Bras, G. L.: Kinetics of the Thermal Decomposition of the CH_3SO_2 Radical and Its Reaction with NO_2 at 1 Torr and 298 K, J. Phys. Chem., 100, 8895 – 8900, 1996.

Régimbal, J.-M. and Mozurkewich, M.: Peroxynitric acid decay mechanisms and kinetics at low pH, J. Phys. Chem. A, 101, 8822–8829, 1997.

Ross, A. B., Mallard, W. G., Helman, W. P., Bielski, B. H. J., Buxton, G. V., Cabelli, D. E., Greenstock, C. L., Huie, R. E., and Neta, P.: NDRL-NIST Solution Kinetics Database: - Ver. 1, National Institute of Standards and Technology, Gaithersburg, MD, 1992.

Ross, A. B., Mallard, W. G., Helman, W. P., Buxton, G. V., Huie, R. E., and Neta, P.: NDRL-NIST Solution kinetics database: Version 3.0, Notre Dame Radiation Laboratory, Notre Dame, and National Institut of Standards and Technology, Gaithersburg, 1998.

Rudich, Y., Talukdar, R. K., Imamura, T., Fox, R. W., and Ravishankara, A. R.: Uptake of NO_3 on KI solutions: Rate coefficient for the $NO_3 + I^-$ reaction and gas-phase diffusion coefficients for NO_3, Chem. Phys. Lett., 261, 467–473, 1996.

Sander, R., Vogt, R., Harris, G. W., and Crutzen, P. J.: Modeling the chemistry of ozone, halogen compounds, and hydrocarbons in the arctic troposphere during spring, Tellus, 49B (5), 522–532, 1997.

Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Huie, R. E., Orkin, V. L., Moortgat, G. K., Ravishankara, A. R., Kolb, C. E., Molina, M. J., and Finlayson-Pitts, B. J.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Tech. Rep. JPL Publication 02-25, Jet Propulsion Laboratory, Pasadena, CA, 2003.

Scheffler, D., Grothe, H., Willner, H., Frenzel, A., and Zetzsch, C.: Properties of Pure Nitryl Bromide. Thermal Behaviour, UV/Vis and FTIR Spectra, and Photoisomerization to *trans*-BrONO in an Argon Matrix, Inorg. Chem., 36, 335 – 338, 1997.

Schwartz, S. E. and White, W. H.: Solubility equilibria of the nitrogen oxides and oxyacids in dilute aqueous solution, in Advances in Environmental Science and Engineering, edited by J. R. Pfafflin and E. N. Ziegler, vol. 4, pp. 1–45, Gordon and Breach Science Publishers, NY, 1981.

Schwarz, H. A. and Bielski, B. H. J.: Reactions of HO_2 and O_2^- with iodine and bromine and the I_2^- and I atom reduction potentials, J. Phys. Chem., 90, 1445–1448, 1986.

Schweitzer, F., Mirabel, P., and George, C.: Uptake of hydrogen halides by water droplets, J. Phys. Chem. A, 104, 72–76, 2000.

Sehested, K., Rasmussen, O. L., and Fricke, H.: Rate constants of OH with HO_2, O_2^-, and $H_2O_2^+$ from hydrogen peroxide formation in pulse-irradiated oxygenated water, J. Phys. Chem., 72, 626–631, 1968.

Sehested, K., Holcman, J., and Hart, E. J.: Rate constants and products of the reactions of e^-_{aq}, O_2^- and H with ozone in aqueous solutions, J. Phys. Chem., 87, 1951–1954, 1983.

Sehested, K., Holcman, J., Bjergbakke, E., and Hart, E. J.: A pulse radiolytic study of the reaction OH + O_3 in aqueous medium, J. Phys. Chem., 88, 4144–4147, 1984.

Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics, John Wiley & Sons, New York, Chichester, Weinheim, 1998.

Shoute, L. C. T., Alfassi, Z. B., Neta, P., and Huie, R. E.: Temperature dependence of the rate constants for reaction of dihalide and azide radicals with inorganic reductants, J. Phys. Chem., 95, 3238 – 3242, 1991.

Takami, A., Kato, S., Shimono, A., and Koda, S.: Uptake coefficient of OH radical on aqueous surface, Chem. Phys., 231, 215–227, 1998.

THALOZ: Final report of the EU project THALOZ: Tropospheric Halogens - effect on ozone, coordinated by R. A. Cox, University of Cambridge, U.K., 2005.

Thomas, K., Volz-Thomas, A., and Kley, D.: Zur Wechselwirkung von NO_3-Radikalen mit wässrigen Lösungen: Bestimmung des Henry- und des Massenakkomodationskoeffizienten, Ph.D. thesis, Institut für Chemie und Dynamik der Geosphäre 2, Forschungszentrum Jülich GmbH, FRG, 1993.

Troy, R. C. and Margerum, D. W.: Non-metal redox kinetics: Hypobromite and hypobromous acid reactions with iodide and with sulfite and the hydrolysis of bromosulfate, Inorg. Chem., 30, 3538 – 3543, 1991.

Tsalkani, N., Mellouki, A., Poulet, G., Toupance, G., and Bras, G. L.: Rate constant measurement for the reactions of OH and Cl with peroxyacetyl nitrate at 298 K, J. Atmos. Chem., 7, 409–419, 1988.

Urbanski, S., Stickel, R. E., Zhao, Z. Z., and Wine, P. H.: Mechanistic and kinetic study of formaldehyde production in the atmospheric oxidation of dimethyl sulfide, J. Chem. Soc. Faraday Trans., 93, 2813 – 2819, 1997.

Urbanski, S., Stickel, R. E., and Wine, P. H.: Mechanistic and kinetic study of the gas-phase reaction of hydroxyl radical with dimethyl sulfoxide, J. Phys. Chem. A, 102, 10 522 – 10 529, 1998.

van Dingenen, R., Jensen, N. R., Hjorth, J., and Raes, F.: Peroxynitrate Formation During the Night-Time Oxidation of Dimethylsulfide: Its Role as a Reservoir Species for Aerosol Formation, J. Atmos. Chem., 18, 211 – 237, 1994.

Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, Nature, 383, 327 – 330, 1996.

von Glasow, R.: Modeling the gas and aqueous phase chemistry of the marine boundary layer, Ph.D. thesis, Universität Mainz, Germany, http://www.rolandvonglasow.de, 2000.

von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 1. Cloud-free MBL, J. Geophys. Res., 107, 4341, doi: 10.1029/2001JD000 942, 2002.

von Gunten, U. and Oliveras, Y.: Advanced oxidation of bromide-containing waters: Bromate formation mechanisms, Environ. Sci. Technol., 32, 63 – 70, 1998.

Wagner, I. and Strehlow, H.: On the flash photolysis of bromide ions in aqueous solution, Ber. Bunsenges. Phys. Chem., 91, 1317 – 1321, 1987.

Wallington, T. J., Andino, J. M., Ball, J. C., and Japar, S. M.: Fourier transform infrared studies of the reaction of Cl atoms with PAN, PPN, CH_3OOH, HCOOH, CH_3COCH_3 and $CH_3COC_2H_5$ at 295±2 K, J. Atmos. Chem., 10, 301 – 313, 1990.

Wang, T. X., Kelley, M. D., Cooper, J. N., Beckwith, R. C., and Margerum, D. W.: Equilibrium, kinetic, and UV-spectral characteristics of aqueous bromine chloride, bromine, and chlorine species, Inorg. Chem., 33, 5872 – 5878, 1994.

Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide and nitrogen dioxide in sunlit continental fair weather clouds, Phys. Chem. Chem. Phys., 1, 5471 – 5483, 1999.

Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, Atmos. Environ., 25A, 1–203, 1991.

Weast, R. C., ed.: CRC Handbook of Chemistry and Physics, 61st Edition, CRC Press, Inc., Boca Raton, FL, 1980.

Weinstein-Lloyd, J. and Schwartz, S. E.: Low-intensity radiolysis study of free-radical reactions in cloudwater: H_2O_2 production and destruction, Environ. Sci. Technol., 25, 791–800, 1991.

Wilhelm, E., Battino, R., and Wilcock, R. J.: Low-pressure solubility of gases in liquid water, Chem. Rev., 77, 219–262, 1977.

Wine, P. H., Wells, J. R., and Nicovich, J. M.: Kinetics of the reactions of F^2P and Cl^2P with HNO_3, J. Chem. Phys., 92, 2223–2228, 1988.

Wine, P. H., Tang, Y., Thorn, R. P., Wells, J. R., and Davis, D. D.: Kinetics of aqueous phase reactions of the SO_4^- radical with potential importance in cloud chemistry, J. Geophys. Res., 94D, 1085–1094, 1989.

Worsnop, D. R., Zahniser, M. S., Kolb, C. E., Gardner, J. A., Watson, L. R., van Doren, J. M., Jayne, J. T., and Davidovits, P.: The temperature dependence of mass accommodation of SO_2 and H_2O_2 on aqueous surfaces, J. Phys. Chem., 93, 1159–1172, 1989.

Wu, D., Wong, D., and Di Bartolo, B.: Evolution of Cl_2^- in aqueous NaCl solutions, J. Photochem., 14, 303–310, 1980.

Yin, F., Grosjean, D., and Seinfeld, J. H.: Photooxidation of Dimethyl Sulfide and Dimethyl Disulfide. I: Mechanism Development, J. Atmos. Chem., 11, 309 – 364, 1990.

Yu, X.-Y.: Kinetics of free radical reactions generated by laser flash photolysis of $OH + Cl^-$ and $SO_4^- + Cl^-$ in the aqueous phase – Chemical mechanism, kinetics data and their implications, Ph.D. thesis, University of Michigan, Ann Arbor, 2001.

Zehavi, D. and Rabani, J.: Oxidation of aqueous bromide ions by hydroxyl radicals. Pulse radiolytic investigation, J. Phys. Chem., 76, 312 – 319, 1972.

Zellner, R., Exner, M., and Herrmann, H.: Absolute OH Quantum Yield in the Laser Photolysis of Nitrate, Nitrite and Dissolved H_2O_2 at 308 and 351 nm in the Temperature Range 278-353 K, J. Atmos. Chem., 10, 411 – 425, 1990.