

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

Regional modelling of tracer transport by tropical convection – Part 1: Sensitivity to convection parameterization

J. Arteta¹, V. Marécal¹, and E. D. Rivière²

¹Laboratoire de Physique et Chimie de l'Environnement, CNRS and Université d'Orléans, 3A avenue de la recherche scientifique, 45071 Orléans cedex 2, France

²Groupe de Spectroscopie Moléculaire et Atmosphérique, Université de Reims, Moulin de la Housse, B.P. 1039, 51687 Reims Cedex, France

Received: 29 December 2008 – Accepted: 27 January 2009 – Published: 4 March 2009

Correspondence to: J. Arteta (Joaquim.Arteta@cnr-orleans.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The general objective of this series of papers is to evaluate long duration limited area simulations with idealised tracers as a tool to assess tracer transport in chemistry-transport models (CTMs). In this first paper, we analyse the results of six simulations using different convection parameterizations. The simulations are using the Grell and Dévényi (2002) mass-flux framework for the convection parameterization with different closures (Grell=GR, Arakawa-Shubert=AS, Kain-Fritsch=KF, Low omega=LO, Moisture convergence=MC) and an ensemble parameterization (EN) based on the other five closures. The simulations are run for one month during the SCOUT-O3 field campaign lead from Darwin (Australia) and have a 60 km horizontal resolution and a fine vertical resolution in the upper troposphere/lower stratosphere. Meteorological results are compared with satellite products, radiosoundings and SCOUT-O3 aircraft campaign data. They show that the model is generally in good agreement with the measurements with less variability in the model. Except for the precipitation field the differences between the six simulations are small with respect to the differences with the meteorological observations. The comparison with TRMM rainrates shows that the six simulations have two different behaviours with the EN, AS and KF parameterizations (Group 1) providing better rain fields than LO, MC and GR (Group 2). The vertical distribution of tropospheric tracers is very different for the two groups showing significantly more transport into the TTL for Group 1 related to the larger average values of the upward velocities. Nevertheless the low values for the Group 1 fluxes at and above the cold point level indicate that the model does not simulate significant overshooting. For stratospheric tracers, the differences between the two groups of parameterizations are small indicating that the downward transport from the stratosphere is more related to the turbulent mixing parameterization than to the convection parameterization.

ACPD

9, 5889–5928, 2009

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

It has long been recognized that air mainly enters the lower stratosphere in the tropics from where it is then distributed at the global scale through the Brewer-Dobson circulation. Although many studies of the troposphere-to-stratosphere transport (TST) have already been published (e.g. reviews by Holton et al., 1995 and Stohl et al., 2003 or e.g. recent work by Ricaud et al., 2007 and Duncan et al., 2007), the detailed processes leading to TST and their quantification are still debated. The Tropical Tropopause Layer (Sherwood and Dessler, 2000), called TTL hereafter, can be defined as the transitional layer between air with typical tropospheric characteristics and air with typical stratospheric characteristics. The TTL is therefore a key layer for TST studies since air masses reaching a height above the zero radiative heating level within the TTL will slowly rise into the lower stratosphere while horizontally advected (Folkins et al., 1999; Sherwood and Dessler, 2001; Fueglistaler et al., 2004). In practice, several definitions of the TTL have been proposed in the literature (Highwood and Hoskins, 1998; Folkins et al., 1999; Gettelman and Forster, 2002; Fueglistaler 2008). In the present paper, we use the recent definition proposed by Fueglistaler (2008). The TTL bottom is set above the top of the main cumulus outflow layer ($z \approx 14 \text{ km} - \theta \approx 355 \text{ K}$). Above this level air is radiatively heated under all sky conditions. The top of the TTL is at $z \approx 18.5 \text{ km}$ ($\theta \approx 425 \text{ K}$) where the most energetic and intense cumulonimbus can reach (overshooting convection). The chemical composition of the TTL is closely linked to tropical convection which can transport vertically and rapidly the lower tropospheric emissions into the TTL altitude range (e.g. Wang et al., 1995; Pickering et al., 1996; Marécal et al., 2006). Convective transport may also have an impact on Stratosphere to Troposphere Transport (STT) from convection induced downdrafts (e.g. Baray et al., 1999; Leclair de Bellevue et al., 2006) and breaking of convectively driven gravity waves (e.g. Rivière et al., 2006).

To study the transport of tracers from the boundary layer into the stratosphere via the TTL, the local convection as well as the large scale advection and the radiative

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

transport processes have to be taken into account. The large scale processes are generally well handled by global chemistry transport models (CTMs) which are forced by dynamical fields from state-of-art weather forecast models. In most current CTMs the subgrid-scale convection is parameterized and the associated tracer transport is taken into account in a consistent manner. Convection is known to be one of the major sources of uncertainty in CTMs through the uncertainty on the convective parameterizations themselves and through the fact that they are applied on off-line dynamical fields. To study TST in the tropics using a CTM it is therefore required to assess the quality of the tracer transport by its convective parameterization. One possibility is to compare with measurements gathered in the TTL or with validated cloud resolving model simulations of observed tropical convection case studies. But the number of case studies available from field campaign or from cloud scale simulations is too small to allow a general evaluation of CTMs. The alternative approach proposed here is to use long duration (~one month) regional (typically 6000 km×4000 km) simulations with a limited-area model using finer vertical (a few hundred meters in the TTL) and horizontal (~20–100 km) resolutions than typical CTM resolutions ($\geq 1^\circ$). Such simulations aim at bridging the gap between the small spatial and temporal scales associated with convection and the CTM global and long time scales. On one hand, the comparison of the regional simulation results with campaign data or cloud scale simulations of convective systems is meaningful thanks to the resolution chosen in the regional run. On the other hand, statistical comparisons with global CTM results are possible since the regional simulations are long enough and over a domain sufficiently large. In this context, the objective of this series of two papers is to evaluate long-duration regional simulations with a limited-area model as a tool to produce realistic tracer transport by tropical convection that could then be used for the assessment of CTMs.

In the framework of tracer transport, several comparative studies of convective parameterizations have been published with different types of models. Using the convective parameterizations proposed by Hack (1994) and Zhang and McFarlane (1995) in a global climate model, Gilliland and Hartley (1998) concluded that the two con-

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

vective schemes have significantly different effects on the tropical circulation and the subsequent interhemispheric tracer transport. Zhang et al. (2008) conducted recently a comparative study on tracer transport of ^{222}Rn in a global climate model. They found large differences in the vertical distribution of the tracer between the cumulus parameterizations from Tiedke (1989) modified by Nordeng (1994) and from Zhang and McFarlane (1995) combined with Hack (1994). Lawrence and Rasch (2005) compared convective mass fluxes based on the plume ensemble formulation (e.g. Arakawa and Schubert, 1974; Grell, 1993) and on the bulk formulation (e.g. Tiedke, 1989; Zhang and McFarlane, 1995) in the MATCH CTM. They showed that the bulk formulation is an adequate approximation for most tracers with lifetimes of a week or longer but not efficient enough for the tracer transport of short-lived species. Folkins et al. (2006) tested four cumulus parameterizations implanted in different global forecast models. The intercomparison was inconclusive since the differences between the model results could be related not only to the convection parameterizations but also to other differences in the model setups. Simulations with the NCAR/MM5 limited area model of a tropical convective system were performed by Wang et al. (1996) who found similar average transport profiles using the Kain and Fritsch (1993) or the Grell (1993) convective schemes. All these studies show that the choice of the convection parameterization is important for the tracer transport in models. This issue is the subject of the present paper (Part 1) that is devoted to the study of the sensitivity of the regional modelling approach to the subgrid scale deep convection parameterization. The second paper (Part 2) of this series of papers is focused on the sensitivity to the model vertical and horizontal resolutions that are known to have a significant effect on the convective tracer transport (e.g. Deng et al., 2004, Wild and Prather 2006).

The present work makes use of the operational limited area CATT-BRAMS (Coupled Aerosol Tracer Transport model to the Brazilian Regional Atmospheric Modeling System) model (Freitas et al., 2007). It is based on the Brazilian version of the RAMS model, tailored to the tropics. The BRAMS includes a deep cumulus parameterization based on the mass-flux approach proposed by Grell and Dévényi (2002) with several

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

possible closures. The CATT-BRAMS has an on-line tracer transport model fully consistent with the simulated atmospheric dynamics including transport by convection. The simulated area is in the Maritime continent known to be a very active region of convection. The simulation period chosen is from mid-November 2005 to mid-December 2005 and corresponds to the SCOUT-O3 field campaign period (Vaughan et al., 2008). During this campaign, convection was very intense and evidence of overshooting events has been identified (Corti et al., 2008). The meteorological data from this experiment are used to validate the model transport by convection, as well as satellite-derived products and radiosoundings. Simulation experiments were run with idealized tracers. This type of tracer cannot be compared to measurements for evaluation but they are useful for understanding the dynamical processes linked to tropical convection driving the tracer spatial distributions. Moreover simulation of real tracers is difficult to analyse due to variations in the intensity, location and time of the emissions and in the background distribution.

In the present paper, the CATT-BRAMS model and the setup of the simulation experiments are described in Sect. 2. The model evaluation of the meteorological fields is presented in Sect. 3. Section 4 is devoted to the analysis and discussion of the model results for the tracers. Concluding remarks are given in Sect. 5.

2 Numerical model

2.1 Model Description

The CATT-BRAMS model (Freitas et al., 2007) used in the present study is an on-line transport model fully consistent with the simulated atmospheric dynamics. The atmospheric model BRAMS (Brazilian RAMS, <http://brams.cptec.inpe.br/>) is based on the Regional Atmospheric Modeling System (RAMS, Cotton et al., 2003). It is tailored to the tropics with several improvements such as cumulus convection parameterization, the soil moisture initialization and the surface scheme.

CATT is a numerical system designed to simulate and to study the transport processes associated with the emission of tracers. This is an Eulerian transport model coupled to BRAMS. The tracer transport is run simultaneously (“on-line”) with the atmospheric state evolution using the same time-step. It is consistent with the BRAMS dynamical and physical parameterizations. The tracer mass mixing ratio, which is a prognostic variable, includes the effects of sub-grid scale turbulence in the planetary boundary layer, convective transport by shallow and deep moist convection in addition to the grid scale advection transport.

2.2 General set-up of the simulations

The series of simulations discussed in the present paper has the same set-up except for the deep-convection parameterizations used. The simulations include one grid covering a domain ranging from 100° E to 160° E and from 20° N to 20° S. Its horizontal grid spacing is 60 km. The geography of the domain and the associated model topography are illustrated in Fig. 1. It includes 56 vertical levels, with a high resolution (300 m depth) between 14.5 km and 19 km, in order to accurately model the upper troposphere and lower stratosphere (UTLS) region. The simulation lasts 30 days from the 15th November 2005 to the 15th December 2005. All radiative calculations were done with the Harrington (1997) scheme. We use the one-moment bulk microphysics parameterization which includes cloud water, rain, pristine ice, snow, aggregates, graupel and hail (Walko et al., 1995). Shallow convection is parameterized as described in Grell and Devenyi (2002). The parameterizations used for deep convection are presented in Sect. 2.3. Initialization fields and domain boundaries came from ECMWF analysis fields and model is forced every 6 h.

The transport of tracers is activated in all the simulations. We chose a set of four idealized tracers to characterize the different pathways of exchange between the troposphere and the stratosphere (see Table 1). The first one is a short-lived tracer designed to focus only on the effect of individual convective events. Its lifetime of 6 h is long enough to be transported by convection but not significantly by large scale advective

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion and diffusion. It is emitted only above land with an arbitrary constant emission rate of $10^{-9} \text{ kg m}^{-2} \text{ s}^{-1}$. It is initialized to 0. The second tracer has the same initial condition and same emission rate but an infinite lifetime in order to analyse TST at the regional scale. Finally, we used two stratospheric tracers to study the effect of convection on Stratosphere to Troposphere Transport (STT). The first one is initialized with a constant mixing ratio of 1 ppt for potential temperatures greater than 380 K (which corresponds approximately to the tropopause level in the tropics and is well into the TTL) and 0 below. Its lifetime is infinite for potential temperatures greater than 380 K and 6 hours below 380 K. The second has the same setup but its lifetime is infinite over the whole atmospheric column.

2.3 Convection parameterizations

In the present paper, we test six convective parameterizations and we analyse their impact on the troposphere-stratosphere transport (TST and STT) of tracers. Convective parameterization schemes are procedures that attempt to account for the collective effect of sub-grid scale convective processes on large-scale model variables. These effects (latent heating, evaporative cooling, generation of cirrus clouds associated with the anvil, etc.) have to be determined from the available model variables. Different cumulus parameterizations were developed during the last decades in order to improve model results in convective areas. The mass-flux approach is generally used in mesoscale models. It attempts to explicitly account for convective processes at each grid point by combining a cloud model with the assumption that convection acts to restore the stratified grid column based on moist parcel stability. The cloud model estimates the properties of the convection and the closure assumption specifies the amount of convection that occurs in order to achieve the desired rate of stabilization.

The parameterizations used are based on the formulation proposed by Grell (1993) and Grell et al. (1994) and modified by Grell and Dévényi (2002) to allow the use of five different closure assumptions: Grell (called GR hereafter) (Grell, 1991), Arakawa Schubert (AS) (Arakawa and Schubert, 1974), Kain-Fritsch (KF) (Kain and Fritsch, 1992),

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



moisture convergence (MC) (Kuo, 1974; Krishnamuti et al., 1983), and Low-Omega (LO) (Frank and Cohen, 1987). The additional convection parameterization used is based on an ensemble approach (EN) (Grell and Dévényi, 2002). The Grell's framework is based on a simple conceptual model; namely, the cloud consists of two steady state circulations caused by an updraft and a downdraft. There is no direct mixing between cloud air and environmental air except at the top and the bottom of the circulation.

In this framework the AS closure uses the quasi-equilibrium assumption which states that the stabilisation of the atmosphere by convection is in quasi-equilibrium with the destabilization by the large scale processes. The GR parameterization is a modified AS closure including moist convective-scale downdrafts. The KF parameterization also uses the stability closure but without any dependence with large scale motion leading to a pure instantaneous stability closure. It assumes that a cloud can rise and then can instantly decay. Thus after subsidence calculations, the convection is supposed to build and decay without a steady-state stage, and the cloud properties are mixed horizontally with the subsided environment. The MC closure assumes that the convective activity is closely related to the total moisture convergence at the base of clouds. LO use the same idea of MC, but introduce a downdraft forcing. This downdraft will cause additional mass-flux convergence, creating subsequent forcing of more convection. EN provides the most probable solution based on statistical methods (Stephenson and Doblas-Reyes 2000) applied to a set of sensitivity calculations using perturbed values in GR, AS, KF, MC and LO parameters. The six simulations run using these parameterizations will be referred hereafter as GR, AS, KF, MC, LO and EN experiments.

3 Evaluation of the model meteorological fields

In the present study we cannot validate the idealised tracers directly using tracer measurement. Rather we evaluate the atmospheric dynamics by comparing the meteorological fields provided by the six simulations against observations. For this purpose we

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

used satellite rainfall estimates from TRMM (Tropical rainfall Measuring Mission), radiosoundings and SCOUT-O3 aircraft measurements. In the Maritime continent area very few radiosoundings from the operational network provide reliable data. Therefore we only used the radiosoundings launched in the frame of the SCOUT-O3 campaign from Darwin in Australia (see Fig. 1) and those launched from Manus (see Fig. 1), a station operating in the frame of the ARM project (Atmospheric Radiation Measurement, <http://www.arm.gov/>). The comparison with the SCOUT-O3 aircraft measurements allows us to make a detailed analysis of the model behaviour on a case study.

3.1 Comparison with TRMM surface rainfall estimates

We compared the surface accumulated rainfall obtained with the six convective schemes to those estimated by TRMM. The dataset used is 3-hourly and $0.25^\circ \times 0.25^\circ$ resolution and was produced by the 3B42 algorithm (Huffman et al., 2007, <http://trmm.gsfc.nasa.gov>). Figure 2a shows the daily mean surface rain rates (in mm day^{-1}) estimated by TRMM during the one-month simulation period.

Almost all the domain experienced significant precipitation (over 0.1 mm day^{-1}) except in three areas located South of 15° S and North of 15° N . We can also identify four major areas with high precipitation rates (above 10 mm day^{-1}) located

- over the New Guinea island (around 140° E ; 5° S), with values reaching 10 to 20 mm day^{-1} in the Southern part of the island.
- from the Eastern coast of the Malaysian peninsula (around 100° E ; 10° N) to the Eastern coast of Thailand (110° E ; 12° N), with very high values above 20 mm day^{-1}
- over all the Indonesian Islands (from 100° E to 115° E ; 0° S to 30° S)
- on the Eastern coast of Philippines (from 110° E to 115° E ; 10° N to 15° N) with values reaching up to 20 mm day^{-1} .

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The TRMM 3B42 product is based on different satellite measurements mainly from passive remote sensing instruments leading to uncertainties on the surface rainrate estimates mainly over land and for very low rainrates. To assess the quality of this product for the chosen area and time period it was compared to the rainrate estimates provided by the Global Precipitation Climatology Project “One-Degree Daily Precipitation DataSet” product (GPCP, Huffman et al., 2001). The TRMM 3B42 product showed a very good agreement for both the precipitation location and the intensity (not shown) giving confidence in the TRMM estimates used here.

The daily mean surface rainrates provided by the 6 experiments are shown in Fig. 2b to g. The six model experiments well locate the areas of high and low precipitation rates compared with the TRMM-based observations. The very convective areas are well simulated by all six model simulations. In other areas TRMM shows more precipitation than all the simulations especially in the centre and in the South over ocean. This could be partly related to a large uncertainty on light precipitation in 3B42 that possibly leads to an overestimation of surface precipitation. But it is also likely due to an uncertainty in light precipitation in the model for all convective parameterizations. Most of the precipitation simulated with GR is convective. There is almost no stratiform precipitation. The EN simulation provides stratiform precipitation but underestimate it.

Comparing the six model experiments we can class the convective closures into two groups providing similar results: AS, KF and EN in Group 1, and GR, MC and LO in Group 2. Group 1 provides results generally closer to TRMM measurements than Group 2. Group 1 simulates more extended areas of light precipitation in better agreement with TRMM. For high rainrates Group 1 is closer to the observations over Malaysia and Indonesian Islands around 20 mm day^{-1} . On the other hand, the lower rates obtained with Group 2 are in better agreement with the measurements over New Guinea where TRMM estimates give values around 10 mm day^{-1} for the South of the island and 5 mm day^{-1} for the North. Group 1 tends to slightly overestimate the values from 1 to 5 mm day^{-1} in this area. Group 1 simulations are also able to capture some of the convection occurring in the North of Australia around Darwin which is missed

by Group 2. Note that the EN parameterization provides the most probable convection characteristics from all the other five closures calculated by varying the main parameters of each closure. This comparison with the TRMM rainrates shows that the EN approach is able to provide as expected the most realistic surface rainrate among the 5 closures by giving more weight to the AS and KF closures.

3.2 Comparison with radiosounding data

Comparisons were done with the radiosounding launched from Darwin during the field campaign of the SCOUT-O3 project and from Manus Island in the North of New-Guinea in the frame of the ARM program. Results provided similar conclusions and are only illustrated here on the 12-hourly Manus (147° E; 2° S) radiosoundings. Note that Manus location is interesting since this island is located in an area where strong convective events are frequent as shown by the TRMM daily-mean rainrate estimates which are above 5 mm day⁻¹ (see Fig. 2a).

Figure 3 shows the mean values/standard deviations and the mean biases between Manus radiosounding data and the six simulations for temperature, wind direction, wind speed and specific humidity. The model results for the 6 experiments are generally close and in good agreement with the radiosonde data for temperature and winds for both the mean and standard deviation. Tropospheric temperatures show a mean warm bias of 1.5°C. A larger positive bias is found around the cold point tropopause reaching 3°C in the model simulations. The very low tropopause temperatures observed in the Western Pacific are related to the intense convective activity in this area characterized by high-reaching cumulo-nimbus. Figures 3a and b show that the model is not able to cool enough around the cold point tropopause in the Manus area. This can be partly related to an underestimation of the convective activity in the model at Manus location (see Fig. 2) and partly to the model horizontal resolution (60 km) that cannot simulate the small scale impact of convection on temperature. The differences between the six simulations are negligible for most altitudes except in the 8–11.5 km range where the Group 1 results are slightly better compared to observations by about 0.3 K.

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Both the simulated wind speed (fig 3c) and direction (Fig. 3e) are in good agreement with the measurements except above 23 km where the strong stratospheric winds are underestimated by the model. The mean bias for the wind speed is around 1 m s^{-1} below 16 km. In the 16–19.5 km range the model shows variations in the easterly wind intensity similar to the measurements but much less pronounced with a bias ranging between -5 to $+4 \text{ m s}^{-1}$. The wind direction presents a mean bias less than 20° over the entire atmospheric column. The radiosoundings data show that the tropopause region over Manus is marked by very strong gradients with a high vertical variability in both the dynamic and the thermodynamical fields. These gradients are produced by all six model simulations thanks to the fine vertical resolution used but smoothed. When comparing model simulations to radiosounding data one has to keep in mind the representativeness issue. Part of the observed gradients and variability may be linked to local effects (Manus Island being ~ 100 km long and ~ 30 km wide) that cannot be captured by the model that uses a 60 km horizontal resolution. The comparison between the six simulations shows differences on the wind speed and direction. For the wind speed (Fig. 3d) they are significant between 10 and 17 km altitude with a reduced bias for the Group 2 up to 0.5 m s^{-1} . For the wind direction there are differences at nearly all the levels with a 8° average mean bias. As for the temperature and the wind speed, the six simulations can be sorted into the same two groups as defined in Sect. 3.1. Depending on the altitude range Group 1 is either better or worse than Group 2 compared to the measurements.

The results for the specific humidity are displayed in Fig. 3g and h. Note that the humidity measurements in the upper troposphere and lower stratosphere should not be considered since they are expected to be not very reliable at very low temperatures and water vapor contents. In Fig. 3g the six model simulations overestimate the specific humidity above 4 km altitude. This can only partly be related to the known remaining small dry bias of the Vaisala RS92 data (Balloon-Borne Sounding System Handbook, http://www.arm.gov/publications/tech_reports/handbooks/sonde_handbook.pdf). This shows that the model, whatever convection parameterization is used, does not

convert enough tropospheric moisture into precipitation leading an overestimation of the water vapour mixing. This is consistent with the model underestimation of the low rainrates linked to a too low production of stratiform precipitation by the model (see Sect. 3.1 and Fig. 2). The differences between the six simulations are small except around 3 to 4 km altitude.

3.3 Comparisons with meteorological data from Falcon and Geophysica flights

During the simulation period several DLR-Falcon and Geophysica (M55) flights (9 for each aircraft) were done around Darwin (Australia), in the framework of the SCOUT-O3 field campaign (Vaughan et al., 2008; Brunner et al., 2008). Most of the flights were done around the Hector convective events regularly occurring over the Tiwi Islands. Some of them were extended flights planned for study of the surrounding regions: survey flights on the 23rd and the 29th November, remote sensing flight on the 5th December. Since the model simulations cover a large area, a comparison with the extended and transit flights were preferred for the model evaluation. A systematic comparison with all these flights was done and similar results were obtained. To illustrate the results we have chosen the flight that took place on the 23rd November. On this date the Geophysica aircraft and the Falcon performed coordinated flights whose objective was the detailed probing of the TTL over the Arafura Sea (see Fig. 1). Both Geophysica and Falcon flew long north-east oriented legs perpendicular to the mean flow expected to be north-westerly in the TTL. Flying back and forth along the same line twice, the Geophysica sampled around the cold point tropopause at four different levels, one significantly below the cold point level at ~ 15.6 km (leg 1), two close to the cold point tropopause at ~ 17.5 km (leg 3) and ~ 16.4 km (leg 4), and one level well above at ~ 18.3 km (leg 2). The flight paths are displayed in Fig. 14 in Brunner et al. (2008).

Figures 4 and 5 show the comparison between the results of the six simulations and the measurements collected respectively by the Geophysica and the Falcon instruments for temperature, wind speed and direction, and specific humidity. The modeled

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



air temperature is in good agreement with the measurements during the Geophysica ascent, leg 1 and the descent and to a lesser extend leg 2. For the two aircraft legs performed around the cold point tropopause level there is an overestimation of the temperature by the model of about 5°C. The comparison with the Falcon temperature measurements shows that modeled temperatures around 12 km altitude are about 2°C degrees lower than the aircraft measurements. This is consistent with the radiosounding comparison showing that the model provides too warm temperatures around the cold point tropopause and too cold temperatures in the troposphere up to 14 km. There are no significant differences between the six simulations for the temperature for both Geophysica and Falcon flights.

The horizontal wind speed and wind direction simulated by the six runs along the aircraft trajectories are generally in good agreement with both the Geophysica (Fig. 4b and c) and the Falcon (Fig. 5b and c) measurements. In the cold point tropopause region (around 17 m altitude) the variations of the wind velocity are well captured by the model compared to the Geophysica measurements but are underestimated. This is consistent with the Manus radiosounding showing a strong increase of the wind speed near the tropopause that is underestimated by the model. The six model runs give very close results for the wind speed and the wind direction with differences of 2 m s⁻¹ and 6 degrees at maximum which are much smaller than the differences between each model simulation and the aircraft observations.

The specific humidity measurements aboard the Falcon aircraft are well simulated by the six model runs with a slight model overestimation for the aircraft leg at 12 km altitude (see Fig. 5d). This overestimation is much lower than that obtained in the comparison with the Manus radiosondes indicating that the Manus sondes likely underwent a significant dry bias in the upper troposphere. The modeled values are also in a fairly good agreement with the Geophysica measurements (Fig. 4d) but with an overestimation of the specific humidity for all legs except leg 2 which was performed significantly above the cold point tropopause. This illustrates the fact that convection parameterizations in all the simulations are not able to modify significantly the specific humidity

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

above the cold point tropopause from its initial smooth water vapour state provided by the ECMWF analysis. Both the Falcon and the Geophysica measurements exhibit strong peaks over short time periods (e.g. around 16 000 and 18 000 s in the Geophysica flight. CO measurements were gathered on board the Geophysica aircraft during the flight. A comparison between the CO and specific humidity data shows that the humidity peaks are correlated with CO peaks indicating their link with deep convection events. The six simulations use a coarse horizontal resolution and therefore do not capture these peaks that are very localized in space and time.

Similarly to Manus radiosonde comparison the model shows a good consistency with the aircraft observations but a lack of variability in the horizontal that can be attributed to the model horizontal resolution. The results of the six simulations provide close results for all the atmospheric variables. When and where differences are found between the model simulations they can be classified into the same two groups as those defined in the TRMM comparison.

3.4 Conclusion and discussion on the meteorological comparison

The simulations give results that are generally consistent with the radiosounding and aircraft meteorological data and exhibit small differences between the 6 convection parameterizations mainly in the upper troposphere and in the TTL. From these differences it is not possible to get guidance on which convection parameterization performs better. But for the surface rainrates the EN, AS, KF (Group 1) closures give significantly better results than LO, MC and GR (Group 2) with a lower underestimation in light precipitation areas. This indicates that the AS and KF closures and the EN closure, the latter seeking the most probable convective characteristics, perform better for the meteorological variables than the other three closures. The fine variations observed in the aircraft data cannot be captured by the model with the chosen horizontal resolution.

In this study we use the Grell's simple mass flux framework with different closure assumptions based on mass-flux parameterizations commonly used in mesoscale models. The five closure assumptions (GR, AS, KF, LO, MC) drive the modulation of con-

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

vection by environment (noted dynamic control in Grell 1993). The Ensemble (EN) also takes into account statistically a variability of the modulation of the environment by the convection and of the cloud model. The similarity of the EN to the AS and KF simulations means that this is the dynamic control dominates the parameterization behaviour in the EN simulation. The assumptions used in the 5 closures have their more important effect on convection triggering and associated rainfall rate. The effect on the temperature, horizontal wind and specific humidity is small on average since convection parameterization is not triggered at all grid points and for each timesteps. In convectively active areas such as Manus Island, there are differences in the meteorological parameters in the upper troposphere and in the TTL corresponding to the top of the deep convection circulation in the Grell's framework where there is direct mixing between cloud air and environment air.

4 Analysis of the tracer transport

Similarly to the results discussed in section 3, the tracer transport with the 6 different convection parameterizations exhibits two types of behaviour. Therefore only the EN and GR simulations will be shown and discussed hereafter since they illustrate the Group 1 and Group 2 results, respectively.

4.1 Tropospheric tracers

Figure 6 shows the tracer mixing ratio profiles averaged over the model domain and over the one month simulation period using 3-hourly model outputs for Tracer 1 (6 h lifetime) and Tracer 2 (infinite lifetime). To interpret these profiles we have displayed the mean temperature and potential temperature profiles in Fig. 7. Note that in Fig. 7 only the profiles for the EN simulations are plotted since the GR results are very close to EN. The mean cold point tropopause (-82°C) and the mean 380 K level are close and located at 17.3 km and 17.1 km altitude respectively. To locate the TTL we use

here the definition proposed by Fueglistaler et al. (2008): TTL top is at the 70 hPa level (425 K) and TTL bottom is located above the levels of main convective outflow at the zero radiative level under all sky conditions (~ 150 hPa, 355 K). This gives for the model simulations the TTL top at 18.9 km altitude and the TTL bottom at 14 km.

In Fig. 6 the shape of the mean mixing ratio profiles for both tracers is typical of convective areas, with large values in the low troposphere, decreasing in the mid-troposphere and increasing in the upper troposphere with a maximum value reached around 15 km altitude. Above there is a rapid decrease reaching very low values around 18 km. The EN parameterization provides for both tracers lower mean mixing ratios in the lower and mid troposphere and larger above ~ 10 km than GR with a ratio of ~ 2 for the maximum values in the upper tropospheric (around 15 km altitude). The GR parameterization gives more frequent convective outflows below 10 km than EN and significantly less transport above. Having a 6-hour lifetime Tracer 1 shows the local effect of convection and is only weakly affected by the model diffusion. This means that the convective EN parameterization is able to drive significant amount of surface tracers into the TTL with a ratio between the TTL bottom and the surface values of 2.2%. The GR parameterization is much less efficient with a ratio of 0.8% and with at least a factor of three less above the cold point tropopause. The Tracer 2 mean profile shows a maximum around 15 km as for Tracer 1 meaning that the tracers lifted by convection in the TTL are not largely transported down during the following days while they travel into the model domain whatever parameterization is used.

Figure 8 displays for EN and GR simulations the meridian mean over the one month period for both tracers and the corresponding vertical velocity. The tracer with a 6 h lifetime (Tracer 1) indicates where the convective transport occurs since it is rapidly removed after uplifting due to its lifetime. In the EN and GR simulations the vertical transport by convection of Tracer 1 occurs at the same longitudes, mainly around 112° E and 143° E. They correspond to emission areas (only islands in the simulation setup) with high vertical velocities where an intense convective activity is modelled as well as observed, mainly Borneo (around 112° E) and New Guinea (around 143° E). For

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tracer 2 the maxima in the TTL for both EN and GR simulations are shifted westward compared to Tracer 1 and located in the 100° E–120° E longitude range. This indicates that a significant part of Tracer 2 mixing ratio that is firstly transported vertically from the low levels to the TTL around 143° E (New Guinea) is then horizontally advected to western longitudes where it adds to the high TTL mixing ratios lifted by local convection (mainly Borneo) and spread in latitudes (Fig. 9a, b). The comparison between Fig. 9a, b with Fig. 9c, d also shows that the tracer meridian distribution in the mid troposphere is also different between Tracer 1 and 2. Therefore the geographical distribution of a long lifetime tracer above the boundary layer depends on both the locations where the main convection occurs and the large scale dynamics that transports horizontally as well as mixes the tracer with its environment.

The major effect of the closure assumptions is on the vertical distribution since they drive the convective updraft and downdraft characteristics in each closure. Grell's formulation gives enhanced tracer amounts at the top of convective outflow than in the mid-troposphere and is on average lower in GR than in EN. To quantify the transport the mean fluxes for tracers 1 and 2 are calculated from vertical wind speeds at several altitudes in the upper troposphere: below the TTL at the level of frequent convective outflow (12 km), at the TTL bottom level (14 km), at the cold point level (17.3 km), at the top of the TTL (18.9 km) and above the TTL (19.5 km). Results are reported in Table 2. The mean surface flux for the two tropospheric tracers that are only emitted over land is $0.182 \times 10^{-9} \text{ kg m}^{-2} \text{ s}^{-1}$. For the EN simulation of Tracer 1 all the fluxes values are positive indicating that there is a mean transport directed upward even up to 19.5 km. At 12 km altitude and at the TTL bottom level, they are similar and around $4 \times 10^{-12} \text{ g m}^{-2} \text{ s}^{-1}$, representing around 2.2% of the emission flux above land. This flux is decreasing rapidly with altitude reaching $5.5 \times 10^{-4} \%$ at the cold point level and negligible values at and above the TTL top. Tracer 1 having a very short lifetime, this means that the EN simulation predicts upward transport in the TTL by overshooting convection but mainly below the cold point dynamic barrier. With the horizontal resolution used the EN simulation is not able to simulate small scale overshooting transport

at very high altitude and would favour the slow radiative ascent for the tracers to reach the stratosphere from the TTL. For the GR simulations Tracer 1 fluxes are ~5-6 times lower at 12 and 14 km altitude. At the cold point level and above the fluxes are negligible, as in EN simulation. This shows that the GR simulation provides dynamical fields that are different from EN simulation when convection occurs and have a large impact on the tracer distribution. Fluxes for Tracer 2 are similar for their sign and variations with altitude to Tracer 1 but with higher absolute values. This means that the large scale transport underwent above the cold point level by Tracer 2 in the EN simulation reinforces the upward tracer fluxes.

4.2 Stratospheric tracers

Figure 10 shows the mean mixing ratio profiles for Tracer 3 and 4 (idealised stratospheric tracer) averaged similarly to Tracers 1 and 2 in Fig. 6. The EN and GR parameterizations provide a similar shape with values close to 1 down to the top of the TTL layer (~19 km altitude), a sharp decrease of the tracer mixing ratio below down to 17 km followed by a smoother decrease down to 15 km where it reaches zero. The comparison with the initial mean profile indicates that stratospheric tracers are partly mixed with the TTL air. More than 0.4 ppt are found for tracer 4 at the cold point tropopause level (17.3 km) showing that the model is able to transport significant amounts of stratospheric tracers below the dynamical barrier of the cold point level. But the very low differences between the EN and GR results suggest that this mixing is likely driven by the subgrid-scale diffusion in the model rather than by the direct effect of the convective parameterization. This is also consistent with the results on the tropospheric tracers showing that convection, even using the EN parameterization, hardly reach the cold point tropopause level.

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 Conclusions

This work is a preliminary step for the CTM/CCM/mesoscale model inter-comparison exercise of idealised tracer simulations done in the frame of the European SCOUT-O3 project. One of the main objectives of this inter-comparison is the evaluation of the tracer transport by tropical convection which is known to be a major source of uncertainty in transport processes of chemical species. In this framework, it was proposed to use regional long-term simulations with a limited area model in tropical regions including similar tracers as in the global models to assess the convective tracer transport in CTMs. Compared to CTMs this type of models uses on-line dynamic fields and finer vertical and horizontal resolutions. The objective if these two papers is to evaluate long-duration regional simulations with the mesoscale model CATT-BRAMS with idealized tracers as a tool to produce realistic tracer transport by tropical convection. In this paper, we analyse the impact of different deep convection parametrizations on the transport of idealised tracer in the TTL. For this purpose a simulation over a 60° longitude×40° latitude domain in the Maritime Continent was run with a 60 km horizontal grid spacing and a 300 m vertical grid spacing in the TTL for one month during the period of the SCOUT-O3 aircraft campaign. The Grell (1993) convection parameterization framework extended by Grell and Dévényi (2002) is used to test the impact on deep convection tracer transport of 5 different closures commonly used in the literature and an ensemble parameterization based on these 5 closures.

Since it is not possible to compare the idealised tracers with measurements there is no direct validation of the tracer fields from the model simulations. The choice of idealised tracers is justified by two reasons: (i) if we had used real instead of idealised tracers the comparison would depend largely in this case on the emissions that are poorly quantified in time and space and (ii) idealised tracers facilitate the analysis and understanding of the impact of convective parameterizations on tracer transport. We used an indirect evaluation of the tracer transport through the assessment of the meteorological fields. The comparison with the TRMM surface rainrate estimates shows that

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the 6 parameterizations exhibit two types of behaviours with AS, KF and EN closures giving significantly better results. They reproduce well rainrates in deep convective areas and tend to underestimate less light stratiform precipitation than the other 3 closures. Comparisons were done with a series of radiosoundings launched from Manus Island during the simulation period and for one case study with SCOUT-O3 aircraft data (mainly gathered around 12 and 15–18 km altitude). The model shows a good agreement with the measurements for temperature and wind speed/direction but underestimate the large variability observed within the TTL. The simulations with the 6 parameterizations tend to overestimate the specific humidity in the mid and upper troposphere and show generally small differences compared one to another. From these differences it is not possible to define which among the parameterizations performs better. This meteorological comparison indicates that the six convective parameterizations provide different convection triggering but on average their mean impact on the large-scale environment on temperature and horizontal winds is very similar. However, from this set of meteorological comparison, we can conclude that the use of AS, KF and EN gives better results than the 3 other closures, because of a better behaviour in reproducing the observed rainrates.

The tracer transport is analysed using four idealised tracers (6 h lifetime and infinite lifetime tropospheric tracers and a infinite lifetime stratospheric tracer) for the EN and GR closures that represent respectively the AS/KF and MC/LO behaviour. For both closures the general shape of the mean profile for both tropospheric tracers is similar with large values near the surface, a general decrease up to 10–11 km altitude, a relative maximum around 15 km and a sharp decrease above. But the EN parameterization transport much larger amounts of tropospheric tracers than GR from the surface into the TTL (14 km to 18.9 km altitude) due the stronger vertical ascending velocities in EN. This shows clearly that the tracer transport by convection of surface emissions depends on the convection parameterization used. Even with the EN parameterization the transport above the cold point tropopause is low indicating that none of the parameterization used in this study are able to simulate significant overshooting convection at

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and above these altitudes in the model. Once the tracer emissions are lifted in the TTL above the emission areas by deep convection, they are redistributed horizontally by large scale circulation if they have a sufficient lifetime. In the EN simulation the major part of the infinite tracer amount is not transported down after a few days below the TTL thanks to large scale slow ascending motions. In the GR simulation less tracers remain in the TTL. The stratospheric tracer is on average significantly mixed with the TTL air but does not reach the mid-troposphere. This mixing is probably linked to the model diffusion rather than to the convection parameterization since only very small differences were found between the EN and GR simulations.

The detailed comparison of the model results with aircraft data from the Falcon and the Geophysica on the 23rd November 2005 shows that the model is not able to simulate the local variations of the meteorological variables that are likely linked to convective activity, in particular of the specific humidity, in the 12 and 18.5 km altitude range. This is due to the 60 km horizontal resolution used in the simulations which does not allow the model to provide the small scale effects of convection that can be of importance in the tracer transport. The important issue of the model horizontal and vertical resolutions is the subject of part 2 of this series of two papers.

In this study, we only used the Grell's formalism which provides, even with different closures, similar response to convective instability with modulations in the flux intensity. A complementary work could be done using mass-flux parameterizations based on a more detailed cloud model. Moreover, to go further in the analysis of the type of simulations done in the present study, the use of real tracers such as carbon monoxide could be considered.

Acknowledgements. This work was supported by the European integrated project SCOUT-O3 (GOCE-CT-2004-505390) and by the program LEFE/INSU in France (projects UTLS-tropicale and Tropopause 2009). The computer resources were provided by CINES (Centre Informatique National de l'Enseignement Supérieur), project pce 2227. The TRMM data were provided by GSFC/DAAC, NASA and the Manus radiosounding data by the ARM program funded by the US Department of Energy. The Falcon meteorological data were provided by the DRL (Deutsches Zentrum für Luft- und Raumfahrt, Germany) and the Geophysica meteorological data by the

CAO (Central Aerological Observatory, Russia) and MDB (Myasishchev Design Bureau, Russia). We acknowledge A. Protat and P. May from the BMRC (Bureau of Meteorology Research Center). CATT-BRAMS is a free software provided by CPTEC/INPI and distributed under the CC-GNU-GPL license.



The publication of this article is financed by CNRS-INSU.

References

- Arakawa, A. and Schubert, W. H.: Interaction of a cumulus cloud ensemble with the large-scale environment, Part I. *J. Atmos. Sci.*, 31, 674–701, 1974.
- Baray, J. L., Ancellet, G., Randriambelo, T., and Baldy, S.: Tropical cyclone Marlene and stratosphere-troposphere exchange, *J. Geophys. Res.*, 104, 13953–13970, 1999.
- Brunner, D., Siegmund, P., May, P. T., Chappel, L., Schiller, C., Müller, R., Peter, T., Fueglistaler, S., MacKenzie, A. R., Fix, A., Schlager, H., Allen, G., Fjaeraa, A. M., Streibel, M., and Harris, N. R. P.: The SCOUT-O3 Darwin aircraft campaign: rationale and meteorology, *Atmos. Chem. Phys.*, 9, 93–117, 2009, <http://www.atmos-chem-phys.net/9/93/2009/>.
- Corti, T., Luo, B. P., de Reus, M., Brunner, D., Cairo, F., Mahoney, M. J., Martucci, G., Matthey, R., Mitev, V., dos Santos, F. H., Schiller, C., Shur, G., Sitnikov, N. M., Spelten, N., Vossing, H. J., Borrmann, S., and Peter, T.: Unprecedented evidence for overshooting convection hydrating the tropical stratosphere, *Geophys. Res. Lett.*, 35, L10810, doi:10.1029/2008GL033641, 2008.
- Cotton, W. R., Pielke Sr., R. A., Walko, R. L., Liston, G. E., Tremback, C. J., Jiang, H., McAnelly, R. L., Harrington, J.-Y., Nicholls, M. E., Carrio, G. G., and McFadden, J. P.: RAMS 2001:

ACPD

9, 5889–5928, 2009

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Current status and future directions, *Meteorol. Atmos. Phys.*, 82, 5–29, doi:10.1007/s00703-001-0584-9, 2003.

Deng, A., Seaman, N., L., Hunter, G. K., and Satuffer, D. R.: Evaluation of interregional transport using the MM5-SCIPUFF system, *J. App. Meteor.* 43, 1864–1886, 2004.

5 Duncan, B. N., Strahan, S. E., Yoshida, Y., Steenrod, S. D., and Livesey, N.: Model study of the cross-tropopause transport of biomass burning pollution, *Atmos. Chem. Phys.*, 7, 3713–3736, 2007,

<http://www.atmos-chem-phys.net/7/3713/2007/>.

10 Folkins, I., Loewenstein, M., Podolske, J., Oltmans, S. J., and Proffitt, M.: A barrier to vertical mixing at 14 km in the tropics: Evidence from ozonesondes and aircraft measurements, *J. Geophys. Res.*, 104, D18, 22095–22102, 1999.

Folkins, I., Bernath, P., Boone, C., Donner, L. J., Eldering, A., Lesins, G., Martin, R. V., Sinnhuber, B.-M., and Walker, K.: Testing convective parameterizations with tropical measurements of HNO₃, CO, H₂O, and O₃: Implications for the water vapour budget, *J. Geophys. Res.*, 111, D23304, doi:10.1029/2006JD007325, 2006.

15 Franck, W. M. and Cohen, C.: Simulation of tropical convective systems. Part 1: A cumulus parameterization, *J. Atmos. Sci.*, 44, 3787–3799, 1987.

Freitas, S. R., Longo K. M., Silva Dias, M. A. F., Chatfield, R., Silva Dias, P., Artaxo, P., Andreae, M. O., Grell, G., Rodrigues, L. F., Fazenda, A., and Panetta, J.: The Coupled Aerosol and Tracer Transport model to the Brazilian developments of the Regional Atmospheric Modeling System (CATT-BRAMS). Part 1: model description and evaluation, *Atmos. Chem. Phys. Discuss.*, 7, 8525–8569, 2007.

20 Fueglistaler, S., Wernli, H., and Peter, T.: Tropical troposphere-to-stratosphere transport inferred from trajectory calculations, *J. Geophys. Res.*, 109, D03108, doi:10.1029/2003JD004069, 2004.

Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., Mote P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009.

Gettelman, A. E. and de F. Forster, P. M.: A climatology of the tropical tropopause layer, *J. Met. Soc. Jpn.*, 80, 911–942, 2002.

30 Gilliland, A. B. and Hartley, D. E.: Interhemispheric transport and the role of convective parameterizations, *J. Geophys. Res.*, 103(D17), 22039–22045, 1998.

Grell, G. A.: Prognostic evaluation of assumptions used by cumulus parameterizations, *Mon. Weather Rev.*, 121, 764–787, 1993.

ACPD

9, 5889–5928, 2009

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Grell, G. A. and Dévényi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation, *Geophys. Res. Lett.*, 29, 1693, doi:10.1029/2002GL015311, 2002.

Grell, G. A., Dudhia, and Stauffer, D. R.: A description of the fifth-generation Penn State /NCAR mesoscale model, Tech. Note, NCAR/TN-398+STR, 138 pp., Natl. Cent. Atmos. Res., 1994.

Hack, J. J.: Parameterization of moist convection in the National Center for Atmospheric Research community climate model (CCM2), *J. Geophys. Res.*, 99(D3), 5551–5568, 1994.

Harrington, J. Y.: The effects of radiative and microphysical processes on simulated warm and transition season Arctic stratus, PhD Diss., Atmospheric Science Paper N° 637, Colorado State University, Department of Atmospheric Science, Fort Collins, CO, USA., 289 pp, 1997.

Highwood, E. J. and Hoskins, B. J.: The tropical tropopause, *Q. J. Roy Meteor. Soc.*, 124, 1579–1604, 1998.

Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-Troposphere exchange, *Rev. Geophys*, 33, 403–439, 1995.

Huffman, G. J., Adler, R. F., Morrissey, M. M., Curtis, S., Joyce, R., McGavock, B., and Susskind, J.: Global precipitation at one-degree daily resolution from multi-satellite observations, *J. Hydrometeorol*, 2, 36–50, 2001.

Huffman, G. J., Adler, R. F., Bolvin, D. T., Gu, G., Nelkin, E. J., Bowman, K. P., Hong, Y., Stocker, E. F., and Wolff, D. B.: The TRMM multi-satellite precipitation analysis: quasi-global, multi-year, combined-sensor precipitation estimates at fine scale, *J. Hydrometeorol.*, 8(1), 38–55, 2007.

Kain, J. S. and Fritsch, J. M.: Convective parameterization for mesoscale models: The Kain-Fritsch scheme, the representation of cumulus convection in numerical models, edited by: Emmanuel, K. and Raymond, D. J., *Am. Meteorol. Soc.*, Boston, Mass., USA, 246 pp., 1993.

Kain, J. S. and Fritsch, J. M.: The role of the convective “trigger function” in numerical forecasts of mesoscale systems, *Meteorol. Atmos. Phys.*, 49, 93–106, 1992.

Krishnamurti, T., N., Low-Nam, S., and Pash, R.: Cumulus parameterizations and rainfall rates, *Mon. Weather. Rev.*, 111, 815–828, 1983.

Kuo, H. L.: Further studies of the parameterization of the effect of cumulus convection on large-scale flow, *J. Atmos. Sci.*, 31, 1232–1240, 1974.

Lawrence, M. G. and Rasch, P. J.: Tracer transport in deep convective updrafts: Plume ensemble versus bulk formulations, *J. Atmos. Sci.*, 62, 2880–2894, 2005.

Leclair De Bellevue, J., Réchou, A., Baray, J. L., Ancellet, G., and Diab, R. D.: Signatures of

- stratosphere to troposphere transport near deep convective events in the southern subtropics, *J. Geophys. Res.*, 111, D24107, doi:10.1029/2005JD006947, 2006.
- Marécal, V., Rivièrè, E. D., Held, G., Cautenet, S., and Freitas, S.: Modelling study of the impact of deep convection on the UTLS air composition. Part I: analysis of ozone precursors. *Atmos. Chem. Phys.*, 6, 1567–1584, 2006,
5 <http://www.atmos-chem-phys.net/6/1567/2006/>.
- Nordeng, T. E.: Extended versions of the convective parameterization scheme at ECMWF and their impact on the mean and transient activity of the model in the tropics, ECMWF research Department Technical Memorandum 2006, European Centre for Medium-range
10 Weather Forecasts, Reading, UK, 618–629, 1994.
- Pickering, K. E., Thompson, A. M., Wang, Y., Tao, W.-K., McNamara, D. P., Kirchhoff, V. W. J. H., Heikes, B. G., Sachse, G. W., Bradshaw, J. D., Gregory, G. L., and Blake, D. R.: Convective transport of biomass burning emissions over Brazil during TRACE A, *J. Geophys. Res.*, 101, D19, 23993–24012, 1996.
- 15 Ricaud, P., Barret, B., Attié, J.-L., Motte, E., Le Flochmoën, E., Teysseèdre, H., Peuch, V.-H., Livesey, N., Lambert, A., and Pommereau, J.-P.: Impact of land convection on troposphere-stratosphere exchange in the tropics, *Atmos. Chem. Phys.*, 7, 5639–5657, 2007,
<http://www.atmos-chem-phys.net/7/5639/2007/>.
- Rivièrè, E., Marécal, V., Cautenet, S., and Larsen, N.: Modelling study of the impact of deep
20 convection on the UTLS air composition. Part II: ozone budget in the TTL, *Atmos. Chem. Phys.*, 6, 1585–1598, 2006,
<http://www.atmos-chem-phys.net/6/1585/2006/>.
- Sherwood, S. C. and Dessler, A. E.: On the control of stratospheric humidity, *Geophys. Res. Lett.*, 27, 2513–2516, 2000.
- 25 Sherwood, S. C. and Dessler, A. E.: A model for transport across the tropical tropopause, *J. Atmos. Sci.*, 58, 765–779, 2001.
- Stephenson, D. B. and Doblas-Reyes, F. J.: Statistical methods for interpreting Monte Carlo ensemble forecasts, *Tellus*, 52A, 300–322, 2000.
- Stohl, A., Bonasoniet, P., Cristofanelli, P., et al.: Stratosphere-troposphere exchange: a review and what we have learned from STACCATO, *J. Geophys. Res.*, 108(D12), 8516,
30 doi:10.1029/2002JD002490, 2003.
- Tiedke, M.: A comprehensive mass flux scheme for cumulus parameterization in large scale models, *Mon. Weather Rev.*, 117, 1779–1800, 1989.

**Regional modelling
of tracer transport by
tropical convection –
Part 1**J. Arteta et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Vaughan, G., Schiller, C., MacKenzie, A. R., Bower, K., Peter, T., Schlager, H., Harris, N. R. P., and May, P. T.: SCOUT-O3/ACTIVE high altitude aircraft measurements around deep tropical convection, *B. Amer. Meteor. Soc.*, 89, 647–661, 2008.

Wang, C., Crutzen, P. J., and Ramanathan, V.: The role of a deep convective storm over the tropical Pacific Ocean in the redistribution of atmospheric chemical species, *J. Geophys. Res.*, 100, D6, 11509–11516, 1995.

Wang, Y., Tao, W.-K., Pickering, K. E., Thompson, A. M., Kain, J. S., Adler, R. F., Simpson, J., Keehn, P. R., and Lai, G. S.: Mesoscale model simulations of TRACE A and Preliminary Regional Experiment for Storm-scale Operational and Research Meteorology convective systems and associated tracer transport, *J. Geophys. Res.*, 101, D19, 24013–24027, 1996.

Walko, R. L., Cotton, W. R., Meyers, M. P., and Harrington, J. Y.: New RAMS cloud microphysics parameterization. Part I: the single-moment scheme, 38, 29–62, 1995.

Wild, O. and Prather, M. J.: Global tropospheric ozone modelling: quantifying errors due to grid resolution, *J. Geophys. Res.*, 111, D11305, doi:10.1029/2005JD0006605, 2006.

Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Centre General Circulation Model, *Atmos. Ocean.*, 33, 407–446, 1995.

Zhang, K., Wan, H., Zhang, M., and Wang, B.: Evaluation of the atmospheric transport in a GCM using radon measurements: sensitivity to cumulus convection parameterization, *Atoms. Chem. Phys.*, 8, 2811–2832, 2008.

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.

Table 1. Characteristics of the idealized tracers used in the simulations.

Tracer	Lifetime	Initial conditions	Emissions
1	6 h	0	$10^{-9} \text{ kg m}^{-2} \text{ s}^{-1}$ over land
2	Infinite	0	$10^{-9} \text{ kg m}^{-2} \text{ s}^{-1}$ over land
3	Infinite if $\theta > 380 \text{ K}$ 6 h if $\theta < 380 \text{ K}$	1 ppt if $\theta > 380 \text{ K}$ 0 ppt if $\theta < 380 \text{ K}$	No emissions
4	Infinite	1 ppt if $\theta > 380 \text{ K}$ 0 ppt if $\theta < 380 \text{ K}$	No emissions

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.

Table 2. Mean tracer fluxes at different levels averaged over the model domain and over the one month simulation period using 3-hourly model outputs.

Altitude (km)	Tracer 1 flux (10^{-12} kg, m $^{-2}$ s $^{-1}$)		Tracer 2 flux (10^{-12} kg m $^{-2}$ s $^{-1}$)	
	EN	GR	EN	GR
12 (below TTL)	3.92	0.82	97.27	57.86
14 (TTL bottom)	4.04	0.59	65.33	22.88

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Regional modelling
of tracer transport by
tropical convection –
Part 1**

J. Arteta et al.




Fig. 1. Model topography of simulated domain. The main islands constituting the Indonesian archipelago are Sumatra, Java, the South part of Borneo, Sulawesi and the West part of the New Guinea.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 2. Daily mean surface rainrate in mm day^{-1} from 15 November 2005 to 15 December 2005 for (a) TRMM and the model simulations using (b) Arakawa-Schubert, (c) Kain-Fritsch, (d) Grell, (e) Moisture Convergence, (f) Low Omega, (g) Ensemble.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 3. Comparison between the Manus radiosounding data and the six model simulations: (a and b) for temperature, (c and d) for horizontal wind speed and (e and f) wind direction, (g and h) for specific humidity. Left panels display the mean (solid line) and standard deviation (dashed line) and right panels the mean bias (model minus observation). Black lines are for the radiosounding data and colored lines for the model simulations. The radiosounding data are averaged over the model vertical levels.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 4. Comparison between the Geophysica meteorological data and the six model simulations. **(a)** Temperature (K), **(b)** horizontal wind speed (m s^{-1}), **(c)** wind direction ($^{\circ}$) and **(d)** specific humidity (g kg^{-1}). The black lines are for the aircraft measurements and the coloured lines for the model results. The dashed line is the model altitude in m.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 5. Same as Fig. 4 but for the Falcon data.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 6. Volumetric tracer mixing ratio profiles (in ppbv) averaged over the model domain and over the one month simulation period using 3-hourly model outputs for **(a)** Tracer 1 and **(b)** Tracer 2. The blue lines correspond to the EN simulation and the red lines to the GR simulation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 7. Mean temperature (top panel) and potential temperature (bottom panel) as a function of altitude for the EN simulation. The mean values are calculated as in Fig. 6. The green, brown and purple lines correspond respectively to the altitudes of the mean TTL bottom and cold point.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 8. Meridian mean of Tracer 1 from EN (a) and GR (b) simulations. (c) and (d) are respectively the same plots but for Tracer 2; and (e) and (f) for vertical velocity. The mean is calculated from the one month period using the 3-hourly outputs.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 9. 15 km height mean of Tracer 2 from EN simulation (a) and GR simulation (b). Wind vector are over plotted. The mean is calculated from the one month period using the 3-hourly outputs.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional modelling of tracer transport by tropical convection – Part 1

J. Arteta et al.




Fig. 10. Same as Fig. 6 but for Tracer 3 and 4 (stratospheric tracers). The dark line is the mean vertical profile at the initial time of the simulation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion