



## Dust deposition and ambient PM<sub>10</sub> concentration in central Asia: Spatial and temporal variability

Xiao-Xiao Zhang<sup>1,2</sup>, Brenton Sharratt<sup>3</sup>, Xi Chen<sup>1</sup>, Zi-Fa Wang<sup>2</sup>, Lian-You Liu<sup>4</sup>, Yu-Hong Guo<sup>2</sup>, Jie Li<sup>2</sup>, Huan-Sheng Chen<sup>2</sup>, and Wen-Yi Yang<sup>2</sup>

5 <sup>1</sup>State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, 830011, China

<sup>2</sup>State Key Laboratory of Atmospheric Boundary Layer and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

<sup>3</sup>USDA-ARS, 215 Johnson Hall, Washington State University, Pullman, WA 99164, USA

10 <sup>4</sup>Key Laboratory of Environmental Change and Natural Disaster, Ministry of Education, Beijing Normal University, Beijing, 100875, China

*Correspondence to:* X. X. Zhang (zhangxx@ms.xjb.ac.cn)

**Abstract.** Aeolian dust transport and deposition are important geophysical processes which influence global biogeochemical cycles. Currently, reliable deposition data are scarce in central Asia. Located in the eastern part of central Asia, Xinjiang Province of northwestern China has long played a strategic role in cultural and economic trade between Asia and Europe. In this paper, we investigated the spatial distribution and temporal variation in dust deposition and ambient PM<sub>10</sub> (particulate matter  $\leq 10 \mu\text{m}$  in aerodynamic diameter) concentration from 2000 to 2013 in Xinjiang Province. This variation was assessed using environmental monitoring records from 14 stations in the province. Over the 14 years, annual average dust deposition across stations in the province ranged from 255.7 to 421.4 t km<sup>-2</sup>. Annual dust deposition was greater in southern Xinjiang (663.6 t km<sup>-2</sup>) than northern (147.8 t km<sup>-2</sup>) and eastern Xinjiang (194.9 t km<sup>-2</sup>). Annual average PM<sub>10</sub> concentration across stations in the province varied from 100 to 156  $\mu\text{g m}^{-3}$  and was 70, 115 and 239  $\mu\text{g m}^{-3}$  in northern, eastern and southern Xinjiang, respectively. The highest annual dust deposition (1394.1 t km<sup>-2</sup>) and ambient PM<sub>10</sub> concentration (582  $\mu\text{g m}^{-3}$ ) were observed in Hotan which is located in southern Xinjiang and at the southern boundary of the Taklamakan Desert. Dust deposition was more intense during the spring and summer than other seasons. PM<sub>10</sub> was the main air pollutant that significantly influenced regional air quality. Annual average ambient PM<sub>10</sub> concentration increased logarithmically with dust deposition ( $R^2 \geq 0.88$ ). The arid climate likely influenced the high dust deposition and PM<sub>10</sub> concentration in the region. While the annual average dust storm frequency remained unchanged from 2000 to 2013, there was a positive relationship between dust storm days and dust deposition and PM<sub>10</sub> concentration across stations. This study suggests that sand storms are a major factor affecting the temporal variability and spatial distribution of dust deposition in central Asia.

**Key words:** Dust deposition; PM<sub>10</sub>; Desert; Atmospheric environment; Central Asia



## 1 Introduction

Airborne dust generated by aeolian activity is an environmental concern in central Asia (Huang et al., 2011). Historically, aeolian activity and airborne dust influenced civilization along the ancient Silk Road which connected Asia and Europe (Zhang, 1984; Dong et al., 2012; Groll et al., 2013). Today, airborne dust is recognized as a factor affecting global radiation and warming (Stanhill, 2005; Carslaw et al., 2013; IPCC, 2013; Huang, et al., 2014) and air quality in distant lands (Tsoar and Pye, 1987; Xu et al., 2007; Uno et al., 2009; Li et al., 2012). Deposition of airborne dust also plays a significant role in soil formation and biological diversity in arid and semi-arid regions (Simonson, 1995). An understanding of atmospheric dust sources, emissions, and deposition is therefore essential to improve regional air quality.

Dust in the atmosphere and its subsequent deposition are vital indicators of aeolian activity and environmental quality. Pye (1987) indicated that reliable dust deposition data are lacking around the world. Annual dust deposition ranges from 10 to 200 t km<sup>-2</sup> on continents and one to two orders of magnitude lower over oceans (Pye, 1987). Observations of dust deposition have been made over deserts with an enhanced awareness of its significance. Table 1 lists observations of dust deposition in desert regions and other regions of the world prone to aeolian activity. High dust deposition occurs in Asia, with an annual deposition of 8365 t km<sup>-2</sup> in the Aral Sea Basin (Wake and Mayewski, 1994; Groll et al., 2013). Shao (2011) estimated that over 2000 megatons of dust is emitted from the Earth's surface into the atmosphere annually. Recent investigations suggest that dust deposition during one dust storm can be 10 to 25 times higher than the annual average (Liu et al., 2004; Goudie, 2014).

Located in the eastern part of central Asia, Xinjiang Province of northwestern China has long played a strategic role in cultural and economic trade between Asia and Europe. Xinjiang Province experiences severe sand and dust storms and is highly susceptible to desertification (Chen, 2010). Xinjiang Province is one of two major source regions of atmospheric dust in China, the other region being western Inner Mongolia (Xuan, 1999; Xuan et al., 2000). Long-range transport of dust from the region strongly affects air quality in eastern Asia (Derbyshire et al., 1998; Uno et al., 2009). In fact, dust from the region can be transported across the Pacific Ocean and thus impact air quality in North America (Husar et al., 2001; Osada et al., 2014). Indeed, particulate matter associated with dust transport can severely deteriorate air quality (Sharratt and Lauer, 2006; Shoemaker et al., 2013). Over the past decades, many observations have been made of processes that govern dust emissions, transport, and deposition in Asia (Shao, 2011; Groll et al., 2013). Little is known, however, concerning dust deposition and concentrations in Xinjiang Province. In fact, temporal and spatial variations in dust deposition and concentration have not been characterized despite the importance of dust transport from the region. To improve our understanding of the fate and transport of airborne dust in central Asia, there is a need for continuous and long-term records of dust deposition and concentration. The purpose of this study is to characterize the spatiotemporal distribution of dust deposition and particulate matter concentration in Xinjiang Province that spans the 21st century. This characterization will strengthen our comprehension on aerosol transport in central Asia and provide aerosol data for modelling dust transport in global desertification regions.



## 2 Methods

### 2.1 Study area

Xinjiang Province is located in northwest China and is the largest inland province which covers an area of more than 1.6 million km<sup>2</sup> (Fig. 1). The Taklamakan and Gurbantunggut Deserts are located in the province. The Taklamakan Desert, located in the southern region, is the world's largest shifting-sand dune desert. The Gurbantunggut Desert, located in the northern region, is the largest fixed-dune desert in China. The province is in part characterized by extreme aridity and aeolian desertification. The average annual precipitation varies from more than 700 mm in high-altitude forests and mountains to less than 50 mm in the deserts. Annual potential evaporation can exceed 2000 mm in desert regions (Chen, 2010). Sand and dust storms occur throughout the year, but are most common in spring. In this study, the province was divided by latitude and longitude into three regions, those being northern Xinjiang, eastern Xinjiang and southern Xinjiang (Table 2).

### 2.2 Experimental Data

Dust deposition and PM<sub>10</sub> concentration were measured at environmental monitoring stations maintained by the Xinjiang Environmental Protection Administration, a division of the Ministry of Environmental Protection (MEP) in China. Data collected at 14 stations (Fig. 1 and Table 2) were used in this study and represent a spatial distribution within this region.

Dust deposition was determined by the gravimetric method and documented at monthly intervals. Glass cylinders were used to monitor dust deposition. Three cylinders (replicates) were installed to monitor dust deposition at each station. The cylinders (0.15 m in diameter, 0.3 m tall, and open at the top) were partly filled with an ethylene glycol (C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>) – water solution prior to deployment. The solution enabled trapping of dust in a liquid media at temperatures below 0°C and also minimized evaporation from the cylinder. Cylinders were mounted vertically on a tower at approximate 10 m above ground. The mass of dust collected by the cylinders was determined after washing the contents out of the cylinders and oven-drying the contents at 105°C. Dust deposition rate was calculated as the mass of dust per unit area per unit time and expressed in units of t km<sup>-2</sup> mon<sup>-1</sup> (MEP, 1994). Monthly and yearly dust deposition data were available through the MEP for the 14 stations from 2000-2013.

Ambient PM<sub>10</sub> concentration was measured with high volume samplers designed to collect particulate matter by filtration. The samplers were installed at 1.5 m above the ground and equipped with fiberglass filters for trapping PM<sub>10</sub>. PM<sub>10</sub> concentration was determined based upon gravimetric filter analysis and flow rate of each sampler. Daily PM<sub>10</sub> concentration data were obtained by the arithmetic mean of four samplers with the sampling time being >18 h for each sampler. PM<sub>10</sub> was expressed in µg m<sup>-3</sup> (MEP, 2011). Annual PM<sub>10</sub> data were available through the MEP for the 14 stations (Xinjiang Statistical Bureau, 2014).

Daily meteorological data, including surface wind speed and direction and dust days, were collected from the China Meteorological Administration. A dust day was defined by visibility according to World Meteorological Organization



protocol; days in which visibility was <10 km at any observation time throughout the day constituted a dust day. The World Meteorological Organization further classifies dust days as dust-in-suspension, blowing dust, and dust storms (Shao and Dong, 2006). Dust-in-suspension constitutes days in which dust is not emitted at the station at the time of observation and visibility is <10 km, blowing dust constitutes days in which dust or sand is emitted at the station and visibility is 1-10 km, and dust storms constitutes days in which dust or sand is emitted at the station and visibility is <1 km. Observations of visibility and wind characteristics at each station were taken at 3 hour intervals throughout the day.

Daily air pollution index (API) data were obtained from air quality monitoring statistics published by the MEP (<http://datacenter.mep.gov.cn>). These data were used to illustrate the impact of airborne dust versus other air pollutants on air quality. The API is calculated according to the daily concentration of three main air pollutants (U.S. EPA, 2006; Wang et al., 2013), namely PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub>. The API is calculated as:

$$\text{API} = \max(\text{API}_i), \quad (1)$$

$$\text{API}_i = \frac{\text{API}_u - \text{API}_L}{C_u - C_L} \times (C_i - C_L) + \text{API}_L, \quad (2)$$

Where API<sub>i</sub> is the index for pollutant i (i.e., PM<sub>10</sub>, SO<sub>2</sub>, and NO<sub>2</sub>), API<sub>u</sub> and API<sub>L</sub> are the upper and lower limits of the index for a specific category of air quality (i.e. excellent, slight, moderate, moderately severe, and severe), C<sub>i</sub> is the observed concentration of pollutant, and C<sub>u</sub> and C<sub>L</sub> are the upper and lower limits of the pollutant for a specific category of air quality.

Information regarding the determination of the API index can be obtained from the MEP (MEP, 2012a, 2012b). Based on the API, air quality was classified as: excellent with an API of 0 to 50, slight pollution with an API of 50 to 100, moderate pollution with an API of 100 to 200, moderately severe pollution with an API of 200 to 300, and severe pollution with an API of 300 to 500. For the purpose of this study, we used only API data collected in 2010 since annual deposition and PM<sub>10</sub> concentration appeared to typify that which occurred between 2000 and 2013 in eastern, northern, and southern Xinjiang Province.

Temporal trends in dust deposition, PM<sub>10</sub> concentration and dust days were evaluated by testing the significance of the slope estimate using a t test at a probability level (P-value) of 0.05.

### 3 Results and discussion

#### 3.1 Dust deposition

Detailed information on dust deposition during 2000-2013 was obtained from 14 environmental monitoring stations (Table 2). Annual average dust deposition across all stations in Xinjiang Province was 301.9 t km<sup>-2</sup>. The highest annual deposition occurred in Hotan and Kashgar in southern Xinjiang while the lowest deposition occurred in Karamay in northern Xinjiang. Based upon spatial characteristics in annual dust deposition, deposition increased from north to south across the province (Fig. 2). The annual average dust deposition was 147.8, 194.9 and 663.6 t km<sup>-2</sup> in northern, eastern and southern Xinjiang, respectively. Although dust deposition was relatively low (<150 t km<sup>-2</sup>) for the majority of stations in northern Xinjiang



Province, dust deposition was at least 50% higher for stations within the industrial belt on the northern slope of the Tianshan Mountains. This industrial belt includes Changji and Urumqi. High dust deposition in the industrial belt was due to industry, coal burning and vehicle exhaust.

Figures 3 and 4 depicted the temporal variation in dust deposition from 2000-2013. The highest annual deposition occurred in 2012 in southern Xinjiang, 2002 and 2012 in eastern Xinjiang, and 2001 in northern Xinjiang. Over the 14 year period, dust deposition varied with time across Xinjiang Province. The slope estimate of the relation between average dust deposition and year ( $-6.4 \pm 0.1 \text{ t km}^{-2} \text{ yr}^{-1}$ ) was significant at  $P=0.05$ . This trend was most apparent in northern Xinjiang (slope estimate was  $-5.6 \pm 0.1 \text{ t km}^{-2} \text{ yr}^{-1}$ ) and least apparent in southern Xinjiang (slope estimate was  $-1.9 \pm 0.1 \text{ t km}^{-2} \text{ yr}^{-1}$ ). High dust deposition occurred in spring in eastern and northern Xinjiang and in spring and summer in southern Xinjiang (Fig. 4). Dust deposition peaked in April in eastern and northern Xinjiang and in May in southern Xinjiang. This corresponds to the onset of wind erosion caused by intensifying zonal flow and rising air temperatures before the arrival of the summer monsoon (Song et al., 2016). The maximum monthly average dust deposition was  $97.5 \text{ t km}^{-2}$  in southern Xinjiang, which was 6.9 and 8 times more than the deposition in northern and eastern Xinjiang, respectively.

### 3.2 $\text{PM}_{10}$ concentration

The annual average  $\text{PM}_{10}$  concentration in Xinjiang was  $125 \mu\text{g m}^{-3}$  based upon data collected at 14 stations from 2000-2013. Ten stations (71 percent) in our study had an annual average  $\text{PM}_{10}$  concentration above the People's Republic of China Class II residential standard of  $70 \mu\text{g m}^{-3}$ . The highest annual average  $\text{PM}_{10}$  concentration ( $582 \mu\text{g m}^{-3}$ ) occurred in Hotan in southern Xinjiang while the lowest average  $\text{PM}_{10}$  concentration ( $46 \mu\text{g m}^{-3}$ ) occurred in Tacheng in northern Xinjiang. The annual average  $\text{PM}_{10}$  concentration appeared to increase from northern to southern regions (Fig. 5). Annual average  $\text{PM}_{10}$  concentration in Xinjiang ranged from 100 to  $156 \mu\text{g m}^{-3}$  (Fig. 6) across years. The annual average  $\text{PM}_{10}$  concentration was 70, 115 and  $239 \mu\text{g m}^{-3}$  in northern, eastern and southern Xinjiang, respectively. The high annual concentration in southern Xinjiang is of the same magnitude as found in other desertification regions of the world such as South Asia, Middle East, and western Sahel desert (WHO, 2015). These high concentrations of suspended particulates, especially finer particulate, may influence the health of sensitive populations who are susceptible to respiratory illness (Goudie, 2014).

Over the period of record (2000-2013), there was a trend for decreasing  $\text{PM}_{10}$  concentration in Xinjiang Province. The slope estimate of the relation between annual  $\text{PM}_{10}$  concentration and year ( $-4.2 \pm 0.1 \mu\text{g m}^{-3} \text{ yr}^{-1}$ ) was significant at  $P=0.05$ . This trend was most apparent in southern Xinjiang (slope estimate was  $-11.8 \pm 0.1 \mu\text{g m}^{-3} \text{ yr}^{-1}$ ). On the contrary,  $\text{PM}_{10}$  concentration appeared to increase with time in eastern and northern Xinjiang (slope estimates were  $1.3 \pm 0.1 \mu\text{g m}^{-3} \text{ yr}^{-1}$  and  $3 \pm 0.1 \mu\text{g m}^{-3} \text{ yr}^{-1}$ , respectively). The slope estimate, however, was not statistically different from zero and indicated no apparent trend in  $\text{PM}_{10}$  concentration with time in northern Xinjiang. A decrease in both dust deposition and  $\text{PM}_{10}$  concentration over 2000 to 2013 suggests a positive relationship between dust deposition and  $\text{PM}_{10}$  concentration. This relationship is supported by data in Fig. 7. A logarithmic function fit the data better than a linear function, suggesting that changes in atmospheric  $\text{PM}_{10}$  concentration are smaller at higher rates of deposition. This trend could be due to deposition of



larger or more massive particles under more severe dust or sand storms. While  $PM_{10}$  concentration may rise under more severe wind erosion events, the limited supply of  $PM_{10}$  in sand (major soil type in the province) will likely suppress a rise in  $PM_{10}$  concentration in the atmosphere under more severe erosion events. Nevertheless, from 2000 to 2013, the decline in both dust deposition and  $PM_{10}$  concentration across Xinjiang could be due to less frequent or intense dust storms because dust deposition in major cities of northern China was found to be closely related to the frequency of sand and dust storms (Zhang et al., 2010).

### 3.3 Influence of atmospheric dust deposition on local air quality

Daily ambient air quality has been reported by MEP since 2000. Airborne dust is one of three pollutants influencing the API, thus the relative contribution of dust to the API was of interest. Accordingly, we made a comparative analysis to identify the impact of airborne dust on air quality in Urumqi and Kuytun in northern Xinjiang, Turpan and Hami in eastern Xinjiang, and Kashgar and Hotan in southern Xinjiang (Fig. 8). In 2010, there were 178, 176, 286, 346, 334, and 363 days in which  $PM_{10}$  was the main constituent of the API in Kuytun, Urumqi, Turpan, Hami, Kashgar and Hotan, respectively (Fig. 8). The  $PM_{10}$  constituent accounted for 48.7%, 48.2%, 78.4%, 94.8%, 91.5%, and 99.5% of the API in the respective above cities. These data suggest that particulate matter is the main air pollutant in Xinjiang. Severe  $PM_{10}$  pollution ( $API > 300$ ) occurred mainly in spring, which was closely associated with the seasonality of strong winds and dust storm activity (Li et al., 2004). Stations in southern Xinjiang (Kashgar and Hotan) had higher API's caused by elevated  $PM_{10}$  concentrations throughout the year. This can be attributed to the violent and persistent eolian activity around the Taklamakan Desert (Pi et al., 2014). Consequently,  $PM_{10}$  is an important pollutant which dominates ambient air quality in Xinjiang.

### 3.4 Factors influencing dust deposition and $PM_{10}$ concentration

Many factors influence ambient particulate concentration and dust deposition, but weather appears to be a dominant factor in arid regions (Zhang et al., 1996; Zhang et al., 2010). The Eurasian atmospheric circulation greatly affects weather in central Asia (Kang, et al., 2013). Strong winds associated with this atmospheric circulation cause large amounts of dust to be emitted into the atmosphere.

Spatial differences in dust deposition and  $PM_{10}$  concentration across Xinjiang Province may be due in part to differences in frequency of dust days in the region. Dust storms normally occurred in all seasons in southern Xinjiang. The magnitude of wind erosion and dust day frequency in southern Xinjiang is nearly twice as large as in northern and the eastern Xinjiang (Wang et al., 2006). Figure 9 displays the variation in dust day frequency in Xinjiang Province from 2000 to 2013. The data indicate that the annual average frequency of dust days fluctuated from 15 to 52 days. The frequency of dust days in the southern region ranged from 41 to 133 days while the frequency of dust days in eastern and northern regions ranged from 2-45 days and 0-3 days, respectively, across years. The slope estimate of the relationship between dust days and years ( $0.11 \text{ day yr}^{-1}$ ) indicated no apparent trend for an increase or decrease in dust days from 2000 to 2013. Thus, despite no temporal trend in dust days, we observed a decline in dust deposition and  $PM_{10}$  concentration across years. This decline in dust





deposition or  $PM_{10}$  concentration could be due to a decrease in frequency of severe dust days versus frequency of dust days from 2000 to 2013 in the region. We are unaware of any previous study which has examined dust day severity in Xinjiang Province, thus we used data available through the China Meteorological Administration to assess trends in dust day severity. Dust days were characterized according to dust-in-suspension, blowing dust, and dust storm events. Although there was no trend in the frequency of blowing dust and dust storm events from 2000 to 2013, there was a trend for fewer dust-in-suspension events from 2000 to 2013 (data not shown). Thus, there appeared to be a close association between frequency of dust-in-suspension events and  $PM_{10}$  concentration and dust deposition. Nevertheless, in examining the relationship between average annual dust days and dust deposition or  $PM_{10}$  concentration across stations, the frequency of dust days was closely related to dust deposition ( $R^2=0.93$ ) (Fig.10) and ambient  $PM_{10}$  concentration ( $R^2=0.89$ ) (Fig.11). There was a significant increase in dust deposition ( $7.91 \text{ t km}^{-2} \text{ day}^{-1}$ ) and  $PM_{10}$  concentration ( $2.06 \mu\text{g m}^{-3} \text{ day}^{-1}$ ) associated with an increase in dust days.

#### 4 Conclusions

The atmospheric environment of central Asia is severely affected by the airborne dust, thus this study was undertaken to quantify dust deposition and ambient  $PM_{10}$  concentration in the region. Data collected at 14 environmental monitoring stations from 2000-2013 in Xinjiang Province, China confirmed that annual average dust deposition ranged from 255.7 to 421.4  $\text{t km}^{-2}$ . Annual average  $PM_{10}$  concentration varied from 100 to 156  $\mu\text{g m}^{-3}$ . The highest dust deposition was observed in Hotan in the southern Taklamakan Desert with 1394.1  $\text{t km}^{-2}$ , which is ten times that in China's Loess Plateau (Liu et al., 2004). The highest ambient  $PM_{10}$  concentration was also observed in Hotan with 582  $\mu\text{g m}^{-3}$ , which far exceeds the World Health Organization's long-term exposure standard (WHO, 2014). These results suggest that dust source in central Asia affect regional air quality and is a potential contributor of global dust.

The spatial distribution and temporal variability in dust deposition and ambient  $PM_{10}$  concentration showed significant variation and a trend for lower deposition and concentration with time. The inner-annual dynamic of dust deposition varied significantly with seasonality. Spring and summer had the highest dust deposition (1.3 times the average), followed by autumn and winter. The highest intensity of dust deposition was observed in May, followed by April, June and July. In dust source areas such as Xinjiang, China, windblown sand and dust affect air quality, especially during the spring season. The analysis of the data indicated no trend in frequency of dust days from 2000 to 2013. A positive relationship existed, however, between dust days and dust deposition as well as airborne  $PM_{10}$  concentration across stations. The effect of weather on dust deposition and ambient air quality cannot be expressed by a simple correlation and should not be extrapolated based on the currently limited evidence. This study provides information on the potential spatial-temporal dust deposition and ambient dust aerosol variation in central Asia. Although longer term datasets are needed to address trends over longer time periods, this work can aid in adjusting model parameters in simulating dry dust deposition or  $PM_{10}$  concentration in desertification regions of central Asia.



*Acknowledgements.* This work was supported by the National Natural Science Foundation of China (No.41301655), the West Light Foundation of the Chinese Academy of Sciences (No.XBBS201104), and the Open Funds (No.LAPC-KF-2013-17) of the State Key Laboratory of Atmospheric Boundary Layer and Atmospheric Chemistry, China.

## References

- 5 Carslaw, K.S., Lee, L.A., Reddington, C.L., Pringle, K.J., Rap, A., Forster, P.M., Mann, G.W., Spracklen, D.V., Woodhouse, M.T., Regayre, L.A., and Pierce, J.R.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503(7474), 67-71, doi: 10.1038/nature12674, 2013.
- Cattle, S.R., McTainsh, G.H., and Elias, S.: Aeolian dust deposition rates, particle-sizes and contributions to soils along a transect in semi-arid New South Wales, Australia, *Sedimentology*, 56, 765-783, doi:10.1111/j.1365-3091.2008.00996.x, 2009.
- Chen, X.: Physical geography of arid land in China, Science Press, Beijing, 124-155, 2010.
- 10 Derbyshire, E., Meng, X., and Kemp, R.A.: Provenance, transport and characteristics of modern aeolian dust in western Gansu Province, China, and interpretation of the quaternary loess record, *Journal of Arid Environments*, 39, 497-516, doi:10.1006/jare.1997.0369, 1998.
- Dong, Z., Lv, P., Qiang, G., Xia, X., Zhao, Y., and Mu G.: Research progress in China's Lop Nur, *Earth-Science Reviews*, 111(1-2), 142-153, doi:10.1016/j.earscirev.2011.11.003, 2012.
- Drees, L.R., Manu, A., and Wilding, L.P.: Characteristics of aeolian dusts in Niger, West Africa, *Geoderma*, 59, 213-233, doi:10.1016/0016-1575(93)90070-2, 1993.
- 15 Gile, L.H. and Grossman, R.B.: The desert project soil monograph: soils and landscapes of a desert region astride the Rio Grande Valley near Las Cruces, New Mexico, USDA-SCS, Washington D.C., 1979.
- Goudie, A.S.: Desert dust and human health disorders, *Environmental International*, 63, 101-113, doi:10.1016/j.envint.2013.10.011, 2014.
- Groll, M., Opp, Chr., and Aslanov, I.: Spatial and temporal distribution of the dust deposition in Central Asia – results from a long term monitoring program, *Aeolian Research*, 9, 49-62, doi:10.1016/j.aeolia.2012.08.002, 2013.
- 20 Huang, J., Wang, T., Wang, W., Li, Z., and Yan, H.: Climate effects of dust aerosols over East Asian arid and semiarid regions, *Journal of Geophysical Research (Atmospheres)*, 119, 11398-11416, doi:10.1002/2014JD021796, 2014.
- Huang, X., Oberhänsli, H., von Suchodoletz H., and Sorrel P.: Dust deposition in the Aral Sea: implications for changes in atmospheric circulation in central Asia during the past 2000 years, *Quaternary Science Reviews*, 30, 3661-3674, doi: 10.1016/j.quascirev.2011.09.011, 2011.
- 25 Husar, R.B., Tratt, D.M., Schichtel, B.A., Falke, S.R., Li, F., Jaffé, D., Gasso, S., Gill, T., Laulainen, N.S., Lu, F., Reheis, M.C., Chun, Y., Westphal, D., Holben, B.N., Gueymard, C., McKendry, I., Kuring, N., Feldman, G.C., McClain, C., Frouin, R.J., Merrill, J., DuBois, D., Vignola, F., Murayama, T., Nickovic, S., Wilson, W.E., Sassen, K., Sugimoto, N., and Malm, W.C.: The Asian dust events of April 1998, *J. Geophys. Res.*, 106(D16), 18317-18330, doi:10.1029/2000JD900788, 2001.
- IPCC Working Group I: Climate change-the physical science basis, Cambridge University Press, New York, 252-271, 2013.
- 30 Kang, L., Ji, M., Huang, J., and Guan, X.: Impact of Eurasian atmospheric circulation on dust events in spring over northern China, *Journal of desert research*, 33(5), 1453-1460, doi:10.7522/j, 2013.
- Khalaf, F. and Al-Hashash, M.: Aeolian sedimentation in the north-western part of the Arabian Gulf, *Journal of Arid Environments*, 6, 319-332, 1983.
- Li, F., Zhao, L., Zhang, H., Zhang, T., and Shirato, Y.: Wind erosion and airborne dust deposition in farmland during spring in the Horqin Sandy Land of eastern Inner Mongolia, China, *Soil and Tillage Research*, 75(2), 121-130, doi:10.1016/j.still.2003.08.001, 2004.
- 35 Li, J., Wang, Z., Luo, G., Luo, G., Sun, Y., and Wang, Y.: Mixing of Asian mineral dust with anthropogenic pollutants over East Asia: a model case study of a super-dust storm in March 2010, *Atmos. Chem. Phys.*, 12, 7591-7607, doi:10.5194/acp-12-7591-2012, 2012.
- Liu, L.Y., Shi, P.J., Gao, S.Y., Zou, X.Y., Erdon, H., Yan, P., Li, X.Y., Ta, W.Q., Wang, J.H., and Zhang, C.L.: Dustfall in China's western loess plateau as influenced by dust storm and haze events, *Atmospheric Environment*, 38(12):1699-1703, doi:10.1016/j.atmosenv.2004.01.003, 2004.
- McTainsh, G.H. and Walker, P.H.: Nature and distribution of Harmattan dust, *Zeitschrift Fur Geomorphologie*, 26, 417-435, 1982.
- 40 Menéndez, I., Díaz-Hernández, J.L., and Mangasa, J.: Airborne dust accumulation and soil development in the North-East sector of Gran Canaria (Canary Islands, Spain), *Journal of Arid Environments*, 71(1), 57-81, doi:10.1016/j.jaridenv.2007.03.011, 2007.
- MEP (Ministry of Environmental Protection, China): Ambient Air Determination of Dustfall Gravimetric Method (GB/T 15265-1994), Standards Press of China, Beijing, 1994.





- MEP (Ministry of Environmental Protection, China): Determination of atmospheric particles PM<sub>10</sub> and PM<sub>2.5</sub> in ambient air by gravimetric method (HJ 618-2011), Standards Press of China, Beijing, 2011.
- MEP (Ministry of Environmental Protection, China): Ambient air quality standards (GB 3095-2012), Standards Press of China, Beijing, 2012a.
- MEP (Ministry of Environmental Protection, China): Technical Regulation on Ambient Air Quality Index (HJ 633-2012), Standards Press of China, Beijing, 2012b.
- 5 Modaihash, A.S.: Characteristics and composition of the falling dust sediments on Riyadh city, Saudi Arabia, *Journal of Arid Environments*, 36, 211-223, doi:10.1006/jare.1996.0225, 1997.
- O'Hara, S.L., Clarke, M.L., and Elatrash, M.S.: Field measurements of desert dust deposition in Libya, *Atmospheric Environment*, 40(21), 3881-3897, doi:10.1016/j.atmosenv.2006.02.020, 2006.
- 10 Osada, K., Ura, S., Kagawa, M., Mikami, M., Tanaka, T.Y., Matoba, S., Aoki, K., Shinoda, M., Kurosaki, Y., Hayashi, M., Shimizu, A., and Uematsu, M.: Wet and dry deposition of mineral dust particles in Japan: factors related to temporal variation and spatial distribution, *Atmos. Chem. Phys.*, 14, 1107-1121, doi:10.5194/acp-14-1107-2014, 2014.
- Péwé, T.L.: Desert dust: origin, characteristics, and effect on man, *Geological Society of America*, 186, 1-10, doi: 10.1130/SPE186-p1, 1981.
- Pi, H., Feng, G., Sharratt, B., Li, X., and Zheng, Z.: Validation of SWEEP for Contrasting Agricultural Land Use Types in the Tarim Basin, *Soil Science*, 15 179(9), 433-445, doi:10.1097/SS.0000000000000083, 2014.
- Pye, K.: *Aeolian dust and dust deposits*, Academic Press, London, 5-63, 1987.
- Saeid, H., Khademia, H., Angel, F., and Ahmad, L.: Characteristics of dust deposited along a transect between central Iran and the Zagros Mountains, *CATENA*, 88(1), 27-36, doi:10.1016/j.catena.2011.09.002, 2012.
- Shao, Y., and Dong, C.: A review on East Asian dust storm climate, modelling and monitoring, *Global and Planetary Change*, 52, 1-22, 20 10.1016/j.gloplacha.2006.02.011, 2006.
- Shao, Y., Wyrwoll, K., Chappell, A., Huang, J., Lin, Z., McTainsh, G., Mikami, M., Tanaka, T., Wang, X., and Yoon, S.: Dust cycle: an emerging core theme in Earth system science, *Aeolian Research*, 2(4), 181-204, doi:10.1016/j.aeolia.2011.02.001, 2011.
- Sharratt, B., and Lauer, D.: Particulate matter concentration and air quality affected by windblown dust in the Columbia Plateau, *Journal of Environmental Quality*, 35, 2011-2016, doi:10.2134/jeq2006.0212, 2006.
- 25 Shoemaker, J.K., Schrag, D.P., Molina, M.J., and Rmanathan, V.: What role for short-lived climate pollutants in mitigation policy, *Science*, 342(6164), 1323-1324, doi:10.1126/science.1240162, 2013.
- Simonson, R.: Airborne dust and its significance to soil, *Geoderma*, 65, 1-43, doi: 10.1016/0016-7061(94)00031-5, 1995.
- Song, H., Zhang, K., Piao, S., and Wan, S.: Spatial and temporal variations of spring dust emissions in northern China over the last 30 years, *Atmospheric Environment*, 126, 117-127, doi:10.1016/j.atmosenv.2015.11.052, 2016.
- 30 Stanhill, G.: Global dimming: a new aspect of climate change, *Weather*, 60(1), 11-14, doi:10.1256/wea.210.03, 2005.
- Tsoar, H., and Pye, K.: Dust transport and the question of desert loess formation, *Sedimentology*, 34, 139-153, doi:10.1111/j.1365-3091.1987.tb00566.x, 1987.
- Uno, I., Eguchi, K., and Yumimoto, K.: Asian dust transported one full circuit around the globe, *Nature Geoscience*, 2, 557-560, doi:10.1038/ngeo583, 2009.
- U.S. EPA: Guideline for reporting of daily air quality - the Air Quality Index (AQI), U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 2006.
- 35 USDA: Dust deposition and weather, *Weatherwise*, 21(2), 65-70, doi:10.1080/00431672.1968.9932796, 1968.
- Wake, C.P. and Mayewski, P.A.: Modern aeolian dust deposition in central Asia, *Tellus*, 46B, 220-233, 1994.
- Wang, X., Oenema, O., Hoogmoed, W.B., Perdok, U.D., and Cai, D.: Dust storm erosion and its impact on soil carbon and nitrogen losses in northern China, *CATENA*, 66:3, 221-227, doi:10.1016/j.catena.2006.02.006, 2006.
- 40 Wang, Y.M., Wang, J.H., Qi, Y., and Yan, C.Z.: China 1:100000 Desert (Sand land) Map, Environmental and Ecological Science Data Centre for West China (<http://westdc.westgis.ac.cn>), doi:10.3972/westdc.006.2013.db, 2005.
- Wang, L., Zhang, P., Tan, S., Zhao, X., Cheng, D., Wei, W., Su, J., and Pan, X.: Assessment of urban air quality in China using air pollution indices (APIs), *Journal of the Air & Waste Management Association*, 63(2), 170-178, doi:10.1080/10962247.2012.739583, 2013.
- WHO: Ambient (outdoor) air quality and health, available at: <http://www.who.int/mediacentre/factsheets/fs313/en/>, 2014.
- 45 WHO: Ambient Air Pollution database, available at: <http://gamapserv.who.int/mapLibrary/app/searchResults.aspx>, 2015.
- Xuan, J.: Dust emission factors for environment of northern China, *Atmospheric Environment*, 33, 1767-1776, doi:10.1016/S1352-2310(98)00339-2, 1999.
- Xuan, J., Liu, G.L., and Du, K.: Dust emission inventory in northern China, *Atmospheric Environment*, 34, 4565-4570, doi:10.1016/S1352-2310(00)00203-X, 2000.



- Yaalon, D.H. and Ganor, E.: Rates of aeolian dust accretion in the Mediterranean and desert fringe environments of Israel, 9th International Sedimentological Congress, Nice, France, 2, 169-174, 1975.
- Xinjiang Statistical Bureau: Xinjiang Statistical Yearbook 2014, China Statistical Press, Beijing, 60-225, 2014.
- Xu, J., Hou, S., Qin, D., Kang, S., Ren, J., and Ming, J.: Dust storm activity over the Tibetan Plateau recorded by a shallow ice core from the north slope of Mt. Qomolangma (Everest), Tibet-Himal region, *Geophys. Res. Lett.*, 34, L17504, doi:10.1029/2007GL030853, 2007.
- 5 Zhang, D.: Preliminary analysis on climatology of dust deposition in China history, *Scientia Sinica (Chinese)*, 3, 278-288, 1984.
- Zhang, X.X., Shi, P.J., Liu, L.Y., Tang, Y., Cao, H.W., Zhang, X.N., Hu, X., Guo, L.L., Lue, Y.L., Qu, Z.Q., Jia, Z.J., and Yang, Y.Y.: Ambient TSP concentration and dustfall in major cities of China: spatial distribution and temporal variability, *Atmospheric Environment*, 44, 1641-1648, doi:10.1016/j.atmosenv.2010.01.035, 2010.
- 10 Zhang, X.Y., Shen, Z.B., Zhang, G.Y., Chen, T., and Liu, H.: Remote mineral aerosols in westerlies and their contributions to the Chinese loess, *Science in China Series D (Earth Science)*, 39 (2), 134-143, 1996.



## Tables

**Table 1.** Observations of dust deposition in desertification regions.

Continent	Location	Period	Dust deposition $\text{t km}^{-2} \text{yr}^{-1}$	Citation
America	Kansas, USA	1964-1966	53.5-62.1	USDA (1968)
	New Mexico, USA	1962-1972	9.3-125.8	Gile and Grossman (1979)
	Arizona, USA	1972-1973	54	Péwé (1981)
Europe	Spain	2002-2003	17-79	Menéndez et al. (2007)
Africa	Nigeria	1976-1979	137-181	McTainsh and Walker (1982)
	Niger	1985	164-212	Drees et al. (1993)
	Libya	2000-2001	420	O'Hara et al. (2006)
Oceania	Australia	2000-2001	5-100	Cattle et al. (2009)
Asia	Israel	1968-1973	57-217	Yaalon and Ganor (1975)
	Kuwait	1982	2600	Khalaf and Al-Hashash (1983)
	Saudi Arabia	1991-1992	4704	Modaihash (1997)
	Lanzhou, China	1988-1991	108	Derbyshire et al. (1998)
	Loess Plateau, China	2003-2004	133	Liu et al. (2004)
	Urumqi, China	1981-2004	284.5	Zhang et al. (2010)
	Iran	2008-2009	72-120	Saeid et al. (2012)
Uzbekistan	2003-2010	8365	Groll et al. (2013)	



**Table 2.** Dust deposition and PM<sub>10</sub> concentrations at 14 stations in Xinjiang.


No.	Station	Region*	Latitude	Longitude	Population** (million)	Annual dust deposition (t km <sup>-2</sup> )	Annual PM <sub>10</sub> concentration (µg m <sup>-3</sup> )
1	Urumqi	NJ	43.832°N	87.616°E	2.26	229.4	141
2	Changji	NJ	44.017°N	87.308°E	0.36	295.7	76
3	Shihezi	NJ	44.306°N	86.080°E	0.62	107.7	61
4	Bole	NJ	44.900°N	82.071°E	0.27	133	48
5	Karamay	NJ	45.580°N	84.889°E	0.29	81.1	54
6	Tacheng	NJ	46.691°N	82.952°E	0.17	84.9	39
7	Yining	NJ	43.912°N	81.329°E	0.53	142.7	78
8	Kuytun	NJ	44.426°N	84.903°E	0.30	108.1	66
9	Hami	EJ	42.818°N	93.514°E	0.48	209.8	84
10	Turpan	EJ	42.957°N	89.179°E	0.28	180.1	145
11	Korla	SJ	41.727°N	86.174°E	0.57	231.8	131
12	Hotan	SJ	37.113°N	79.922°E	0.33	1394.1	352
13	Kashgar	SJ	39.471°N	75.989°E	0.57	516.9	236
14	Aksu	SJ	41.170°N	80.230°E	0.51	511.5	238

\* Xinjiang Province was classified into three regions: northern Xinjiang (NJ), eastern Xinjiang (EJ), and southern Xinjiang (SJ).


\*\*Population in 2013 as reported by the Xinjiang Statistical Bureau.



## Figures




**Figure 1.** Location of Xinjiang Province in China (gray area outlined on inset map). Dust deposition and concentrations were measured at stations signified by small triangles. Land use types are identified across the province according to Wang et al., 2005.




**Figure 2.** Annual average dust deposition ( $t\ km^{-2}$ ) reported at 14 stations in Xinjiang Province from 2000-2013. Land use types across the province are identified according to Wang et al. (2005).







**Figure 3.** Annual average dust deposition in Xinjiang Province from 2000 to 2013. Dust deposition in northern, eastern and southern Xinjiang is the average deposition at 8, 2 and 4 stations, respectively.




**Figure 4.** Monthly average dust deposition in Xinjiang Province from 2000-2013. Dust deposition in northern, eastern and southern Xinjiang is the average deposition at 8, 2 and 4 stations, respectively.




**Figure 5.** Annual average  $PM_{10}$  concentration ( $\mu\text{g m}^{-3}$ ) reported for 14 stations in Xinjiang Province from 2000-2013. Land use types are identified across the province according to Wang et al., 2005.



**Figure 6.** Annual average PM<sub>10</sub> concentration in Xinjiang Province from 2000 to 2013.



**Figure 7.** Relationship between annual dust deposition and PM<sub>10</sub> concentration in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.



**Figure 8.** Daily air pollution index for Kuytun and Urumqi in northern Xinjiang, Hami and Turpan in eastern Xinjiang, and Kashgar and Hotan in southern Xinjiang in 2010. The main air pollutant contributing to the daily API is identified for each station. Not detected indicates excellent air quality (API<50).








Figure 9. Dust day frequency in Xinjiang Province from 2000 to 2013.



**Figure 10.** Relationship between annual dust deposition and dust day frequency in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.



**Figure 11.** Relationship between PM<sub>10</sub> concentration and dust day frequency in Xinjiang Province. Each point represents data averaged across 2000 to 2013 at one station.