



Spatiotemporal patterns of the fossil-fuel CO₂ signal in central Europe: Results from a high-resolution atmospheric transport model

Yu Liu^{1,2}, Nicolas Gruber^{1,2}, and Dominik Brunner³

¹Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, Zurich, Switzerland

²Center for Climate Systems Modeling (C2SM), ETH Zurich, Switzerland

³Laboratory for Air Pollution/Environmental Technology, Swiss Federal Laboratories for Materials Science and Technology, Empa, Duebendorf, Switzerland

Correspondence to: Nicolas Gruber (nicolas.gruber@env.ethz.ch)

Abstract.

The emission of CO₂ from the burning of fossil fuel is a prime determinant of variations in atmospheric CO₂. Here, we simulate this fossil fuel signal together with the natural and background components with a regional high-resolution atmospheric transport model for central and southern Europe considering separately the emissions from different sectors and countries on the basis of emission inventories and hourly emission time functions. The simulated variations in atmospheric CO₂ agree very well with observation-based estimates, although the observed variance is slightly underestimated, particularly for the fossil fuel component. Despite relatively rapid atmospheric mixing, the simulated fossil fuel signal reveals distinct annual mean structures deep into the troposphere reflecting the spatially dense aggregation of most emissions. The fossil fuel signal accounts for more than half of the total (fossil fuel + biospheric + background) temporal variations in atmospheric CO₂ in most areas of northern and western central Europe, with the largest variations occurring on diurnal timescales owing to the combination of diurnal variations in emissions and atmospheric mixing/transport out of the surface layer. Their co-variance leads to a fossil-fuel diurnal rectifier effect with a magnitude as large as 9 ppm compared to a case with time-constant emissions. The spatial pattern of CO₂ from the different sectors largely reflects the distribution and relative magnitude of the corresponding emissions, with power plant emissions leaving the most distinguished mark. An exception is southern and western Europe, where the emissions from the transportation sector dominate the fossil fuel signal. Most of the fossil fuel CO₂ remains within the country responsible for the emission, although in smaller countries, up to 80% of the fossil fuel signal can come from abroad. A fossil fuel emission reduction of 30% is clearly detectable for a surface-based observing system for



atmospheric CO₂, while it is beyond the edge of detectability for the current generation of satellites with the exception of a few hotspot sites. Changes in variability in atmospheric CO₂ might open an additional door for the monitoring and verification of changes in fossil fuel emissions, primarily for surface based systems.

1 Introduction

With annual CO₂ emissions from fossil fuel burning and cement production having soared in the recent decades and approaching 10 Pg C yr⁻¹ (Raupach et al., 2007; Friedlingstein et al., 2014; Le Quéré et al., 2016), these fluxes have reached the same order of magnitude as the natural exchange fluxes between the atmosphere and land surface and between the atmosphere and the ocean, respectively (Sarmiento and Gruber, 2002; Le Quéré et al., 2016). Thus, the fossil fuel emissions have become a key driver for the spatiotemporal dynamics of atmospheric CO₂, not only close to major sites of emissions, but also far downstream (Peylin et al., 2011; Keppel-Aleks et al., 2013; Nassar et al., 2013). This represents simultaneously a challenge and an opportunity. It is an opportunity since the substantial and growing size of this fossil fuel CO₂ signal facilitates the use of variations in atmospheric CO₂ to monitor and verify changes in fossil fuel emissions (Bovensmann et al., 2010; Velasco et al., 2011; McKain et al., 2012; Ciais et al., 2014; Shiga et al., 2014). At the same time, the large fossil fuel CO₂ signal complicates the use of atmospheric CO₂ observations to determine sources and sinks of CO₂ driven by the land biosphere through atmospheric inverse modeling methods. This requires the separation of the biospheric signal in atmospheric CO₂ from the total signal, which is usually accomplished by subtracting an estimate of the fossil fuel component from the measured atmospheric CO₂ concentration. This implies that any error in the fossil fuel component tends to be projected directly onto the inversely estimated biospheric fluxes (Nassar et al., 2013; Peylin et al., 2011). Thus, in order to benefit from the monitoring and verification opportunity as well as to minimize the magnitude of the challenge associated with atmospheric inversions, it is paramount to well characterize the fossil fuel component in atmospheric CO₂ in time and space.

Two sets of approaches have been developed to determine this fossil fuel component in atmospheric CO₂. A first set of approaches relies on concurrent observations of carbon monoxide (CO) and/or radiocarbon to determine the fossil fuel component in the observed atmospheric CO₂ variations (Bréon et al., 2015; Ciais et al., 2013; Levin and Karstens, 2007; Van Der Laan et al., 2010; Turnbull et al., 2011; Newman et al., 2013; Vogel et al., 2013; Lindenmaier et al., 2014; Vardag et al., 2015; Oney et al., 2016, in review). A major advantage of these observation-based methods is that they do not require any atmospheric transport modeling, and thus are not sensitive to any errors in the modeled atmospheric transport. A major disadvantage is that these observation-based estimates are available only at a relatively small set of observing sites, providing a very limited picture of the spatiotemporal dynamics of the fossil fuel signal for larger areas. Further complications may arise



from e.g., poorly known and varying ratios of the emissions of CO and CO₂ in the case of CO-based methods (Oney et al., 2016, in review), or the emission of radiocarbon from nuclear power and reprocessing plants in the case of radiocarbon-based methods (Graven and Gruber, 2011).

60 In the second set of approaches the fossil fuel CO₂ signal is modeled, starting from the specification of fossil fuel emissions as a bottom boundary condition in an atmospheric transport model, and then running this model forward in time (Peylin et al., 2011; Ogle et al., 2015). A key advantage of this set of approaches is that the spatiotemporal dynamics is fully resolved. But this comes at the disadvantage that the resulting accuracy of the modeled fossil fuel CO₂ signal not only depends on
65 the quality of the fossil fuel emissions data, but also on that of the transport model. The latter disadvantage is well illustrated by the results of a recent model intercomparison study, where inter-model differences in the simulated spatiotemporal pattern of the fossil fuel CO₂ were 2-3 times larger than the differences resulting from the use of different emission inventories (Peylin et al., 2011). Of particular relevance is the resolution of the atmospheric model, as this is key to better resolve the
70 topography and land surface contrasts that govern much of the atmospheric circulation and mixing in the lower atmosphere.

The challenge associated with the modeling of atmospheric transport is particularly acute for the fossil fuel component, since fossil fuel emissions are distributed in time and space in a highly heterogeneous and non-Gaussian manner (Ray et al., 2014). This reflects the nature of the processes
75 underlying these emissions, ranging from the point source nature of the emissions from coal-fired power plants, whose emissions vary in response to changing needs for electricity, to the strong diurnal fluctuations of the dispersed emissions associated with road transportation (Nassar et al., 2013). This strong spatial and temporal patterning of the fossil fuel emissions interacts with the spatiotemporal variability of atmospheric transport, forming distinct patterns of the fossil fuel signal in atmospheric CO₂ (Feng et al., 2016). Of particular relevance are the diurnal and the seasonal changes
80 in emissions, since they tend to co-vary with atmospheric transport, which can lead to atmospheric CO₂ concentration gradients due to a rectification effect (Denning et al., 1995; Zhang et al., 2016). Such unaccounted for variations in the fossil fuel signal would bias the biospheric signal in atmospheric inversion frameworks, hindering us from developing a better understanding of the role of the
85 land biosphere as a carbon sink. At the same time, this strong temporal patterning of the emissions creates also distinct signals that might be used to detect or track the fossil fuel signal.

In fact, several studies already explored the possibilities to detect the fossil fuel signal (Ciais et al., 2014; Nassar et al., 2013). These include a range of methods and systems, including bottom up methods based on surface observation systems (Shiga et al., 2014; McKain et al., 2012; Keller
90 et al., 2016), CO and radiocarbon based methods (Levin and Karstens, 2007; Van Der Laan et al., 2010; Vogel et al., 2013), airborne measurements (Turnbull et al., 2011), satellite constraints (Kort et al., 2012), and top-down approaches on the basis of atmospheric inversions (Ogle et al., 2015; Lauvaux et al., 2016; Brioude et al., 2013). Spatially, the focus ranges from point scale emissions



(Bovensmann et al., 2010; Velazco et al., 2011; Turnbull et al., 2016), or urban-scale (Newman et al.,
95 2013; Bréon et al., 2015; Turnbull et al., 2015; Pillai et al., 2016) to regional and global (Keppel-
Aleks et al., 2013; Basu et al., 2016).

A necessity to successfully deploy any of these different detection approaches is a good under-
standing of the spatiotemporal dynamics of the fossil fuel signal over a scale that is sufficiently large
in order to avoid an unacceptably high sensitivity to the lateral boundary conditions, i.e., over scales
100 exceeding a few 100 km. A successful detection also requires a good understanding of the contribu-
tion of the other processes affecting atmospheric CO₂ variations, namely the exchange fluxes with
the land biosphere and with the ocean, respectively. Further, often it would be quite useful to know
the source processes responsible for the fossil-fuel CO₂ signature, i.e., what fraction of the signal
stems from emissions from a coal-fired power plant and what part from road transportation. This
105 helps, e.g., with the assessment of how the implementation of a particular policy affects the fossil
fuel signature, such as e.g., the shutting down of coal- fired power plants.

Few studies have taken a continental to global perspective on the fossil fuel signal (Keppel-Aleks
et al., 2013), as the focus in the last few years had been on urban areas (McKain et al., 2012; New-
man et al., 2013; Kort et al., 2012), or just whether the emissions in the city be detected or not (Hase
110 et al., 2015; Pillai et al., 2016). In addition, comparatively less work has been carried out in Europe
(Schneising et al., 2008), and the majority of those used relatively coarse resolution atmospheric
transport models, resulting in relatively washed-out gradients of the fossil fuel signal over Europe
(Keppel-Aleks et al., 2013; Peylin et al., 2011), or few of them focused on whether the potential re-
duced emissions could be discerned by current observation methods in this region or not (Levin et al.,
115 2011). Furthermore, little consideration has been given to the temporal variations of the emissions.

The main objective of this work is to fill these gaps, and to develop a quantitative understanding
of the fossil fuel CO₂ signal in Europe. To this end, we employ a forward modeling approach using a
high resolution atmospheric transport model for Europe, forced with finely resolved fossil fuel emis-
sion fluxes in time and space. In this paper, we will (i) investigate the magnitude of the contribution
120 of the fossil fuel CO₂ signal to the variations in total CO₂; (ii) understand how the high temporal
resolution considered in the fossil fuel emissions affect the fossil CO₂ signal; and (iii) determine the
detectability of a reduction of fossil fuel emissions from different sources through changes in the
column mean CO₂ as seen, e.g., by a satellite-based observing system. We first describe the model
and methods, followed by the evaluation of the model in the third part. We then present the results,
125 followed by a discussion of each of the aforementioned three main topics, and then conclude with a
summary and an outlook.



2 Methods and Data

To simulate the fossil-fuel CO₂ over central and southern Europe in the context of the variations in total atmospheric CO₂, we employ a regional high-resolution atmospheric transport model for the European domain and prescribe lateral and boundary conditions for the various components that constitute atmospheric CO₂. These include the fossil fuel emissions, the CO₂ exchange fluxes with the land and ocean surfaces, and the lateral atmospheric CO₂ boundary conditions. The simulations cover the period April 2008 until April 2009. The following subsections describe the methods and data in more detail.

2.1 Atmospheric transport model

The simulations were undertaken with the limited-area atmospheric prediction model COSMO (Consortium for Small-scale Modeling) (Baldauf et al., 2011) Version 4.23. We employed the COSMO-7 setup developed by the Swiss Federal Office for Meteorology and Climatology (MeteoSwiss) for the purpose of providing boundary conditions for the inner COSMO-2 grid used for forecasting the weather in Switzerland. The COSMO-7 setup has a grid spacing of 6.6 km and its domain covers central and southern Europe (35.16°N/9.80°E (lower left) to 56.84°N/23.02°E (upper right) (see Fig. 1).

The COSMO model is based on the primitive hydro-thermodynamical equations describing compressible non-hydrostatic flow in a moist atmosphere without any scale approximations. The model equations are solved numerically on a rotated latitude-longitude grid, with terrain-following coordinates in the vertical (60 vertical levels, and lowest level at 10 meters), using an Eulerian finite difference method. Parameterization schemes are used to resolve the sub-grid scale physical processes such as vertical diffusion (turbulence), convection, radiation, and soil processes. A tracer transport module was recently added to the COSMO model, permitting the online transport of passive tracers in a manner that is fully consistent with the physics of the model (Roches and Fuhrer, 2012). In our setup, advective transport was accomplished with a 3-dimensional semi-Lagrangian scheme. The tracers are transported in the model as moist air mass mixing ratios q_{CO_2} . Values reported here are provided as dry air mole fractions χ_{CO_2} , calculated as $\chi_{CO_2} = q_{CO_2} / (1 - q_{H_2O}) M_{dry} / M_{CO_2}$, where q_{H_2O} is the specific humidity and M_{dry} and M_{CO_2} are the molar masses of dry air and CO₂, respectively. The dry air column average mixing ratio is calculated as $X_{CO_2} = (\sum_{k=1}^K (p(k+1/2) - p(k-1/2)) q_{CO_2}(k)) / (\sum_{k=1}^K (p(k+1/2) - p(k-1/2)) (1 - q_{H_2O}(k))) * M_{dry} / M_{CO_2}$. K is the vertical level, and p pressure, which is at the staggered level (Baldauf et al. (2011) is recommended for more information).



2.2 Fossil fuel emissions

160 The fossil fuel emissions for CO₂ were generated by merging a relatively coarse emission inventory
for the regions outside Switzerland (EDGAR v4.2_FT2010, approx. 10 km, (Janssens-Maenhout
et al., 2012)) with a high-resolution (0.5 km) emission inventory for Switzerland. The latter was pro-
duced by the company MeteoTest specifically for the CarboCount CH project. The annual emissions
from this merged product for the year 2008 amount to 2.54 Pg CO₂ over the domain, representing
165 about 10% of the global emissions of that year (Le Quéré et al., 2016). We merged the emission
categories from the two inventories to 5 large emission categories, i.e., power generation, residential
heating, road transportation, industrial processes, and others. Even though each of these different
categories have a distinct emission pattern, many of them co-occur in the large metropolitan areas,
leading to a very patchy emission pattern with strong emission hotspots, and extensive regions with
170 relatively low emission densities (Fig. 1).

These emission inventories are given for each emission category as annual totals for each grid
cell, requiring us to multiply them with time functions to generate hourly timeseries of the fossil fuel
emissions at each location (Nassar et al., 2013). The time functions we employed were originally
generated by the University of Stuttgart (Institute für Energiewirtschaft und Rationelle Energiean-
wendung, IER) for the GENEMIS project (Friedrich and Reis, 2004) and have been used since in
175 several air quality modeling studies. The time functions are comprised of diurnal, weekly and sea-
sonal components and are specific to each of the main economic sectors (activities collected in the
Selected Nomenclature for Air Pollution (SNAP) codes) (Kuenen et al., 2014). The time functions
(except for the daily one) vary also by country, and are locally adjusted to reflect local time. Some
180 reassignments were necessary to align the categories used in EDGAR v4.2 and the CarboCount CH
inventory (both following IPCC guidelines) with the SNAP categories as described in the supple-
mentary material.

The time functions differ greatly between the various categories, reflecting their very different
time course of activities over the average day, week or year (see Fig. 2a,b). Among all diurnal time
185 functions, road transportation has the largest diurnal variability and is characterized by two peaks
during the day reflecting the rush hour periods (local time 8:00-9:00 and 17:00-18:00). Also residen-
tial/commercial combustion has a distinct diurnal cycle with two peaks. In contrast, the emissions
from industrial processes and fossil- fuel fired power plants vary less over the course of the day and
also have only one peak. The time functions for the day-of-week reflect primarily the lower industrial
190 activities and traffic during the week end, while most other sectors continue to emit at only slightly
smaller rates (see Fig. 2a). Combining all the sectors together, emissions during the weekend are
15-20% lower than during the week. The seasonal time functions depend primarily on the local cli-
matic conditions (see Fig. 2b), with northern and central European countries having a maximum in
winter due to their heating requirement, while there is little seasonality in emissions in the southern
195 European countries.



In order to be able to trace the fossil fuel signature in atmospheric CO₂ back to the emitters, we consider separate fossil-fuel tracers for ten different countries (or groups of countries) in our atmospheric transport model (see Fig. 1). Each of these tracers receives only the emissions from its respective country or group of countries, while elsewhere, the emissions are set to zero. Due to the
200 linearity of atmospheric transport and the absence of any transformation of CO₂ in the atmosphere, the individual country-based tracers can then be summed to obtain the total fossil fuel CO₂ signal. In addition, in order to determine the contribution of the different CO₂ emission categories to the total fossil fuel CO₂, we also included five additional fossil fuel tracers, one each for the five categories we consider, i.e., power generation, residential heating, road transportation, industrial processes, and
205 others. For these 5 tracers, we used time-invariant emissions, permitting us to assess also the role of the time variations in emissions on the fossil fuel CO₂ signal. In total, we included 17 fossil fuel tracers (10 countries, 5 sectors, and total fossil fuel CO₂ with time varying emission, and total fossil fuel CO₂ with time constant emission) in our high-resolution simulation study.

2.3 Other CO₂ component fluxes

210 In order to simulate the distribution of total atmospheric CO₂, we also include in our model three other CO₂ components, namely background CO₂, the terrestrial biospheric CO₂ and the oceanic CO₂ components. The background CO₂ represents that part of the atmospheric CO₂ that enters the domain through its boundaries. These boundary concentrations are provided by the post-assimilation results of CarbonTracker Europe (Peters et al., 2010). For the terrestrial biospheric CO₂
215 component, we used the hourly terrestrial biospheric fluxes from the Vegetation Photosynthesis and Respiration Model (VPRM) (Mahadevan et al., 2008). For the oceanic CO₂ component, we combined the monthly air-sea CO₂ flux estimates for the Atlantic from Landschützer et al. (2013) with the annual mean flux estimates for the Mediterranean by D'Ortenzio et al. (2008). As the oceanic flux contribution is small, no attempt was made to add higher frequency variability.

220 2.4 Simulations

The hindcast simulation started on March 1, 2008, with the initial and boundary conditions for meteorology taken from the operational hourly COSMO-7 analyses of MeteoSwiss and the initial and boundary conditions for atmospheric CO₂ provided by CarbonTracker Europe (Peters et al., 2010). The model was then run for 13 months until April 30, 2009. No assimilation of any meteorological
225 data was performed. The lateral and boundary conditions for the total of 18 CO₂ tracers considered (15 fossil fuel, 3 other components) were prescribed as described above. We consider the first month as a spinup, and use the subsequent 12 months for our analyses.



3 Evaluation

3.1 Total atmospheric CO₂

230 We evaluate our COSMO-based results for the total atmospheric CO₂ concentration (computed by summing the fossil fuel component with the three others) by comparing them to the measurements from four sites in central Europe, namely Mace Head (MHD, 3.33°N, 9.90°W, 5m above ground, coastal site, 15 m a.s.l.), Cabauw (CBW, 51.97°N, 4.92°E, 20m, 60m, 200m above ground, flatland, near urban site, 0 m a.s.l.), Hegyhatsal (HUN, 6.95°N, 16.65°E, 10m, 48m, and 115m above ground, 235 continental site, 248 m a.s.l.) (Geels et al., 2007), Puy de Dome (PUY, 45.46°N, 2.58°E, mountain site, 1480 m a.s.l.). In order to minimize the impact of local influences, we use the average CO₂ concentrations between 12:00 and 18:00 local time, i.e., the time of day of maximum vertical mixing.

The modeled atmospheric CO₂ records at these four sites agree well with the observed ones (see Table 1). The correlation between the modeled and observed values exceed 0.7 at all sites and 240 heights. The highest correlation is found at Mace Head (MHD) (>0.81). This is due to the relatively steady conditions that characterize this relatively clean coastal site. Influence from air pollution is only observed during episodes of transport from the United Kingdom and continental Europe, which are very well captured by the model. The correlations are somewhat lower at the more polluted and more continental sites, i.e., between 0.72 and 0.78 at the coastal tall tower station Cabauw (CBW) in 245 the Netherlands, and around 0.8 at the continental tall tower station Hegyhatsal (HUN) in Hungary. Even the atmospheric CO₂ variations at the mountain top site Puy de Dome in France are well captured ($r = 0.75$).

COSMO tends to systematically underestimate the observed CO₂ concentration at most of the stations and levels, except at the coast of Ireland (MHD), where it is overestimated by 0.3 ppm (Table 250 1). The biases tend to get larger with increasing continentality of the sites, and the associated higher complexity of the surrounding terrain and other influencing factors. At the Cabauw site (CBW), the biases amount to between -0.8 and -1.6 ppm, while in central Hungary (HUN) the biases are already more than -4 ppm at all vertical levels. In general, this may be related to COSMO-7 ventilating the planetary boundary layer too strongly, particularly in winter time under weakly stratified conditions. 255 This is especially acute for the HUN site, because the air in the lowest atmospheric levels tends to get trapped at this site owing to the winter-time prevalence of anticyclonic conditions in the Carpathian Basin (Haszpra et al., 2012). An alternative explanation is that the biospheric sink simulated by VPRM is too strong, as discussed later.

Even though COSMO exhibits some biases in the mean, it captures the observed variability generally well (Table 1). In particular, COSMO reproduces the strong gradient in variability between 260 the coastal site Mace Head (~6 ppm) and the continental site in central Hungary (~12 ppm), reflecting primarily differing contributions of synoptic variations on atmospheric CO₂. However, the



absolute magnitude of the variations are not matched by our simulations, with COSMO consistently underestimating the observed variability.

265 Overall, the evaluation of the total atmospheric CO₂ concentration reveals a good agreement with the observations, both in terms of mean and variability. The low and positive bias at the Mace Head site, where the contribution of the background CO₂ component dominates, suggests that this component is overall well modeled and likely not responsible for the bias at the other sites. This bias is likely due to the superposition of biases in atmospheric transport (as argued for the HUN site) and
270 biases in the underlying boundary conditions for the fossil fuel emissions and/or terrestrial fluxes. Since the contribution of the oceanic fluxes is very small, this component can be excluded as an explanation. Unfortunately, we do not have observationally-based estimates of the fossil fuel or terrestrial biosphere components at the four sites discussed so far, requiring us to use data from other sites for further evaluation.

275 3.2 Fossil fuel CO₂ component

Estimates of the fossil fuel component in atmospheric CO₂ are available for our model simulation period from Lutjewad in the Netherlands (LUT, 6.35° E, 53.4° N, 1 m a.s.l.) (Van Der Laan et al., 2010; Bozhinova et al., 2014) and from Heidelberg (HEI, 49.417°N, 8.675°E, 116 m a.s.l.) (Levin and Karstens, 2007). Both estimates are based on a combination of concurrent CO and ¹⁴C-
280 measurements and represent the fossil fuel induced offset relative to a regional background. They are thus comparable to our modeled fossil fuel component, as this reflects the offset relative to the domain-wide background induced by the lateral boundary conditions. Lutjewad is located on the Waddensea dike in the north of the Netherlands, influenced by the highly populated and industrialized areas in the Netherlands and in northwestern Germany (Ruhr area). The Heidelberg station is located near an
285 urban center with considerable fossil fuel emissions.

At the Dutch site LUT, the daily average fossil fuel CO₂ component simulated by our model compares well with the observations ($r=0.73$, mean bias -4 ppm) (see Fig. 3a). Generally, the model reproduces the observed variability, especially in summer, when the fossil fuel CO₂ component is low owing to the deep mixing in the atmosphere. But the model underestimates the estimated fossil
290 fuel CO₂ component substantially in winter. This may be due to several reasons. First, the model may transport signals too quickly out of the planetary boundary layer, which is a known problem of many atmospheric transport models under stratified conditions typical of wintertime (see also above) (Holtslag et al., 2013). Second, our wintertime emission inventory in the region might be too small, owing to, for example, our underestimating the strength of the seasonal signal in the
295 time functions. Third, the observations might be biased high. One reason is that these reconstruction rely on a constant ratio between CO and ¹⁴C-CO₂, which may lead to an underestimation of the ¹⁴C-CO ratio compared to real values at some time of the year, and subsequently overestimation of the inferred fossil fuel CO₂ (Van Der Laan et al., 2010; Bozhinova et al., 2014).



At Heidelberg, our model captures the fossil fuel CO₂ component even better, particularly since
300 the model has a very small mean bias of 0.75 ppm. Also the day- to-day and the seasonal variations
are well represented with a correlation coefficient of 0.72. The model's (small) overestimation of
the fossil fuel component may be due to our prescribing all emissions at the surface, while the
fossil-fuel fired power plants that contribute substantially to the fossil fuel CO₂ at this site tend to
have an effective emission height quite some distance above the ground due to the height of the
305 stacks and the additional rise of the buoyant plumes (Vogel et al., 2013). Another reason might be
an overestimation of the emissions in our emission inventory EDGAR - an explanation furthered
by EDGAR's emission being higher than those of IER (Peylin et al., 2011). Especially assuring,
and particularly so in comparison to the situation at LUT, is the COSMO model's ability at HEI to
capture most of the variability and amplitude of the fossil fuel component in winter. An exception
310 are the observations from late December and early January, where the data include a number of
exceptionally high peaks. These peaks may be the result of very strong local trapping of the emitted
fossil fuel CO₂ by e.g., a local inversion situation, i.e., a process that our model cannot properly
resolve.

Despite these discrepancies, the good to excellent evaluation results provide us with good con-
315 fidence to use our COSMO-7 based system to investigate the spatio- temporal variability of the
fossil-fuel CO₂ in central and southern Europe. It is particularly encouraging to note the good agree-
ment not only for the fossil fuel CO₂ component, but also for total atmospheric CO₂. The presence
of an overall negative bias in the total atmospheric CO₂ in the absence of such a bias in the fossil
fuel component suggests that the bias comes from the terrestrial biospheric component. This could
320 be due to our VPRM-based estimates of the net fluxes being too negative as suggested by Oney
et al. (2016, in review), i.e., suggesting a too strong sink for central and southern Europe, or for our
model simulating a too small diurnal and/or seasonal rectification effect (Denning et al., 1995), i.e.,
a too small correlation between the time variations in the terrestrial exchange fluxes and atmospheric
transport/mixing. This deficiency does not impact our results much, since our focus will be on the
325 spatio-temporal variability of the fossil fuel CO₂ signal.

4 The spatiotemporal pattern of the fossil fuel CO₂

4.1 The spatial pattern

In the annual mean, computed from data from all times of the day, the fossil fuel component of
atmospheric CO₂ in the surface layer (~10 m above ground) amounts to more than 10 ppm across
330 wide swaths of central Europe (Fig. 4a). In large metropolitan areas, such as in western Germany
(Ruhrgebiet), Berlin, London, Paris, and Milan, the annual mean fossil fuel component exceeds
even 30 ppm. To first order, the distribution of the surface fossil fuel CO₂ reflects the distribution
of the emissions (see Fig. 1), suggesting a somewhat limited efficiency of atmospheric transport



and mixing to disperse the signal laterally. In mountainous regions this is clearly a consequence of
335 topographic constraints; elsewhere this is largely a result of the strong spatial gradients in emissions,
which remain conserved in the annual mean due to the overall diffusive nature of the dispersion.
Nevertheless, a substantial amount of the emitted CO₂ is being transported away, leading to a size-
able fossil fuel CO₂ signal extending far into the oceans surrounding Europe, especially the North
Sea.

340 Despite this lateral transport, the relatively good conservation of the spatial gradients in emissions
sets our results distinctly apart from previous studies, where the fossil fuel CO₂ signal was modeled
to be very smooth in space and on average also substantially smaller. For example, compared to the
results obtained with the medium-resolution (0.5°) Regional Model (REMO) (Peylin et al., 2011),
one can detect in our simulations nearly all major metropolitan regions and other fine-scale features,
345 such as individual fossil-fuel fired power plants (e.g., in eastern Germany). This is primarily the
result of the high horizontal and vertical resolution of COSMO permitting this model to conserve
the spatial gradients well. This good conservation is particularly well illustrated when considering
snapshot distributions of the fossil fuel CO₂ for individual seasons (Fig. 5). This figure also shows
the strong impact of the transport and dilution by the diurnal variations of the planetary boundary
350 layer, whose impact is particularly strong in summer.

For much of Europe, the fossil fuel component is the dominant contributor to the spatial gradients
in annual mean atmospheric CO₂ (Fig. 4b-d). In many places it accounts for nearly all of the spatial
gradients, with the contribution of the background and the terrestrial biospheric component being
substantially smaller. The latter shows gradients up to 10 ppm (Fig. 4c), while the background signal
355 does not exceed a few ppm (Fig. 4d). In the big cities, the fossil fuel CO₂ component represents
even a sizeable fraction (10%) of the total CO₂ concentration. This dominance of the fossil fuel
component together with its highly patterned nature owing to the many point sources leads to a
hotspot pattern in the near surface map of total atmospheric CO₂ over much of Europe (Fig. 4b).
However, due to lower emissions in southwestern Europe, the fossil fuel CO₂ signal is less strikingly
360 visible there compared to central Europe, while the biospheric signal is stronger. This results in a
relatively uniform spatial pattern of atmospheric CO₂ across Europe (Fig. 4b). Also the relatively
low CO₂ concentrations in the mountain regions, such as the Alps, Apennines, Pyrenees and central
France, reflect the much lower contribution from the fossil fuel component.

Naturally, when investigating the column averaged dry air mole fractions (X_{CO_2}), i.e., the prop-
365 erty typically measured by remote sensing from a satellite, the annual mean gradients of the fossil
fuel component are much smaller than those seen at the surface (see Fig. 6a). This is a consequence
of the lateral gradients being much weaker aloft, owing to a more effective transport and mixing.
As a result, most of the hotspot nature seen in the surface concentration pattern is blurred in X_{CO_2} .
Also the magnitude of the signal is much weaker. While the surface signal of the fossil fuel CO₂
370 signal amounted to more than 30 ppm in strong emissions regions, the signal in the column aver-



aged annual mean XCO_2 hardly exceeds 2 ppm. The impact of the predominant westerly air-flow becomes much more obvious in the column averaged dry air mole fraction XCO_2 , with the fossil fuel component revealing a clear eastward increase that is substantially stronger than the gradient in the underlying emissions.

375 The relative dominance of the fossil fuel component over the other components of atmospheric CO_2 is much weaker when considering the column averaged dry air mole fraction of CO_2 (see Fig. 6b-d). As a result, the total XCO_2 is made up of three relatively equally sized contributions, with the fossil fuel CO_2 signal continuing to dominate the XCO_2 variations in the major metropolitan areas. Contrary to the annual surface pattern, where CO_2 tends to increase eastward, the highest
380 XCO_2 are found in southwestern Europe with a trend toward lower values going eastward. This is partly a consequence of the lateral boundary conditions for atmospheric CO_2 , which tend to lead to the advection of high background CO_2 into the domain from the southwest. But the most important reason is the strong negative terrestrial biosphere signal over Europe, reflecting the sizeable carbon sink in European forests in the last decade (Reuter et al., 2016, in press). Interestingly, the relatively
385 uniform negative distribution for XCO_2 contrasts with a more patterned biospheric signal in the lowest atmosphere (Figure 4c), where the strong negative signal is restricted to central Europe, while much of southern Europe has a positive annual mean biospheric signal. The likely reason for this difference is the biospheric rectification effect (Denning et al., 1995), which tends to lead a vertical redistribution of CO_2 , i.e. positive values in the lower atmosphere and negative ones aloft.
390 In most of Europe, this rectification signal is relatively small in comparison to the annual mean biospheric component, so that the latter determines the overall signal. But in southern Europe, where the biospheric fluxes tend to be smaller in magnitude and in the annual mean to be near zero, the rectifier effect can dominate, explaining the positive signals in the surface layer (Figure 4c) and simultaneously the negative signals when the biospheric signal is integrated vertically (Figure 6c).

395 4.2 The temporal variability

The temporal variability of the fossil fuel CO_2 signal at the surface is very large, leading to a standard deviation around the annual mean of 30 ppm or more in the hotspot regions (Figure 7a). These hotspots correspond largely to the regions of highest emissions (Figure 1). But this high variability is not only a result of the temporal variability of the emissions, but arises also from the interaction of
400 variability in atmospheric transport and mixing with the strong lateral gradients seen in the snapshot figures (see Fig. 5).

A similar pattern of variability is seen in surface atmospheric CO_2 (Figure 7b), suggesting that the fossil fuel CO_2 is a major determinant not only of the annual mean spatial distribution of atmospheric CO_2 , but also of its temporal variability. This is confirmed by Figure 8a, which shows the relative
405 contribution of the fossil fuel CO_2 signal to the temporal standard deviations of atmospheric CO_2 . In many places, particularly in Europe's major metropolitan areas, but also in many urban areas



across Europe, the fossil fuel signal dominates the variability in atmospheric CO₂. But the high fossil fuel contribution is not limited to the urban areas. In fact, the region delineated by having a 50% contribution or more extends over much of northern central Europe, including the North Sea
410 (see Fig. 8a).

In order to better understand the origin of the strong variability, we decomposed the variability into seasonal, synoptic and diurnal contributions. The seasonal variation component was derived by averaging the data on a monthly basis and by subtracting the annual mean. The synoptic component was then computed by subtracting from the data the time series of the monthly means and then
415 forming daily averages of these deseasonalized data. Finally, the diurnal variability was derived by subtracting the seasonal and synoptic components from the data.

This decomposition reveals that the contribution of the fossil fuel CO₂ to the total variability of atmospheric CO₂ varies greatly depending on the temporal scale considered (Figure 8). While the fossil fuel contribution is comparably small on seasonal timescales (Figure 8b), the contribution on
420 synoptic and particularly on diurnal timescales is actually very large, exceeding 60% across nearly the entire northern part of central Europe (Figure 8c-d). The small contribution on the seasonal timescales is the result of the dominance of the seasonal cycle of the biospheric fluxes in most of Europe. An exception are a few places in northern Europe and in the very south of our European domain. We interpret this to be caused primarily by the relatively strong seasonality of the fossil fuel
425 emissions in these regions, owing to the strong summer-time requirement for cooling in the south and the strong winter-time demand for heating in the north.

The pattern of the fossil fuel contribution on synoptic timescales is very similar to that of the total contribution, meaning its contribution is one of the dominant contributions to the total temporal variability. This is consistent with synoptic variations also being among the strongest contributors to
430 atmospheric variability, owing to baroclinic waves and frontal systems being formed out of the strong baroclinicity that characterize the mid-latitudes. These synoptic weather events transport the emitted CO₂ also quite efficiently outside the main metropolitan areas, explaining the widespread signal of the fossil fuel contribution to the total variance of atmospheric CO₂. Even larger than the fossil fuel contribution to synoptic variability is the contribution on the diurnal timescale, with the fossil fuel
435 CO₂ contributing more than half of the variability over most of Europe. This high variability comes from the interaction of the diurnal variability of the fossil fuel emissions, with the strong diurnal variability of atmospheric transport, particularly the diurnal mixing of the planetary boundary layer. This co-variability between fossil fuel emissions and atmospheric transport exceeds that between the biospheric fluxes and atmospheric transport over the entire year, owing to the latter fluxes being large
440 and relevant only during the spring/summer period, while the fossil fuel emissions are relatively high during most of the months of the year, particularly close to the sources.



5 Discussion

The analyses of the results raise a number of questions that we would like to discuss next. First, why is the diurnal variability so high, and in particular, what is the contribution of our consideration of diurnal (and seasonal) variations in CO₂ emissions on the simulated fossil fuel CO₂ signal? Further, is there an impact beyond the variability, e.g., on the mean fossil fuel CO₂ signal? Second, what is the contribution of the various sectors on the fossil fuel CO₂ signal and in what way do emissions from one country influence the fossil fuel CO₂ signal in another country? Third, how can we use the insights gained from the study of the fossil fuel CO₂ signal to develop optimal strategies for detecting changes in fossil fuel CO₂ emissions? We discuss each of these three questions next.

5.1 The impact of variations in fossil fuel emissions on atmospheric CO₂

In order to elucidate the role of the temporal variations in fossil fuel emissions on the fossil fuel CO₂, we contrast the results of our standard simulation with time-varying emissions with those where the fossil fuel emissions were kept constant over time. The annual emissions are identical for the two cases, but the time constant case has, on average, considerably higher emissions in summer and at night.

The contrast between these two cases shows only a small change in the high diurnal variability of atmospheric CO₂ seen in Figure 8d, implying that the contribution of the diurnal variations in fossil fuel emissions is less important than other factors (results not shown). The largest contributions are found around some of the large metropolitan areas (e.g., London, Paris, Milan), but they do not exceed 10%. Thus the majority of the diurnal variability in the fossil fuel CO₂ stems from the diurnal variations in atmospheric transport and mixing acting on the strong horizontal gradients in emissions.

While not contributing much to the diurnal variability in the fossil fuel CO₂, the consideration of the time-varying emission matters quite substantially for the annual mean distribution of the fossil CO₂ signal. Figure Fig. 9a reveals that the annual mean fossil CO₂ signal in the simulation with time varying emissions is substantially lower over wide swaths of Spain, Italy, the Benelux countries, (western) Germany and the UK compared to the simulation where fossil fuel emissions were kept constant. The strongest negative signals are found close to the strongest emitters in these countries, with magnitudes exceeding several ppm. But the magnitude of the signal does not correspond to the magnitude of emissions, since regions with comparably low emissions such as Spain, have signals that are as large as those in high emission regions of the Netherlands. The relatively large signals in southern Europe are likely due to the stronger PBL dynamics in these regions throughout the year in comparison to central and northern Europe. Some regions also have a positive signal from the time-varying emissions, such as parts of France and northeastern Germany. Thus the interaction between the variations in fossil fuel emissions and the variations in atmospheric transport and mixing leads



to a substantial net signal in atmospheric CO₂, even though the total emissions in both cases are identical.

This net signal represents a fossil fuel-driven rectification effect (Zhang et al., 2016) in analogy
480 to the rectification effect associated with the terrestrial biosphere (Denning et al., 1996; Larson and
Volkmer, 2008), i.e., a signal that is due to the co-variance of emissions and atmospheric trans-
port/mixing. Its (mostly) negative sign emerges from the fact that when the emissions are large, e.g.,
during the day, the transport and mixing away from the surface is strong, diluting the fossil fuel
signal in atmospheric CO₂. In contrast, when the emissions are small, e.g., during the night, the
485 transport and mixing tends to be weak. Taken together, this results in a more efficient dilution of the
emissions in the time-varying emission case compared to the time- invariant case, thus explaining
the mostly negative sign of the fossil fuel rectification effect.

This explanation is supported by the mostly positive correlation between the height of the plan-
etary boundary layer (PBL) and the fossil fuel emissions, since the height of the PBL is a direct
490 measure of the magnitude of the dilution in the lowest levels of the atmosphere (Figure Fig. 9b).

But there are a number of notable exceptions. For example, wide swaths of northeastern Germany
and Poland and some places in central France have a positive rectification signal. Further, there are
places where the co-variation of fossil fuel emission and the PBL is negative, yet the fossil-fuel rec-
tification effect is still negative (e.g. the Ruhr valley region in western Germany), suggesting that our
495 explanation does not cover all aspects. In response, one first needs to recognize that not only PBL
but also other temporally varying phenomena, such as local atmospheric circulation patterns (e.g.
mountain winds, sea-breezes) can lead to co- variability between emissions and transport/mixing,
creating a rectification signal that can differ in sign. The contribution of the sea-breeze can be iden-
tified quite clearly by the strong negative sign along most of the coastline between southern Europe
500 and the Mediterranean. Second, the local timing between the growth and decay of the PBL and the
emissions can be quite different, owing in part, to the substantially different time functions for the
different emission categories and their different local contributions (Figure 1). For example, in re-
gions with a large contribution from road transportation, the local emissions have a strong peak in the
early morning hours, when the PBL is still shallow, leading to a high signal there, while emissions
505 are lower when the PBL is at its maximum in the early afternoon. This would create a positive rec-
tification signal. Finally, in certain places, also the seasonal rectification appears to play a role, i.e., the
seasonal co-variations of the emissions with the PBL height. In fact, in many places the magnitude
of the correlation between emission and PBL height on seasonal timescales exceeds that on diurnal
timescales. This seasonal variation is particularly large for residential heating, which is maximum in
510 winter when the PBL is low, leading to a positive seasonal rectification. This effect likely contributes
to the negative correlations between emissions and PBL height in large urban centers such as Paris
(Figure Fig. 9b). We suspect that such seasonal effects are also the primary reason for the positive
rectification signal in northeastern Germany and northern Poland. In southern Europe, these seasonal



co- variations tend to lead to a negative fossil-fuel rectification effect, since the emissions peak in
515 summer (Figure 2b), when the PBL height is at its seasonal maximum.

The magnitude of the fossil fuel rectification effect is smaller than the rectifier effect induced by
the exchange fluxes with the terrestrial biosphere (Zhang et al., 2016), but still quite substantial.
Thus, the fossil fuel rectification effect clearly needs to be taken into consideration when model-
ing the atmospheric fossil fuel CO₂ signal, highlighting the need to use and apply accurate time
520 functions. Our results thus clearly support the results of Nassar et al. (2013), who demonstrated the
substantial impact of the consideration of time-varying emissions on atmospheric CO₂. We extend
their result by demonstrating an effect on the annual mean fossil fuel CO₂, suggesting that special
attention needs to be given to the relative timing of variations in atmospheric transport and mixing
and fossil fuel emissions. Our results confirm the recent findings by Zhang et al. (2016) who demon-
525 strated the fossil fuel rectification effect for the first time in a global model. Their signal is locally
smaller than ours, owing to their using a much coarser resolution model, but they also show that the
sign of the fossil fuel rectification effect tends to vary between timescales, with the diurnal being pri-
marily negative, while the seasonal rectification effect being positive. This supports our explanation
for the positive signals in northeastern Germany and northern Poland.

530 5.2 Fossil fuel CO₂ signal from different sources

Near the surface, the fossil fuel emissions from a particular region create a distribution that stays
mostly within the region of origin (see Fig. 10 a,b). The fossil fuel CO₂ is highly concentrated near
the localized areas of high emissions and then drops off quickly by distance with an e-folding spatial
scale of a few hundred kilometers. As a result, the fossil fuel signal tends to be relatively small
535 outside the region of origin, rarely exceeding 1 ppm in contrast to the > 20 ppm signal close to the
sources. The different magnitudes of the fossil fuel CO₂ signals from different regions largely reflects
the total emissions, but also the emission intensity, i.e., the emission per unit area. For example, with
a total emission of 0.59 Pg CO₂ yr⁻¹, Germany is the biggest source of fossil fuel CO₂ within
Europe, nearly double that of the second biggest emitter, i.e., France, yet Germany is almost half the
540 size of France, resulting in a considerably higher emission intensity over Germany.

A different picture emerges when considering XCO₂, i.e., the column averaged dry air mole
fraction CO₂. After having been transported aloft, where the fossil fuel signal can be much more
readily dispersed, the imprint of the emissions of any particular region to the fossil fuel CO₂ within
another region is actually quite large (Figure 10 c,d). In a small country, such as Switzerland, only
545 20% of the fossil fuel signature in XCO₂ above its territory stems from emissions within, while the
contribution of Germany alone is 21% and that of France 18% (Figure 10). A similar distribution of
sources is found for other small countries, such as Austria. In contrast, the fraction of the territorial
emissions to the total fossil fuel signal is quite a bit larger for large countries/regions, such as France
or Germany. In the latter case, more than 50% of its total fossil fuel CO₂ signal stems from emissions



550 within, with 4 countries contributing most of the remainder. The countries/regions with high overall
emissions contribute, of course, also most strongly to the fossil fuel CO₂ signal in other countries,
with Germany contributing 18% to the signal in France, 11% to that in Italy and 20% to that in the
Netherlands. Owing to its lower total emissions, France just contributes 9% to the signal in Germany
and 8% to that in Italy. Thus, as is the case with classical air pollution, the fossil fuel CO₂ does not
555 stop at the national borders, but extends to continental scales.

Among all the processes, the CO₂ emissions from power plants dominate the fossil fuel distribu-
tion, with concentrations reaching up to 16 ppm in the northern part of the domain (see Fig. 12).
The point-source nature of this emission sector is clearly visible in the surface distribution, as is
the spatially distinct distribution owing to the large differences in power production in the different
560 countries of central Europe. While France has very few fossil-fuel fired power plants as a result
of its high reliance on nuclear and hydroelectric power plants, Germany, Italy, the Netherlands and
Poland rely strongly on coal- and gas-fired power plants for their electricity production. This leads
to a highly heterogeneous fossil fuel CO₂ signals of the power plant sector. In total, this sector con-
tributes 31.8% to the total fossil fuel CO₂ signal in central Europe, which is slightly smaller than
565 its contribution to emissions (32.8%). This small difference emerges from the somewhat stronger
loss of the signal across the lateral boundaries from this sector relative to the signal from the other
sectors.

The second largest fossil fuel CO₂ signal is generated by the emissions from the road transporta-
tion sector (22.0%) (Fig. 12d), with this share actually being somewhat larger than its share in total
570 emissions (21.1%). The transportation sector signal is very smooth, owing to the distributed nature
of the emissions from this sector (see also Fig. 1).

The CO₂ signal from the industrial and residential sectors are more granular than that from the
transportation sector, but still not as distinct as the power plant sector, as there are less country
specific policies impacting the CO₂ emissions from these sectors. The emissions and consequently
575 the CO₂ signal largely follow population density. The residential sector (mostly heating) contributes
18.1% to the total fossil fuel signal in atmospheric CO₂, slightly larger than the emissions from
the industrial sector (17.4%). These two shares in the signal very nearly reflect their shares in total
emissions. The emissions from the ‘other’ sectors (e.g., shipping, waste incineration, etc) is smaller,
in comparison (10.7%), but not negligible.

580 The relative contribution of the emissions from the different sectors to the fossil fuel CO₂ vary
strongly by region (Fig. 13). Clearly, close to major fossil-fuel fired power plants, this sector domi-
nates, but elsewhere, any of the four major sources can take the leading role. For example, in Switzer-
land, Paris, and London, the emissions from the residential sector dominate the signal, while over
much of southern and western Europe, the transportation sector dominates. The industrial sector
585 dominates the signal in a few hotspot areas, where its emissions are high, but where there is no
major fossil fuel fired power plants nearby.



These high spatial variations in the relative contribution puts the findings of Vogel et al. (2013) into a spatial context, as they reported for the Heidelberg site a dominance for emissions from power plants (28%), while the transportation sector contributed only 15%. This is a typical value for much of western Germany, reflecting the relative contribution of the different emission sectors (see also Fig. 1). But the contributions are very different, for example, for the CarbCount CH sites in Switzerland (Oney et al., 2015). At Beromünster, the transportation sector dominates over the other sectors, with nearly 70% stemming from this sector alone, while the contribution from power plant emissions is very low at this site, since Switzerland does not operate any fossil fuel power plants.

These large differences in the relative contribution from the different emission sectors have major implications for the analysis of the fossil fuel CO₂ and how it may change in response to mitigation measures. For example, these large differences will lead to substantial spatial gradients in the CO to CO₂ ratio in the fossil fuel signal, as the different emission sectors have very different CO to CO₂ emission ratios. Since CO is often used to identify the fossil fuel component from atmospheric CO₂ observations, these variations need to be carefully disentangled in order to properly diagnose the fossil fuel component. The strong variations in the contributions from the different sectors thus adds a substantial amount of uncertainty to the CO method (Oney et al., 2016, in review; Vardag et al., 2015). A second consequence concerns the detection of changes in emissions from the different sectors. Thus, with the transportation sector contributing little to the very large fossil fuel signal in much of the northeastern part of our domain, reductions in this sector will be difficult to discern in that region. In contrast, the high relative contribution of the transportation sector to the total signal in southwestern Europe makes it actually quite feasible to detect mitigation measures in this sector in that part of Europe, even though the overall signal might not be that high.

An important caveat of our simulations is the fact that the effective height of the emissions above surface was not considered, but rather all CO₂ was released into the lowest model level. As a consequence, the surface CO₂ signals from elevated stack emissions from power plants and residential heating are likely biased high relative to those from the transportation sector. Given the large contribution from power plant emissions, it will be important to accurately consider the effective emission height (including plume rise) in future simulations, a point that was also raised by Vogel et al. (2013).

5.3 The response of atmospheric CO₂ to an emission reduction

According to their intended nationally determined contributions filed with the United Nations Framework Convention on Climate Change (UNFCCC) in late 2015, the European Union and its member states have agreed to a binding target of an at least 40% domestic reduction in greenhouse gases emissions by 2030 compared to 1990 (<http://www4.unfccc.int/Submissions/INDC/Published%20Documents/Latvia/1/LV-03-06-EU%20INDC.pdf>). A major question driving international policy making is to what degree such a reduction can be verified through independent means, such as through the monitoring of atmospheric CO₂ (Ciais et al., 2014, 2015). To address this question, we conducted several



sensitivity experiments to investigate how various reductions in the magnitude and types of emissions affect not only the annual mean fossil fuel CO₂ signal, but also its variability. The goal is to
625 determine whether reduced fossil fuel emissions might be detectable by current and future observing systems, especially satellites.

Since CO₂ is a conservative tracer in the atmosphere at the time scales considered here, a uniform reduction in the emissions leads to a uniform and directly proportional reduction of its current distribution, i.e., a 30% reduction of total fossil fuel emission would simply lead to a 30% reduction of the fossil fuel CO₂ signal at the surface (Fig. 4a) and throughout the atmospheric column (Fig. 6a). Concretely, the fossil fuel CO₂ would be reduced by more than 4 ppm near the surface for vast stretches of central and northern Europe, with maximum reductions of 10 ppm or more in the emission hotspots (Figure 14a). This contrasts with the reduction in the averaged column annual mean X_{CO_2} amounting to just over 0.2 ppm in the regions where the surface decreases by 4 ppm
635 or more (Figure 14b). A reduction of 0.5 ppm is reached in just a few isolated locations, generally characterized by a high density of point sources, primarily fossil-fuel fired power plants. Thus, given current measurement accuracies of better than 0.1 ppm for a ground-based observing network (Zellweger et al., 2016), a 30% reduction in the fossil fuel emissions is fundamentally easily detectable for such a system, although one needs to bear in mind the non-trivial task to separate the signal from
640 the background variability. In contrast, such a reduction in the fossil fuel emissions is not trivial to detect by satellite observations for most regions (except around the big power plants) as it is very challenging to obtain and maintain accuracies better than 0.5 ppm by current space-based observing systems (Buchwitz et al., 2015). Furthermore, such high accuracies are only achieved when the data are averaged over large scales, i.e., order of 1000 km or more. Nevertheless, taking 0.5 ppm as the
645 threshold for detection within a single pixel, a 30% reduction in fossil fuel emissions thus appears to be beyond the detectability, except for a few hotspot regions (Figure 14b). Even a 50% reduction would not be trivial to detect for a satellite-based system on the basis of changes in the column averaged dry air mole fraction .

Given these challenges, a potentially attractive second avenue for determining changes in fossil
650 fuel emissions is the reduction in temporal variability of atmospheric CO₂ that goes alongside the reduction in the mean signal. This is particularly promising given the very high contribution of the fossil fuel CO₂ signal to the variability in atmospheric CO₂ (see Fig. 8). As is the case for the mean, the conservative nature of atmospheric CO₂ implies that a uniform reduction of the emissions will lead to a uniform and proportional reduction of the variability of the fossil fuel signal as well.
655 However, this is not the case for the variability in total atmospheric CO₂, since co-variations between the fossil fuel signal and the signal from e.g., the terrestrial biosphere can lead to non-linear effects. For example, a negative correlation between the two components would lead to a situation where the variability of atmospheric CO₂ was smaller than that of the individual components. In such a case, a reduction of the fossil fuel emission would lead to a smaller decrease in variability than expected.



660 If the two components were positively correlated, the opposite would occur, i.e., the variability in atmospheric CO₂ would decrease more than expected.

Near the surface, the reduction in the temporal standard deviation and in the mean have nearly the same amplitude for most places (Figure 14c). This makes the analysis of changes in the temporal variability indeed an attractive option to enhance the detectability of changes in fossil fuel emissions.

665 This is much less the case for the annual mean XCO_2 , where the standard deviation changes are in general much smaller than the changes in the mean, with just a few isolated places revealing changes in the standard deviation of 0.5 ppm or more that might be discerned by the current generation of satellites.

But in these isolated places, the analysis of the temporal variability might be an interesting option
670 even for satellite-based measurement systems (Fig. 15). In those places, indicated by the green circles in (Figure 14c), the changes in the temporal standard deviation are very large. Even for changes in emissions of around only 30%, the changes would be detectable for current satellites (Fig. 15). But the number of such sites is very low across Europe, making this not a general, but rather a specialized option.

675 The detection challenge is not simpler for other potential emission reduction scenarios, as outlined, for example in the EU roadmap (http://ec.europa.eu/clima/policies/strategies/2050/index_en.htm). A 50% reduction in the emissions from power plants alone (representing a reduction of the overall emissions by 16%), results in the mean surface concentration of atmospheric CO₂ going down by more than 2 ppm over large parts of northwestern Europe, following the pattern of the
680 surface signal of this sector (cf. Fig. 12a). Alongside we find a substantial reduction of the standard deviation of surface atmospheric CO₂ by more than 2 ppm in these regions, with the hotspots of power plant emissions seeing a reduction in the standard deviation of atmospheric CO₂ of 5 ppm or more. The reduction of the average annual mean column XCO_2 is much smaller than that of atmospheric CO₂ at the surface, amounting to little more than 0.2 ppm over wide swaths of northern Europe. The maximum reductions are of the order of 0.5 ppm in the proximity of large clusters
685 of fossil-fuel fired power plants, i.e., generally too small to be detected. But, in these regions, the changes in the variability in XCO_2 is quite high, making this method again potentially attractive for detecting changes. In fact, in several regions, including some major cities, a 19% reduction of the fossil fuel emissions would result in a change of more than 0.5 ppm in the standard deviation, i.e.,
690 above detection level. This thus supports the findings of Pillai et al. (2016) that changes in fossil fuel emissions are fundamentally detectable over major cities or major point sources, but it also shows that this detection is very challenging.

The signals get even more difficult to discern if the emission reductions occur in individual sectors other than the power plants. For example, detectable signals by current generation satellites occur
695 only if industrial emissions are cut by more than 80% or if residential emissions are cut by more than 90%. Also country level emissions are not trivial to be clearly detected. A reduction in Germany by



50% is potentially detectable by current satellites, with a maximum reduction of $X\text{CO}_2$ by 0.95 ppm. For most other countries, however, a 50% reduction in emissions is difficult to be detected.

All the analyses here relied on using the model output on all available days, i.e., we assumed
700 perfect temporal coverage. This is overly optimistic, since cloud cover and other complicating factors (e.g., aerosol layers) will cause the coverage to decrease considerably, complicating the detection.

But regardless of this additional challenge, there is much additional information contained in high frequency observations of atmospheric CO_2 . As we demonstrated above, the temporal variations are potentially highly useful for detecting fossil fuel emissions changes from various sources, especially
705 those with a strong spatial granularity such as power plants or individual cities. For a routine monitoring of strong point sources, Velazco et al. (2011) therefore proposed a constellation of 5 satellites of type CarbonSat that combine imaging capability with a relatively wide swath (Bovensmann et al., 2010). Such a constellation would offer daily global coverage, though the presence of clouds would reduce the effective coverage considerably. As the precision and accuracy of satellite retrieved $X\text{CO}_2$
710 will improve in the future, that minimum change will go down as well.

6 Summary and conclusions

We have investigated the fossil fuel signal in atmospheric CO_2 over southern and central Europe using a regional high-resolution atmospheric model forced with temporally and spatially highly resolved variations in the fossil fuel emissions. The assessment of the modeled atmospheric CO_2 with
715 in-situ measurements across multiple sites across Europe reveals good to excellent agreement on all timescales considered with biases of less than 1 ppm, with the exception of the tall tower site Hegyhatsal in central Hungary. The model is also able to capture the reconstructed fossil fuel component at two sites quite successfully. Although the model tends to underestimate the amplitude of the daily averaged fossil fuel CO_2 in winter, the simulation matches fossil fuel CO_2 from both sites very well
720 most of the time, revealing the high quality of the transport model and reasonable time profiles of the fossil fuel emissions used as input.

Over much of Europe, the fossil fuel CO_2 is a dominant component of the spatial variability of atmospheric CO_2 , particularly near the surface. In some places, it even contributes significantly to the total (including background) CO_2 , up to 110% in large urban centers and power plant plumes.
725 Also the contribution to the temporal variability is very substantial. Fossil fuel CO_2 makes a particularly large contribution at synoptic and diurnal time scales whereas the seasonal variability is dominated by biospheric activity. The influence is not only large over the hot spot regions of fossil fuel emissions, but also over large areas downstream. In case of diurnal variability, fossil fuel CO_2 is the dominant component over wide areas of northern and western Europe.

730 Temporal variability of the emissions has a non-negligible influence on annual mean fossil fuel CO_2 mole fractions near the surface, due to diurnal and seasonal rectifier effects. Differences be-



tween annual mean values with temporally variable and constant emissions can be up to a few ppm in the hot spot regions, but are mostly below 1 ppm elsewhere. This implies that temporal variability of fossil fuel emissions needs to be accurately represented for realistic simulations, confirming the results of Zhang et al. (2016). It is also important for reliably detecting fossil fuel emission changes from specific sources since different sources have different temporal profiles.

Simulating fossil fuel emissions from different countries and sectors suggests that the major part of the signal near the surface remains in the country of origin. Ground-based in situ observations are thus most sensitive to fossil fuel emissions from the country where they are located. A different picture emerges for column averaged dry air mole fractions (X_{CO_2}) as measured by satellites, for which the signal is much more dispersed. Only over Germany, the contribution from emissions within the country is larger than 50%, whereas over France the signal from neighboring countries dominates (66%). An important reason for these contrasting results seems to be the differences in electricity production, which mostly relies on nuclear power in France but on fossil fuels in neighboring countries including Germany, UK and Italy. Over small countries such as Switzerland or the Netherlands, the contribution from abroad is typically the dominating component. Among all the processes, fossil fuel emissions from power plants contributes the most (approx. one third) to the total fossil fuel signal of CO_2 both at the surface and in the column. However, the power plant signal at the surface is likely overestimated in our simulations, since all emissions were released into the lowest model level without considering the true elevation of the source. The signal from power plant emissions has a pronounced and distinct spatial pattern that provides us an opportunity to discern changes in from power plant emissions from changes in other sources.

Based on a number of sensitivity studies, we show that reductions in fossil fuel emissions not only leave a distinct signal in the time mean distribution of atmospheric CO_2 , but also in its temporal variability. This opens potentially additional ways to detect and verify emission reductions. But this opportunity exists primarily for surface based measurement networks, while the satellite based systems that measure the column-averaged X_{CO_2} will see too small changes, in general, relative to their current measurement capabilities. An important exception are a few hotspot sites, where the satellites will be able to detect fairly modest changes of about 30% when assuming an accuracy of the satellite observations of 0.5 ppm.

As both satellite and surface measurements have advantages and disadvantages, combining surface measurements with satellite data and increasing the frequency and coverage of the latter will be the optimal path forward to enhance the possibility of detecting future changes in fossil fuel emissions.

Acknowledgements. This study was funded by the Swiss National Foundation (SNF) as part of the CarboCount CH Sinergia Project (Grant Number: CRSII2 136273). Additional funding was provided by ETH Zürich (YL and NG) and Empa (DB). We thank the Center for Climate Systems Modeling (C2SM) and especially Anne



Roches for providing extensive support. We acknowledge MeteoSwiss for the provision of their operational
COSMO analysis products. The computations were done on the supercomputers of the Swiss National Super-
770 computing Centre (CSCS). We are very thankful to the principal investigators responsible for the Mace Head
(MHD) station in Ireland, for the SNO-ICOS-France station Puy de Dome (PUY) in France, for the Hegyhatsal
station (HUN) in Hungary, and for the Cabauw station (CBW) in the Netherlands for making their data publi-
cally available. We are also grateful to Dr. Sander van Der Laan and Dr. Felix Vogel for sharing their estimates
of the fossil fuel component at Lütjewed and Heidelberg, respectively. Furthermore, we are very grateful to
775 Prof. Wouter Peters, Wageningen University, for providing the CO₂ global model boundary conditions and to
Dr. Christoph Gerbig for providing the VPRM biospheric flux data.



References

- Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational convective-scale numerical weather prediction with the COSMO model: description and sensitivities, *Mon. Weather Rev.*, 139, 3887–3905, doi:http://dx.doi.org/10.1175/MWR-D-10-05013.1, 2011.
- 780 Basu, S., Miller, J. B., and Lehman, S.: Separation of biospheric and fossil fuel fluxes of CO₂ by atmospheric inversion of CO₂ and ¹⁴CO₂ measurements: Observation System Simulations, *Atmos. Chem. Phys.*, 16, 5665–5683, doi:10.5194/acp-16-5665-2016, 2016.
- Bovensmann, H., Buchwitz, M., Burrows, J., Reuter, M., Krings, T., Gerilowski, K., Schneising, O., Heymann, J., Tretner, A., and Erzingler, J.: A remote sensing technique for global monitoring of power plant CO₂ emissions from space and related applications, *Atmos. Meas. Tech.*, 3, 781–811, doi:10.5194/amt-3-781-2010, 2010.
- 785 Bozhinova, D., Van Der Molen, M. K., Van Der Velde, I. R., Krol, M. C., Van Der Laan, S., Meijer, H. A. J., and Peters, W.: Simulating the integrated summertime $\Delta^{14}\text{CO}_2$ signature from anthropogenic emissions over Western Europe, *Atmos. Chem. Phys.*, 14, 7273–7290, doi:10.5194/acp-14-7273-2014, 2014.
- 790 Bréon, F., Broquet, G., Puygrenier, V., Chevallier, F., Xueref-Remy, I., Ramonet, M., Dieudonné, E., Lopez, M., Schmidt, M., Perrussel, O., et al.: An attempt at estimating Paris area CO₂ emissions from atmospheric concentration measurements, *Atmos. Chem. Phys.*, 15, 1707–1724, doi:10.5194/acp-15-1707-2015, 2015.
- Brioude, J., Angevine, W. M., Ahmadov, R., Kim, S.-W., Evan, S., McKeen, S. A., Hsie, E.-Y., Frost, G. J., Neuman, J. A., Pollack, I. B., et al.: Top-down estimate of surface flux in the Los Angeles Basin using a mesoscale inverse modeling technique: assessing anthropogenic emissions of CO, NO_x and CO₂ and their impacts, *Atmos. Chem. Phys.*, 13, 3661–3677, doi:10.5194/acp-13-3661-2013, 2013.
- 795 Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenstock, T., et al.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): Comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, *Remote Sens. Environ.*, 162, 344–362, 2015.
- 800 Ciais, P., Gasser, T., Paris, J. D., Caldeira, K., Raupach, M. R., Canadell, J. G., Patwardhan, A., Friedlingstein, P., Piao, S. L., and Gitz, V.: Attributing the increase in atmospheric CO₂ to emitters and absorbers, *Nat. Clim. Chang.*, 3, 926–930, doi:10.1038/nclimate1942, 2013.
- 805 Ciais, P., Dolman, A., Bombelli, A., Duren, R., Peregón, A., Rayner, P., Miller, C., Gobron, N., Kinderman, G., Marland, G., et al.: Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11, 3547–3602, doi:10.5194/bg-11-3547-2014, 2014.
- Ciais, P., Crisp, D., Van Der Gon, H., Engelen, R., Heimann, M., Janssens-Maenhout, G., and Scholze, M.: Towards a European operational observing system to monitor fossil CO₂ emissions (Final Report from the expert group), European Commission, JRC98161, Brussels, 2015.
- 810 Denning, A. S., Fung, I. Y., Randall, D., et al.: Latitudinal gradient of atmospheric CO₂ due to seasonal exchange with land biota, *Nature*, 376, 240–243, doi:10.1038/376240a0, 1995.
- Denning, A. S., Randall, D. A., Collatz, G. J., and Sellers, P. J.: Simulations of terrestrial carbon metabolism and atmospheric CO₂ in a general circulation model, *Tellus B*, 48, 543–567, doi:10.1034/j.1600-0889.1996.t01-1-00010.x, 1996.



- D'Ortenzio, F., Antoine, D., and Marullo, S.: Satellite-driven modeling of the upper ocean mixed layer and air-sea CO₂ flux in the Mediterranean Sea, *Deep. Res. Part I Oceanogr. Res. Pap.*, 55, 405–434, doi:10.1016/j.dsr.2007.12.008, 2008.
- Feng, S., Lauvaux, T., Newman, S., Rao, P., Ahmadov, R., Deng, A., Díaz-Isaac, L. I., Duren, R. M., Fischer, M. L., Gerbig, C., et al.: Los Angeles megacity: a high-resolution land–atmosphere modelling system for urban CO₂ emissions, *Atmos. Chem. Phys.*, 16, 9019–9045, doi:10.5194/acp-16-9019-2016, 2016.
- 820 Friedlingstein, P., Andrew, R. M., Rogelj, J., Peters, G. P., Canadell, J. G., Knutti, R., Luderer, G., Raupach, M. R., Schaeffer, M., van Vuuren, D. P., and Le Quéré, C.: Persistent growth of CO₂ emissions and implications for reaching climate targets, *Nat. Geosci.*, 7, 709–715, doi:10.1038/ngeo2248, 2014.
- 825 Friedrich, R. and Reis, S.: Emissions of Air Pollutants, doi:10.1787/data-00598-en, 2004.
- Geels, C., Gloor, M., Ciais, P., Bousquet, P., Peylin, P., Vermeulen, A., Dargaville, R., Aalto, T., Brandt, J., Christensen, J., et al.: Comparing atmospheric transport models for future regional inversions over Europe–Part 1: mapping the atmospheric CO₂ signals, *Atmos. Chem. Phys.*, 7, 3461–3479, doi:10.5194/acp-7-3461-2007, 2007.
- 830 Graven, H. D. and Gruber, N.: Continental-scale enrichment of atmospheric ¹⁴CO₂ from the nuclear power industry: Potential impact on the estimation of fossil fuel-derived CO₂, *Atmos. Chem. Phys.*, 11, 12 339–12 349, doi:10.5194/acp-11-12339-2011, 2011.
- Hase, F., Frey, M., Blumenstock, T., Groß, J., Kiel, M., Kohlhepp, R., Mengistu Tsidu, G., Schäfer, K., Sha, M., and Orphal, J.: Application of portable FTIR spectrometers for detecting greenhouse gas emissions of the major city Berlin, *Atmos. Meas. Tech.*, 8, 3059–3068, doi:10.5194/amt-8-3059-2015, 2015.
- 835 Haszpra, L., Ramonet, M., Schmidt, M., Barcza, Z., Pátkai, Z., Tarczay, K., Yver, C., Tarniewicz, J., and Ciais, P.: Variation of CO₂ mole fraction in the lower free troposphere, in the boundary layer and at the surface, *Atmos. Chem. Phys.*, 12, 8865–8875, doi:10.5194/acp-12-8865-2012, 2012.
- Holtslag, A., Svensson, G., Baas, P., Basu, S., Beare, B., Beljaars, A., Bosveld, F., Cuxart, J., Lindvall, J., 840 Steeneveld, G., et al.: Stable atmospheric boundary layers and diurnal cycles: challenges for weather and climate models, *B. Am. Meteorol. Soc.*, 94, 1691–1706, doi:http://dx.doi.org/10.1175/BAMS-D-11-00187.1, 2013.
- Janssens-Maenhout, G., Dentener, F., Van Aardenne, J., Monni, S., Pagliari, V., Orlandini, L., Klimont, Z., Kurokawa, J., Akimoto, H., Ohara, T., et al.: EDGAR-HTAP: a harmonized gridded air pollution emission dataset based on national inventories, European Commission Publications Office, Ispra (Italy). JRC68434, EUR report No EUR, 25, 299–2012, doi:10.2788/14069, 2012.
- 845 Keller, E. D., Turnbull, J. C., and Norris, M. W.: Detecting long-term changes in point-source fossil CO₂ emissions with tree ring archives, *Atmos. Chem. Phys.*, 16, 5481–5495, doi:10.5194/acp-16-5481-2016, 2016.
- Keppel-Aleks, G., Wennberg, P. O., O'Dell, C. W., and Wunch, D.: Towards constraints on fossil fuel emissions 850 from total column carbon dioxide, *Atmos. Chem. Phys.*, 13, 4349–4357, doi:10.5194/acp-13-4349-2013, 2013.
- Kort, E. A., Frankenberg, C., Miller, C. E., and Oda, T.: Space-based observations of megacity carbon dioxide, *Geophys. Res. Lett.*, 39, 1–5, doi:10.1029/2012GL052738, 2012.



- 855 Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier Van Der Gon, H. A. C.: TNO-MACC-II emission inventory; A multi-year (2003-2009) consistent high-resolution European emission inventory for air quality modelling, *Atmos. Chem. Phys.*, 14, 10963–10976, doi:10.5194/acp-14-10963-2014, 2014.
- Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T. P., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793–7815, doi:10.5194/bg-10-7793-2013, 2013.
- 860 Larson, V. E. and Volkmer, H.: An idealized model of the one-dimensional carbon dioxide rectifier effect, *Tellus B*, 60, 525–536, doi:10.1111/j.1600-0889.2008.00368.x, 2008.
- Lauvaux, T., Miles, N. L., Deng, A., Richardson, S. J., Cambaliza, M. O., Davis, K. J., Gaudet, B., Gurney, K. R., Huang, J., O’Keefe, D., et al.: High-resolution atmospheric inversion of urban CO₂ emissions during the dormant season of the Indianapolis Flux Experiment (INFLUX), *J. Geophys. Res-Atmos.*, 121, 5213–5236, doi:10.1002/2015JD024473, 2016.
- 865 Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Liener, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., O’Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2016, *Earth System Science Data*, 8, 605–649, doi:10.5194/essd-8-605-2016, 2016.
- 875 Levin, I. and Karstens, U.: Inferring high-resolution fossil fuel CO₂ records at continental sites from combined ¹⁴CO₂ and CO observations, *Tellus B*, 59, 245–250, doi:10.1111/j.1600-0889.2006.00244.x, 2007.
- Levin, I., Hammer, S., Eichelmann, E., and Vogel, F. R.: Verification of greenhouse gas emission reductions: the prospect of atmospheric monitoring in polluted areas, *Philos. T. Roy. Soc. A.*, 369, 1906–1924, doi:10.1098/rsta.2010.0249, 2011.
- 880 Lindenmaier, R., Dubey, M. K., Henderson, B. G., Butterfield, Z. T., Herman, J. R., Rahn, T., and Lee, S.-H.: Multiscale observations of CO₂, ¹³CO₂, and pollutants at Four Corners for emission verification and attribution, *P. Natl. Acad. Sci. USA*, 111, 8386–8391, doi:10.1073/pnas.1321883111, 2014.
- Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., Gerbig, C., Munger, J. W., 885 Chow, V. Y., and Gottlieb, E. W.: A satellite-based biosphere parameterization for net ecosystem CO₂ exchange: Vegetation Photosynthesis and Respiration Model (VPRM), *Global Biogeochemical Cycles*, 22, doi:10.1029/2006GB002735, 2008.
- McKain, K., Wofsy, S. C., Nehrkorn, T., Eluszkiewicz, J., Ehleringer, J. R., and Stephens, B. B.: Assessment of ground-based atmospheric observations for verification of greenhouse gas emissions from an urban region, *P. Natl. Acad. Sci. USA*, 109, 8423–8428, doi:10.1073/pnas.1116645109, 2012.
- 890 Nassar, R., Napier-Linton, L., Gurney, K. R., Andres, R. J., Oda, T., Vogel, F. R., and Deng, F.: Improving the temporal and spatial distribution of CO₂ emissions from global fossil fuel emission data sets, *J. Geophys. Res-Atmos.*, 118, 917–933, doi:10.1029/2012JD018196, 2013.



- Newman, S., Jeong, S., Fischer, M. L., Xu, X., Haman, C. L., Lefter, B., Alvarez, S., Rappenglueck, B., Kort,
895 E. a., Andrews, a. E., Peischl, J., Gurney, K. R., Miller, C. E., and Yung, Y. L.: Diurnal tracking of anthropogenic CO₂ emissions in the Los Angeles basin megacity during spring 2010, *Atmos. Chem. Phys.*, 13, 4359–4372, doi:10.5194/acp-13-4359-2013, 2013.
- Ogle, S. M., Davis, K., Lauvaux, T., Schuh, A., Cooley, D., West, T. O., Heath, L. S., Miles, N. L., Richardson,
900 S., Breidt, F. J., Smith, J. E., McCarty, J. L., Gurney, K. R., Tans, P., and Denning, A. S.: An approach for verifying biogenic greenhouse gas emissions inventories with atmospheric CO₂ concentration data, *Environ. Res. Lett.*, 10, 34 012, doi:10.1088/1748-9326/10/3/034012, 2015.
- Oney, B., Henne, S., Gruber, N., Leuenberger, M., Bamberger, I., Eugster, W., and Brunner, D.: The CarboCount
CH sites: characterization of a dense greenhouse gas observation network, *Atmos. Chem. Phys.*, 15, 11 147–
11 164, doi:10.5194/acp-15-11147-2015, 2015.
- 905 Oney, B., Gruber, N., Henne, S., Leuenberger, M., and Brunner, D.: A CO₂-based method to determine the regional biospheric signal in atmospheric CO₂, *Tellus B*, 2016, in review.
- Peters, W., Krol, M., Van Der Werf, G., Houweling, S., Jones, C., Hughes, J., Schaefer, K., Masarie, K., Jacobson,
A., Miller, J., et al.: Seven years of recent European net terrestrial carbon dioxide exchange constrained
by atmospheric observations, *Glob. Chang. Biol.*, 16, 1317–1337, doi:10.1111/j.1365-2486.2009.02078.x,
910 2010.
- Peylin, P., Houweling, S., Krol, M. C., Karstens, U., Rödenbeck, C., Geels, C., Vermeulen, A., Badawy, B.,
Aulagnier, C., Pregger, T., Delage, F., Pieterse, G., Ciais, P., and Heimann, M.: Importance of fossil fuel
emission uncertainties over Europe for CO₂ modeling: Model intercomparison, *Atmos. Chem. Phys.*, 11,
6607–6622, doi:10.5194/acp-11-6607-2011, 2011.
- 915 Pillai, D., Buchwitz, M., Gerbig, C., Koch, T., Reuter, M., Bovensmann, H., Marshall, J., and Burrows, J. P.:
Tracking city CO₂ emissions from space using a high resolution inverse modeling approach: A case study for
Berlin, Germany, *Atmos. Chem. Phys. Discussions*, 16, 9591–9610, doi:10.5194/acp-16-9591-2016, 2016.
- Raupach, M. R., Marland, G., Ciais, P., Le Quéré, C., Canadell, J. G., Klepper, G., and Field, C. B.:
Global and regional drivers of accelerating CO₂ emissions., *P. Natl. Acad. Sci. USA*, 104, 10 288–10 293,
920 doi:10.1073/pnas.0700609104, 2007.
- Ray, J., Yadav, V., Michalak, A. M., van Bloemen Waanders, B., and McKenna, S. A.: A multiresolution spatial
parameterization for the estimation of fossil-fuel carbon dioxide emissions via atmospheric inversions,
Geoscientific Model Development, 7, 1277–1315, doi:10.5194/gmdd-7-1277-2014, 2014.
- Reuter, M., Buchwitz, M., Hilker, M., Heymann, J., Bovensmann, H., Burrows, J. P., Houweling, S., Liu, Y. Y.,
925 Nassar, R., Chevallier, F., Ciais, P., Marshall, J., and Reichstein, M.: How much CO₂ is taken up by the European
terrestrial biosphere?, *Bull. Am. Meteorol. Soc.*, doi:10.1175/BAMS-D-15-00310.1, 2016, in press.
- Roches, A. and Fuhrer, O.: Tracer module in the COSMO model, *COSMO Tech. Rep. No. 20*, 2012.
- Sarmiento, J. L. and Gruber, N.: Sinks for anthropogenic carbon, *Phys. Today*, 55, 30–36,
doi:10.1063/1.1510279, 2002.
- 930 Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Bergamaschi, P., and Peters, W.: Three years
of greenhouse gas column-averaged dry air mole fractions retrieved from satellite-Part 2: Methane, *Atmos.
Chem. Phys.*, 8, 8273–8326, doi:10.5194/acpd-8-8273-2008, 2008.



- Shiga, Y., Michalak, A., Gourdji, S. M., Mueller, K. L., and Yadav, V.: Detecting fossil fuel emissions patterns from sub-continental regions using North American in-situ CO₂ measurements, *Geophys. Res. Lett.*, 41, 1–8, doi:10.1002/2014GL059684, 2014.
- 935
- Turnbull, J. C., Tans, P. P., Lehman, S. J., Baker, D., Conway, T. J., Chung, Y. S., Gregg, J., Miller, J. B., Southon, J. R., and Zhou, L. X.: Atmospheric observations of carbon monoxide and fossil fuel CO₂ emissions from East Asia, *J. Geophys. Res.-Atmos.*, 116, 1–14, doi:10.1029/2011JD016691, 2011.
- Turnbull, J. C., Sweeney, C., Karion, A., Newberger, T., Lehman, S. J., Tans, P. P., Davis, K. J., Lauvaux, T., Miles, N. L., Richardson, S. J., et al.: Toward quantification and source sector identification of fossil fuel CO₂ emissions from an urban area: Results from the INFLUX experiment, *J. Geophys. Res.-Atmos.*, 120, 292–312, doi:10.1002/2014JD022555, 2015.
- 940
- Turnbull, J. C., Keller, E. D., Norris, M. W., and Wiltshire, R. M.: Independent evaluation of point source fossil fuel CO₂ emissions to better than 10%, *P. Natl. Acad. Sci. USA*, 113(37), 10 287–10 291, doi:10.1073/pnas.1602824113, 2016.
- 945
- Van Der Laan, S., Karstens, U., Neubert, R. E. M., Van Der Laan-Luijkx, I. T., and Meijer, H.: Observation-based estimates of fossil fuel-derived CO₂ emissions in the Netherlands using $\Delta^{14}\text{C}$, CO and ²²²Radon, *Tellus B*, 62, 389–402, doi:10.1111/j.1600-0889.2010.00493.x, 2010.
- Vardag, S., Gerbig, C., Janssens-Maenhout, G., and Levin, I.: Estimation of continuous anthropogenic CO₂: Model-based evaluation of CO₂, CO, $\delta^{13}\text{C}(\text{CO}_2)$ and $\Delta^{14}\text{C}(\text{CO}_2)$ tracer methods, *Atmos. Chem. Phys.*, 15, 12 705–12 729, doi:10.5194/acp-15-12705-2015, 2015.
- 950
- Velazco, V. a., Buchwitz, M., Bovensmann, H., Reuter, M., Schneising, O., Heymann, J., Krings, T., Gerilowski, K., and Burrows, J. P.: Towards space based verification of CO₂ emissions from strong localized sources: Fossil fuel power plant emissions as seen by a CarbonSat constellation, *Atmos. Meas. Tech.*, 4, 2809–2822, doi:10.5194/amt-4-2809-2011, 2011.
- 955
- Vogel, F. R., Thiruchittampalam, B., Theloke, J., Kretschmer, R., Gerbig, C., Hammer, S., and Levin, I.: Can we evaluate a fine-grained emission model using high-resolution atmospheric transport modelling and regional fossil fuel CO₂ observations?, *Tellus B*, 65, 1–16, doi:10.3402/tellusb.v65i0.18681, 2013.
- Zellweger, C., Emmenegger, L., Firdaus, M., Hatakka, J., Heimann, M., Kozlova, E., Spain, T. G., Steinbacher, M., van der Schoot, M. V., and Buchmann, B.: Assessment of recent advances in measurement techniques for atmospheric carbon dioxide and methane observations, *Atmos. Meas. Tech. Discussions*, 9, 4737–4757, doi:10.5194/amt-9-4737-2016, 2016.
- 960
- Zhang, X., Gurney, K. R., Rayner, P., Baker, D., and Liu, Y.-P.: Sensitivity of simulated CO₂ concentration to sub-annual variations in fossil fuel CO₂ emissions, *Atmos. Chem. Phys. Discuss.*, 15, 20 679–20 708, doi:10.5194/acpd-15-20679-2015, 2016.
- 965




Figure 1: Map of the fossil fuel emissions used in this study. Also depicted is the domain of the COSMO-7 setup employed here. Shown in transparent color are the fossil fuel CO₂ emissions for different sectors in units of gC m⁻² yr⁻¹. The colors from the different sector blend to a darker color when they are co-located as shown by the color mixing star at the bottom right.




Figure 2: Time dependence of fossil fuel CO₂ emissions for different sectors and countries. (a) Time functions for the diurnal and weekly emissions for four sectors. (b) Annual evolution of the CO₂ emission intensity for three different countries or group of countries. Shown are the daily minima and maxima for each country.




Figure 3: Comparison between modeled and observation-based estimates of the fossil fuel CO₂ component. (a) Comparison at the Lutjewad site in the Netherlands (LUT, 6° 21'E, 53° 24'N, 1 m a.s.l.) (Van Der Laan et al., 2010; Bozhinova et al., 2014). (b) Comparison at Heidelberg (HEI, 49.417° N, 8.675° E, 116m a.s.l.) (Levin and Karstens, 2007). The observational estimates are based on concurrent observations of CO and ¹⁴CO₂.




Figure 4: Maps of the model simulated annual mean components of atmospheric CO₂ in the surface layer (10 m above ground). (a) fossil fuel component, (b) total atmospheric CO₂, (c) terrestrial biosphere component, and (d) background CO₂ component. The results are shown as dry air mole fraction with units of ppm.




Figure 5: Instantaneous snapshot of the model simulated fossil fuel CO_2 in the surface layer. (a) Snap shot on July 1st at 06 00 GMT, (b) as (a) but at 18 00 GMT, (c) snapshot on January 1st at 06 00 GMT, (d) as (c) but at 18 00 GMT.




Figure 6: As Figure 4, but for whole air column averaged dry air mole fraction in units of ppm .




Figure 7: Maps of the annual standard deviation of (a) the fossil fuel component and (b) atmospheric CO₂ in the surface layer.




Figure 8: Maps of the contribution of fossil fuel CO₂ variability to total atmospheric CO₂ variability on various timescales in percent. (a) Contribution over all timescales; (b) contribution for the seasonal timescale only; (c) contribution for the synoptic timescale only; (d) contribution for the diurnal timescale only.




Figure 9: Maps of the impact of the consideration of time-varying fossil fuel emissions. (a) Difference in annual mean surface CO_2 between the case with time varying and time-constant fossil fuel emissions. This difference represents the fossil fuel rectification effect. (b) Linear correlation between the fossil fuel emissions and the height of the planetary boundary layer height in the COSMO-7 model. Pixels with emissions smaller than $0.06 \text{ gC m}^{-2} \text{ yr}^{-1}$ are not plotted. The positive correlation implies high emissions when the PBL is deep, and vice versa. Most of this correlation stems from the diurnal time-scale, but the correlation is enhanced through the (mostly) positive correlation also on seasonal timescales.




Figure 10: Maps of the annual mean fossil fuel CO_2 generated by different countries/regions. (a) Surface pattern created by the emissions from Germany, (b) as (a), but for the France. (c) Column averaged pattern created by the emissions from Germany, and (d) as (c), but for France.




Figure 11: Pie charts depicting the origin of the fossil fuel CO₂ signal for each country/region. The percentages represent the contribution of each country/region of origin to the total fossil fuel signal in the averaged over the air column. The pie chart for Switzerland reveals, for example, that only 20% of the fossil fuel CO₂ signal over its territory stems from its territorial emission. Here, CH: Switzerland; DE: Germany; FR: France; IT: Italy; AT: Austria; NL: Netherlands; SW: countries in southwest of the domain; UK: United Kingdom; EA: countries in eastern domain; OT: the rest of countries.




Figure 12: Maps of the annual mean surface fossil fuel CO₂ stemming from different sectors in units of ppm. (a) fossil-fuel fired power plants, (b) residential heating, (c) industrial processes, and (d) road transportation.

Table 1: Evaluation of COSMO-7 based simulations of the atmospheric CO₂ concentration at 4 European sites (locations are shown in Figure 1). The comparison are shown for the 3 hourly means between 12 to 18 PM local time for the period April 2008 through April 2009. m.s.a.g.is the height above ground or relative height.

Station	characteristics	m.s.a.g(m)	S.T.D. obs(ppm)	S.T.D. mod(ppm)	Correlation	Bias(ppm)
Cabauw (CBW, Netherlands)	tower	20	11.46	11.80	0.78	0.72
Cabauw (CBW, Netherlands)	tower	60	10.86	11.06	0.77	0.27
Cabauw (CBW, Netherlands)	tower	200	9.35	8.19	0.74	0.88
Puy de Dome (PUY, France)	mountain top	10	7.83	7.65	0.75	0.85
Hegyhatsal (HUN, Hungary)	continental	10	12.08	9.42	0.8	4.0
Hegyhatsal (HUN, Hungary)	continental	48	11.51	9.32	0.8	4.04
Hegyhatsal (HUN, Hungary)	continental	115	10.69	8.72	0.8	3.86
Mace Head, (MHD, Ireland)	coastal	15	6.26	3.87	0.81	-0.34




Figure 13: Maps of the annual mean relative contribution of each sector to the total surface fossil fuel CO₂. a) fossil-fuel fired power plants, (b) residential heating, (c) industrial processes, and (d) road transportation.




Figure 14: Changes in annual mean atmospheric CO_2 and its standard deviations resulting from a 30% reduction in the fossil fuel emissions from all sectors. (a) Change in surface mean CO_2 . (b) Change in the column averaged CO_2 , i.e., $X\text{CO}_2$. (c) Change in the standard deviation of surface CO_2 (all seasons). (d) Change in the standard deviation of the column averaged CO_2 , i.e., $X\text{CO}_2$. The standard deviation refers to the differences of the afternoon data (at 1:00 PM) to the annual afternoon average.




Figure 15: Impact of reductions in power plant emissions on the mean and standard deviation of the fossil fuel CO₂ signal. (a) probability density distribution of the surface atmospheric CO₂ for the present and for a case when the power plant emissions were reduced by 50% at a site in eastern Germany (50.32°N,13.19°E). (b) Relationship between the changes in the mean and the standard deviation of the column averaged CO₂ for a given reduction in power plant emissions, with different color representing representing different sites with different characteristics in their response to this reduction in emission: Blue (50.32°N, 13.19°E), Cyan (50.32°N, 6.59°E), Red (42.48°N, 6.51°W), Orange (49,28°N, 6.14°E) (Locations shown in Figure 14b with green circles).