



## 1 Long-term series of surface solar radiation at Athens, Greece

2

3 Stelios Kazadzis<sup>1,2</sup>, Dimitra Founda<sup>2</sup>, Vassilios E. Psiloglou<sup>2</sup>, Harry Kambezidis<sup>2</sup>, Nikolaos  
4 Mihalopoulos<sup>2,3</sup>, Arturo Sanchez-Lorenzo<sup>4</sup>, Charikleia Meleti<sup>5</sup>, Panagiotis I. Raptis<sup>1,2</sup>, Frangiskos  
5 Pierros<sup>2</sup>, Pierre Nabat<sup>6</sup>

6

7 [1] {Physikalisch-Meteorologisches Observatorium Davos, World Radiation Center (PMOD/WRC)  
8 Dorfstrasse 33, CH-7260 Davos Dorf, Switzerland}

9 [2] {Institute of Environmental Research and Sustainable Development, National Observatory of Athens,  
10 Greece}

11 [3] {Department of Chemistry, Univ. of Crete, Heraklion, Crete}

12 [4] {Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza,  
13 Spain}

14 [5] {Physics Department, Aristotle University of Thessaloniki, Greece}

15 [6] {CNRM UMR 3589, Météo-France/CNRS, Toulouse, France}

16 Corresponding author: S. Kazadzis, kazadzis@noa.gr

## 17 Abstract

18 We present a long-term series of solar surface radiation (SSR) for the city of Athens, Greece. The  
19 SSR measurements were performed from 1953 to 2012, and before that (1900-1952) sunshine  
20 duration (SD) records have been used in order to reconstruct monthly SSR. Analysis from the whole  
21 dataset (1900-2012) mainly showed: a decrease of 2.9% per decade in SSR from 1910 to 1940  
22 assuming a linear change in SSR. For the dimming period (1955-1980), a -2% change per decade  
23 has been observed, that matches various European long-term SSR measurement related studies.  
24 This percentage for Athens is in the lower limit, compared to other studies for the Mediterranean  
25 area. For the brightening period (1980-2012) we have calculated a +1.5% per decade which is also  
26 in the lower limit of the reported positive changes in SSR around Europe. Comparing the 30-year  
27 periods (1954-1983 and 1983-2012) we have found a difference of 4.5%. The difference was  
28 observed for all seasons except winter. Using an analysis of SSR calculations of all sky and clear  
29 sky (cloudless) conditions/days, we report that most of the observed changes in SSR after 1954 can  
30 be attributed partly to cloudiness and mostly to aerosol load changes.

31



## 1 **1 Introduction**

2 In the past decades surface solar radiation (SSR) and the transmission of the atmosphere have been  
3 of increasing interest because of the related impacts on climate. Most of the energy in the Earth-  
4 atmosphere system is introduced by solar radiation as it provides heating, which creates pressure  
5 gradients and ultimately wind, as well as it triggers water, carbon and oxygen cycles through  
6 evaporation and photosynthesis. These processes define the climatological conditions, and changes  
7 of incoming solar radiation rapidly affect the energy balance (Wild et al., 2015). Interest on the  
8 solar radiation changes has also been raised after the development of solar energy applications,  
9 which are continuously growing in number over the recent years. Changes in SSR have been  
10 recorded over the last century and can be caused either by natural events such as volcanic eruptions  
11 or human-related activities, mainly in polluted regions (Wild, 2016). At larger scales (thousands of  
12 years) changes in SSR, might have been caused by changes in the Earth's orbit and Sun solar output  
13 (Lean, 1997; Ohmura, 2006).

14 Systematic continuous measurements of SSR were established in the middle of the 20th century at  
15 selected meteorological observatories. Solar variations have been investigated in several studies  
16 using ground based SSR measurements from various monitoring networks worldwide (e.g.,  
17 Ohmura, 2009) and also by satellite-derived estimations (e.g. Kambezidis et al., 2010). Overall,  
18 most of these studies (Gilgen et al., 1998; Noris and Wild ,2009; Wild, 2009 and 2016 and  
19 references therein) have reported a worldwide decrease of solar incoming radiation in the period  
20 1960-1985 (known as dimming period), followed by an increase (brightening period) thereafter.  
21 These changes were larger in more polluted and urban areas but have also been recorded in isolated  
22 regions such as the Arctic (Stanhill, 1995) and Antarctica (Stanhill and Cohen 1997). Changes in  
23 atmospheric transmission due to variations in cloudiness and aerosol concentration are the main  
24 factors to be investigated in order to determine the possible causes of such trends in SSR (Wild,  
25 2009).

26 The cloud and aerosol radiative effects on solar radiation variations over the past decades have been  
27 investigated by numerous studies during the last years. The inter-annual variations in cloudiness is  
28 crucial for studying SSR time series, but its decadal variability is not always connected with the  
29 widespread dimming and brightening effects (Wang et al., 2012; Wild, 2016). Aerosols play  
30 significant role in incoming radiation, by scattering and absorbing light and by acting as cloud-  
31 condensation nuclei. Over the 20-year dimming phase (from 1960 to 1980) and the 15-year  
32 brightening phase (from 1990 to 2005), it was found that the aerosol effects (direct and indirect)  
33 played the most important role in SSR variation (Dudok de Wit et al., 2015). Concerning Central



1 Europe, Ruckstuhl et al. (2008) suggested that the brightening phase under cloud-free conditions is  
2 in line with decreasing anthropogenic aerosol emissions (Streets et al., 2006). Nabat et al., 2013  
3 using a blending of remote sensing and model products showed that a decreasing Aerosol Optical  
4 Depth (AOD) trend of 0.05 per decade has been calculated for Europe. In addition, Nabat et al.,  
5 2014 reported that anthropogenic aerosol decline in Europe from 1980 to 2012 statistically explains  
6 explain  $81 \pm 16\%$  of the observed brightening. Overall, changes in anthropogenic aerosol emissions  
7 are now considered as the major cause of brightening and dimming effects (Wild, 2016). The  
8 gaseous and particulate air pollutants may reduce solar radiation by up to 40% during air pollution  
9 episodes (Jauregui and Luyando, 1999). This attenuation is much larger during forest fires, dust  
10 events and volcanic eruptions. Vautard et al., (2009), have also reported on a decline of the  
11 frequency of low-visibility conditions such as fog, mist and haze in Europe over the past 30 years,  
12 suggesting a significant contribution of air-quality improvements

13 Long-term series of SSR measurements are essential for such studies. One of the main constraints in  
14 studying SSR temporal changes is the small number of sites with reliable long-term records, even  
15 over areas with high density of stations such as Europe, Japan or the USA. In Europe for example,  
16 there are currently less than 80 stations with more than 40-years homogeneous data (Sanchez-  
17 Lorenzo et al., 2015), with very few of them operating over Southern Europe. Recently, a high-  
18 quality dataset of SSR has been set up over Italy (Manara et al., 2016), but there is still lack of high  
19 quality long-term trends in other countries around the Mediterranean Basin.

20 In addition, even more sporadic measurements are available before the 1950s (Stanhill and  
21 Achiman, 2016); the few studies of them have pointed out an SSR increase in the first decades of  
22 the 20<sup>th</sup> century and a maximum around 1950 (Ohmura, 2006). This topic is still controversial due  
23 to the few long-term series available (Antón et al., 2014). Recently, there have been efforts to  
24 reconstruct SSR series in periods with no direct measurements available, using other variables such  
25 as sunshine duration (SD), which is available in a large number of sites since the late 19<sup>th</sup> century  
26 (e.g., Stanhill and Cohen, 2005, for USA; Sanchez-Lorenzo and Wild 2012, for Switzerland;  
27 Matuszko 2014, for Poland). For example, Sanchez-Lorenzo and Wild (2012) used data from 17  
28 stations in Switzerland, considered SD as a proxy and successfully reconstructed SSR time series  
29 since the late 19<sup>th</sup> century. Thus, they calculated that the variability in SSR monthly anomalies can  
30 be explained by SD anomalies in a range of 76%-96%, and a monthly root mean squared error of  
31  $4.2 \text{ W m}^{-2}$  between recorded and estimated SSR for all-sky conditions and of  $5.5 \text{ W m}^{-2}$  for clear-  
32 sky conditions. Other studies have tried to use pan evaporation as a proxy of SSR, for the first half  
33 of the 20<sup>th</sup> century (Stanhill and Möller, 2008). Kambezidis et al. (2016) used monthly re-analysis



1 datasets from the Modern Era Retrospective-Analysis for Research and Applications (MERRA) and  
2 calculated shortwave radiation trends from 1979-2012 for the Mediterranean basin. They reported  
3 an increase in MERRA by  $+0.36 \text{ W m}^{-2}$  per decade, with higher rates over the western  
4 Mediterranean ( $+0.82 \text{ W m}^{-2}$  per decade).

5 For the Southeastern Mediterranean, there have been a few studies discussing the  
6 brightening/dimming effect. Zerefos et al. (2009) have studied the Ultraviolet A (UVA) changes for  
7 the area of Thessaloniki (Greece) from 1984 to 2008. They calculated a 5% positive trend per  
8 decade linked to a negative trend in aerosol optical depth (AOD) for the area due to air pollution  
9 abatement strategies. Founda et al., (2014) have studied the SD long-term variability over Athens  
10 area. They reported a 7% decline in the annual SD from 1951-1982 and a 3% increase from 1983-  
11 2011 under all sky conditions. Although, under near clear sky conditions, these percentages are -7%  
12 and + 9% for the dimming and brightening periods respectively. Similarly, Founda et al. (2016a)  
13 analyzed long-term SD and total cloud cover time series over 15 sites in Greece (the oldest one  
14 beginning on 1897). They have shown an increase in SD almost at all stations since the mid-1980s,  
15 which in certain areas of Southeastern Greece amounts to an increase of 20 h per year. This increase  
16 is not accompanied with synchronous decrease in total cloud cover, possibly evidencing to  
17 decreasing aerosols loads, despite the fact that their impact on SD should be lower than on SSR  
18 (Sanchez-Romero et al., 2014). Yildirim et al. (2014) have analyzed 41 years of SD measurements  
19 in 36 stations in Turkey. They reported a decreasing trend (between 1970 to about 1990) for most of  
20 the stations. After 1990 they observed either zero trend variation or a reduction in the decreasing  
21 rate of SD for most of the locations. They concluded that the decreasing period might be attributed  
22 to human-induced air pollution. Founda et al. (2016b) have investigated the visibility trends over  
23 Athens area from 1931 to 2013. They reported a deterioration in the visibility up to 2004 and a  
24 slight recovery afterwards, negatively/positively correlated with relative humidity/wind speed and  
25 positively correlated with AOD from 2000 to 2013.

26 In this work, measurements of SSR, recorded for 60 years at the center of Athens, are presented. In  
27 addition, with the use of the SD measurements that are conducted in Athens since 1900, we could  
28 reconstruct the time series of SSR during the first half of the 20th century. These time series (1900-  
29 2012) are the oldest, uninterrupted and high quality SSR time series in the SE Mediterranean and  
30 one of the oldest in Europe, providing unique information about the variations and trends in the area  
31 for the past decades. Time-series of SSR over Athens are presented to try answering questions such  
32 as:



- 1 Are the dimming–brightening patterns observed in Europe over the past century also observed, at  
2 the same extent, over the Eastern Mediterranean?  
3 Is SSR variability during the first decades of the 20th century in Athens in line with the other few  
4 locations reporting trends over this period?  
5 Can we verify that anthropogenic aerosols play the most important role on the brightening/dimming  
6 observed SSR after 1950 in agreement with other European regions?  
7

## 8 **2 Data and Methodology**

### 9 **2.1 Data collection and analysis**

10 The SSR data used in this study cover the period from December 1953 to December 2012 and were  
11 measured by a series of pyranometers that are mentioned in Table 1. These instruments have been  
12 operating continuously at the Actinometric Station of the National Observatory of Athens  
13 (ASNOA) (Hill of Pnyx, Thissio), that is located near the center of Athens, Greece (38.00° N,  
14 23.73° E, 110 m above mean sea level). Table 1 presents the instruments and the period of  
15 operation, as well as the maximum error in the calculation of the daily values. References  
16 mentioned in Table 1 describe the exact type of errors and uncertainties related to the sensors. In the  
17 period 1953-1986, the maximum daily error was about 5%, and 2% afterwards. The spectral  
18 response of the sensors is in the range of 285-2800 nm; since 1986 a first-class Eppley PSP  
19 pyranometer (WMO, 1983) is operating at ASNOA. Since 1992, frequent calibrations (every two  
20 years) have been performed by the NOAA's Laboratory of Meteorological Device Calibration  
21 (LMDC, 2016) in order to ensure the high quality of measurements. LMDC follows the standard  
22 calibration procedure for thermopile pyranometers (ISO 9847, 1992), with exposure to real sunlight  
23 conditions and comparison with a standard thermopile (Secondary Standard) pyranometer. LMDC's  
24 reference pyranometer, Kipp & Zonen CM21, is regularly calibrated in PMOD/WRC, Davos,  
25 Switzerland.

26

27 Table 1: History of SSR instruments used at ASNOA. SSR measurements refer to the total solar  
28 radiation on a horizontal surface.

	Instrument	Period	Class	Maximum error (daily integral)	Reference	Class	Comments	Resolution
1	Solarigraph GOREZYNSKI	1953- 1959	2nd	5%	Coulson (1975)	B	One instrument being used	1 hour



2	Eppley 180° pyranometer (No. 3604)	1960-1966	2nd	5%	Coulson (1975), Drummond (1965)	B	Manual measurements archiving with mvoltmeter	1 hour
3	Eppley 180° pyranometer (No. 3604) coupled with a Leeds-Northup recorder, Speedomax, type G	1966-1968	2nd	5%	Coulson (1975), Drummond (1965)	B	Same instrument as #2 with Speedomax recorder	1 hour
4	Eppley 180° pyranometer (No. 3034) coupled with a Leeds-Northup recorder, Speedomax, type G	1968-1973	2nd	5%	Coulson (1975), Drummond (1965)	B	New instrument, same recorder	1 hour
5	Eppley pyranometer, type 8-48 and type 8-48A coupled with a Leeds-Northup recorder, Speedomax, type G	1974-1986	2nd	3-5%	Hulstrom (1989)	B	Type 8-48 and type 8-48A Instruments were measuring alternatively for three years each	1 h
6	Eppley Precision Spectral Pyranometer (PSP)	1986-now	1st	1-2%	Hulstrom (1989)	A	Regular recalibrations. Coupled with a, A/D recorder (Campbell Scientific Ltd. Datalogger, type CR-21X at the beginning until 2003, a CR10X until 2012	1 min

1

2 SSR data are processed using a set of quality-control (QC) tests in order to ensure the quality of the  
 3 data set. The QC procedures include rejection of:


- 4
- Measurements for solar elevation angles less than 5 deg;
  - 5 • SSR values equal to or less than  $5 \text{ W m}^{-2}$ , during sunrise and sunset, due to the  
 6 pyranometers' sensitivity;
  - 7 • SSR values greater than 120% of the seasonally corrected solar constant.

8 After the initiation of diffuse horizontal radiation measurements at ASNOA in 1991, the following  
 9 quality criteria were added for rejection:

- 10
- diffuse horizontal values greater than the corresponding SSR ones;
  - 11 • diffuse horizontal values greater than 80% of the seasonally correct solar constant;
  - 12 • direct-beam solar component exceeding the extraterrestrial solar irradiance.



1 Also, both total and diffuse horizontal measurements are corrected for the night-time dark-signal  
2 offset of the pyranometers.  
3 Mean daily SSR values were calculated from the data set of this study (December 1953 – December  
4 2012); only months with more than 20 days of measurements were considered in the analysis. Over  
5 the 60 years of measurements, only three months (January and February of 1998 and March of  
6 2012) did not fulfill this criterion.  
7 Figure 1 shows the intra-annual variability of SSR at ASNOA based on the measurements from all  
8 instruments during the period 1953-2012. Mean daily SSR at Athens ranges between approximately  
9 6 to 27 MJ m<sup>-2</sup> during the year. Mean and standard deviations were calculated using the 60 year  
10 record for each day.



11

12 **Figure 1.** Average Intra-annual variability of Surface Solar Radiation (SSR) at Actinometric Station of the  
13 National Observatory of Athens (red), along with the inter-annual variability for a given day ( $\pm 1$  standard  
14 deviation, blue), calculated over the period 1953-2012.  
15

16 The results of figure 1 show the average yearly pattern of SSR at ASNOA. The day to day  
17 variability that is shown as “noise” in the plotted blue line comes from the 60 year averaging of  
18 each day and is mostly related with the amount of cloudiness for each of the averaged days.  
19 Minimum and maximum SSRs at solstices, compared to a cloudless sky aerosol free model, are also  
20 related with the highest probability of the presence of clouds during winter months. For the



1 calculation of each of the daily averages the available data points vary from 55 to the possible  
2 maximum of 60.

3 In addition, collocated measurements of SD recorded at ASNOA have been used. According to  
4 WMO (2010), the SD during a given period is defined as the sum of the sub-periods for which the  
5 direct solar irradiance exceeds  $120 \text{ W m}^{-2}$ . In Athens, SD has been recorded using classical  
6 Campbell-Stokes heliographs (since 1894) and been replaced by electronic instrumentation in 1998  
7 (EKO, MS-091 analog SD sensor). Monthly SD values since January 1900 have been used in this  
8 study. A more analytical study of these time series can be found in Founda et al. (2014).

9 Complementary to this study, cloud-cover observations from the Hellenic National Meteorological  
10 Service (HNMS) from 1954 have also been used. These observations are recorded at a site 7 km  
11 away from ASNOA. All cloud observations at HNMS are conducted every 3 hours and are  
12 expressed in octas.

13 Concerning the data availability for SSR and SD data: SSR monthly means calculated here have  
14 been retrieved from daily calculated SSRs. Over the 59 years (708 months), 98% of the months had  
15 none or one day missing, 3 months had from 10-20 missing days and 2 months from 20-30 missing  
16 days. For SD, 1931-1940 monthly data have been used taken from the NOAA measurement annals.  
17 From 1940 on, hourly measurements have been used in order to derive daily and monthly  
18 measurements. The SD time series have no gaps with only six missing days during December 1944  
19 (Founda et al., 2014).

20 In order to examine the AOD impact on SSR, we have used the longest satellite based AOD series  
21 available for the area. This is the AOD time series from Advanced Very High Resolution  
22 Radiometer (AVHRR). AOD retrievals at 630 nm over global oceans at spatial resolution of  $0.1^\circ \times$   
23  $0.1^\circ$  and one overpass per day, have been used. Data used were downloaded from NOAA Climate  
24 Data Record (CDR) version 2 of aerosol optical thickness (Zho and Chan, 2014), and cover the  
25 period from August 1981 to December 2009. AVHRR AOD embodies a large variety of  
26 uncertainties, including radiance calibration, systematic changes in single scattering albedo and  
27 ocean reflectance (Mishchenko et al, 2007). Current dataset radiances have been recalibrated using  
28 more accurate MODIS data (Chan et al, 2013). We used daily data at the region around Athens  
29 (longitude:  $37.5^\circ$ - $38.2^\circ$ N, latitude:  $23.2^\circ$ - $24.4^\circ$ E) which includes 50 active available (ocean) grid-  
30 points. The above region was selected based on data availability on each grid with the distance up to  
31 50 km from ASNOA.



1 To complement the analysis on the evolution of aerosols, the recent climatology developed by  
2 Nabat et al., 2013 has been considered over the period 1979-2012. This product provides monthly  
3 averages of AOD at 550 nm over the Mediterranean region at 50 km resolution. It is based on a  
4 combination of satellite-derived (MODIS instrument) and model-simulated products (MACC  
5 reanalysis and RegCM-4 simulations), which have been selected among many available datasets,  
6 from an evaluation against ground-based measurements of the AERONET network. Thus this  
7 climatology is able to give an estimation as best as possible of the atmospheric aerosol content over  
8 the period 1979-2012. For the present work, the AOD time series over the grid cell of the ASNOA  
9 (38.00° N, 23.73° E) has been extracted and is referred to as the ChArMEx data thereafter.

10

## 11 **2.2 Clear-sky SSR**

12 For the determination of the clear sky (defined here as the cloudless) days, we have used both the  
13 cloud octas and SD data. Daily observations have been used for this analysis. We have defined as a  
14 clear sky day each day that fulfills the following criteria:

- 15 - the mean daily cloudiness (in octas) should be less than 1.5, and
- 16 - the total daily SD should be higher than 90% of its theoretical (astronomical) value.

17 The procedure for calculating a single mean cloud octa value for each day was the following:

18 We have first excluded night-time cloud observations; then, we have weighted each observation  
19 based on the hour of the observation. Weights have been calculated based on the solar radiation  
20 contribution of the specific time slot and day of the month, compared with the daily clear sky SSR  
21 integral, of the particular day and month.

22

## 23 **2.3 Reconstruction of SSR from SD**

24 We have used the 1900-2012 SD time series in order to extend our SSR time series back to 1900.  
25 For that purpose we have used a recent period (1983-2012) in order to derive a function between  
26 SD and SSR and a testing period (1953-1982) to verify the validity of this function and results.  
27 Finally we applied this derived and tested formula for the 1900-1952 period where SD  
28 measurements were available while SSR were not. In order to derive a relationship between SD and  
29 SSR, we used the broadly accepted formula of Ångström (1924):

$$30 \quad \text{SSR}/\text{SSR}_{\text{max}} = a + b(\text{SD}/\text{SD}_{\text{max}}) \quad (1)$$

31 where  $\text{SSR}_{\text{max}}$  and  $\text{SD}_{\text{max}}$  refer to the theoretical extra-terrestrial value of SSR and the astronomical  
32 value of SD, respectively, while  $a$  and  $b$  are constants usually defined monthly. This formula can




1 only be used in large data sets as a statistical approach. That is because for different cloud height,  
2 thickness and positioning, the constants can show a large variability (Angell, 1990).

3

### 4 **3 Results**

#### 5 **3.1 Relationship between SD and SSR measurements**

6 In order to benefit from the over-century long time-series of SD at NOAA we used Eq. (1) for the  
7 period of synchronous SD and SSR measurements to estimate constants  $a$  and  $b$  for the period  
8 1983-2012. The 1983-2012 period was chosen for determining the SSR vs. SD relationship as  
9 mainly SSR measurements have lower uncertainties compared with the 1953-2012 one. We thus  
10 calculated  $a=0.237$  and  $b=0.458$  (as derived from the linear equation shown in Figure 2) and used  
11 them for validation in the period 1953-1982. Figure 2 shows the correlation of the monthly  
12  $SSR/SSR_{max}$  and  $SD/SD_{max}$  ratios for the 1983-2012 period, together with the derived coefficients  
13 ( $a$  and  $b$ ) and the coefficient of determination ( $R^2$ ). In addition, we have included the testing period  
14 statistics, together with the coefficients  $a$  and  $b$  that can be determined using the whole period  
15 (1953-1982), to show that each of the two periods could provide similar results.




16

17 **Figure 2.** Plot of  $Y=SSR/SSR_{max}$  vs  $X=SD/SD_{max}$ . Blue dots represent the 1953-1982 period and  
18 red ones 1983-2012. The red and blue lines represent the respective linear regression lines.

19 Corresponding regression relations are given in the inner boxes.

20 For testing the method we have applied the retrieved coefficients and calculated the SSR for the  
21 1953-1982 period. Comparing normalized simulated with the observed SSR values an  $R^2=0.79$  has  
22 been found (see Figure 3).



1  
2  
3  
4  
5

**Figure 3.** Plot of the ratios of mean monthly SSR values to SSR maximum, of measurements in the period 1983-2012 vs the estimated ratios from Eq (1) for the same period. Blue is the 1:1 and black is the regression line.

6 We also followed the same procedure to calculate the coefficients of the Ångström formula  
7 separately for each month and for each season during the control period 1983-2012. For individual  
8 months, calculated SSR/SSRmax vs SDU/SDUmax coefficients of determination ranged from 0.5  
9 to 0.65 for winter months, 0.32 to 0.67 for spring months, 0.47 to 0.53 for autumn months and 0.1  
10 to 0.38 for summer months. The low coefficients for the summer period are related with the small  
11 range of values of SDU/SDU max and SSR/SSRmax that are related with the absence of clouds.  
12 When we have calculated seasonal based Ångström formulas for winter, spring, summer, and  
13 autumn months we have found  $a=[0.22, 0.22, 0.34, 0.27]$ ,  $b=[0.48, 0.49, 0.45, 0.34]$  and  $R^2=[0.6,$   
14  $0.74, 0.2, 0.63]$  respectively. In order not to include statistical uncertainties introduced from the  
15 correlations of individual months and seasons that are reported, we decided to use the Ångström  
16 formula derived using all months in the same dataset. Such assumption could introduce a season  
17 dependent trend to the extrapolation of SSR back to 1900 but it is considered more safe than using  
18 other least trusted seasonal Ångström formulas.

19

### 20 3.2 Long-term variations and trends (1900-2012)


21 Based on the results shown in Section 3.1, we have reconstructed monthly SSR from 1900  
22 to 1953. Using the full dataset of reconstructed (1900-1952) and measured SSR (1953-  
23 2012) we have calculated the mean monthly SSR values and used them for de-seasonalising  
24 the results shown in Figure 4. The de-seasonalizing was determined by: a. calculating the



- 1 average SSR ( $SSR_{mi}$ ) for each month (i) out of the 12 months of the given year, for all 1900-
- 2 2012 years, b. calculating the changes in % in SSR ( $SSR\%(i,y)$ ) for each month (i) of each
- 3 year (y) as:

$$4 \quad SSR\%(i,y) = \frac{SSR_{iy} - SSR_m}{SSR_m} * 100$$

5



6

7

8 **Figure 4.** Full time series of de-seasonalised SSR per cent changes (using the 1900-2012 monthly  
9 averages). Upper panel: different colors represent seasonal analysis, lower panel: black bold line  
10 represents the annual series, and light grey line the mean monthly values

11

12 According to Figure 4, the month-to-month variation (shown with light grey line) can reach more  
13 than 30% in comparison with the mean monthly average of the whole data set. Annual means show



1 a 10%-12% (peak to peak) decrease in SSR from 1910 to late1930's and then an increase of 15%-  
2 17% from 1940 to early 1950's. Subsequently, there is a decrease during the 1960's and then a  
3 positive change of the order of 20% till today with an episode in the early 1990's that show low  
4 SSR values. The latest can be directly linked with the Pinatubo volcanic eruption and its known  
5 effect in the SSR (e.g. Zerefos et al., 2012). Analytical linear trends of each of the sub-periods and  
6 for every season are presented in Table 2.


7 **Table 2.** Annual and seasonal SSR trends in percent per decade over the period 1900-2012 and  
8 different sub-periods. Percentages in parenthesis show the limits of the 95% confidence bounds.  
9

Season	1900-2012	1900-1952	1953-1982	1983-2012
Winter	-0.15 ( $\pm 0.46$ )	-0.68 ( $\pm 1.26$ )	-6.43 ( $\pm 3.83$ )	+0.52 ( $\pm 3.26$ )
Spring	+1.05 ( $\pm 0.38$ )	+0.15 ( $\pm 1.08$ )	-0.60 ( $\pm 3.10$ )	+2.77 ( $\pm 3.10$ )
Summer	+0.54 ( $\pm 0.31$ )	+0.43 ( $\pm 0.78$ )	-1.14 ( $\pm 2.90$ )	+1.38 ( $\pm 2.55$ )
Autumn	+0.14 ( $\pm 0.34$ )	+0.14 ( $\pm 1.02$ )	-1.28 ( $\pm 3.42$ )	-1.50 ( $\pm 1.83$ )
Year	<b>+0.40</b> ( $\pm 0.26$ )	<b>+0.02</b> ( $\pm 0.73$ )	<b>-2.33</b> ( $\pm 2.28$ )	<b>+0.80</b> ( $\pm 1.96$ )

10

11 Looking at the 1900-2012, period the seasonal and annual trends in SSR are less than 1% per  
12 decade. A positive change of 0.40% per decade has been calculated from annual values. For the  
13 whole data set, all seasons show positive trends, except for winter. For the periods with simulated  
14 SSR values (1900-1952), even smaller trends have been detected for spring and summer. The  
15 measuring period of 1954-2012 has been split into two sub-periods of 1954-1982 and 1983-2012.  
16 The first sub-period shows a negative annual change of -2.33% per decade in SSR, which is also  
17 reflected in all seasons with predominant changes during winter (-6.43% per decade). The second  
18 sub-period shows a positive trend of +0.80% per decade with highest ones in spring (+2.77% per  
19 decade) and summer (+1.38% per decade) and negative in autumn (-1.50% per decade). Looking at  
20 the trend significance described by the 95% confidence bounds, we can see significant positive  
21 trends for 1900-2012 (yearly, summer and spring) and significant negative trends for yearly  
22 analysis and winter of 1953-1982.

23 In order to have a better understanding of the SSR changes over the 112-year period (1900-2012),  
24 we have calculated the decadal SSR trends for different time-windows (15 to 40 years). Figure 5  
25 shows the results of this analysis.



1

2 **Figure 5.** Trends in SSR (% per decade) calculated for different sliding time windows. The value of  
3 the trend has been calculated at the central year of each time window.  
4

5 For the first two decades of the 20<sup>th</sup> century there appears a decrease in SSR, in line with other  
6 long-term SD series as recently shown by Stanhill and Achiman (2016). Then, in all calculations an  
7 increase is shown from mid 1930's to late 1940's, in line with the early brightening effect pointed  
8 out by other authors (Ohmura, 2009; Sanchez-Lorenzo et al., 2008). It should be reminded that this  
9 period is based on estimations of SSR from SD measurements, which thus include additional  
10 measurement uncertainties. Nevertheless, both the early dimming and brightening periods reported  
11 in Stanhill and Achiman (2016) and this study seem to be in line with trends in anthropogenic black  
12 carbon (McConnell et al., 2007; Lamarque et al., 2010) and biomass-burning (Lamarque et al.,  
13 2010) emissions peaking in the 1920's and then decreasing. The dimming period from 1950's to  
14 1970's can be observed in all time windows with a brightening effect after late 1970s.

15 The 40-year and 30-year time windows in the analysis presented in Figure 5 show the maximum  
16 rate of increase in early 1940's (resulting in an increase of 2% per decade and 3% per decade,  
17 respectively). Then a maximum rate of decrease is observed in early-mid 1960's, followed by a  
18 positive rate of increase after 1990's. Shorter time windows (15 years) are also interesting as they  
19 are able to capture the Pinatubo effect in early 1990's.


20



### 1 3.3 Variations and trends in SSR for the 1953-2012 measurement period

2 In order to further analyze the whole 59-yr SSR data set of this study, we have divided it in two 30-  
3 yr climatological sub-periods: 1954-1983, and 1983-2012 (the common year is meant to have equal  
4 duration for both periods). Investigating a possible seasonal dependence, the relative difference in  
5 SRR for every month from its mean monthly value of the whole measurement (1953-2012) period  
6 was calculated.

7 Figure 6, shows the mean daily insolation for each month for the two sub-periods and the whole 59-  
8 year period. Examining the monthly average differences between the two periods, we observe that  
9 for spring and summer months these are of the order of 6%. In addition, for all months SSR  
10 differences of the 1983-2012 period compared to the 1954-1983 period are positive with an  
11 exception of November (-1.9%) and December (-1.2%). In general, the second measurement period  
12 shows a 3% to 8% larger monthly SSR than the first measurement period.




13  
14 **Figure 6.** Intra-annual variability of monthly mean daily SSR over the sub-periods of 1954-1983  
15 (blue line) and 1983-2012 (red line) and the entire period of 1954-2012 (black line). The green line  
16 (right axis) represents the monthly relative difference between the two 30-year sub-periods.  
17

18 We have also calculated decadal trends in time windows of 15 to 30 years for the entire SSR  
19 measurement period (see Figure 7). In the left panel of Figure 7 each of the points used for  
20 constructing the colored lines represents the percent change per decade of the SSR. Mostly positive  
21 trends are detected using any time window centered after 1975. Larger trends are calculated for time



1 windows centered at 1975 to 1980 and after 2000 (in the order of 5% per decade using the 15-year  
2 time window). For the period 1954 to 1970 mainly negative trends are shown.



3 **Figure 7.** Left panel: calculation of SSR changes in % per decade for different time scales (15 to 30  
4 years) used. Each point on the line represents the middle year of the time window used for the trend  
5 calculation Right panel: % change per decade for different time scales and different months using  
6 1954 as the starting year (the last 30 year period is magnified with changes presented as % per  
7 decade).  
8

9 The right panel of Figure 7 shows the SSR change per decade for the months of January, April,  
10 July, October and yearly (all months). The figure is showing a trend analysis for the entire data set  
11 with time windows from 10 to 59 years, where each time window starts from 1953. For all months  
12 SSR changes become positive for time windows of 35 years and higher (1953-1988 time window  
13 and any larger window starting from 1953). Negative trends calculated from 1954 to any given year  
14 up to 1989 are mainly due to the large negative changes during the winter period. Especially during  
15 the 1954-1974 period, winter SSR changes show a 18% per decade decrease. Linear trends in SSR  
16 from 1954-2012 showed a positive trend of the order of 1% per decade, while individual months  
17 vary from 0.5% per decade to 1.5% per decade.

18

#### 19 **4. Comparison between all-sky and clear-sky SSR records variation**


20 We have used the 59-year data set (1953-2012) in order to quantify the factors controlling the SSR  
21 variations in Athens, Greece, focusing mainly on two known dominant factors, clouds and aerosol  
22 load.

##### 23 **4.1 The role of clouds**

24 Figure 8 shows the 1954-2012 time series of the monthly and yearly anomalies based on daily SSR,  
25 together with yearly total cloud coverage in weighted octas. The yearly de-seasonalised SSR values  
26 for all-sky conditions show a drop of ~14% from 1960 to 1970 and then a continuous increase




1 excluding the Pinatubo period in the early 1990's. Most pronounced positive changes can be seen  
2 during the last 15 years with a change of the order of about 15%.  
3



4  
5 **Figure 8.** De-seasonalised monthly mean (i.e. deviation from the respective monthly mean  
6 calculated over the whole period; dash line) and yearly mean SSR (black line), along with mean  
7 annual total cloud cover (green line).

8  
9 Figure 8 suggests anti-correlation between variations in SSR and cloud cover. Figure 9 shows the  
10 correlation between annual mean SSR and cloud cover. From the best-fit linear regression line it is  
11 seen that a  $-1.54 \text{ MJ m}^{-2}$  (or  $-9.6\%$ ) change in mean daily insolation accounts for a change of 1 octa  
12 in cloud cover.

13



1

2 **Figure 9.** Correlation between annual means of daily insolation and cloud cover over the period  
3 1954-2012. The straight line represents the best-fit regression line to the data points. The year 1953  
4 has not been included in the analysis since it does not include measurements for all months.

5

6 However, the great scatter of the data points and the low correlation of the two parameters in Figure  
7 9 ( $R^2=0.229$ ) indicate that the presence of cloud cover can only partly explain the changes in SSR.

8

9 In addition, there is no significant change in cloudiness over the 59 year period for Athens, Greece.

9

10 Calculating linear changes of cloudiness from data shown Figure 8, shows a non significant change  
11 of -0.4% per decade which can practically have a limited effect on SSR changes during the  
12 examined period.

12

13 Nevertheless, it is worth mentioning that different cloud properties like cloud optical thickness and  
14 cloud phase, not described by the measurements of cloud cover, can influence SSR.

14

15

## 16 **4.2 Clear sky records**

17

18 In order to minimize the cloud influence and investigate the possible role of direct aerosol effects on  
19 Athens SSR series, we had to select clear-sky (or cloudless) days. We have used daily SSR  
20 measurements from 1953 to 2012 and we have separated the cloudless days according to the criteria  
21 mentioned in Section 2.2.

21

22 For considering the SSR seasonality, we have calculated a five-degree polynomial derived from the  
23 maximum daily SSR (for all years of the data set), as a function of the day of the year (Figure 10).

23

24 Afterwards we have calculated the ratio of the daily SSR to the SSR calculated by this function.


24

25 Seasonal and yearly means of this ratio have been estimated and have been used to describe  
cloudless-sky SSR percentage changes on a seasonal and yearly basis. This approach has been



1 chosen since averaging a random set of cloudless days, within each month during the 59-year  
2 period, could cause solar elevation-related (due to the change of maximum solar elevation within  
3 each month) discrepancies, when calculating the monthly average SSR.

4




5

6 **Figure 10.** Clear-sky SSR measurements (red dots) and all-sky SSR measurements (blue dots)  
7 derived with the cloud octa (cloudiness<1.5) and sunshine duration (SD>0.9) related criteria. The  
8 black line represents the polynomial fit to the daily SSR<sub>max</sub> values.

9

10 Using the clear sky conditions seasonal and yearly averages of SSR have been calculated. The use  
11 of seasonal instead of monthly SSR has been introduced in order to improve the averaging SSR-  
12 related statistics, since the average number of cloudless days (per year) can be relatively low  
13 especially during the winter months. For all cases the ratios of the mean daily cloudless SSR to the  
14 SSR<sub>max</sub> derived from the daily best-fit curve in Figure 10 has been calculated and deviations of this  
15 ratio from its 59-yr mean have been calculated for each year.

16



1  
2 **Figure 11.** Changes in yearly mean SSR to relative to the 1954-2012 average for cloudless sky (in  
3 %; blue), AVHRR AOD series (red) and ChArMEEx AOD climatology (blue; Nabat et al., 2013) for  
4 Athens area is shown in the right axis.


5  
6 Figure 11 shows that most of the SSR variation observed for the measuring period has to be  
7 explained by other factors than changes in cloudiness (see figure 8 for variations due to cloudiness).  
8 Different seasons with the exception of wintertime show similar patterns to the year-to-year  
9 variability. Individual seasonal calculated SSRs do not exceed by more than  $\pm 5\%$ , the SSR  
10 variability of all sky data, with the exception of the winter season. Comparing clear sky and all sky  
11 yearly mean SSR, we find a high correlation ( $R^2 = 0.71$ ), which can be explained as a combination  
12 of: aerosol changes driving the SSR changes and by the number of clear sky days during the year.  
13 There is a decrease of more than 15% in the clear sky SSR from the start of the series to the end of  
14 1960's. A decline after 1983 could possibly be related with El Chichon volcanic eruption.

15 The Pinatubo-related drop of 10% from the early 1990's to the mid 1990's, can also be seen in both  
16 cloudless and all-sky datasets and also to the increase in AOD in the AVHRR dataset (Figure 11).  
17 Finally, the  $\sim 13\%$  change from 1995 to 2012 shown for all skies (Fig. 8) and clear skies (Fig. 11) is  
18 accompanied with a drop of  $\sim 25\%$  in AOD measured by AVHRR. The year to year variations of  
19 clear sky SSR series and the AVHRR-related AOD show an anti-correlation with  $R = -0.78$  ( $N = 29$ ),  
20 verifying the hypothesis that SSR clear sky changes are associated with aerosol load changes, at  
21 least within the common AVHRR/measurement period (1982-2009).

22 Similar to the AVHRR data the ChArMEEx 4-D aerosol climatology is shown in figure 11, providing  
23 similar conclusions, namely the AOD negative trend of 0.03 or 14% per decade from 1979 to 2012.



1 Differences between the AVHRR and ChaArMEx data can be explained in part by the different  
2 AOD wavelengths presented here (630 vs 550 nm) and also by a general negative bias of AVHRR  
3 over the Mediterranean compared to AERONET (Nabat et al., 2014). The smooth decline in the  
4 ChArMEx AOD data is due to the method used to build this product and uses the trend and not the  
5 interannual variability which is not included in the global model that was used.



6  
7 Figure 12. Ratio of cloudless vs all days, per season and yearly

8  
9 In addition to figure 11 we have included figure 12 showing the ratio of cloudless days to all  
10 available days for each season and for each year. Figure 12 shows a minimum (less than 30%  
11 during a year) of the number cloudless days from mid-1970's to early 1990's. It is mostly linked  
12 with the decrease of cloudless days during summer months. The figure provides a hint on the SSR  
13 relative changes observed during this period, but it can not directly interpret year to year SSR  
14 changes as they depend also on cloud fraction and properties for cloudy days. In addition, it can  
15 only partly be linked with fig. 11 as aerosol effects on cloudless sky calculated SSRs depend mostly  
16 on AOD levels and not on the number of days included in the calculations.

17  
18 In Table 3 we have calculated the linear trends for the 1953-2012 period and for both clear sky and  
19 all sky measurements and the 1953-1982 and 1983-2012 sub-periods for clear sky measurements.  
20 Results show comparable changes per decade (2% for the clear sky and 1.5% for the all sky cases).  
21 Seasonal analysis show that clear sky trends for summer, autumn and winter months are higher than  
22 the ones derived for all skies. Such differences are linked with the seasonal variability and long-  
23 term changes of cloudiness for the specific seasons.



1  
2 Table 3: Clear sky and all sky data trends comparison for the whole 1953-2012 period and the two  
3 30-yr sub-periods (% per decade). Percentages in parenthesis show the limits of the 95% confidence  
4 bounds.

Season	Clear sky 1953-2012	All skies 1953-2012	Clear sky 1953-1982	Clear sky 1983-2012
Winter	0.91 ( $\pm 2.31$ )	-6.43 ( $\pm 3.83$ )	-7.01 ( $\pm 3.16$ )	0.55 ( $\pm 2.41$ )
Spring	1.22 ( $\pm 1.12$ )	-0.60 ( $\pm 3.10$ )	-0.92 ( $\pm 1.11$ )	2.62 ( $\pm 1.97$ )
Summer	2.03 ( $\pm 0.78$ )	-1.14 ( $\pm 2.90$ )	-0.36 ( $\pm 0.83$ )	1.31 ( $\pm 0.81$ )
Autumn	2.74 ( $\pm 1.37$ )	-1.28 ( $\pm 3.42$ )	-1.03 ( $\pm 1.84$ )	-1.48 ( $\pm 1.73$ )
Year	<b>2.17 (<math>\pm 1.21</math>)</b>	<b>-2.33 (<math>\pm 2.28</math>)</b>	-1.44 ( $\pm 2.35$ )	1.94 ( $\pm 2.08$ )

5  
6 Clear sky results for the 1953-2012 period show significant positive changes in SSR for all seasons  
7 except winter. Looking individually at the 1953-1982 and 1983-2012 periods we have calculated  
8 significant negative trends only for the winter over the first and for summer and spring over the  
9 second.

10  
11 The effect of various parameters on SSR has been discussed by Kambezidis et al. (2016) in their  
12 study about the global dimming/brightening effect over the Mediterranean in the period 1979-2012;  
13 they show that the influence of parameters related to the atmospheric transparency, like water  
14 vapor, aerosols and trace gases, as well as changes in the surface albedo on SSR have been larger in  
15 the southern parts of the Mediterranean, over the Balkan countries and central Turkey. This  
16 outcome is in agreement with the conclusion of the present study that other factors than cloudiness  
17 play significant role in the SSR variations.

18  
19

## 20 5 Conclusions

21 Surface solar radiation (SSR) at National Observatory of Athens, in the center of the city, is  
22 presented using a unique dataset covering a period of 59 years (1954-2012). Sunshine duration (SD)  
23 records for another 54 years have been used as a proxy to reconstruct SSR time series for the period  
24 from 1900 to 2012.

25 A comparison of the SSR results in Athens with visibility observations since 1931 (Founda et al.,  
26 2016b) did not show any correlation among SSR and horizontal visibility. The steep visibility



1 decrease from 1931 till early 90's is not accompanied by a relative SSR decrease excluding  
2 individual sub-periods. Studying the literature for similar cases, similar conclusions have been  
3 drawn by Liepert and Kukla (1997) showing an SSR decrease over 30 years of measurements  
4 accompanied by a visibility increase and no significant changes in the cloud cover conditions, in  
5 Germany. This Athens SSR vs visibility relationship can be partly explained by the fact that: SSR  
6 and visibility have different response on cloud conditions, water vapor and rainfall and also by the  
7 fact that visibility is affected by aerosols only in the first few hundred meters above the surface,  
8 while SSR is affected by the columnar AOD, which in the case of Athens can be significantly  
9 different due to aerosol long-range transport in altitude (e.g. Saharan dust; Léon et al., 1999). The  
10 data accuracy of such historic radiation measurements is generally not well established, at least  
11 compared with the measurements after the 1990s. Quality assessment procedures in the presented  
12 time series have been applied based on criteria based on instrument characteristics and the  
13 availability of additional collocated measurements. Year to year fluctuations of the measured SSR  
14 in addition to the reversal of the downward tendencies at the Athens site adds credibility to the  
15 measured variations. That is because a typical radiometer behavior is to lose sensitivity with time  
16 indicating spurious downward, but not upward trends. The more recent (after 1989) SSR  
17 measurements can be characterized as high-quality radiation data with known accuracy.

18 De-seasonalized SSR data analysis from 1900 to 2012 showed high month to month variability that  
19 could reach up to 25%, mainly related with monthly cloudiness variations. During the period 1910-  
20 mid-1930s where only few datasets have reported worldwide SSR results, we observe a 2.9% per  
21 decade or a total of 8.7 % decrease in SSR, assuming linear changes in SSR during this period. This  
22 early dimming was followed by a 5% per decade increase from 1930 to the 1950s. Similar results  
23 have been found at Washington DC and at Potsdam, Germany (Stanhill and Achiman, 2016). They  
24 have reported an early brightening at both locations in the 1930's. For the SSR measurement period  
25 of 1953 to 1980, European related studies presented in Wild (2009) showed a -1% down to -7%  
26 change per decade in SSR measurements over various European sites (dimming period). For the  
27 Mediterranean region, Manara et al. (2016) showed a decrease in the order of -2% to -4% per  
28 decade in Italy. We are reporting a change in SSR of -2% per decade in Athens. Finally, for the  
29 brightening (1990-2012) phase again Wild et al. (2009) reported a 1.6% up to 4.7% per decade  
30 positive change in SSR while we have calculated a +1.5% per decade, which is lower than the 3-6%  
31 per decade reported in Manara et al. (2016) for Italy. A summary of the above findings can be seen  
32 in table 4.

33



1  
2  
3  
4

Table 4: Summary of per cent SSR changes per decade for various locations

Period	Location	Trend % per decade	Reference
1893-2012	Potsdam, Germany	0.71	Stanhill and Achiman, 2014
1900-2012	Athens, Greece	0.40 ( $\pm 0.26$ )	This work
1959-1988	Europe	-2.0	Ohmura and Lang, 1989
1971-1986	Europe	-2.3	Norris and Wild, 2007
1959-1985	Italy	-6.4( $\pm 1.1$ ) / -4.4( $\pm 0.8$ )	Manara et al, 2016
1953-1982	Athens, Greece	-2.33( $\pm 2.28$ )	This work
1985-2005	Europe	2.5	Wild, 2009
1990-2012	Italy	6.0 ( $\pm 1.1$ ) / 7.7 ( $\pm 1.1$ )	Manara et al, 2016
1986-2013	Athens, Greece	0.80 ( $\pm 1.96$ )	This work

5

6 The decadal variations of SSR measured since 1954 at Athens, Greece, originate from the  
7 alterations in the atmosphere's transparency (namely by clouds and aerosols). Using an analysis of  
8 SSR calculations of all sky and clear sky (cloudless) days we end up that since cloud cover changes  
9 during the 59 period were very small, most of the observed decadal changes might be related with  
10 changes in the aerosol load of the area. An additional hint in support of this conclusion is the high  
11 correlation of clear sky and all sky yearly SSR. We also found an anti-correlation between either  
12 clear sky and all sky SSR measurements and AOD time series from AVHRR (1981-2009) or  
13 ChArMex (1979-2012). Looking at linear trends over the 59 year period clear sky changes per  
14 decade were 2% while it was 1.5% for all sky conditions. The most pronounced changes have been  
15 calculated for summer and autumn seasons (2% and 2.7% respectively).

16

## 17 Acknowledgements

18 This study contributes to the Chemistry-Aerosol Mediterranean Experiment (ChArMEX) Work  
19 package 6 on trends. The work was partly funded by the national project "Aristotelis", work package  
20 1: "Study of long term variations of Solar Radiation in the region of Athens". The authors wish to  
21 thank all the past and present NOAA staff members who carefully collected and archived the long-  
22 term data used in this study.

23



## 1 **References**

- 2 Angell, J.K.: Variation in United States cloudiness and sunshine duration between 1950 and the  
3 drought year of 1988, *Journal of Climate* 3, 296:308, 1990.
- 4 Ångström, A.: Solar and terrestrial radiation. Report to the international commission for solar  
5 research on actinometric investigations of solar and atmospheric radiation, *Quarterly Journal of the*  
6 *Royal Meteorological Society* 50.210: 121-126, 1924.
- 7 Anton M., Vaquero J.M., Aparicio A.J.P.: The controversial early brightening in the first half of  
8 20th century: A contribution from pyrheliometer measurements in Madrid (Spain),  
9 10.1016/j.gloplacha.2014.01.013, 2014.
- 10 Chan, P. K., Zhao, X. P., and Heidinger, A. K.: Long-Term Aerosol Climate Data Record Derived  
11 from Operational AVHRR Satellite Observations, *Dataset Papers in Geosciences*, 140791,  
12 doi:10.7167/2013/140791, 2013.
- 13 Coulson, K.L.: *Solar and Terrestrial Radiation: Methods and Measurements*, Academic Press, 1975.
- 14 Drummond, A.J. and Roche, J.J.: Corrections to be applied to measurements made with Eppley (and  
15 other) spectral radiometers when used with Schott colored glass filters, *Journal of Applied*  
16 *Meteorology*, 4(6), pp.741-744, 1965.
- 17
- 18 Dudok de Wit, T., Ermolli, I., Haberreiter, M., Kambezidis, H., Lam, M., Lilensten, J., Matthes, K.,  
19 Mironova, I., Schmidt, H., Seppälä, A., Tanskanen, E., Tourpali, K. & Yair, Y. (Eds.): Earth's  
20 climate response to a changing sun, Les Ulis Cedex: EDP Sciences , doi:10.1051/978-2-7598-1733-  
21 7, 2015.
- 22 .
- 23



- 1 Founda, D., Kalimeris A., Pierros F.: Multi annual variability and climatic signal analysis of sun-  
2 shine duration at a large urban area of Mediterranean (Athens). *Urban Climate*,  
3 <http://dx.doi.org/10.1016/j.uclim.2014.09.008>, 2014.
- 4
- 5 Founda, D., Kazadzis, S., Mihalopoulos, N., Gerasopoulos, E., Lianou, M. and Raptis, P.I.: Long-  
6 term visibility variation in Athens (1931–2013): a proxy for local and regional atmospheric aerosol  
7 loads. *Atmospheric Chemistry and Physics*, 16(17), pp.11219-11236, 2016.
- 8 Founda, D., Pierros, F. and Sarantopoulos, A.: Evidence of Dimming/Brightening over Greece from  
9 long-term observations of Sunshine Duration and Cloud Cover, *Perspectives in Atmospheric*  
10 *Sciences*, pp. 753-759, Springer, ISBN 978-3-319-35094-3, 2016.
- 11
- 12 Gilgen, H., Wild, M. and Ohmura, A., 1998. Means and trends of shortwave irradiance at the  
13 surface estimated from global energy balance archive data. *Journal of Climate*, 11(8), pp.2042-  
14 2061.
- 15
- 16 Hulstrom, R. L.: “Solar resources”, is the volume 2 in the Series: “Solar Heat Technologies:  
17 Fundamentals and Applications”, edited by Charles A. Bankston, The MIT Press, Cambridge, 1989.
- 18
- 19 ISO 9847: Solar energy - Calibration of field pyranometers by comparison to a reference  
20 pyranometer, International Organization for Standardization, 1992,
- 21
- 22 Jauregui, E., and E. Luyando.: Global radiation attenuation by air pollution and its effects on the  
23 thermal climate in Mexico City, *International Journal of Climatology* 19, no. 6: 683-694, 1999.
- 24 Kambezidis, H., Demetriou D., Kaskaoutis, D., Nastos, P.: Solar dimming/brightening in the  
25 Mediterranean EGU General Assembly 2010, held 2-7 May, 2010 in Vienna, Austria, p.10023,  
26 2010.
- 27 Kambezidis, H., Kaskaoutis, D., Kalliampakos, G., Rashki, A. and Wild, M.: The solar  
28 dimming/brightening effect over the Mediterranean Basin in the period 1979 - 2012, *Journal of*  
29 *Atmospheric and Solar-Terrestrial Physics*, <http://dx.doi.org/10.1016/j.jastp.2016.10.006>, 2016.
- 30
- 31 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C.,  
32 Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J.,



- 1 Cooper, O. R., Kainuma, M., Mahowald, N., Mc-Connell, J. R., Naik, V., Riahi, K., and van  
2 Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of  
3 reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039,  
4 doi:10.5194/acp-10-7017-2010, 2010.
- 5
- 6 Lean, J.: The Sun’s variable duration and its relevance for earth, *Annual Review of Astronomy and*  
7 *Astrophysics* 35.1 33-67, 1997.
- 8
- 9 Léon, J.F., Chazette, P., and Dulac, F.: Retrieval and monitoring of aerosol optical thickness over an  
10 urban area by spaceborne and ground-based remote sensing, *Appl. Opt.*, 38, 6918-6926, 1999.
- 11
- 12 Liepert B. and Kukla G., Decline in Global Solar Radiation with Increased Horizontal Visibility in  
13 Germany between 1964 and 1990, *Journal of Climate*, 2391-2401, September, 1997.
- 14 LMDC,: Laboratory of Meteorological Device Calibration, <http://www.meteo.noa.gr/lmdc.html>, B.  
15 Psiloglou, Scientific Responsible, personal contact, November 2016.
- 16
- 17 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J.  
18 R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th Century industrial black carbon  
19 emissions altered Arctic climate forcing, *Science*, 317, 5843, doi:10.1126/science.1144856, 2007.
- 20
- 21 Manara, V., Brunetti, M., Celozzi, A., Maugeri, M., Sanchez-Lorenzo, A., and Wild, M.: Detection  
22 of dimming/brightening in Italy from homogenized all-sky and clear-sky surface solar radiation  
23 records and underlying causes (1959–2013), *Atmos. Chem. Phys.*, 16, 11145-11161,  
24 doi:10.5194/acp-16-11145-2016, 2016.
- 25 Matuszko, D.: Long-term variability in solar radiation in Krakow based on measurements of  
26 sunshine duration, *Int. J. Climatol.* 34: 228 – 234, 2014.
- 27 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J.  
28 R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th Century industrial black carbon  
29 emissions altered Arctic climate forcing, *Science*, 317, 5843, doi:10.1126/science.1144856, 2007.



- 1 Mishchenko, M. I., Geogdzhayev, I. V., Rossow, W. B., Cairns, B., Carlson, B. E., Lacis, A. A.,  
2 Liu, L., and Travis, L. D.: Long-term satellite record reveals likely recent aerosol trend, *Science*,  
3 315, p. 1543, doi:10.1126/science.1136709, 2007.  
4
- 5 Nabat, P., Somot, S., Mallet, M., Chiapello, I., Morcrette, J.J., Solmon, F., Szopa, S., Dulac, F.,  
6 Collins, W., Ghan, S. and Horowitz, L.W.: A 4-D climatology (1979-2009) of the monthly  
7 tropospheric aerosol optical depth distribution over the Mediterranean region from a comparative  
8 evaluation and blending of remote sensing and model products, *Atmospheric Measurement*  
9 *Techniques*, 6(5), p.1287, 2013.  
10
- 11 Nabat, P., Somot, S., Mallet, M., Sanchez - Lorenzo, A. and Wild, M.: Contribution of  
12 anthropogenic sulfate aerosols to the changing Euro - Mediterranean climate since 1980,  
13 *Geophysical Research Letters*, 41(15), pp.5605-5611, 2014.
- 14 Norris, J.R. and Wild, M.: Trends in aerosol radiative effects over Europe inferred from observed  
15 cloud cover, solar “dimming,” and solar “brightening”. *Journal of Geophysical Research:*  
16 *Atmospheres*, 112(D8),2007.
- 17
- 18 Norris, J. R., and M. Wild : Trends in aerosol radiative effects over China and Japan inferred from  
19 observed cloud cover, solar “dimming,” and solar “brightening”, *J. Geophys. Res.*, 114, D00D15,  
20 doi:10.1029/ 2008JD011378, 2009.
- 21 Ohmura, A., Observed long-term variations of solar irradiances at the Earth’s surface, *SpaceSci.*  
22 *Rev.*, 125, 111–128, doi:10.1007/s11214- 006-9050-9, 2006.  
23
- 24 Ohmura, A.: Observed decadal variations in surface solar radiation and their causes, *J. Geophys.*  
25 *Res.*, 114, D00D05, doi:10.1029/2008JD011290, 2009.  
26
- 27 Ohmura, A. and H. Lang: Secular variation of global radiation over Europe, in *Current Problems in*  
28 *Atmospheric Radiation*, edited by J. Lenoble and J. F. Geleyn, pp. 98–301, Deepak, Hampton, Va. ,  
29 1989.



- 1
- 2 Ruckstuhl C, Philipona R, Behrens K, Collaud Coen M, Durr B, Heimo. A, Matzler C, Nyeki S,  
3 Ohmura A, Vuilleumier L, Weller M, Wehrli C, Zelenka A.: Aerosol and cloud effects on solar  
4 brightening and the recent rapid warming ,Geophysical Research Letters 35:L12708. DOI:  
5 10.1029/2008gl034228, 2008.
- 6
- 7 Sanchez-Lorenzo, A., Calbó J, Martin-Vide J : Spatial and temporal trends in sunshine duration  
8 over Western Europe (1938-2004), Journal of Climate 21 (22), 6089-6098, 2008.
- 9
- 10 Sanchez-Lorenzo, A., and Wild, M.: Decadal variations in estimated surface solar radiation over  
11 Switzerland since the late 19th century, Atmospheric Chemistry and Physics 12.18: 8635-8644,  
12 2012.
- 13
- 14 Sanchez-Lorenzo, A., Wild M., Brunetti M., Guijarro J. A., Hakuba, M. Z. , Calbó, J. , Mystakidis,  
15 S. and Bartok, S.: Reassessment and update of long-term trends in downward surface shortwave  
16 radiation over Europe (1939–2012), J. Geophys. Res. Atmos., 120, 9555–9569,  
17 doi:10.1002/2015JD023321, 2015.
- 18
- 19 Sanchez-Romero, A., A. Sanchez-Lorenzo, J. Calbó, J. A. González, and C. Azorin-Molina : The  
20 signal of aerosol-induced changes in sunshine duration records: A review of the evidence, J.  
21 Geophys. Res. Atmos., 119, 4657–4673, doi:10.1002/2013JD021393, 2014
- 22 Stanhill, G. and Achiman, O.: Early global radiation measurements: a review. Int. J. Climatol..  
23 doi:10.1002/joc.4826, 2016
- 24 Stanhill, G.: Global irradiance, air pollution and temperature changes in the Arctic, Philos. Trans. R.  
25 Soc. A, 352, 247 –258, doi:10.1098/rsta.1995.0068, 1995.
- 26
- 27 Stanhill, G., and Cohen S. : Recent changes in solar irradiance in Antarctica, J. Clim., 10, 2078–  
28 2086, doi:10.1175/1520-0442, 1997.



- 1 Stanhill, G. and Cohen, S.: Solar radiation changes in the United States during the twentieth  
2 century: Evidence from sunshine duration measurements. *Journal of Climate*, 18(10), pp.1503-  
3 1512, 2005.  
4
- 5 Stanhill, G. and Möller, M.: Evaporative climate change in the British Isles. *International Journal of*  
6 *Climatology*, 28(9), pp.1127-1137, 2008.  
7
- 8 Stanhill, G., and O. Ahiman: Radiative forcing and temperature change at Potsdam between 1893  
9 and 2012, *J. Geophys. Res. Atmos.*, 119, 9376–9385, doi:10.1002/2014JD021877, 2014.  
10
- 11 Streets, D. G., Ye W., and Mian C.: Two-decadal aerosol trends as a likely explanation of the global  
12 dimming/brightening transition, *Geophysical Research Letters* 33.15, 2006.  
13
- 14 Vautard, R. and Yiou, P.: Control of recent European surface climate change by atmospheric flow.  
15 *Geophysical Research Letters*, 36(22),2009.  
16
- 17 Wang, K. C., Dickinson, R. E., Wild, M., and Liang, S.: Atmospheric impacts on climatic  
18 variability of surface incident solar radiation, *Atmos. Chem. Phys.*, 12, 9581-9592,  
19 doi:10.5194/acp-12-9581-2012, 2012.
- 20 Wild M.: Global dimming and brightening: a review, *Journal of Geophysical Research* 114. DOI:  
21 10.1029/2008JD011470, 2009.  
22
- 23 Wild M, Folini D, Hakuba MZ, Schär C, Seneviratne SI, Kato S, Rutan D, Ammann C, Wood EF,  
24 König-Langlo G.: The energy balance over land and oceans: an assessment based on direct  
25 observations and CMIP5 climate models, *ClimDyn* 44:3393–3429. doi:10.1007/s00382-014-2430-  
26 z, 2015.
- 27 Wild, M.: Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for  
28 global warming. *WIREs Clim Change*, 7: 91–107. doi:10.1002/wcc.372, 2016



- 1 WMO: “Measurement of radiation”, in Guide to Meteorological Instrument and Observing  
2 Practices, Chapter 9, fifth ed., WMO-No. 8, 1983.
- 3 WMO: Scientific Assessment of Ozone Depletion: 2010, report 52, World Meteorological  
4 Organization (WMO), Global Ozone Research and Monitoring Project, Geneva, Switzerland;  
5 National Oceanic and Atmospheric Administration (NOAA), Washington, DC, USA; National  
6 Aeronautics and Space Administration (NASA), Washington, DC, USA; United Nations  
7 Environment Program (UNEP), Nairobi, Kenya; and the European Commission, Research  
8 Directorate General, Brussels, Belgium, 2010.  
9
- 10 Yildirim, U., Ismail, O. Y. and Akinoglu B. G.: Trend analysis of 41 years of sunshine duration  
11 data for Turkey, Turkish Journal of Engineering and Environmental Sciences 37.3: 286-305, 2014.  
12
- 13 Zerefos, C.S., K. Eleftheratos, C. Meleti, S. Kazadzis, A. Romanou, C. Ichoku, G. Tselioudis, and  
14 A. Bais: Solar dimming and brightening over Thessaloniki, Greece, and Beijing,  
15 China. Tellus, 61B, 657-665, doi:10.1111/j.1600-0889.2009.00425.x, 2009.  
16
- 17 Zerefos, C.S., Tourpali, K., Eleftheratos, K., Kazadzis, S., Meleti, C., Feister, U., Koskela, T. and  
18 Heikkilä, A.: Evidence of a possible turning point in solar UV-B over Canada, Europe and Japan,  
19 Atmospheric Chemistry and Physics, 12(5), pp.2469-2477, 2012.  
20
- 21 Zhao X., Chan, P., and NOAA CDR Program: NOAA Cli-mate Data Record (CDR) of AVHRR  
22 Daily and Monthly Aerosol Optical Thickness over Global Oceans, Version 2.0.AOT1, NOAA  
23 National Centers for Environmental Information, doi:10.7289/V5SB43PD, 2014.  
24