



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Consistent circulation differences in the Southern Hemisphere caused by ozone changes: a chemistry-climate model and observational study

P. Braesicke<sup>1</sup>, J. Keeble<sup>2</sup>, X. Yang<sup>1</sup>, G. Stiller<sup>3</sup>, S. Kellmann<sup>3</sup>, N. L. Abraham<sup>1</sup>, A. T. Archibald<sup>1</sup>, P. Telford<sup>1</sup>, and J. A. Pyle<sup>1</sup>

<sup>1</sup>NCAS/University of Cambridge, Chemistry Department, Cambridge, UK

<sup>2</sup>University of Cambridge, Chemistry Department, Cambridge, UK

<sup>3</sup>Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, Karlsruhe, Germany

Received: 30 January 2013 – Accepted: 12 March 2013 – Published: 28 March 2013

Correspondence to: P. Braesicke (pb261@cam.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

We report results from two pairs of chemistry-climate model simulations using the same climate model but different chemical perturbations. In each pair of experiments an ozone change was triggered by a simple change in the chemistry. One pair of model experiments looked at the impact of Polar Stratospheric Clouds (PSCs) and the other pair at the impact of short lived halogenated species on composition and circulation. The model response is complex with both positive and negative changes in ozone concentration, depending on location. These changes result from coupling between composition, temperature and circulation. Even though the causes of the modelled ozone changes are different, the high latitude Southern Hemisphere response in the lower stratosphere is similar. In both pairs of experiments the high latitude circulation changes, as evidenced by N<sub>2</sub>O differences, suggesting a slightly longer-lasting/stronger stratospheric descent in runs with higher ozone destruction. We contrast the idealised model behaviour with interannual variability in ozone and N<sub>2</sub>O as observed by the MIPAS instrument on ENVISAT, highlighting the similarity of the modelled climate equilibrium changes to the year 2006/2007 in observations. We conclude that the climate system can respond quite sensitively to small chemical perturbations, that circulation adjustments seen in the model can occur in reality, and that coupled chemistry-climate models are needed for an assessment of future ozone and climate changes.

## 1 Introduction

The recent United Nations Environment Programme/World Meteorological Organization (UNEP/WMO) ozone assessment (WMO, 2011) highlights the importance of chemistry-climate interactions, emphasising the strong link between composition differences and climate change. For example, to first order, when the concentrations of halogen radical species decrease in the stratosphere, we expect to see an increase in

ACPD

13, 8455–8487, 2013

### Consistent circulation differences in the SH

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ozone. However, that ozone change will be coupled to changes in temperature and circulation, feeding back further onto the initial ozone change. Changes in stratospheric temperature and circulation are also projected under increased greenhouse gas loading. Stratospheric temperatures will decline which, due to the temperature dependence of the controlling gas phase reactions, should lead to ozone increases away from polar latitudes. In addition, climate model simulations with increasing long-lived greenhouse gases consistently show a strengthening (e.g. enhanced tropical upwelling at 70 hPa) of the stratospheric overturning circulation, the so-called Brewer–Dobson Circulation (BDC; after Brewer, 1949; Dobson, 1956), which will affect the meridional distribution of ozone in the stratosphere. In consequence modelled stratospheric age-of-air<sup>1</sup> decreases in the extratropical stratosphere (Butchart et al., 2006). However, age-of-air trends derived from trace gas observations (Engel et al., 2009) have not been able to confirm this model result. Recently, Stiller et al. (2012) highlighted the inhomogeneous spatial behaviour of age-of-air trends derived from satellite observed SF<sub>6</sub>. This could be a possible reason why Engel et al. (2009) in their spatially aggregated data set where unable to detect a trend. It also challenges modellers to understand better the causes of regional stratospheric circulation changes and age-of-air differences.

A number of studies have emphasised chemistry-climate interactions in the Southern Hemisphere (SH). There is clear evidence for the Southern Hemisphere ozone hole having caused a change in the SH polar vortex dynamics (Thompson et al., 2011) and a shift in the tropospheric jets (Son et al., 2010). Furthermore, different levels of ozone depletion, in different models, tend to modify surface climate modes (e.g. Gillett and Thompson, 2003; Morgenstern et al., 2008), including the Southern and Northern Annual Modes. Changes in the stratospheric vortex/jet strength and position are also reflected in the overturning speed of the BDC.

Note that the BDC is not just one single overturning cell. It can be envisaged as a hemispheric to global scale circulation with three distinct height regimes as illus-

<sup>1</sup>Age-of-air or, for short, age is a measure for the time it takes an air mass to travel from the troposphere to a particular point in the stratosphere (see, e.g. Waugh and Hall, 2002).

**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



trated in Fig. 1, following Plumb (2002): a global scale branch with air flowing from the summer to the winter pole in the upper stratosphere and mesosphere (red), an upper/middle stratospheric branch with air flowing from the tropical region to middle and high latitudes in the winter hemisphere (orange), and a lower stratospheric/upper tropospheric branch flowing from tropical to middle and high latitudes in both hemispheres at all seasons, notwithstanding some seasonal modulation (green). All three branches are relevant to determining the composition and mean age of SH polar air in the lower stratosphere. We will investigate in idealised model experiments how a chemical change impacts the balance of the three BDC branches in the SH and, in turn, also affects composition. We will present evidence that such modulation of the BDC can be detected in observations.

We report results from two pairs of chemistry-climate model simulations using the same climate model with different chemistry schemes. Differences between the two runs in each pair of experiments show some commonalities regarding composition and circulation changes in the SH. We aim to characterise and explain the qualitatively similar changes and to put them into context with observed interannual variability. We do not provide a detailed analysis of the individual model runs (this is left to other publications in the near future from the co-authors Alex Archibald, James Keeble and Xin Yang). Here, we point out the common features in the differences between the two integrations in each of the two pairs of experiments that illustrate a robust chemistry-climate feedback between changes in ozone depletion and circulation differences.

We start with discussing details of our runs and the analysis strategy (Sect. 2), followed by the detailed description of the modelled response (Sect. 3). Observational considerations and a case study supporting the model results are presented in Sect. 4. Section 5 summarises and presents a generalisation of the findings.

## 2 Method

The model is an update of the model described in Morgenstern et al. (2009). The underlying Unified Model (UM) atmospheric climate model has been updated from version 6.1 to 7.3 (Hewitt et al., 2011). The integrations are performed with a horizontal resolution of 3.75° in longitude and 2.5° in latitude on an Arakawa-C grid. A hybrid sigma-geometric height coordinate is used to resolve the vertical range from the surface up to ~84 km with 60 levels, resulting in a vertical resolution of just above 1 km in the lower stratosphere. All model integrations are performed for year 2000 boundary conditions. Averaged sea surface temperatures (SSTs) and sea ice concentrations (SICs) are prescribed. There is an annual cycle in the SST and SIC boundary conditions, but no interannual variability. The model can use a number of different chemistry schemes. Here a *Chemistry of the Stratosphere* (CheS) scheme based on Morgenstern et al. (2009), including some minor updates, is used within UM7.3 in experiment pair A. In experiment pair B the chemistry scheme includes a more detailed tropospheric volatile organic compound mechanism, and additional halogenated short lived species and related reactions (CheST+; *Chemistry of the Stratosphere and Troposphere with additional halogenated short lived species (+)*)<sup>2</sup>. Both model set-ups use trace gases interactively in the calculation of heating rates allowing for feedbacks between changing circulation and composition.

For each model set-up a pair of experiments is performed, as summarized in Fig. 2. One pair of model experiments (Pair A using the CheS model set-up) looks at the impact of PSCs. The second pair (Pair B using the CheST+ model set-up) considers the impact of short lived halogenated species on composition and circulation. In the following we simply refer to pair A and pair B. Pair A consists of one model integration allowing chlorine activation on Polar Stratospheric Clouds (PSCs) and one inhibiting the activation on PSCs. In this case, we expect a large impact in polar latitudes, with

<sup>2</sup>Effectively this scheme is an extended merger of CheS (see above) and the standard tropospheric chemistry CheT as described in O'Connor et al. (2010).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the integration that includes activation on PSCs leading to much lower ozone there. Pair B consists of one model integration where short lived halogenated species are emitted in addition to the usual long-lived halogenated species and one not emitting these short-lived species. In this case we anticipate that the ozone differences will be largest in the very low stratosphere and have a more global extent than for pair A.

For each pair of integrations, differences are calculated by subtracting the run in which we expect higher ozone (lower ozone depletion) from the run with potentially higher ozone depletion, following the general rule of thumb, above, that a larger availability of halogenated species will lead to larger ozone destruction. This should lead to consistent signs in the model differences (Fig. 2). All runs are spun-up for 10 yr and are integrated for 20 (pair A) and 10 (pair B) yr, respectively.

To characterise the changes in circulation and composition we will consider model fields of nitrous oxide ( $N_2O$ ; lifetime of around 120 yr, with its source in the troposphere), ozone ( $O_3$ ; lifetime of weeks to month in the lower stratosphere, with its major source in the stratosphere), mean age of air (an idealised tracer increasing linearly with time at the model surface, e.g. Waugh and Hall, 2002), tropopause height and temperature.

### 3 Model response

We start our discussion by looking at the changes in total ozone (Sect. 3.1). Subsequently we discuss the changes in tropopause height (Sect. 3.2), before detailing the zonal mean ozone and temperature changes in December (Sect. 3.3). Using high latitude  $N_2O$  we will discuss changes in the large scale circulation and support these conclusions by looking at the age-of-air differences for December (Sect. 3.4).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.1 Total ozone changes

Figure 3 shows the latitude-time evolution of zonal mean total ozone differences for the two experiment pairs A (left) and B (right). Stippled areas mark out regions where differences exceed one standard deviation of the run subtracted, thus indicating a likely significant change between runs in a pair. Negative differences dominate in high southern latitudes in both experiment pairs. As expected Pair A, with PSCs switched on/off, and hence with the largest expected changes in activated chlorine, shows the larger negative change on the SH.

The response in the NH is quite complex, and in both cases some areas of ozone increase can be seen. Based simply on chemical arguments, for one experiment in each pair we would expect just to see an ozone decrease. The increases must be related to circulation changes, e.g. possibly enhanced poleward transport during NH winter in pair A and an earlier vortex break-up in pair B, but the statistical significance of the modelled features is small.

In the SH the following picture emerges: the modelled timing of the maximum ozone deficit in the SH differs between the experiment pairs. The largest negative anomaly occurs in pair A between October and December, whereas the smaller negative anomaly in pair B occurs slightly later. In addition to the chemically induced ozone differences, dynamical changes also occur, as we will show below. For example, the lower stratospheric branch of the BDC acts as a year-round supply route for ozone from low to high latitudes. In contrast, the middle stratospheric branch of the BDC fulfils this role during winter (e.g. Grewe, 2006). As a consequence the relative effectiveness of these transport routes has a strong seasonality that plays a role in maintaining (or wiping out) high latitude ozone anomalies from spring to summer, when the supply of ozone from the middle stratospheric branch of the BDC decays. Depending on the strength of the lower stratospheric BDC the lifetime of the SH springtime “ozone mass deficit” (Bodeker et al., 2005) can either be enhanced or diminished. In our experiments the negative high-latitude SH ozone anomalies are long-lived from austral spring to sum-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mer, indicative of a weakened lower stratospheric BDC, similar to the pattern seen in Braesicke et al. (2006) for a doubling of CO<sub>2</sub> in a predecessor model. In addition, observational evidence from SF<sub>6</sub> derived age-of-air data supports this statement (Stiller et al., 2012), showing “old” polar air masses until late summer. Given the proposed importance of circulation changes, we explore these next. In the following subsection we will link the ozone changes to the modelled tropopause height changes and, then, more generally to BDC changes by discussing the changes in modelled N<sub>2</sub>O distributions.

### 3.2 Tropopause height changes

Figure 4 shows the latitude-time evolution of zonal mean tropopause height differences for both experiment pairs. Positive differences dominate high SH latitudes, maximising in southern polar latitudes during austral summer when the ozone differences are also highest. Even though the SH responses are very similar in both experiment pairs, the Northern Hemisphere spring evolution differs between Pairs A and B.

The total ozone differences shown in Fig. 3 are consistent with the tropopause behaviour seen in Fig. 4. Generally, a higher than average troposphere will result in lower than average column ozone in most latitudes. Dynamical control produces a shallower region of stratospheric ozone mixing ratios in the lowermost stratosphere above a high tropopause, leading to lower column ozone values when integrated vertically.

However, there are some differences in timing between ozone deficits (Fig. 3) and tropopause heights (Fig. 4). In pair A the largest ozone deficit precedes the period of highest tropopause; in pair B the largest ozone deficit occurs simultaneously with the highest tropopause. This difference is related to the more seasonal and regional change of ozone in A where chemical ozone loss is suppressed in one integration by neglecting halogen activation on PSCs during Antarctic spring, leading to changes in the thermal structure and thus an elevated tropopause, which by its very existence prolongs the existence of a low ozone episode. Due to the strong chemical contribution the largest ozone deficit precedes the highest elevation of the tropopause. In experiment pair B the ozone change is more modest and stays in synchronisation with

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the tropopause change. Even though the initial “disturbance” to the system is the enhanced availability of reactive halogens which leads to increased ozone destruction, the dynamical signature (e.g. high tropopause/low ozone during December/January in the SH) dominates the new equilibrium climate state.

To explore the common features during austral summer in more detail, we will next investigate the latitude-height structure of ozone and temperature changes for December.

### 3.3 Ozone and temperature changes in December

Figure 5 shows long-term monthly mean latitude-height cross-sections of ozone differences for December, when peak column ozone differences occur in the Southern Hemisphere (Fig. 3). Pairs A and B both show a consistent decrease in lower stratospheric ozone in high southern latitudes in response to the changes implemented in each pair of model integrations.

As expected, in pair A the magnitude of ozone loss is larger in response to the suppression of halogen activation on PSCs. Pair B has the maximum loss slightly higher in altitude; in this case, halogen activation on PSCs is present in both runs but the abundance of reactive halogen is increased relative to A. In pair A the ozone difference south of 75° S and between 15 to 20 km exceeds 60%. In pair B only up to 30% difference is modelled in the Antarctic stratosphere. Even though the general expectation for the experiments is increased ozone loss, it should be noted that there are some areas in the mid and upper stratosphere where ozone is slightly increased in response to complex composition-climate for which it is difficult to disentangle cause and effect.

The large tropospheric changes of ozone in pair A are in part due to the simple tropospheric chemistry in CheS, and also reflect changes in ozone exchange between the lowermost stratosphere and the uppermost troposphere; summer tropospheric ozone production does not mitigate the downward transport of a stratospheric ozone deficit. CheST+ (pair B) includes a comprehensive tropospheric chemistry and together with the smaller ozone changes in the lowermost stratosphere, tropospheric changes are

**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



also much smaller and less significant. The negative ozone anomalies in Fig. 5 are largely triggered by increased chemical ozone destruction, but they happen in conjunction with transport changes, in particular in high latitudes. To link the ozone changes more closely to transport and climate changes we will discuss the change in temperature during austral summer next.

Figure 6 shows the temperature differences for experiment pairs A and B as a latitude-height cross-section. Note that we focus our discussion here on SH early summer (December), when the high latitude lower stratospheric ozone changes established in spring are still significant and will impact on the radiation in the model. In the SH the temperature response relates directly to the ozone differences shown in Fig. 5, with SH polar temperatures being lower where ozone is depleted (between 10 to 22 km) due to lower short-wave heating rates. Above 25 km higher temperatures indicate largely a dynamical response with enhanced descent causing adiabatic warming. Due to feedbacks the situation is more complicated in lower latitudes. In some areas ozone increases where temperatures are lower, thus highlighting the slow-down of the gas-phase ozone loss reactions (e.g. the tropical stratosphere above 25 km in B). In other areas signatures are far more complex and are determined by ozone differences above altering photolysis rates below, in conjunction with thermal and circulation changes.

For both model pairs the South Polar responses are similar, with a larger magnitude in the case with a more severe ozone loss (pair A). The tropical and NH polar changes are very different between the two pairs of experiments and are not significant, due to high dynamical variability during NH winter. Despite the SH high latitude similarities the tropical changes also differ between the pairs of experiments, with a negative temperature anomaly around and above the tropopause in pair A and a positive anomaly in pair B. This is also evident in the magnitude of the transport changes modelled in  $N_2O$  that are discussed in the next section.

## Consistent circulation differences in the SH

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.4 Circulation changes

The use of tracer data to characterise transport changes is well established (e.g. Ray et al., 2002, 2010). Here we use  $N_2O$  to visualise changes in the modelled BDC. Figure 7 shows the seasonal progression of  $N_2O$  (isolines) and  $N_2O$  differences (shaded) in high southern latitudes (averaged over all latitudes south of  $70^\circ S$ ) from the surface to 40 km for both pairs of model integrations.

The absolute  $N_2O$  values (from the “high ozone” run that is subtracted in each pair of experiments) are shown as isolines;  $N_2O$  differences between runs in each experiment pair are shaded. Starting in SH late summer (February) the downward slope of the isolines is indicative of the high-latitude descending branch of the Brewer–Dobson Circulation. Following Plumb (2002) and as indicated in Fig. 1, during autumn and winter the two upper vertical regions of the BDC contribute to the pronounced descent visualised by the downward slope of the  $N_2O$  isolines. During spring a strong vertical gradient in  $N_2O$  is maintained. Note how rapidly  $N_2O$  increases again in early summer (November/December) above 25 km when the upper part of the BDC undergoes significant seasonal change. Positive anomalies in autumn (persisting until winter) indicate stronger meridional transport and decreased descent in the “low ozone” run, which can be interpreted as a strengthening of the meridional drift of the middle BDC branch bringing more  $N_2O$  to high latitudes and a weakening of the uppermost descending BDC branch, which would otherwise bring low  $N_2O$  down. Subsequently the negative anomalies below 25 km during spring/summer indicate an enhanced/prolonged descent bringing low  $N_2O$  down in altitude over a longer period during which ozone concentrations are decreased. In addition, a weaker middle/lower stratospheric BDC branch does not replenish high latitude  $N_2O$  as effectively from the tropical entry point. The BDC changes are also reflected in a change of stratospheric vortex lifetime. The vortex lasts 15 to 20 days longer in integrations with more severely depleted ozone (more details are discussed below).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Positive anomalies around 15 km indicate a strengthening of the lowermost stratospheric BDC branch and at 25 km a strengthening of the middle stratospheric BDC branch, an assertion that is verified by the transformed Eulerian mean circulation diagnostics (Andrews et al., 1987) available from pair A (not shown), in which the residual mean velocities are generally more poleward. In contrast, positive N<sub>2</sub>O anomalies just above the Antarctic tropopause result from enhanced mixing with tropospheric air (see below).

Figure 8 shows latitude-height cross-sections of relative mean age-of-air differences above the tropical tropopause for each pair of experiments during December. For both pairs, the mean age-of-air difference shows a dipole structure in high southern latitudes with younger air just above the tropopause and older air around or slightly above 15 km. This pattern suggests more efficient mixing slantwise across the tropopause in high latitudes (younger air anomaly, blue) and a weakened lower to middle stratospheric BDC branch, advecting less young air from lower latitudes (older air anomaly, orange). This is consistent with the longer-lived SH polar vortex in runs with more ozone depletion, because a more pronounced vortex edge suppresses meridional transport. In both pairs of experiments the general structure of the high latitude SH pattern is similar. The amplitude of the feature is slightly larger in pair B, where we expect a more year-round global change of chemistry, compared to the more regional and seasonal change in pair A (the PSC experiments). Note that the negative ozone anomaly south of 60° S in A is at lower altitudes compared with B (Fig. 5). Consequently the change in ozone gradient across the Antarctic tropopause is different in both pairs of experiments. The larger vertical extent of the area of younger air above the Antarctic tropopause in B is consistent with the longer duration of a positive anomaly in N<sub>2</sub>O (Fig. 7), both indicating a more effective exchange of air between the upper troposphere and the upper lowermost stratosphere during December in B as compared to A, where the positive N<sub>2</sub>O anomaly starts earlier and does not last throughout December.

Changes in age north of 60° S differ between experiment pairs A and B. In A age is generally decreased whereas in B age is generally higher (B being the experiment

**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pair with less seasonality). Note that the change in A is nearly everywhere significant (stippled area, as defined in Sect. 3.1), whereas the change in B is only significant in high southern latitudes and, to some degree, in northern middle latitudes. Note, also, that the differences in age-of-air between pairs A and B away from the SH polar region are consistent with the modelled changes in zonal mean temperature, with younger air (blue) in A where colder temperatures at the tropopause indicate a strengthened upwelling and older air (yellow) in B where warmer temperatures at the tropopause indicate a weakened upwelling (Fig. 6). The ozone change is more complex, because it depends on a coupled chemistry-transport response. In A the tropical lower stratospheric decrease in ozone is consistent with the increased uplift. In both experiment pairs the ozone changes in the upper stratosphere are anti-correlated with the temperature changes, as expected for ozone in photochemical steady state. In B, the decrease in ozone in the tropical lower stratosphere is driven by the change in available bromine from the short lived halogenated species that occurs at all latitudes and during all months.

In summary, the changes in ozone and circulation are intimately linked with temperature and tropopause height differences, with a higher tropopause indicating a stronger tropospheric influence on the lowermost stratosphere and hence younger air. Interpreting our results it is important to appreciate the more seasonal and regional trigger in A in contrast to the year-round, global trigger in B. By definition the changes in each pair are different, but they are embedded in the naturally occurring seasonal cycle. Despite many differences, some of the impact manifests itself in similar changes in hemispheric transport patterns, especially near the South Pole. The similarities in SH N<sub>2</sub>O anomalies (Fig. 7) between A and B are indicative of similar changes in the seasonal progression of the BDC, and in the summer change in age-of-air (Fig. 8) that is linked to an elevated SH high-latitude tropopause. This links directly to the evidence chain presented in conjunction with Fig. 3 (above), relating the lifetime of a SH polar ozone mass deficit to the strength of the BDC.

**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 MIPAS observations

For nearly a decade (2002–2012) the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) instrument on ENVISAT (“Environmental Satellite”) has successfully observed atmospheric trace gas distributions. Here we use SF<sub>6</sub> derived age-of-air (Stiller et al., 2012) to validate the seasonality of the modelled high latitude circulation. The subsequent case study for 2006/2007 uses a recent version (V5R.x.220) of MIPAS N<sub>2</sub>O and ozone data processed at IMK/IAA (updated from von Clarmann et al., 2009) to highlight interannual circulation changes and accompanying ozone differences.

### 4.1 Seasonality

As a way of validation of the modelled seasonal progression of the BDC we discuss the observed and modelled high latitude age-of-air gradients. Figure 9 shows the seasonal progression of age-of-air differences between 80° S and 70° S as a function of height for the observations (left) and experiment pair A (right). In observations, air at 70° S is older than the air at 80° S below 16 km during spring. This is presumably related to horizontal mixing processes below the vortex linked to meridional drift in the lowermost stratospheric branch of the BDC and the different descent rates between the vortex core and a vortex region more towards the edge.

Above 16 km, air more inside the vortex is older compared to air more outside the vortex. The resulting strong gradient during late winter/early spring is artificially enhanced in the observations (Fig. 9, left) due to very old mean age in the vortex, caused by mesospheric SF<sub>6</sub> loss that cannot fully be accounted for in the derivation of the mean age. This structure is not seen in the model (Fig. 9, right), and the gradient reversal at lower altitudes is less pronounced. The signature of older air in higher latitudes becomes more prominent at lower altitudes during summer in both observations and model. The seasonality of the age gradient is a robust feature of the atmospheric circulation; neither in the observed case study or climatology (left, isolines and shadings)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nor in the climatological means of both experiments in pair A (right, shadings and iso-lines) is the overall seasonality very different. Nevertheless small changes can occur on top of the overall mean seasonality and are observed and modelled. We will discuss next these small changes, merging the ozone, N<sub>2</sub>O and circulation differences into a consistent framework, diagnosing a change in the seasonality of the BDC.

## 4.2 Case study for 2006/2007

Isolines in Fig. 10 (left) show the climatological mean of N<sub>2</sub>O from MIPAS observations for the period 2005–2012 using all months where data is available. The structural agreement between the observed and modelled climatological N<sub>2</sub>O is high (see Fig. 7), indicating that the model captures the seasonal changes in high latitude circulation well, as is also indicated by the age-of-air gradient shown in Fig. 9. Screening all annual N<sub>2</sub>O anomalies, we find a particular year (2006/2007; shadings in Fig. 10, left) that shows clear similarities with the transport change patterns identified in our idealised model studies (Fig. 7). In addition, the ozone change observed for January 2007 (shaded in Fig. 10, right) shows a similar pattern to the modelled structure (c.f. the SH polar dipole in Fig. 5).

This good correspondence with observations provides us with confidence that the modelled BDC change is relevant for the real atmosphere; the change modelled manifests itself in the observed interannual variability, even though it does not imply a causal relationship between ozone and N<sub>2</sub>O anomalies. It does suggest that halogen changes in the atmosphere can cause a shift in the circulation, towards a state of the atmosphere more similar to the SH condition of 2006/2007.

Note that we are not, of course, concluding that the change seen in the observations originates from the same initial disturbance we imprint on the model (e.g. the ozone difference in January 2006 was different). Note further that we compare modelled differences for climate equilibrium states, e.g. Fig. 7, A and B, with a particular event in the real atmosphere. However, the similarity between the modelled and observed anomalies seems to indicate that our model results can be realised in the “real world”.

### Consistent circulation differences in the SH

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As pointed out by Salby et al. (2012), both the change in stratospheric ozone and circulation in a particular month may have the same cause, and we just observe/model a consistent response of the system.

## 5 Summary and conclusions

We report results from two pairs of chemistry-climate model simulations using the same climate model but different chemistries. In each pair of experiments an ozone change was triggered by a simple change in the chemistry. One pair of model experiments looked at the impact of PSCs and the other at the impact of short lived halogenated species on composition and circulation. In both pairs of experiments, although the run with higher active chlorine has less ozone globally, there are regions of the atmosphere, and particular periods, where ozone is increased locally. The response to the perturbations is not purely chemical but is an example of a chemical-dynamical-radiative feedback (or chemistry-climate feedback). In tropical and northern latitudes, the model responses differ between pairs A and B, although the temperature and ozone changes are consistent within each pair and reduced column ozone is consistently associated with an increased tropopause height. In the SH, the two pairs of experiments have a consistent response. In both pairs of integrations the Antarctic polar ozone destruction is increased due to the higher availability of reactive chlorine (and bromine) in one of the integrations. Even though the magnitude of the chemical change differs between experiments, a common response pattern in circulation could be detected. We now focus our discussion on this region.

### 5.1 Schematic summary of model results

A schematic of the robust results emerging from experiment pairs A and B is shown in Fig. 11. The x-axis represents the seasonal progression of the year, the y-axis indicates latitude with high southern latitudes being to the front, and the z-axis indicates altitude

regions with around 15 km and 25 km highlighted. The +, – and o symbols indicate positive, negative and neutral anomalies between runs within a pair of integrations (pair A or B).  $\Delta\text{TO}_3$  is the difference in total ozone, and  $\Delta\text{N}_2\text{O}$  is the difference in nitrous oxide mixing ratio. The arrows indicate the general seasonal cycle of the BDC found in each run, colour coded as in Fig. 1. Green arrows show the mean meridional drift of the lower stratospheric BDC branch, orange arrows indicate the mean seasonal evolution of the middle stratospheric BDC branch. Red arrows indicate descent due to the polar convergence of the uppermost branch of the BDC.

Within each pair of experiments, the fundamental, consistent difference is that in the run with the increased amount of active halogen, the SH spring ozone depletion is more severe (negative  $\Delta\text{TO}_3$ ). This negative anomaly carries through to summer and maximises around December, because of a delayed vortex breakdown in integrations with larger amounts of ozone depleted in spring. In addition the lower stratospheric branch of the BDC is slightly weakened (negative  $\Delta\text{N}_2\text{O}$  around 15 km), thus inhibiting an efficient replenishment of ozone from the tropical lower stratospheric ozone production region. Consequently the negative total ozone anomaly is very long lived and persists into autumn.

In autumn the middle stratospheric branch of the BDC strengthens in the runs with higher ozone depletion (positive  $\Delta\text{N}_2\text{O}$  around 25 km). This is followed by a positive  $\text{N}_2\text{O}$  anomaly lower down in winter, which occurs because the positive anomaly from the previous season is advected downwards with the onset of descent linked to the uppermost BDC branch and is maintained by a strengthening of the lower stratospheric BDC branch (also evidenced by increased mid-latitude meridional heat flux at 150 hPa from June to August; not shown), which also increases ozone transport from low to high latitudes as well.

The positive  $\text{N}_2\text{O}$  anomaly is maintained throughout winter by the strengthened lower stratospheric branch and a decrease in descent rates. Consistently, in the following spring season negative  $\text{N}_2\text{O}$  anomalies around 25 km indicate a weakening (related to a later onset) of the middle stratospheric branch of the BDC (the September meridional

**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



heat flux anomaly in middle latitudes is negative; not shown) and slightly increased descent from above. This starts the seasonal cycle of changes again: in conjunction with the increased ozone depletion in spring this leads to a delayed break-up of the polar vortex during December.

As a caveat it should be mentioned that the Antarctic vortex in all model integrations presented here is slightly longer lived than observations would suggest. In many reanalyses data sets (see e.g. Fig. 4.2 of the CCMVal-2 Report, <http://www.sparc-climate.org/publications/sparc-reports/sparc-report-no5/>) the transition to easterlies at 60° S and 10 hPa occurs in late November. The previous model version (labelled UMUKCA-UCAM in the CCMVal-2 Report) captured this timing very well. In the integrations here the earliest break-up is modelled without halogen activation on PSCs using the CheS chemistry and occurs in early December. With halogen activation on PSCs the break-up is delayed by ~ 20 days (at 10 hPa, experiment pair A). The more comprehensive CheST+ chemistry (which always includes halogen activation by PSCs) has its transition in between the two dates for experiment pair A. And including additional halogen containing compounds increases the ozone deficit and delays the break-up by ~ 18 days (at 10 hPa, experiment pair B).

## 5.2 Implications of the results

Much of what we describe does not allow inference of causality. It merely points to a consistent change of the seasonal BDC development in both model integrations with enhanced ozone depletion in Antarctic spring. The delay of the vortex break-up in response to enhanced polar ozone depletion has been described before, and is one of the important drivers of the change during other seasons. Certainly other aspects are also changing, e.g. planetary wave propagation in conjunction with the delayed vortex break-up (at 60° S meridional heat transport anomalies are negative during September to November; not shown), but in addition because generally ozone and thermal gradients have changed.

---

**Consistent  
circulation  
differences in the SH**P. Braesicke et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 What we have presented here is a robust chemistry-climate feedback; one of many that require further investigation. Many IPCC models do not have interactive ozone and so do not allow readjustments of stratospheric ozone and circulation (see the discussion in Son et al., 2010). If we would like to predict future climate change reliably and with narrowing uncertainties, our results suggest that we need to understand more about fairly small composition changes that can contribute to systematic changes in the hemispheric circulation. As always there is a caveat regarding the general applicability of our results as all models exhibit some bias with respect to observations. Nevertheless we are able to identify similar consistent changes in MIPAS data of the recent past as part of the interannual variability of the stratospheric climate system. So we believe that a future change in composition could lead to a systematic shift in the mean seasonal cycle of the BDC in the long-term, beyond the general strengthening postulated for increasing long-lived greenhouse gases.

15 Feedbacks between composition and circulation are important in shaping our future climate. The above is just one particular example in which an initial composition change (increased availability of halogens) leads to a readjustment of the equilibrium state of the coupled chemistry-climate system. The overall ozone changes are determined by the initial composition change and the readjusted circulation (because the ozone change has changed heating rates and in conjunction winds). Understanding these processes in models and observations will lead to improvements in our ability to model the past climate and to project future changes more reliably.

25 *Acknowledgement.* We thank NCAS-CMS for modelling support. Model integrations have been performed using the UK National Supercomputing Service HECToR. PB and JK thank NERC for support through grant NE/H024778/1. JAP and XY thank ERC for support through the ACCI grant and the EU for support through SHIVA.

## References

- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle Atmosphere Dynamics, Academic Press, Orlando, FL, 1987.
- 5 Birner, T. and Bönisch, H.: Residual circulation trajectories and transit times into the extratropical lowermost stratosphere, *Atmos. Chem. Phys.*, 11, 817–827, doi:10.5194/acp-11-817-2011, 2011.
- Bodeker, G. E., Shiona, H., and Eskes, H.: Indicators of Antarctic ozone depletion, *Atmos. Chem. Phys.*, 5, 2603–2615, doi:10.5194/acp-5-2603-2005, 2005.
- 10 Braesicke, P. and Pyle, J. A.: Changing ozone and changing circulation in northern mid-latitudes: possible feedbacks?, *Geophys. Res. Lett.*, 30, 1059, doi:10.1029/2002GL015973, 2003.
- Braesicke, P., Hurwitz, M. M., and Pyle, J. A.: The stratospheric response to changes in ozone and carbon dioxide as modelled with a GCM including parameterised ozone chemistry, *Meteorol. Z.*, 15, 343–354, doi:10.1127/0941-2948/2006/0124, 2006.
- 15 Brewer, A. W.: Evidence for a world circulation provided by the measurements of helium and water vapour distribution in the stratosphere, *Quarterly J. R. Meteorol. Soc.*, 75, 351–363, 1949.
- Butchart, N., Scaife, A. A., Bourqui, M., de Grandpre, J., Hare, S. H. E., Kettleborough, J., Langematz, U., Manzini, E., Sassi, F., Shibata, K., Shindell, D., and Sigmond, M.: Simulations of anthropogenic change in the strength of the Brewer-Dobson, circulation, *Clim. Dynam.*, 27, 727–741, doi:10.1007/s00382-006-0162-4, 2006.
- 20 CCMVal-2 Report: SPARC Report on the Evaluation of Chemistry-Climate Models, WCRP-132, WMO/TD No. 1526, SPARC Report No. 5, [http://www.sparc-climate.org/fileadmin/customer/6\\_Publications/SPARC\\_reports.PDF/5\\_CCMval\\_SPARCreportNo5\\_Jun2010\\_preamble.pdf](http://www.sparc-climate.org/fileadmin/customer/6_Publications/SPARC_reports.PDF/5_CCMval_SPARCreportNo5_Jun2010_preamble.pdf), 2010.
- 25 Dobson, G. M. B.: Origin and distribution of the polyatomic molecules in the atmosphere, *Proc. R. Soc. London*, A236, 187–193, 1956.
- Engel, A., Moebius, T., Boenisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nat. Geosci.*, 2, 28–31, doi:10.1038/ngeo388, 2009.

## Consistent circulation differences in the SH

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Garcia, R. R. and Randel, W. J.: Acceleration of the Brewer-Dobson circulation due to increases in greenhouse gases, *J. Atmos. Sci.*, 65, 2731–2739, doi:10.1175/2008JAS2712.1, 2008.
- Gillett, N. P. and Thompson, D. W. J.: Simulation of recent Southern Hemisphere climate change, *Science*, 302, 273–275, doi:10.1126/science.1087440, 2003.
- 5 Grewe, V.: The origin of ozone, *Atmos. Chem. Phys.*, 6, 1495–1511, doi:10.5194/acp-6-1495-2006, 2006.
- Hewitt, H. T., Copsey, D., Culverwell, I. D., Harris, C. M., Hill, R. S. R., Keen, A. B., McLaren, A. J., and Hunke, E. C.: Design and implementation of the infrastructure of HadGEM3: the next-generation Met Office climate modelling system, *Geosci. Model Dev.*, 4, 223–253, doi:10.5194/gmd-4-223-2011, 2011.
- 10 Morgenstern, O., Braesicke, P., Hurwitz, M. M., O'Connor, F. M., Bushell, A. C., Johnson, C. E., and Pyle, J. A.: The World Avoided by the Montreal Protocol, *Geophys. Res. Lett.*, 35, L16811, doi:10.1029/2008GL034590, 2008.
- Morgenstern, O., Braesicke, P., O'Connor, F. M., Bushell, A. C., Johnson, C. E., Osprey, S. M., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model – Part 1: The stratosphere, *Geosci. Model Dev.*, 2, 43–57, doi:10.5194/gmd-2-43-2009, 2009.
- 15 O'Connor, F. M., Johnson, C. E., Morgenstern, O., and Collins, W. J.: Interactions between tropospheric chemistry and climate model temperature and humidity biases, *Geophys. Res. Lett.*, 36, L16801, doi:10.1029/2009GL039152, 2009.
- 20 Plumb, R. A.: Stratospheric transport, *J. Meteorol. Soc. Jpn.*, 80, 793–809, doi:10.2151/jmsj.80.793, 2002.
- Ray, E. A., Moore, F. L., Elkins, J. W., Hurst, D. F., Romashkin, P. A., Dutton, G. S., and Fahy, D. W.: Descent and mixing in the 1999–2000 northern polar vortex inferred from in situ tracer measurements, *J. Geophys. Res.-Atmos.*, 107, 8285, doi:10.1029/2001JD000961, 2002.
- 25 Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Boenisch, H., Morgenstern, O., Smale, D., Rozanov, Eugene, Hegglin, M., Pitari, G., Mancini, E., Braesicke, P., Butchart, N., Hardiman, S., Li, F., Shibata, K., and Plummer, D. A.: Evidence for changes in stratospheric transport and mixing over the past three decades based on multiple data sets and tropical leaky pipe analysis, *J. Geophys. Res.-Atmos.*, 115, D21304, doi:10.1029/2010JD014206, 2010.
- 30 Salby, M. L., Titova, E. A., and Deschamps, L.: Changes of the Antarctic ozone hole: controlling mechanisms, seasonal predictability, and evolution, *J. Geophys. Res.-Atmos.*, 117, D10111, doi:10.1029/2011JD016285, 2012.

## Consistent circulation differences in the SH

P. Braesicke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Son, S.-W., Gerber, E. P., Perlwitz, J., Polvani, L. M., Gillett, N. P., Seo, K.-H., Eyring, V., Shepherd, T. G., Waugh, D., Akiyoshi, H., Austin, J., Baumgaertner, A., Bekki, S., Braesicke, P., Bruehl, C., Butchart, N., Chipperfield, M. P., Cugnet, D., Dameris, M., Dhomse, S., Frith, S., Garny, H., Garcia, R., Hardiman, S. C., Joeckel, P., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Nakamura, T., Morgenstern, O., Pitari, G., Plummer, D. A., Pyle, J., Rozanov, E., Scinocca, J. F., Shibata, K., Smale, D., Teyssedre, H., Tian, W., and Yamashita, Y.: Impact of stratospheric ozone on Southern Hemisphere circulation change: a multimodel assessment, *J. Geophys. Res.-Atmos.*, 115, D00M07, doi:10.1029/2010JD014271, 2010.

Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12, 3311–3331, doi:10.5194/acp-12-3311-2012, 2012.


Thompson, D. W. J. and Solomon, S.: Recent stratospheric climate trends as evidenced in radiosonde data: global structure and tropospheric linkages, *J. Climate*, 18, 4785–4795, doi:10.1175/JCLI3585.1, 2005.

Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nat. Geosci.*, 4, 741–749, doi:10.1038/NGEO1296, 2011.

von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H<sub>2</sub>O, O<sub>3</sub>, HNO<sub>3</sub>, CH<sub>4</sub>, N<sub>2</sub>O, ClONO<sub>2</sub> and ClO from MIPAS reduced resolution nominal mode limb emission measurements, *Atmos. Meas. Tech.*, 2, 159–175, doi:10.5194/amt-2-159-2009, 2009.

Waugh, D. W. and Hall, T. M.: Age of stratospheric air: theory, observations, and models, *Rev. Geophys.*, 40, 1010, doi:10.1029/2000RG000101, 2002.

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project-Report No. 52, 516 pp., Geneva, Switzerland, available at: <http://www.esrl.noaa.gov/csd/assessments/ozone/2010/citations.html>, 2011.



**Fig. 1.** Conceptual structure of the Brewer-Dobson Circulation (BDC) following Plumb (2002). The three main circulation branches are highlighted in different colours, which are used in summary Fig. 11 to distinguish the height regions. Blue shadings indicate the general altitude gradient of  $N_2O$  with near constant  $N_2O$  in the troposphere (Trop.). The complex meridional  $N_2O$  gradient structure in the stratosphere is shaped by the different BDC branches and is not detailed here.

**Consistent circulation differences in the SH**

P. Braesicke et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Consistent circulation differences in the SH

P. Braesicke et al.



**Fig. 3.** Long-term, monthly and zonal mean total ozone differences (Dobson Units, DU) for experiment pairs **(A)** (left) and **(B)** (right). Negative differences are shown in cold colours. See Fig. 2 for definition of the differences.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Consistent circulation differences in the SH

P. Braesicke et al.



**Fig. 4.** Long-term, monthly and zonal mean WMO (lapse rate) tropopause height differences (m) for experiment pairs **(A)** (left) and **(B)** (right). Positive differences are shown in warm colours. See Fig. 2 for definition of the differences.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.



**Fig. 5.** December zonal and monthly mean ozone differences in percent for experiment pair **(A)** (left) and experiment pair **(B)** (right). Negative differences are shown in cold colours. The black dashed line indicates the position of the tropopause in the base cases.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.



**Fig. 6.** December zonal and monthly mean temperature differences (K) for experiment pair **(A)** (left) and experiment pair **(B)** (right). Negative differences are shown in cold colours. The black dashed line indicates the position of the tropopause in the base cases.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 7.** Time-height cross sections of N<sub>2</sub>O and N<sub>2</sub>O differences. Isolines: absolute values of long-term monthly mean (LTMM) N<sub>2</sub>O (parts per million by volume, ppbv) averaged over all latitudes south of 70° S for runs subtracted in pair (B) (left) and pair (B) (right). Shadings: absolute differences in N<sub>2</sub>O (ppbv) for experiment pair (A) (left) and experiment pair (B) (right). Positive differences are shown in warm colours.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**Consistent  
circulation  
differences in the SH**

P. Braesicke et al.



**Fig. 8.** December zonal and monthly mean age-of-air differences in percent for experiment pair **(A)** (left) and experiment pair **(B)** (right). Positive differences are shown in warm colours. The black dashed line indicates the position of the tropopause in the base cases.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Consistent circulation differences in the SH

P. Braesicke et al.



**Fig. 10.** Time-pressure cross section of climatological MIPAS  $N_2O$  (ppbv, isolines, left) and  $N_2O$  differences between 2006/2007 and the climatology (ppbv, shaded, left), both averaged over all latitudes south of  $70^\circ S$ . Latitude-pressure cross section of climatological January MIPAS ozone (ppmv, isolines, right) and ozone differences between January 2007 and the climatology (ppbv, shaded, right). Positive differences are shown in warm colours.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Consistent circulation differences in the SH**

P. Braesicke et al.



**Fig. 11.** Sketch summarising the time evolution of the changes modelled (LS: lower stratospheric, MS: middle stratospheric). Arrows indicate a mean state. Deviations are indicated with + (positive), - (negative) and o (neutral).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

