



## Abstract

Organic aerosol particles play a key role in climate by serving as nuclei for clouds and precipitation. Their sources and composition are highly variable, and their phase state ranges from liquid to solid under atmospheric conditions, affecting the pathway of activation to cloud droplets and ice crystals. Due to slow diffusion of water in the particle phase, organic particles may deviate in phase and morphology from their thermodynamic equilibrium state, hampering the prediction of their influence on cloud formation. We overcome this problem by combining a novel semi-empirical method for estimation of water diffusivity with a kinetic flux model that explicitly treats water diffusion. We estimate timescales for particle deliquescence as well as various ice nucleation pathways for a wide variety of organic substances, including secondary organic aerosol (SOA) from the oxidation of isoprene,  $\alpha$ -pinene, naphthalene, and dodecane. The simulations show that in typical atmospheric updrafts, glassy states and solid/liquid core-shell morphologies can persist for long enough that heterogeneous ice nucleation in the deposition and immersion mode can dominate over homogeneous ice nucleation. Such competition depends strongly on ambient temperature and relative humidity as well as humidification rates and particle sizes. Due to relatively high glass transition temperature and low hygroscopicity, naphthalene SOA particles have a higher potential to act as heterogeneous ice nuclei than the other investigated substances. Our findings demonstrate that kinetic limitations of water diffusion into organic aerosol particles strongly affect their ice nucleation pathways and require advanced formalisms for the description of ice cloud formation in atmospheric models.

## 1 Introduction

Atmospheric aerosol particles influence climate through affecting the earth's radiation budget directly by scattering and absorbing light, and indirectly by acting as nuclei for cloud droplets and ice crystals (Yu et al., 2006; Andreae and Rosenfeld, 2008; IPCC,

ACPD

14, 16451–16492, 2014

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2013). Ice nucleation is an important pathway for high-altitude cirrus cloud formation and it occurs either homogeneously in liquid aerosol particles or heterogeneously in the presence of active ice nuclei (IN), which are solid particles that facilitate nucleation. Homogeneous ice nucleation generally requires high supersaturations in aqueous aerosol droplets, occurring at ice saturation ratios of  $S_{\text{ice}} \geq 1.4$  (Koop et al., 2000). Only a small fraction of atmospheric aerosol particles act as IN below this homogeneous ice nucleation threshold (DeMott et al., 2003; Cziczo et al., 2013). Heterogeneous ice nucleation can occur via several pathways such as deposition nucleation, i.e. deposition of gaseous water molecules to form crystalline ice on a solid IN, or immersion freezing, which describes nucleation induced by IN immersed in supercooled aqueous droplets (Pruppacher and Klett, 1997; Hoose and Möhler, 2012).

Organic aerosol particles are ubiquitous and abundant in the atmosphere, but traditionally are not referred to as effective IN when compared to dust or biological particles (see Hoose and Möhler (2012) and references therein). More recently, however, several laboratory studies have shown that glassy organic particles can act as IN at low-temperature cirrus conditions in the deposition mode (Murray et al., 2010; Wagner et al., 2012; Wilson et al., 2012) or at slightly elevated temperatures in the immersion mode (Wang et al., 2012; Baustian et al., 2013; Schill et al., 2014), in agreement with inferences from field data (Froyd et al., 2010). This IN ability has been observed for a number of different types of particles composed of pure organic substances such as simple sugars and acids (Murray et al., 2010; Wagner et al., 2012; Wilson et al., 2012; Baustian et al., 2013) and biomass burning marker compounds (Wagner et al., 2012; Wilson et al., 2012), for (phase-separated) organic-inorganic mixtures (Wagner et al., 2012; Wilson et al., 2012; Baustian et al., 2013; Schill and Tolbert, 2013), as well as for SOA particles derived from aromatic VOCs (Wang et al., 2012) or emerging from aqueous phase reactions (Schill et al., 2014). It has also been proposed recently that formation of highly porous structures upon atmospheric freeze-drying could enhance the IN ability of organic aerosol particles (Adler et al., 2013).

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



These observations suggest a connection between particle phase state and the resulting predominant ice nucleation pathway (Murray et al., 2010). Organic aerosol particles can adopt liquid, semisolid or solid states, or may even exhibit mixed phases, depending on composition and ambient conditions (Mikhailov et al., 2009; Koop et al., 2011; Vaden et al., 2011; Kuwata and Martin, 2012; Perraud et al., 2012; Song et al., 2012; You et al., 2012; Renbaum-Wolff et al., 2013; Kidd et al., 2014). SOA particles are expected to be liquid at high temperature and high humidity, but are very likely to exhibit a highly viscous semisolid or even glassy state at low temperature and low humidity (Virtanen et al., 2010; Saukko et al., 2012; Renbaum-Wolff et al., 2013; Shiraiwa et al., 2013). Glassy states are characterized by viscosities greater than  $10^{12}$  Pa s, corresponding to diffusion time scales within these particles that can exceed days or even years (Shiraiwa et al., 2011; Koop et al., 2011; Zhou et al., 2013). Water uptake into glassy aerosols has been shown to occur slowly and to proceed gradually with increasing relative humidity (Mikhailov et al., 2009; Tong et al., 2011; Zobrist et al., 2011; Bones et al., 2012; Price et al., 2014).

Hence, several competing processes can occur in glassy organic aerosol particles during updraft of an air parcel: heterogeneous ice nucleation in the deposition mode onto the glassy solid aerosol surface; diffusion of water into the particle, inducing a gradual phase transition towards the liquid state; and immersion freezing during the transition between both states. In order to determine those atmospheric conditions at which one of these processes dominates, we employ a numerical aerosol diffusion model based on the kinetic multi-layer model for gas-particle interactions in aerosols and clouds (KM-GAP), which explicitly treats mass transport of water molecules in the gas and particle phases (Shiraiwa et al., 2012). Due to experimental constraints associated with very long observation times, parameterizations for water diffusivity in glassy organic material are sparse and hence are only known for a few model compounds. Therefore, water diffusivity in SOA materials from various biogenic and anthropogenic precursors are deduced from water diffusivity parameterizations of model

compounds using a semi-empirical physico-chemical model of water diffusion in glass-forming aqueous organics.

## 2 Modelling approach

### 2.1 Numerical diffusion model

5 The numerical diffusion model employed in this study is based on the kinetic multi-layer model for gas-particle interactions in aerosols and clouds, KM-GAP (Shiraiwa et al., 2012). KM-GAP consists of multiple model compartments and layers, respectively: gas phase, near-surface gas phase, sorption layer, surface layer, near-surface bulk, and a number of  $n$  bulk layers (cf. Fig. S1 in the supplementary material). The follow-  
10 ing processes are considered in KM-GAP: gas phase diffusion, gas-surface transport, surface-bulk transport, and bulk diffusion. The bulk layers can either grow or shrink in response to mass transport. The initial bulk layer sizes are chosen small enough to ensure numerical convergence (usually 100–750 layers), but are not allowed to fall below the molecular length scale ( $\sim 0.3$  nm).

15 The model was complemented by modules predicting homogeneous ice nucleation as a function of water activity according to Koop et al. (2000), heterogeneous ice nucleation at a pre-defined ice supersaturation level, and it considers Kelvin effects. Moreover, a few further conceptual changes have been introduced to the original KM-GAP, including a more explicit treatment of gas diffusion, composition-based bulk diffusion and a mechanism of surface-to-bulk transport facilitated by surface-adsorbed water, as detailed in the following sections. Parameterizations of composition-dependent density,  
20 water activity and bulk diffusivity for the sucrose/water system have been adopted from Zobrist et al. (2011). A detailed description of the gas diffusion scheme and a list of all employed parameterizations are provided as supplementary material.

25 In this study, the model is used to simulate an atmospheric updraft situation by following a preselected trajectory in temperature, relative humidity and pressure. It tracks

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the chemical composition of an amorphous aerosol particle as a function of time and depth below the particle surface in discretized layers, providing concentration profiles of water and organics at any given time. The equilibrium composition is calculated through a water activity parameterization that translates ambient relative humidity into equilibrium mass fractions of the bulk constituents. Mass fluxes from the far-surface into the near-surface gas phase, onto the particle surface, into as well as between bulk layers are coupled in flux-based differential equations, which are solved with an ordinary differential equation solver using Matlab software (ode23tb).

### 2.1.1 Ice nucleation modules

Besides water diffusion, the model is able to simulate ice nucleation and growth. However, the initial numerical solution of the differential equations treats merely water uptake into the particle. The model registers an ice nucleation event when all necessary conditions in ambient relative humidity and water activity are satisfied. From this point onwards, the model simulates ice crystal growth by deposition of water molecules from the gas phase.

For homogeneous ice nucleation, a stochastic approach based on classical nucleation theory has been chosen. An ice nucleation event is triggered when the probability of the particle being liquid ( $P_{liq}$ ) falls below 50%.  $P_{liq}$  is the product of the individual probabilities in all  $n$  layers, using the homogeneous nucleation rate coefficient for each layer  $J_{hom,n}$  as parameterized by Koop et al. (2000). The nucleation rate then translates into  $P_{liq}$  by multiplication with layer volume  $V_n$  and (numerical) integration time step  $dt$ :

$$P_{liq,tot}(t) = \int_{t_0}^t \prod_{n=1}^L (1 - J_{hom,n}(t) \cdot V_n(t)) dt \quad (1)$$

Heterogeneous nucleation is assumed to occur once a certain freezing threshold is exceeded. In this work, we distinguish between heterogeneous ice nucleation thresholds

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for sucrose and SOA, which have been shown to occur at different ice supersaturations, as summarized by Schill et al. (2014). For sucrose, we apply a linear fit to nucleation data from Baustian et al. (2013), whereas for SOA we fit the nucleation data of naphthalene SOA from Wang et al. (2012) and those of aqSOA from Schill et al. (2014). The fit results are shown in Fig. A1 in Appendix A.

To distinguish between deposition and immersion freezing, additional criteria are employed. For deposition nucleation, the necessary condition is solidness of the outermost layer of the particle, requiring the water activity to be below the quasi-equilibrium glass transition point. In case of immersion mode nucleation, a 1 nm thick region in the near-surface bulk is required to be entirely liquefied before nucleation can occur in the immersion mode. For this purpose, a 2 nm thick region below the particle surface is finely resolved by multiple bulk layers (cf. Fig. S1).

### 2.1.2 Bulk diffusion and bulk layer mixing

Bulk diffusion of water is treated as kinetic flux  $J_{bk,bk\pm 1}$  from one bulk layer ( $bk$ ) to the next ( $bk \pm 1$ ). Because layer thickness is not allowed to fall below molecular resolution, concentrations in adjacent layers can differ significantly. As in Zobrist et al. (2011), this heterogeneity is accounted for with a virtual mixing scheme for the determination of bulk diffusivities between layers. In this scheme, the composition of a mixture of two subsequent bulk layers is determined and the bulk diffusion coefficient calculated according to the effective composition along the diffusion path. Scenarios with very low diffusivities and hence steep concentration gradients thus lead to situations in which a liquefied layer (high bulk diffusivity of water,  $D_{H_2O}$ ) “softens” the subsequent glassy layer (low  $D_{H_2O}$ ), facilitating further diffusion. Such a process can be seen analogously to a dissolution process, in which the glassy matrix dissolves into nearby water-rich regions.

Diffusion of the organic matrix has been neglected for this study, because the organic molecules investigated here can be expected to diffuse much slower than water molecules. Also, in the glassy state, the organic molecules diffuse on a much longer

timescale compared to the experimental time scale of minutes to hours (cf. Shiraiwa et al., 2011; Koop et al., 2011).

### 2.1.3 Surface monolayers and surface softening

The original KM-GAP uses a double monolayer approach to describe the particle surface, comprising a sorption layer and a quasi-static surface layer. In this study the quasi-static surface layer was replaced by a near-surface volume layer similar to that used in Shiraiwa et al. (2013), which is more suitable for systems with low diffusivity.

Surface-adsorbed water can lead to softening of the solid surface (Koop et al., 2011), thereby facilitating exchange between surface and first near-surface bulk layer. In the model, this is accounted for by introducing a surface softening scheme that estimates the surface-to-bulk transport rate by mixing a hypothetical water monolayer with a hypothetical bulk monolayer containing water and bulk material. Using the momentary molar fractions of water ( $x_{b1,H_2O}$ ) and organics ( $x_{b1,org}$ ) of the near-surface bulk layer, the effective surface coverages of water ( $\theta_{ss,H_2O}$ ) and organics ( $\theta_{ss,org}$ ) at the surface bulk layer can be described as:

$$\theta_{ss,i} = \frac{x_{b1,i} \cdot \sigma_i}{x_{b1,org} \cdot \sigma_{org} + x_{b1,H_2O} \cdot \sigma_{H_2O}} \quad (2)$$

where  $\sigma_i$  is the molecular cross section of species  $i$  (i.e. water (H<sub>2</sub>O) or organics (org)), respectively. The weight fraction of organics in the “softened” surface is then given by:

$$w_{ss,org,mix} = \frac{\frac{\theta_{ss,org}}{\sigma_{org}} \cdot M_{org}}{\frac{\theta_{ss,org}}{\sigma_{org}} \cdot M_{org} + \left(\frac{\theta_{ss,H_2O} + 1}{\sigma_{H_2O}}\right) \cdot M_{H_2O}} \quad (3)$$

where  $M_{org}$  and  $M_{H_2O}$  are the molar mass of organics and water. This process facilitates the initial water uptake into a glassy particle and leads (in most cases) to a sub-surface layer that is in equilibrium with the surrounding gas phase. In the temperature range

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



relevant for immersion freezing, liquefaction of the surface was always obtained at the quasi-equilibrium glass transition point due to the surface softening mechanism. At lower temperatures however (deposition regime), the particle surface was not always in quasi-equilibrium with ambient humidity.

## 2.2 Estimation of water diffusivity in SOA

For model systems other than sucrose/water, no direct parameterization of water diffusivity in the full atmospherically relevant temperature and composition range is available to date. For compounds chemically similar to sucrose (i.e. organic polyols and acids), we present a scheme that enables estimation of bulk diffusivity data from glass transition and hygroscopicity data. Bulk diffusivity of water is parameterized using a Vogel-Fulcher-Tammann (VFT) approach (Vogel, 1921; Fulcher, 1925; Tammann and Hesse, 1926). The estimation scheme utilizes the structure of the VFT equation, Eq. (S9), and the physical interpretation of its parameters. The method can be described by the following set of assumptions:

1. Two similar organic substances act similar in the way they approach the glass transition and thus have a similar fragility:  $B_{\text{org},1} \approx B_{\text{org},2}$ .
2. The same two substances have a similar diffusion coefficient in the high temperature limit:  $A_{\text{org},1} \approx A_{\text{org},2}$ .
3. A difference in glass transition temperatures between the two substances indicates a difference in Vogel temperatures of same direction and (relative) magnitude

$$\frac{T_{0,\text{org},1}}{T_{0,\text{org},2}} \approx \frac{T_{\text{g},\text{org},1}}{T_{\text{g},\text{org},2}} \quad (4)$$

Thus, diffusivities within an organic substance can be estimated by knowledge of its glass transition curve relative to a known standard with similar chemical functionality.

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



“quasi-equilibrium glass transition“, we denote the conditions under which a binary organic-water system would undergo amorphous deliquescence when humidification occurs sufficiently slow so that equilibrium between ambient RH and water activity is always maintained. Humidification may be fast enough to cause a difference in phase state from equilibrium: water activity, colour-coded from dark blue (low water activity) to light blue (high water activity), trails behind ambient RH due to kinetic limitations in water diffusivity (Koop et al., 2011). Note that when using a constant  $D_{\text{H}_2\text{O}}$ , diffusion gradients appear less pronounced (cf. Fig. S2 and Movie S2). Hence, self-accelerating water diffusion leads to a sharpening of the diffusion gradient that can be close to the molecular length scale (Zobrist et al., 2011).

Several morphological stages can be distinguished during the humidification process. Starting from a homogeneous, glassy particle, an increase in RH first leads to liquefaction of a thin outer layer and emergence of a core-shell morphology. This liquid outer layer grows in equilibrium with ambient relative humidity and also extends towards the particle centre by diffusion of water into the glassy organic matrix, leading to shrinkage of the residual glassy core until the particle is fully deliquesced. Thus, during the continuous amorphous deliquescence process two characteristic instants can be distinguished, each occurring at a different humidity: we define the *Partial Deliquescence Relative Humidity* (PDRH) as the point where a thin aqueous outer shell of the particle is homogeneously mixed and the shell’s water activity is larger than that of the quasi-equilibrium glass transition. In this study we set the thickness of this surface shell to 1 nm, corresponding to about 5 monolayers of water. We define the *Full Deliquescence Relative Humidity* (FDRH) as the point where the entire particle is homogeneously mixed and its water activity corresponds to that of a liquid (i.e. it is larger than that of the quasi-equilibrium glass transition). Note that in the case of a sufficiently slow updraft, both PDRH and FDRH would occur at  $\text{RH}_g$ . In fact, the KM-GAP simulations suggest that, with updraft velocities typical for atmospheric conditions (e.g.  $0.01\text{--}10\text{ m s}^{-1}$ ), PDRH often coincides with  $\text{RH}_g$ . In contrast, FDRH often extends far into the liquid region of the phase diagram, indicating the importance of kinetic limita-

tions and implying that particles can contain glassy cores even at relative humidities above  $RH_g$  due to slow water diffusion.

### 3.2 Ice nucleation regimes

Next, we investigate by kinetic model simulations the competition between amorphous deliquescence and ice nucleation during an atmospheric updraft. For our initial calculations we use sucrose as a proxy for organic aerosols since detailed physico-chemical parameterizations for water diffusivity, the RH-dependent equilibrium composition as well as glass transition data are available (Zobrist et al., 2011). The heterogeneous ice nucleation onset ( $RH_{het}$ ) for sucrose was obtained from ice nucleation experiments by Baustian et al. (2013) and is shown as brown dashed lines in Fig. 2. Here we use the ice saturation ratio  $S_{ice}$  as an indicator of humidity because it scales with RH according to  $S_{ice} = \rho_{liq,0}(T)/\rho_{ice}(T) \cdot RH$ , but is also a more direct indicator of the supersaturation of ice.

Figure 2a shows results obtained with KM-GAP simulating the updraft of 100 nm sucrose particles for a wide range of temperatures. Each simulated trajectory started at ice saturation ( $S_{ice} = 1$ ), as is often the case for cloud chamber or environmental cell experiments (Murray et al., 2010; Wang et al., 2012). Temperature was decreased so that the resulting humidification rate was constant at  $1\% RH \text{ min}^{-1}$ , corresponding to an atmospheric updraft of about  $0.2 \text{ m s}^{-1}$ , typical of atmospheric gravity waves.

As expected FDRH of sucrose particles, indicated by the red solid line, occurs significantly above  $RH_g$  at all temperatures. The intersection of  $RH_{het}$  with  $RH_g$  defines the upper temperature limit for deposition nucleation. Below this temperature, a sucrose particle is a glassy solid when  $RH_{het}$  is reached, and hence deposition ice nucleation may occur. Above this temperature, the particle is partially deliquesced when approaching  $RH_{het}$  and the glassy core of the particle may act as an IN for immersion freezing. The upper limit of the immersion freezing regime is given by the intersection of  $RH_{het}$  with the FDRH line. Above this temperature, particles are already fully deliquesced once  $RH_{het}$  is reached. Hence, these particles do not nucleate ice hetero-

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



geneously and freeze only at the homogeneous ice nucleation limit (green dashed line; Koop et al., 2000). Finally at  $\sim 232$  K, the homogeneous ice nucleation limit coincides with water saturation (solid black line) and above this temperature the aerosol particles activate into cloud droplets consisting of supercooled water, thus representing the upper limit of the homogeneous ice nucleation regime.

The delay between the nominal quasi-equilibrium glass transition  $RH_g$  and the actual full deliquescence at FDRH is governed by the competition between humidification rate (synonymous to updraft velocity) and timescale for water diffusion within the particle bulk. FDRH will shift towards higher relative humidities when higher humidification rates are employed, as shown in Fig. 2b. For example, increasing the rate of humidification to  $10\% RH \text{ min}^{-1}$ , a value corresponding to an updraft velocity of about  $2 \text{ m s}^{-1}$  and commonly reached in convective updrafts, shifts the FDRH line upwards (solid dark blue line) and thus its intersection with the  $RH_{\text{het}}$  line towards higher temperatures. Accordingly, decreasing the updraft velocity to  $0.02 \text{ m s}^{-1}$ , a value found in large-scale, synoptic updrafts, leads to FDRH (solid light blue line) much closer to the quasi-equilibrium glass transition  $RH_g$ . Moreover, an increase in particle size delays the deliquescence process (indicated by the solid purple line), since it increases the timescale of diffusion. The range of the immersion freezing regime thus strongly depends on ambient conditions and is extended towards higher temperatures in fast updrafts and for large particles.

Laboratory ice nucleation measurements of sucrose particles ( $4 \mu\text{m}$  diameter,  $1\% RH \text{ min}^{-1}$ ) (Baustian et al., 2013) are used to validate our model calculations of ice nucleation regimes in Fig. 2c, for details see Appendix B. The modelled FDRH curve (solid blue line) correctly confines the region below which heterogeneous ice nucleation is observed in the experiments. Based on our calculations, the experimental data points below  $RH_g$  (full brown circles) can be assigned to the deposition nucleation regime, whereas points between  $RH_g$  and FDRH (open brown circles) can be assigned to immersion freezing. Additional analyses for validation have been performed for other types of organic particles (Figs. B1 and B2).

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

### 3.3 Biogenic and anthropogenic SOA

In order to apply our kinetic model to ice nucleation in secondary organic aerosol (SOA), estimates of  $D_{\text{H}_2\text{O}}$  in SOA material have been inferred. Four major SOA precursors were chosen to represent biogenic and anthropogenic origin, respectively:  $\alpha$ -pinene and isoprene, as well as naphthalene and dodecane. Each of these SOA is represented by a choice of marker compounds taken from the literature (cf. Table S1). Water diffusivities are estimated utilizing the scheme described in Sect. 2.2. The heterogeneous ice nucleation onset ( $\text{RH}_{\text{het}}$ , brown dashed line) for SOA was obtained from laboratory measurements by Wang et al. (2012) and Schill et al. (2014) as derived in Fig. A1. In all simulations, particles of 100 nm diameter were humidified at a rate of  $1\% \text{ RH min}^{-1}$ .

Figure 3a shows the simulation results of FDRH for all four precursor types. Naphthalene SOA is observed to be fully deliquesced latest due to the high estimated glass transition temperature and low hygroscopicity (cf. Table A1), followed by  $\alpha$ -pinene and isoprene. Dodecane SOA showed the earliest deliquescence, reflecting the low glass transition temperature of pure dodecane SOA of  $\sim 210 \text{ K}$ . By comparison of FDRH with measured  $\text{RH}_{\text{het}}$  on SOA, compound-specific upper temperature limits for heterogeneous ice nucleation on SOA particles can be determined (arrows on x-axis, values are given in Table S2). Uncertainty estimates for FDRH and  $\text{RH}_{\text{g}}$  of all four precursor classes are given in Fig. S4.

For the calculations in Fig. 3a, we chose an average oxidation state typically observed for SOA from the respective precursor. The atomic oxygen to carbon ratio (O/C) increases upon chemical ageing, thereby affecting hygroscopicity (Lambe et al., 2011) and glass transition temperature (Fig. A2). The resulting effects of chemical ageing on modelled FDRH are shown exemplarily for  $\alpha$ -pinene and dodecane SOA in Fig. 3b and c, respectively. For  $\alpha$ -pinene SOA (B), a higher O/C results in hardening of the organic material with ageing, leading to a FDRH increase, whereas for dodecane SOA (C)

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a higher O/C results in softening, thus leading to earlier deliquescence and a FDRH decrease.

The observed effects can be explained by the competition between a simultaneous increase of hygroscopicity with O/C and an increasing glass transition temperature of the pure organic matrix due to stronger molecular interactions in the highly oxidized organic material. A higher glass transition value enhances the rigidity of the pure organic matrix, whereas a higher hygroscopicity enhances the amount of water taken up by the aqueous organic mixture at a given humidity and thus its plasticizing effect.

Figure 4 illustrates this competition by displaying estimated characteristic timescales of water diffusion in 100 nm diameter SOA particles at 220 K as a function of hygroscopicity ( $\kappa_{\text{org}}$ ) and glass transition temperature of the pure organic matrix ( $T_{\text{g,org}}$ ). Dotted contour lines show characteristic mass transport times associated with the diffusion coefficient  $D_{\text{H}_2\text{O}}$  (Shiraiwa et al., 2011). Coloured oval shapes indicate estimated ranges of  $\kappa_{\text{org}}$  and  $T_{\text{g,org}}$  for the four SOA precursor classes, for three different oxidation states each (cf. Table A1). The arrows pointing from the lowest to the highest oxidation state reveal that both  $\kappa_{\text{org}}$  and  $T_{\text{g,org}}$  increase with O/C. The slope of these arrows when compared to the slope of the contour lines indicates whether a compound undergoes hardening (steeper slope) or softening (more shallow slope) during the ageing process. Apparently, both biogenic SOA types undergo hardening upon ageing, whereas the two anthropogenic SOA types undergo softening, with the strongest effects for pinene and dodecane SOA.

The area between 1 s and 1 h represents the time scale of atmospheric updraft processes. For SOA in this range, diffusion processes occur on the same time scales as typical air parcel updrafts and the predominant cloud formation process depends strongly on atmospheric conditions. All four SOA types fall within or beneath this range, indicating the importance of the actual updraft velocity for ice nucleation on glassy aerosols. But it is also obvious that SOA particles from naphthalene are most likely to be subject to kinetic effects and may thus act as IN.

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Atmospheric implications of glassy organic IN

Organic aerosols can induce cloud formation via many different pathways depending on ambient conditions and composition. At high temperature and high humidity, liquid organic particles can act as cloud condensation nuclei (CCN). At lower temperatures, they facilitate formation of ice crystals. Figure 5 summarizes how the phase state and morphology of atmospheric organic aerosol particles may vary upon changes in ambient relative humidity (humidity-induced phase transitions). Upon humidifying, the phase state changes from amorphous solid (glassy) over a partially-deliquesced state with a solid core residual coated by a liquid shell to a fully-deliquesced liquid. Upon drying the transition may occur via an inverse core-shell morphology, i.e. a liquid coated by a solid shell. Consequently, the particle phase state determines the active ice nucleation pathway: glassy solids can nucleate ice in the deposition mode, partially deliquesced particles with core-shell morphologies may act as IN in the immersion mode and liquid particles nucleate ice homogeneously, at significantly higher ice supersaturation.

From the SOA types investigated in this study, aromatic SOA or highly aged  $\alpha$ -pinene SOA may persist in a glassy state to the highest temperatures and humidities and may thus facilitate heterogeneous ice nucleation at temperatures of up to 225 K. Below 210 K, SOA from all precursors are expected to be in the glassy state required for heterogeneous ice nucleation. Our microphysical simulations suggest a potential anthropogenic influence of IN from emission of aromatic VOCs and by providing high oxidative capacities in urban areas leading to an increase of ice nucleation in and on glassy organic particles.

Compared to typical atmospheric IN such as dust, soot and biological particles, glassy organic particles require temperatures below  $\sim 230$  K to nucleate ice heterogeneously (Hoose and Möhler, 2012). This restriction confines their atmospheric activity range to the upper troposphere – lower stratosphere region since the glassy state is

ACPD

14, 16451–16492, 2014

### Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



prevalent only up to temperatures of about 200–240 K under typical atmospheric humidities ( $S_{\text{ice}} \approx 1$ ), depending on composition.

In this study we show a strong interplay between diffusion time scales in the atmosphere and atmospheric updraft speeds: the stronger the updraft and the larger the particle size, the more kinetic limitations delay the liquefaction of glassy particles. These findings also imply that an ice nucleation onset determined in laboratory studies needs to be interpreted carefully in order to apply it to realistic atmospheric parameters, i.e. humidification rate, particle size and starting humidity. Kinetic limitations are already pronounced at the smallest atmospherically relevant updraft velocities of  $0.02 \text{ m s}^{-1}$ . When humidification is fast (e.g. in convective updrafts), the glassy state may persist well above its quasi-equilibrium boundaries. In situations where particles equilibrated in dry air ( $S_{\text{ice}} < 0.9$ ) are elevated quickly, upper temperature limits for immersion freezing on glassy organics might reach much higher values than the conservative estimates given in this study. Thus, also mid-latitude cloud formation may be affected by this heterogeneous ice nucleation pathway. Such ice nucleation potential of organic aerosols challenges traditional views and requires new formalisms in regional and global models for cloud formation and precipitation, climate and air quality.

## Appendix A: Details on the estimation of bulk diffusivities from glass transition and hygroscopicity data

### A1 Justification of the method

Even though the estimation scheme described in Sect. 2.2 represents a rather crude estimation of water diffusivities, it builds on basic physical principals: in solutions of chemically similar organic substances (like the mixture of highly functionalized organic species in SOA), the types of molecular interactions are mostly hydrogen bonds and dispersion interactions, irrespective of the actual composition. Differences in diffusive properties are to a substantial degree due to factors such as molar mass and shape,

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



both of which directly affect the glass transition temperature (Koop et al., 2011). The way by which the glass transition is approached is not affected strongly by the substance type, as all organic compounds relevant for SOA are *fragile* glass-formers (Angell, 1985). The proposed method is consistent with the following previous studies.

Rampp et al. (2000) used NMR spectroscopy to determine water diffusion coefficients in different carbohydrate matrices (sucrose, allosucrose, leucrose, trehalose) and fitted VFT parameters to the temperature and concentration-dependent data sets. Overall, similar VFT parameters  $A$  and  $B$  were found for these chemically similar substances, even though  $D_{\text{H}_2\text{O}}$  seemed to depend strongly on organic mass fraction, thus supporting assumptions 1 and 2 above. The observed concentration dependence was described almost exclusively by a change in  $T_0$ , with only small trends in  $A$  and minor variation in  $B$ , possibly due to experimental error, thus supporting assumption 3.

Angell (1997) investigated the correlation of Kauzmann temperatures  $T_k$  with Vogel temperatures  $T_0$  and found their ratio to be close to unity. The ratio of  $T_g$  to  $T_0$  has been shown to be confined to a narrow range between  $1.07 < T_g/T_0 < 1.82$  for a wide variety of strongly different substances. This ratio seems to be correlated in magnitude to the substances fragility (i.e. VFT parameter  $B$ ), with high fragilities implying high  $T_g/T_0$  ratios. Reversely the assumption of similar fragilities (assumption 2) directly points towards similar  $T_g/T_0$  ratios (assumption 3). Accordingly, deducing Vogel temperatures  $T_0$  from glass transition properties seems reasonable.

## A2 Estimation of glass transition temperatures $T_{g,\text{org}}$

The proposed estimation scheme enables the prediction of bulk diffusion coefficients only from knowledge of glass transition values for the desired RH range. The glass transition curve can be described by three parameters: the glass transition temperature of the pure molecular compound  $T_{g,\text{org}}$ ; the Gordon-Taylor constant  $k_{\text{GT}}$  of the aqueous organic mixture; and the hygroscopicity  $\kappa_{\text{org}}$  for translating composition into water activity.  $T_{g,\text{org}}$  exhibits a linear correlation with melting point  $T_m$ , also known as the Boyer-Beaman rule (Koop et al., 2011).  $T_m$  can be estimated by group contribution

16468

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



models with knowledge of its chemical structure. We use the melting point prediction model of UPPER (Unified Physical Property Estimating Relationships) as presented by Jain et al. (Jain et al., 2004; Jain and Yalkowsky, 2006).

Table S1 shows our choice of marker substances for four different types of SOA along with molar mass, melting points predicted with UPPER and predicted glass transition values based on the Boyer-Beaman rule. The SOA groups were chosen to include SOA from the most commonly studied precursors and are derived from one specified precursor substance each. The groups “ $\alpha$ -Pinene” and “Isoprene” represent SOA from biogenic origin, whereas “Naphthalene” and “Dodecane” are our choice for precursors of anthropogenic origin.

The group “ $\alpha$ -Pinene” contains compounds characteristic for photo-oxidation of the biogenic SOA precursor  $\alpha$ -pinene, which has been chosen as proxy for the different monoterpene VOCs responsible for biogenic SOA formation. The list contains compounds with the highest yields according to MCM-based simulations of Shilling et al. (2009) as well as of Zuend and Seinfeld (2012), who also included two dimer substances. Furthermore, we included 3-MBTCA, a highly oxidized pinene derivative found in ambient samples (Szmigielski et al., 2007) as well as terpenylic acid, a tracer for rather fresh SOA, along with two of its derivatives (Claeys et al., 2009).

The group “Isoprene” contains isoprene-derived compounds found in ambient and laboratory aerosol as suggested by Surratt et al. (2006) and references therein. These authors also proposed a high contribution of esterification products with 2-methylglyceric acid as monomeric unit to SOA mass. Table S1 lists these oligomers up to the tetramer level, where predicted glass transition values start to level off.

The group “Naphthalene” represents typical products originating from the oxidation of anthropogenic aromatic precursors. Note that for highly functionalized aromatic compounds, UPPER predicts unusually high values for  $T_m$ , which are inconsistent with observations. For example, phthalic acid melts under decomposition (presumably anhydrate formation) at 403 K (Lide, 2005), whereas UPPER suggests a melting point of about 539 K. For this reason, we used only those naphthalene oxidation products

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for which literature melting points are known, such as the substances given in Saukko et al. (2012) and a number of compounds listed in Kautzman et al. (2010). Note that for the same reason we did not include oligomerization products to the “Naphthalene” group. Oligomerization is however also expected for aromatic SOA, shown e.g. by Kalberer et al. (2004), which would lead to higher  $T_{g,org}$  (Koop et al., 2011). For these reasons, our estimates for aromatic SOA materials may be regarded as a conservative estimate.

The group “Dodecane” in Table S1 lists oxidized organics derived from the C12 straight chain alkane to represent the family of similar compounds originating from aliphatic VOCs of anthropogenic origin. The list is a selection from the comprehensive chemical mechanism in Yee et al. (2012) and three compounds from those suggested by Zhang et al. (2014).

The resulting glass transition values are presented in Fig. A2 as a function of atomic O/C ratio and a clear positive correlation is observed within each group of compounds. Solid lines are obtained by linear regressions using a bisquare weighting function and shaded areas are confidence intervals at the  $1\sigma$  level. The chosen marker compounds occupy compound-specific ranges of O/C values, which is in part due to a different carbon number in the precursor molecule. To estimate a value characteristic for a mixture of the single compounds, we choose three values of O/C ratios that are typical for the respective group and take at each of those values the corresponding  $T_{g,org}$  that arises from the linear fit. The errors are then given by the extension of confidence bands at each point. The results are shown in Table A1.

### A3 Estimation of Gordon-Taylor constants $k_{GT}$

Gordon-Taylor constants are necessary to estimate the glass transition temperatures of compound mixtures. Zobrist et al. (2008) determined Gordon-Taylor constants for a variety of atmospherically relevant substances and SOA proxies. However, data are sparse when compared to the wide structural variety of compounds in SOA and no clear correlation can be drawn from the molecular structure. For this reason,

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Koop et al. (2011) recommended the usage of a mean Gordon-Taylor constant of  $k_{GT} = 2.5 \pm 1$  (cf. Table A1). Figure S6 shows the temperature dependence of FDRH in calculations similar to Figs. 2 and 3, this time using the best guess parameters recommended in Koop et al. (2011). The uncertainty in FDRH that arises from the given input parameter ranges is shown (grey shaded), but also the specific uncertainty from varying  $k_{GT}$  between 1.5 and 3.5 is highlighted (orange shaded).

### A4 Estimation of hygroscopicities $\kappa_{org}$

The hygroscopicity of a compound can be expressed by a single parameter  $\kappa_{org}$ , which is strongly correlated to its degree of oxidation (Petters and Kreidenweis, 2007; Lambe et al., 2011). A typical value for  $\kappa_{org}$  in biogenic SOA particles collected in pristine rainforest environments is 0.1 (Gunthe et al., 2009), which was also used by Koop et al. (2011) for their estimation of glass transition values in biogenic SOA.

For estimation of  $\kappa_{org}$ , we use the parameterization of Lambe et al. (2011) that correlates the O/C ratio of secondary organic material to its hygroscopicity, Eq. (A1).

$$\kappa_{org} = (0.18 \pm 0.04) \cdot O/C + 0.03 \quad (A1)$$

Each SOA precursor class is assigned a typical O/C value from previous investigations of marker compounds (cf. Fig. A2) and results are shown in Table A1. With the knowledge of  $T_{g,org}$ ,  $k_{GT}$  and  $\kappa_{org}$ , the entire glass transition curves for the four SOA types can be calculated, as visualized in Fig. S4. Dashed lines and grey shaded areas indicate ranges of uncertainty.

### A5 Evaluation of the method

For evaluation of the performance of the diffusivity estimation scheme, we compare estimated diffusivity values with values obtained in experiments by Price et al. (2014). In these experiments,  $D_2O$ - $H_2O$  exchange in an organic matrix at constant temperature and humidity is investigated by Raman spectroscopy. Figure S3 shows the experimen-

tally determined  $D_{\text{H}_2\text{O}}$  values for sucrose and levoglucosan in Price et al. (blue and purple markers) as well as the  $D_{\text{H}_2\text{O}}$  parameterization from Zobrist et al. (2011) (blue solid line).  $D_{\text{H}_2\text{O}}$  in levoglucosan has also been estimated with the diffusivity estimation scheme (purple solid line), utilizing input parameters from Zobrist et al. (2008) ( $T_{\text{g,org}} = 283.6 \text{ K}$ ,  $k_{\text{GT}} = 5.2$ ). Water activity has been parameterized using the parameters in Table S4.

Experimental and estimated values coincide for the highest and lowest water activities but differ under medium conditions due to the different curvature of the base parameterization from Zobrist et al. that underlies all calculations. However, diffusivities differ only within at most two orders of magnitude, which is a considerably small deviation compared to the large set of approximations made here and the difference between experimental techniques.

## Appendix B: Application of the model to ice nucleation experiments in the literature

### B1 Sucrose experiments

Baustian et al. (2013) investigated sucrose particles deposited on a quartz substrate and humidified inside an experimental flow cell. After cooling and drying below the glass transition, particles with an average diameter of  $4 \mu\text{m}$  were humidified by cooling at a rate of  $0.1 \text{ K min}^{-1}$ . Humidification was initialized below ice saturation ( $S_{\text{ice}} < 0.9$ ). The resulting heterogeneous ice nucleation onsets (brown circles) are shown in Fig. 2c along with the full deliquescence relative humidity (FDRH, blue solid line) from multiple model runs (spacing:  $2 \text{ K}$ ) mimicking the experimental conditions. Simulations below  $215 \text{ K}$  (left black square marker) are found to nucleate in the deposition mode, whereas particles in runs between about  $215 \text{ K}$  and  $238 \text{ K}$  (right black square marker) are assumed to undergo immersion freezing. This result is compliant with the experimental values, none of which exceeds a nucleation temperature of  $235 \text{ K}$ . Above  $238 \text{ K}$  full

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

deliquescence occurs before the ice supersaturation required for heterogeneous ice nucleation (brown dashed line) is reached. Also, homogeneous ice nucleation is not possible anymore below the water saturation limit according to Koop et al. (2000; green dashed line), leaving no remaining ice nucleation pathway.

## 5 B2 Naphthalene SOA experiments

Wang et al. (2012) generated SOA by oxidation of naphthalene by OH in a potential aerosol mass (PAM) reactor, deposited the particles on glass slides and investigated the onsets of water uptake and ice nucleation inside an ice nucleation cell that was mounted on a microscope. Experimental results are shown in Fig. B1 for three different SOA oxidation states: Low O/C (0.27) given in red, medium O/C (0.54) in green and high O/C (1.0) in blue. For the comparing model simulations, we employ our diffusivity estimation scheme with the glass transition parameterization for naphthalene given above. A humidification rate of  $1\% \text{ RH min}^{-1}$  was employed and temperature varied accordingly to maintain a constant dew point. In Fig. B1, the lines of full deliquescence relative humidity (FDRH) divide the measured heterogeneous ice nucleation onsets into two groups, irrespective of the degree of oxidation used in the simulation. Heterogeneous nucleation at or below 225 K is consistent with simulation results (closed diamonds), whereas ice nucleation at or above 230 K cannot be explained with the estimated water diffusion properties (open diamonds). According to the model simulation, naphthalene SOA should be already deliquesced at temperatures and RH where ice nucleation is still experimentally observed. The model simulations thus suggest that a different ice nucleation mechanism than heterogeneous ice nucleation on glassy particles was active in these experiments.

The reliability of the method is confirmed by comparing experimental and modelled water uptake onsets that show very good correlation. The modelled water uptake onset was defined as the point where the particle diameter had increased by 100 nm to take into account the fact that experimental onsets were determined by visible inspection under a light microscope.

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### B3 Citric acid experiments

Murray et al. (2010) observed the process of heterogeneous ice nucleation on glassy aerosols by investigating citric acid particles in the AIDA cloud chamber. The experimentally determined onsets of heterogeneous (orange diamonds) and homogeneous ice nucleation (green circles) are shown in Fig. B2 along with results of simulations mimicking the experimental conditions. In the calculations, we assumed a particle diameter of 150 nm and a humidification rate of 12 % RH min<sup>-1</sup>, corresponding to a cooling rate of 1–2 K min<sup>-1</sup>. Humidification was initiated at  $S_{\text{ice}} = 1$  since the cloud chamber walls were covered with ice during the initial cooling process. We performed two series of simulations for two different water activity parameterizations available in the literature. According to the parameterization in Lienhard et al. (2012) (dashed lines), heterogeneous nucleation occurs exclusively above the (equilibrium) glass transition relative humidity  $RH_g$  and thus in the immersion freezing regime. With the parameterization from Koop et al. (2011) (solid lines), equilibrium glass transition and full deliquescence occur at later stages in the humidification process. According to this data, only the experimental data point at about 206 K would have occurred in the immersion mode.

At 212 K, ice nucleation occurs only homogeneously in Murray's experiments, indicated by the much later ice nucleation onset. The humidification run started with liquid aerosol particles that showed retarded deliquescence, but were not able to nucleate ice heterogeneously.

**The Supplement related to this article is available online at doi:10.5194/acpd-14-16451-2014-supplement.**

*Acknowledgements.* This work was funded by the Max Planck Society (MPG), the Deutsche Forschungsgemeinschaft through the Ice Nuclei Research Unit INUIT (FOR1525, Grant KO 2944/2-1), and the European Commission under the PEGASOS project (grant no. 265148). T. B. was supported by the Max Planck Graduate Center with the Johannes Gutenberg-

16474

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chan, A. W. H., Herrmann, H., Seinfeld, J. H., and Maenhaut, W.: Terpenylic acid and related compounds from the oxidation of alpha-pinene: implications for new particle formation and growth above forests, *Environ. Sci. Technol.*, 43, 6976–6982, doi:10.1021/es9007596, 2009.

Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M. H., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the dominant sources and mechanisms of cirrus cloud formation, *Science*, 340, 1320–1324, doi:10.1126/science.1234145, 2013.

DeMott, P. J., Cziczo, D. J., Prenni, A. J., Murphy, D. M., Kreidenweis, S. M., Thomson, D. S., Borys, R., and Rogers, D. C.: Measurements of the concentration and composition of nuclei for cirrus formation, *P. Natl. Acad. Sci. USA*, 100, 14655–14660, doi:10.1073/pnas.2532677100, 2003.

Froyd, K. D., Murphy, D. M., Lawson, P., Baumgardner, D., and Herman, R. L.: Aerosols that form subvisible cirrus at the tropical tropopause, *Atmos. Chem. Phys.*, 10, 209–218, doi:10.5194/acp-10-209-2010, 2010.

Fulcher, G. S.: Analysis of recent measurements of the viscosity of glasses, *J. Am. Ceram. Soc.*, 8, 339–355, doi:10.1111/j.1151-2916.1925.tb16731.x, 1925.

Gunthe, S. S., King, S. M., Rose, D., Chen, Q., Roldin, P., Farmer, D. K., Jimenez, J. L., Artaxo, P., Andreae, M. O., Martin, S. T., and Pöschl, U.: Cloud condensation nuclei in pristine tropical rainforest air of Amazonia: size-resolved measurements and modeling of atmospheric aerosol composition and CCN activity, *Atmos. Chem. Phys.*, 9, 7551–7575, doi:10.5194/acp-9-7551-2009, 2009.

Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817–9854, doi:10.5194/acp-12-9817-2012, 2012.

IPCC: Climate Change 2013, The Physical Science Basis, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.

Jain, A. and Yalkowsky, S. H.: Estimation of melting points of organic compounds-II, *J. Pharm. Sci.*, 95, 2562–2618, doi:10.1002/jps.20634, 2006.

Jain, A., Yang, G., and Yalkowsky, S. H.: Estimation of melting points of organic compounds, *Ind. Eng. Chem. Res.*, 43, 7618–7621, doi:10.1021/ie049378m, 2004.

Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004.

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S., Dalleska, N. F., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of gas- and aerosol-phase products from the photooxidation of naphthalene, *J. Phys. Chem. A*, 114, 913–934, doi:10.1021/jp908530s, 2010.
- 5 Kidd, C., Perraud, V., Wingen, L. M., and Finlayson-Pitts, B. J.: Integrating phase and composition of secondary organic aerosol from the ozonolysis of  $\alpha$ -pinene, *P. Natl. Acad. Sci. USA*, 111, 7552–7557, doi:10.1073/pnas.1322558111, 2014.
- Koop, T., Luo, B. P., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, doi:10.1038/35020537, 2000.
- 10 Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238–19255, 2011.
- Kuwata, M. and Martin, S. T.: Phase of atmospheric secondary organic material affects its reactivity, *P. Natl. Acad. Sci. USA*, 109, 17354–17359, doi:10.1073/pnas.1209071109, 2012.
- 15 Lambe, A. T., Onasch, T. B., Massoli, P., Croasdale, D. R., Wright, J. P., Ahern, A. T., Williams, L. R., Worsnop, D. R., Brune, W. H., and Davidovits, P.: Laboratory studies of the chemical composition and cloud condensation nuclei (CCN) activity of secondary organic aerosol (SOA) and oxidized primary organic aerosol (OPOA), *Atmos. Chem. Phys.*, 11, 8913–8928, doi:10.5194/acp-11-8913-2011, 2011.
- 20 Lide, D. R.: *CRC Handbook of Chemistry and Physics*, CRC Press, National Institute of Standards and Technology, Boca Raton, FL, 2005.
- Lienhard, D. M., Bones, D. L., Zuend, A., Krieger, U. K., Reid, J. P., and Peter, T.: Measurements of thermodynamic and optical properties of selected aqueous organic and organic-inorganic mixtures of atmospheric relevance, *J. Phys. Chem. A*, 116, 9954–9968, doi:10.1021/jp3055872, 2012.
- 25 Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491–9522, doi:10.5194/acp-9-9491-2009, 2009.
- 30 Murray, B. J., Wilson, T. W., Dobbie, S., Cui, Z. Q., Al-Jumur, S., Mohler, O., Schnaiter, M., Wagner, R., Benz, S., Niemand, M., Saathoff, H., Ebert, V., Wagner, S., and Karcher, B.: Heterogeneous nucleation of ice particles on glassy aerosols under cirrus conditions, *Nat. Geosci.*, 3, 233–237, doi:10.1038/ngeo817, 2010.

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Perraud, V., Bruns, E. A., Ezell, M. J., Johnson, S. N., Yu, Y., Alexander, M. L., Zelenyuk, A., Imre, D., Chang, W. L., Dabdub, D., Pankow, J. F., and Finlayson-Pitts, B. J.: Nonequilibrium atmospheric secondary organic aerosol formation and growth, *P. Natl. Acad. Sci. USA*, 109, 2836–2841, doi:10.1073/pnas.1119909109, 2012.

5 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.

Price, H. C., Murray, B. J., Mattsson, J., O'Sullivan, D., Wilson, T. W., Baustian, K. J., and Benning, L. G.: Quantifying water diffusion in high-viscosity and glassy aqueous solutions using a Raman isotope tracer method, *Atmos. Chem. Phys.*, 14, 3817–3830, doi:10.5194/acp-14-3817-2014, 2014.

10 Pruppacher, H. and Klett, J.: *Microphysics of Clouds and Precipitation*, 2 edn., Kluwer Academic Publishers, Boston, MA, 1997.

Rampp, M., Buttersack, C., and Ludemann, H. D.: c,T-dependence of the viscosity and the self-diffusion coefficients in some aqueous carbohydrate solutions, *Carbohydr. Res.*, 328, 561–572, 2000.

15 Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, K., Sellier, M., Murray, B. J., Schilling, J. E., Martin, S. T., and Bertram, A. K.: Viscosity of  $\alpha$ -pinene secondary organic material and implications for particle growth and reactivity, *P. Natl. Acad. Sci. USA*, 110, 8014–8019, doi:10.1073/pnas.1219548110, 2013.

Saukko, E., Lambe, A. T., Massoli, P., Koop, T., Wright, J. P., Croasdale, D. R., Pedernera, D. A., Onasch, T. B., Laaksonen, A., Davidovits, P., Worsnop, D. R., and Virtanen, A.: Humidity-dependent phase state of SOA particles from biogenic and anthropogenic precursors, *Atmos. Chem. Phys.*, 12, 7517–7529, doi:10.5194/acp-12-7517-2012, 2012.

25 Schill, G. P. and Tolbert, M. A.: Heterogeneous ice nucleation on phase-separated organic-sulfate particles: effect of liquid vs. glassy coatings, *Atmos. Chem. Phys.*, 13, 4681–4695, doi:10.5194/acp-13-4681-2013, 2013.

Schill, G. P., De Haan, D. O., and Tolbert, M. A.: Heterogeneous ice nucleation on simulated secondary organic aerosol, *Environ. Sci. Technol.*, 48, 1675–1682, doi:10.1021/es4046428, 2014.

30 Shilling, J. E., Chen, Q., King, S. M., Rosenoern, T., Kroll, J. H., Worsnop, D. R., DeCarlo, P. F., Aiken, A. C., Sueper, D., Jimenez, J. L., and Martin, S. T.: Loading-dependent elemental

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

composition of  $\alpha$ -pinene SOA particles, *Atmos. Chem. Phys.*, 9, 771–782, doi:10.5194/acp-9-771-2009, 2009.

Shiraiwa, M., Ammann, M., Koop, T., and Pöschl, U.: Gas uptake and chemical aging of semisolid organic aerosol particles, *P. Natl. Acad. Sci. USA*, 108, 11003–11008, doi:10.1073/pnas.1103045108, 2011.

Shiraiwa, M., Pfrang, C., Koop, T., and Pöschl, U.: Kinetic multi-layer model of gas-particle interactions in aerosols and clouds (KM-GAP): linking condensation, evaporation and chemical reactions of organics, oxidants and water, *Atmos. Chem. Phys.*, 12, 2777–2794, doi:10.5194/acp-12-2777-2012, 2012.

Shiraiwa, M., Yee, L. D., Schilling, K. A., Loza, C. L., Craven, J. S., Zuend, A., Ziemann, P. J., and Seinfeld, J. H.: Size distribution dynamics reveal particle-phase chemistry in organic aerosol formation, *P. Natl. Acad. Sci. USA*, 110, 11746–11750, doi:10.1073/pnas.1307501110, 2013.

Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation and morphology of internally mixed dicarboxylic acids/ammonium sulfate/water particles, *Atmos. Chem. Phys.*, 12, 2691–2712, doi:10.5194/acp-12-2691-2012, 2012.

Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical composition of secondary organic aerosol formed from the photooxidation of isoprene, *J. Phys. Chem. A*, 110, 9665–9690, doi:10.1021/jp061734m, 2006.

Szmigielski, R., Surratt, J. D., Gomez-Gonzalez, Y., Van der Veken, P., Kourtchev, I., Vermeylen, R., Blockhuys, F., Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Edney, E. O., Seinfeld, J. H., Maenhaut, W., and Claeys, M.: 3-methyl-1,2,3-butanetricarboxylic acid: an atmospheric tracer for terpene secondary organic aerosol, *Geophys. Res. Lett.*, 34, L24811, doi:10.1029/2007gl031338, 2007.

Tammann, G. and Hesse, W.: The dependancy of viscosity on temperature in hypothermic liquids, *Z. Anorg. Allg. Chem.*, 14, 245–257, doi:10.1002/zaac.19261560121, 1926.

Tong, H.-J., Reid, J. P., Bones, D. L., Luo, B. P., and Krieger, U. K.: Measurements of the timescales for the mass transfer of water in glassy aerosol at low relative humidity and ambient temperature, *Atmos. Chem. Phys.*, 11, 4739–4754, doi:10.5194/acp-11-4739-2011, 2011.

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vaden, T. D., Imre, D., Beranek, J., Shrivastava, M., and Zelenyuk, A.: Evaporation kinetics and phase of laboratory and ambient secondary organic aerosol, *P. Natl. Acad. Sci. USA*, 108, 2190–2195, doi:10.1073/pnas.1013391108, 2011.

Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., YliPirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824–827, doi:10.1038/nature09455, 2010.

Vogel, H.: The temperature dependence law of the viscosity of fluids, *Phys. Z.*, 22, 645–646, 1921.

Wagner, R., Möhler, O., Saathoff, H., Schnaiter, M., Skrotzki, J., Leisner, T., Wilson, T. W., Malkin, T. L., and Murray, B. J.: Ice cloud processing of ultra-viscous/glassy aerosol particles leads to enhanced ice nucleation ability, *Atmos. Chem. Phys.*, 12, 8589–8610, doi:10.5194/acp-12-8589-2012, 2012.

Wang, B. B., Lambe, A. T., Massoli, P., Onasch, T. B., Davidovits, P., Worsnop, D. R., and Knopf, D. A.: The deposition ice nucleation and immersion freezing potential of amorphous secondary organic aerosol: pathways for ice and mixed-phase cloud formation, *J. Geophys. Res.-Atmos.*, 117, D16209, doi:10.1029/2012jd018063, 2012.

Wilson, T. W., Murray, B. J., Wagner, R., Möhler, O., Saathoff, H., Schnaiter, M., Skrotzki, J., Price, H. C., Malkin, T. L., Dobbie, S., and Al-Jumur, S. M. R. K.: Glassy aerosols with a range of compositions nucleate ice heterogeneously at cirrus temperatures, *Atmos. Chem. Phys.*, 12, 8611–8632, doi:10.5194/acp-12-8611-2012, 2012.

Yee, L. D., Craven, J. S., Loza, C. L., Schilling, K. A., Ng, N. L., Canagaratna, M. R., Ziemann, P. J., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from low-NO<sub>x</sub> photooxidation of dodecane: evolution of multigeneration gas-phase chemistry and aerosol composition, *J. Phys. Chem. A*, 116, 6211–6230, 2012.

You, Y., Renbaum-Wolff, L., Carreras-Sospedra, M., Hanna, S. J., Hiranuma, N., Kamal, S., Smith, M. L., Zhang, X. L., Weber, R. J., Shilling, J. E., Dabdub, D., Martin, S. T., and Bertram, A. K.: Images reveal that atmospheric particles can undergo liquid-liquid phase separations, *P. Natl. Acad. Sci. USA*, 109, 13188–13193, doi:10.1073/pnas.1206414109, 2012.

Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Belouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., and Zhou, M.: A review of measurement-based assessments of the

aerosol direct radiative effect and forcing, *Atmos. Chem. Phys.*, 6, 613–666, doi:10.5194/acp-6-613-2006, 2006.

Zhang, X., Schwantes, R. H., Coggon, M. M., Loza, C. L., Schilling, K. A., Flagan, R. C., and Seinfeld, J. H.: Role of ozone in SOA formation from alkane photooxidation, *Atmos. Chem. Phys.*, 14, 1733–1753, doi:10.5194/acp-14-1733-2014, 2014.

Zhou, S., Shiraiwa, M., McWhinney, R., Pöschl, U., and Abbatt, J. P. D.: Kinetic limitations in gas-particle reactions arising from slow diffusion in secondary organic aerosol, *Faraday Discuss.*, 165, 391–406, doi:10.1039/C3FD00030C, 2013.

Zobrist, B., Marcolli, C., Pedernera, D. A., and Koop, T.: Do atmospheric aerosols form glasses?, *Atmos. Chem. Phys.*, 8, 5221–5244, doi:10.5194/acp-8-5221-2008, 2008.

Zobrist, B., Soonsin, V., Luo, B. P., Krieger, U. K., Marcolli, C., Peter, T., and Koop, T.: Ultra-slow water diffusion in aqueous sucrose glasses, *Phys. Chem. Chem. Phys.*, 13, 3514–3526, doi:10.1039/c0cp01273d, 2011.

Zuend, A. and Seinfeld, J. H.: Modeling the gas-particle partitioning of secondary organic aerosol: the importance of liquid-liquid phase separation, *Atmos. Chem. Phys.*, 12, 3857–3882, doi:10.5194/acp-12-3857-2012, 2012.

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Table A1.** Assumed physical properties of SOA classes for use in conjunction with diffusivity estimation scheme.

SOA Class	O/C	$T_{g,org}$ (K)	$k_{GT}$	$\kappa_{org}$
$\alpha$ -Pinene	0.3	$228.9 \pm 10.6$	2.5	$0.084 \pm 0.012$
	0.5	$278.5 \pm 7.0$	2.5	$0.120 \pm 0.020$
	0.7	$328.1 \pm 12.8$	2.5	$0.156 \pm 0.028$
Isoprene	0.6	$258.2 \pm 22.2$	2.5	$0.138 \pm 0.024$
	0.8	$287.2 \pm 11.9$	2.5	$0.174 \pm 0.032$
	1.0	$316.3 \pm 19.1$	2.5	$0.210 \pm 0.040$
Naphthalene	0.3	$294.2 \pm 5.7$	2.5	$0.084 \pm 0.012$
	0.5	$313.1 \pm 8.8$	2.5	$0.120 \pm 0.020$
	0.7	$332.0 \pm 15.0$	2.5	$0.156 \pm 0.028$
Dodecane	0.1	$210.3 \pm 9.7$	2.5	$0.048 \pm 0.004$
	0.3	$216.8 \pm 5.1$	2.5	$0.084 \pm 0.012$
	0.5	$223.4 \pm 11.4$	2.5	$0.120 \pm 0.020$
Koop SOA	–	$270 \pm 21$	$2.5 \pm 1$	$0.1008$ $-0.0504$



**Figure 1.** Schematic temporal evolution of particle morphology along a trajectory of an atmospheric updraft (grey arrow). Humidification of ambient air upon adiabatic expansion leads to liquefaction of initially glassy particles (dark blue) to liquid particles (light blue). Whereas partial deliquescence (PDRH) coincides with  $RH_g$ , full deliquescence (FDRH) is delayed to much higher RH, indicating that diffusion processes occur on much longer time scales than humidification. The speed of the displayed trajectory corresponds to that typical for cloud chamber or environmental cell experiments.

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.




**Figure 3.** (A) Simulated humidification of SOA particles from the four different precursors  $\alpha$ -pinene, isoprene, dodecane and naphthalene. Naphthalene SOA (dark red) shows the latest deliquescence, whereas dodecane SOA (light red) liquefied rather early in the simulations. The two biogenic SOA estimates lie between both extremes with pinene SOA (dark green) showing slightly later deliquescence than isoprene SOA. Intercepts (square markers) with a heterogeneous nucleation onset typical for SOA (brown dashed line) indicate upper temperature limits for immersion freezing (arrows on x-axis). The effect of particle ageing also depends on precursor type: Pinene SOA (B) shows hardening upon increase in O/C (indicated by higher FDRH), whereas dodecane SOA (C) exhibits softening (indicated by lower FDRH). Similarly, isoprene and naphthalene SOA show only moderate hardening and softening, respectively (Fig. S5).

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

Water diffusion time scales in SOA (220 K,  $S_{ice} = 1.45$ ,  $d_p = 100$  nm)



**Figure 4.** Characteristic time scales of water diffusion in SOA as function of hygroscopicity  $\kappa_{org}$  and glass transition temperatures of the pure organic matrix  $T_{g,org}$ . Calculations have been performed at 220 K,  $S_{ice} = 1.45$  and for 100 nm particles. Oval shapes confine estimated ranges in  $\kappa_{org}$  and  $T_{g,org}$  for the four SOA types in three different oxidation states (Appendix A and Table A1). The grey area indicates the time scale of typical atmospheric updrafts (1 s to 1 h) and thus divides the plot in areas of quasi-equilibrium and non-equilibrium water diffusion. Within the grey area, the relative speed of both processes depends upon the actual atmospheric conditions. The aging process is indicated by arrows pointing from regions of low O/C to regions of high O/C.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 5.** Overview of processes in organic aerosol particles affecting atmospheric cloud formation. Particles form by oxidation of volatile organic compounds (VOCs) originating from anthropogenic and biogenic emissions. The dominating cloud formation process depends on particle phase state, which is a function of temperature and humidity. Humidity-induced phase transitions between phase states may be kinetically limited and occur under formation of partially deliquesced particles with core-shell morphologies. Glassy or partially deliquesced particles are able to undergo heterogeneous ice nucleation, occurring at lower relative humidity or higher temperature than homogeneous ice nucleation of liquid particles.

**Water uptake and ice nucleation by glassy aerosols**

T. Berkemeier et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Figure A1.** Determination of heterogeneous ice nucleation onsets. For sucrose, data from Baustian et al. (2013) (blue circles) are fitted. For SOA, deposition freezing data on Naphthalene SOA from Wang et al. (2012) (red downward triangles) as well as nucleation data on aqSOA from Schill et al. (2014) (pink upward triangles) are used. The resulting linear regression fits (blue and red dashed lines) lie significantly below the homogeneous nucleation limit and are displayed along with their parameterizations.

## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.



**Figure A2.** Predicted glass transition values of SOA marker substances as function of O/C ratio. The predicted  $T_{g,org}$  exhibit a linear correlation with O/C for each of the four SOA systems. Solid lines are robust linear regressions using a bisquare weighting function and shaded areas are confidence intervals at the  $1\sigma$  level. Anthropogenic aliphatic SOA constituents show the lowest values of  $T_{g,org}$  and a weak dependence on O/C. In contrast, aromatic SOA shows the highest glass transition values despite a rather low average O/C ratio.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.



**Figure B1.** Comparison between calculation results of naphthalene SOA deliquescence and experimental ice nucleation and water uptake data from Wang et al. (2012). For the numerical simulations, aerosol particles are assumed to be 1  $\mu\text{m}$  in diameter and are humidified at a rate of 1 % RH  $\text{min}^{-1}$ , corresponding to a cooling rate of about 0.1 K  $\text{min}^{-1}$  used by Wang et al. (2012).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Water uptake and ice nucleation by glassy aerosols

T. Berkemeier et al.



**Figure B2.** Comparison between calculation results of citric acid aerosol deliquescence (orange lines) and experimental ice nucleation data from Murray et al. (2010) (orange diamonds, green circles). In the numerical simulations, 150 nm diameter aerosol particles are humidified at a rate of  $12\% \text{ RH min}^{-1}$ , corresponding to a cooling rate around  $1\text{--}2 \text{ K min}^{-1}$  typical for cloud chamber experiments. Black lines and shaded areas confine the region where a glass is the favored thermodynamic state. The dashed lines were obtained using the water activity parameterization provided by Lienhard et al. (2012), whereas the solid lines were obtained with the parameterization in Koop et al. (2011).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

