

Response to Reviewer #1

(1) This manuscript simulates atmospheric aerosol formation over Europe using a regional modeling framework. The investigated research topic is very important, yet very few investigations like this have published before. As a result, I consider the paper very welcome to the scientific community. The paper itself is clearly written and easy to follow. The presented analysis appears scientifically sound, but it should be expanded a bit in order to get a better idea how robust the obtained results really are.

We do appreciate the positive comments.

Major issues:

(2) As the authors point out in section 2 (the top of page 13589), organic vapors do not assist the growth of <100 nm in the current model implementation. This is understandable considering the current complexities in simulating secondary organic aerosol formation. However, since organics probably do play an important role in ultrafine aerosol growth in many parts of Europe, the authors should bring out this thing in a bit concrete manner when discussing the results and their implications. The model evaluation is based only on comparing particle number concentration. The capability of the model to reproduce observed growth rate could be investigated as well, at least for the sites for which such information has been published in the literature. Such an analysis, if possible carry out based on model output information, would immediate give some hints whether "missing organic condensation" is a serious problem and where.

This is a valid concern. As suggested, we have calculated predicted growth rates (from the model output, based on the method describe in Hirsikko et al. 2005) and compared with observed values that are reported in Manninen et al. 2010. We have added a new figure showing this comparison and a new section (new paragraph 5.2) discussing the results as well as the possible effect of limited organic condensation on the results. As expected, the model underpredicts the growth rates in all studied sites, most probably due to insufficient organic condensation in the model. This is now stated in the paper more clearly as suggested by the reviewer.

(3) Concerning the sensitivity analysis, is there any possibility to investigate how sensitive the results are on the availability of condensable vapors (other than sulfuric acid)? For example, could one think of artificially enhancing the ultrafine particle growth rate by a certain factor to mimic what organics might do for these particles.

We have performed a new sensitivity run to quantify the effect of organics condensation as suggested by the reviewer. The secondary OA contribution to the particle growth rate was artificially enhanced by assuming that a fraction of the new secondary organics has zero volatility. Enhancing the organics condensation resulted in an increase of N_{100} by 10 percent on average over the whole domain. However, during nucleation event days the predicted increase was larger. In Melpitz for example, where frequent nucleation events are predicted (and observed), N_{100} increased by 30 percent in this test. This was the largest increase predicted in N_{100} among the 7 sites studied. At the same the total OA concentration in the new simulation increased on average by approximately a factor of 1.7. Text has been added discussing the results in the new section 5.2 (second paragraph).

Minor/technical issues:

(4) The kinetic approach (equation 2) was suggested already by McMurry and Friedlander (1979, Atmos. Environ., p 1635) and should therefore be cited here.

We have added the reference to the original work.

(5) Page 13597, lines 9-10: organics contributing to the growth of fresh particles are considered to be low-volatile rather than semi-volatile.

Corrected.

References

Hirsikko, A., Laakso, L., Horrak, U., Aalto, P. P., Kerminen, V.-M., and Kulmala, M.: Annual and size dependent variation of growth rates and ion concentrations in boreal forest, *Boreal Env. Res.*, 10, 357–369, 2005.

Manninen, H. E., Nieminen, T., Asmi, E., Gagne, S., Hakkinen, S., Lehtipalo, K., Aalto, P., Vana, M., Mirme, A., Mirme, S., Horrak, U., Plass-Dulmer, C., Stange, G., Kiss, G., Hoffer, A., Toro, N., Moerman, M., Henzing, B., de Leeuw, G., Brinkenberg, M., Kouvarakis, G. N., Bougiatioti, A., Mihalopoulos, N., O'Dowd, C., Ceburnis, D., Arneth, A., Svenningsson, B., Swietlicki, E., Tarozzi, L., Decesari, S., Facchini, M. C., Birmili, W., Sonntag, A., Wiedensohler, A., Boulon, J., Sellegri, K., Laj, P., Gysel, M., Bukowiecki, N., Weingartner, E., Wehrle, G., Laaksonen, A., Hamed, A., Joutsensaari, J., Petaja, T., Kerminen, V.-M., and Kulmala, M.: EUCAARI ion spectrometer measurements at 12 European sites – analysis of new particle formation events, *Atmos. Chem. Phys.*, 10, 7907–7927, doi:10.5194/acp-10-7907-2010, 2010.

Response to Reviewer #2

This paper discusses a modeling study of aerosol particle formation and evolution with a sectional 3-D model. The paper is generally well written and has important new information. Below are some requested edits and calculations.

(1) Introduction. Mention that secondary particles also form by in cloud processing of aerosols followed by cloud drop evaporation, releasing a modified aerosol.

This is now mentioned in the revised manuscript.

(2) P. 13588. "...assuming a density of 1.4 g/cm³." Is this a constant density assumed for all particles of all sizes and for all times? What about when soil dust particles are present in larger size bins?

No, the density is not constant. The aerosol density is calculated and updated continuously in each size bin based on the particle composition. The value of 1.4 g/cm³ mentioned in the original text was indicative to estimate the corresponding size range. To avoid confusion we have deleted this statement.

(3) P. 13588. "For the integration of the differential equations during nucleation..." Does this mean that nucleation is operator split from condensation? If so, please state explicitly.

Nucleation and condensation rates are simulated together using the pseudo-steady-state approximation (PSSA) for sulfuric acid vapor (Pierce and Adams, 2009). We have re-phrased this sentence to avoid any misunderstandings.

(4) P. 13588. "Condensation of ammonia..." Ammonia is more likely to dissolve in solution than condense. Is its dissolution treated? What about crystallization to ammonium nitrate or ammonium sulfate?

Ammonia dissolution as well as formation of ammonium salts is treated by making use of the ISORROPIA (Nenes et al., 1998) aerosol thermodynamic model.

(5) P. 13588. "We limit the ammonia condensation flux in each time step to prevent condensing more than what is available..." Shouldn't the flux also be constrained by the pH of solution in drops of each size, which itself is a function of all other acids and the condense or dissolve (e.g., Jacobson, *Aerosol Science, and Technology* 39, 92-103, 2005). Even if this is not included, it would be useful to clarify what the ammonium concentration in solution is a function of.

This is actually constrained by the ISORROPIA model. We have now deleted this sentence to avoid any misunderstandings.

(6) P. 13592. "The first two days of each simulation were excluded..." Did you check whether results from the first two days are really inaccurate compared with other days or were they just automatically excluded? I would suggest to check whether there really was much error.

We did check the results from the first two days and no major problems were evident compared to the other days. Despite that, we decided to be cautious and exclude these first 2 days to minimize any influence of the rather arbitrary initial conditions on our results.

(7) P. 13597. What was the operator-splitting time step used and what was the order of calculation?

Based on results of Jung et al. (2006) the operator-splitting time step used here is 300 s (which corresponds to the master time step of PMCAMx-UF). The calculations follow in general the order of gas-phase chemistry – nucleation – coagulation – condensation. More specifically, in the operator-splitting scheme, the condensation of NH₃ is simulated for $\Delta t/2$ (where Δt is the operator time step) and then particles are allowed to coagulate for $\Delta t/2$. In the next step, the gas-phase chemistry and nucleation equations are integrated. Finally the overall step ends with calculation of coagulation for $\Delta t/2$ and condensation for $\Delta t/2$.

(8) P. 13597. Since the authors are operator splitting nucleation and growth, I think it is important to do the following two sensitivity tests:

1) Solve condensation before rather than after nucleation each time step to see whether the depletion of sulfuric acid (and ammonia) during condensation reduces the nucleation rate. 2) Reduce the operator-split time step between nucleation and condensation to see whether increased competition reduces the nucleation rate.

This is not applicable to the current structure of the model (see also reply to previous comment for the order of calculation). As stated in the text, the two species (i.e. sulfuric acid and ammonia) are treated independently in the aerosol microphysics scheme. For sulfuric acid vapor, nucleation and condensation rates are calculated simultaneously. We have included some text explaining this and also stating that a different model structure (e.g. Jacobson, 2002) could also be used.

(9) Table 2. Use the same number of significant digits for the model and observations.

Done.