

## ***Interactive comment on “Local short-term variability in solar irradiance” by Gerald M. Lohmann et al.***

**Gerald M. Lohmann et al.**

gerald.lohmann@uni-oldenburg.de

Received and published: 20 April 2016

Thank you very much for reviewing our manuscript. We appreciate the feedback and feel that the comments have helped us to improve the quality of the paper.

Below, each comment is quoted in italics and followed by its respective author response. A corresponding revised version of the manuscript is attached to this response as a supplement. It has been prepared by means of latexdiff and highlights all differences between the original and revised versions of the paper. All page and line numbers quoted below refer to this supplement file.

*This paper examines clear sky irradiance at 1 Hz. Given the response time of solar photovoltaics and solar photothermal, is this frequency neces-*

Printer-friendly version

Discussion paper



*sary? I suspect an argument for this could go either way; please include a discussion in the discussion/conclusions.*

On the one hand, typical solar thermal energy (STE) systems are intentionally designed with storage capacities, and 1 Hz irradiance data is probably not necessary for the majority of applications involving STE.

On the other hand, the response of a single photovoltaic cell to changes in its illumination is orders of magnitude faster than 1 Hz and thus virtually instantaneous within the scope of our analyses. Of course, when aggregating many cells in a PV module, and then connecting a large number of modules in a PV system, spatial smoothing increases the system's response time. For very many inter-connected PV systems in a very large area, e.g. all of Europe, the necessary temporal resolution of data is hence appreciably reduced (the European Energy Exchange, for example, uses 15 minute time steps for electricity trading). However, Marcos et al. (2011b) find rare power fluctuations of e.g. up to  $\pm 50\%$  from one second to the next (and changes of more than 90 % for a time lag of 20 s) in 1 Hz power data from a single, relatively small, 48 kWp PV plant (typical rooftop systems are smaller yet and experience less smoothing). Yordanov et al. (2013b) even argue that the optimal temporal resolution of single point measurements should be around 10 Hz, in order not to miss extremely short but relatively high magnitude changes. Thus, a temporal resolution of 1 Hz may not be necessary for large-scale analyses, but it is key to characterize local short-term variability in solar irradiance on the spatio-temporal scales that we investigated.

We have extended the corresponding paragraphs in section 6 (Discussion) on page 18 (lines 30–32) and page 19 (lines 1–11) in line with the above answer.

*The finding that of a very low spatial autocorrelation at  $\tau=1$  is not surprisingly, given the physical nature of clouds. Indeed, previous papers cited by the author suggest that the spatial autocorrelation should be low even at*

*5 or 15 minute time steps. Why is this important to have discovered, and what does it show beyond what we already know?*

We agree that very low spatial autocorrelation coefficients of clearsky index increments  $\rho_{ij}^{\Delta k_{\tau}^*}$  are indeed no surprise for  $\tau = 1$  s. However, we rather see the reason for this in the smallest inter-sensor separation bin (ranging from 28 m to 51 m) being much greater than typical cloud speeds (between  $2 \text{ ms}^{-1}$  and  $10 \text{ ms}^{-1}$ ). The cloud-induced shadows would frequently have to cover the shortest of the sensor pair distances within a second, in order to yield high values of  $\rho_{ij}^{\Delta k_{\tau}^*}$  at short distances. For a robust characterization of the decrease of  $\rho_{ij}^{\Delta k_{\tau}^*}$  from 1 to 0, the pyranometer network would have to be reconfigured to feature much shorter inter-sensor distances. We have inserted a short paragraph along these lines at the appropriate place in section 7 (Conclusions) on page 20 (lines 21–25) to make ourselves more clear.

We disagree, however, that previously published papers would generally suggest  $\rho_{ij}^{\Delta k_{\tau}^*}$  to yet be low for time lags of 300 s or even 900 s on the small spatial scales considered by our analysis. For example, the virtual networks studied by Perez et. al. (2012) on average suggest positive station pair correlation coefficients of 1.0 at distances of 100 m for these time lags, and correlation coefficients of  $\sim 0.6$  ( $\sim 0.75$ ) for distances of 1000 m for 300 s (900 s) increments (see their Fig. 4). Likewise, the satellite-derived model by Hoff and Perez (2012) used for comparison in our Fig. 6 predicts correlation coefficients for e.g. 1000 m distances to range from 0.375 through 0.75 for time lags of 300 s, and from 0.643 through 0.9 for 900 s (using effective wind speeds between  $2 \text{ ms}^{-1}$  and  $10 \text{ ms}^{-1}$ ). In good agreement with the above, Hinkelman (2013) shows 300 s increments to be associated with  $\rho_{ij}^{\Delta k_{\tau}^*} \simeq 1$  for distances of about 100 m and  $\rho_{ij}^{\Delta k_{\tau}^*} \simeq 0.4$  for distances of about 1000 m, based on data from a small pyranometer network during 13 days of broken clouds.

*A similar question with the decorrelation... Wouldn't it make sense that*

*the decorrelation distances would be a function of clouds? In an area of almost entirely one climatic zone, cloud conditions in one location will (almost by definition) be correlated with other locations, and thus the distance needed to obtain decorrelation will increase. It is not clear this is novel; at a minimum, the authors should describe why this finding agrees entirely with what one would expect from real time data.*

We agree that decorrelation distances of  $\Delta k_{\tau}^*$  greatly depend on cloud field properties. In previous analyses for example, Hoff and Perez (2010 and 2012) argue that (de)correlation of  $k^*$  increments should mainly be a function of (1) distance, (2) time scale, and (3) effective cloud speed, while Perpignan et al. (2013) show spatial autocorrelation structures of PV power fluctuations to depend on (1) distance, (2) time scale, and (3) one of three daily 'fluctuation categories' (which are comparable to our sky types). The distinction between sky types is thus not what is novel about our study. What is novel, however, is the detailed characterization of  $\rho_{ij}^{\Delta k_{\tau}^*}$  at very high spatio-temporal resolutions, including – but not limited to – decorrelation distances under different sky conditions. We have removed the single sentence emphasizing the varying decorrelation distances under different sky conditions from section 7 (Conclusions) on page 20 (lines 20–21) in order to avoid potential misunderstandings in terms of what's novel.

Please also note the supplement to this comment:

<http://www.atmos-chem-phys-discuss.net/acp-2016-2/acp-2016-2-AC2-supplement.pdf>

---

Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2016-2, 2016.

Printer-friendly version

Discussion paper

