

A comprehensive biomass burning emission inventory with high spatial and temporal resolution in China

Ying Zhou^{1,2}, Xiaofan Xing^{1,2}, Jianlei Lang^{1,2}, Dongsheng Chen^{1,2}, Shuiyuan Cheng^{1,2,3}, Lin Wei^{1,2}, Xiao Wei⁴, Chao Liu⁵

¹Key Laboratory of Beijing on Regional Air Pollution Control, Beijing University of Technology, Beijing 100124, China

²College of Environmental & Energy Engineering, Beijing University of Technology, Beijing 100124, China

³Collaborative Innovation Center of Electric Vehicles, Beijing 100081, China

⁴Beijing Municipal Research Institute of Environmental Protection, Beijing 100037, China

⁵Environmental Meteorological Center of China Meteorological Administration, Beijing 100081, China

Correspondence to: Ying Zhou (y.zhou@bjut.edu.cn) and Shuiyuan Cheng (chengsy@bjut.edu.cn)

Abstract. Biomass burning injects many different gases and aerosols into the atmosphere, which could have a harmful effect on air quality, climate change and human health. In this study, a comprehensive biomass burning emission inventory including crop straw domestic combustion and in-field crop residue burning, firewood and livestock excrement combustion, forest and grassland fire was developed for mainland China in 2012 based on county-level activity data, satellite data, and updated source-specific emission factors (EFs). The emission inventory within 1 × 1 km grid was generated using geographical information system (GIS) technology according to source-based spatial surrogates. A range of key information related to emission estimation (e.g., province-specific proportion of crop straw domestic ~~burning-combustion~~ and in-field crop residue burning~~open burning~~, detailed firewood combustion quantities, uneven temporal distribution coefficient) was obtained from field investigation, systematic combing of the latest research and regression analysis of statistical data. The established emission inventory includes the major precursors of complex pollution, greenhouse gases and heavy metal released from biomass burning. The results show that the emissions of SO₂, NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, NH₃, CO, EC, OC, CO₂, CH₄ and Hg in 2012 ~~were 332.8 Gg, 972.5 Gg, 3676.0 Gg, 3479.4 Gg, 3429.6 Gg, 395.8 Gg, 33987.9 Gg, 367.1 Gg, 1151.7 Gg, 665989.0 Gg, 2076.5 Gg and 3.65 Mg, respectively. Indoor and outdoor burning of straw are 336.8 Gg, 990.7 Gg, 3728.3 Gg, 3526.7 Gg, 3474.2 Gg, 401.2 Gg, 34380.4 Gg, 369.7 Gg, 1189.5 Gg, 675299.0 Gg, 2092.4 Gg and 4.12 Mg, respectively. Straw domestic burning, in-field crop residue burning,~~ and firewood combustion are identified as the dominant biomass burning sources. The largest contributing source is different for various pollutants. Straw ~~indoor-domestic~~ burning is the major source of SO₂, CO, CH₄ and ~~Hg-NMVOC~~ emission; firewood contributes most to EC and NH₃ emission. ~~Corn, rice and wheat represent the major crop straws, with their total emission contribution exceeding 80% for each pollutant. Corn straw burning has the greatest contribution to EC, NO_x and SO₂ emissions; rice straw burning is dominant contributor to CO₂, VOC, CH₄ and NH₃ emissions.~~ Corn, rice and wheat represent the major crop straws. The combined emissions of these three straw types account for 80% of the total straw burned emissions for each specific pollutants mentioned in this study. As for the straw burning

emission of various crops, corn straw burning has the largest contribution to EC, NO_x and SO₂ emissions; rice straw burning has higher contribution to CO₂, NMVOC, CH₄ and NH₃ emissions; wheat straw burning has higher contribution to CO and Hg emissions. Heilongjiang, Shandong, and Henan provinces located in northeast and central-south region of China have higher emissions. Gridded emissions, which were obtained through spatial allocation based on the gridded rural population and fire point data from emission inventory at county resolution, could better represent the actual situation. Higher biomass burning emissions are concentrated in the areas with greater agricultural and rural activity. The temporal distribution shows that April, May, June and October are the top four months with higher emissions, due to the in-field crop residue burning. While as for EC, the emission in February, January, October and December are relatively higher due to the biomass domestic burning in heating season. ~~The temporal distribution shows that higher emissions occurred in April, September, and October during the whole year.~~ There's regional difference in monthly variation due to the diversity of main planted crop and the climate conditions. Furthermore, PM_{2.5} component results showed that OC, Cl⁻, EC, K⁺, NH₄⁺, K element and SO₄²⁻ are the main PM_{2.5} species accounting for 80% of the total emissions. The species with relatively higher contribution to ~~VOCNMVOCs~~ emission include ~~ing~~ ethylene, propylene, toluene, ~~mp-xylene and and ethyl benzene, halocarbons~~ which are key species for the formation of secondary air pollution. The detailed biomass burning emission inventory generated by this study could provide useful information for air quality modelling and support the development of appropriate pollution control strategies.

Keywords: Biomass burning; Emission inventory; High resolution; Species

1 Introduction

Biomass burning is considered a significant source of gas and particulate matter (PM), resulting in a major impact on atmospheric chemistry, climate change and human health. Active trace gases (e.g., SO₂, NO_x, ~~VOCNMVOCs~~, NH₃) released from biomass burning are the major precursors of secondary inorganic/organic aerosols and tropospheric ozone (O₃) in the atmosphere (Penner et al., 1992; Kaufman and Fraser, 1997; Koppmann et al., 2005; Langmann et al., 2009). Several studies have indicated that observed local and regional air pollution could be attributed to the chemical species emitted from biomass burning (Huang et al., 2012b; Zha et al., 2013; Cheng et al., 2014; Yan et al., 2014; Zong et al., 2016). The emission factor (EF) of some biomass burning pollutants is even greater than coal burning, which is widely recognized as a major pollution source (Zheng et al., 2009; Fu et al., 2013). Primary particles (e.g., BC and OC) discharged by biomass burning not only impact visibility, but also have an influence on climate change due to the positive effects of the absorption of light and cloud condensation (IPCC, 2011). Biomass burning is also a ~~critical-significant~~ source of greenhouse gases such as methane (CH₄) and carbon dioxide (CO₂) (Andreae and Merlet, 2001), which contribute to global warming (Sun et al., 2016). Moreover, several reports (Fernandez et al., 2001; Huang et al., 2012b; Shi and Yamaguchi, 2014) reveal that the long-term or short-term exposure to PM (e.g., BC emitted from indoor biomass burning) can cause adverse effects to human health, such as

decreased lung function, increased respiratory diseases and lung cancer mortality. Furthermore, studies have identified that indoor biomass burning could bring adverse health effects on residents (Jiang and Bell, 2008; Fullerton et al., 2008).

Prior to its rapid economic development, China was a largely agricultural country and thus once consumed a large amount of biofuels (e.g., crop residues and firewood). With the dramatic urbanization that accompanied the economic development, the pattern of energy consumption in rural areas has been gradually transformed. In particular, in some agricultural areas with relatively high income, crop residues were more burned directly in the field (Sun et al., 2016). Beginning in 1999, the Chinese government has issued a series of laws and regulations to ban the ~~open-in-field~~ burning of straw ~~residues~~ and to encourage straw comprehensive utilization, such as returning to field, livestock feeding, industrial raw materials manufacturing, briquette fuel processing, etc. (MEP, 1999). However, the effect of this legislation was not satisfactory because the processes of straw comprehensive utilization not only required high ~~labor~~labour costs but also delayed sowing of the next crop. Thus, the phenomenon of straw ~~outdoor-in-field~~ burning continued to occur. ~~The amount of straw outdoor burning in China in 2009 is 0.215 billion tons (MA, 2014) The amount of in-field crop residue burning in China in 2009 was estimated as 0.215 billion Mg. The data is obtained from the governmental report on the investigation and evaluation of crop straw resources in various provinces in China (MA, 2011).~~ Accordingly, a comprehensive and detailed emission inventory of biomass burning representing the current status in China, is important to provide valuable information for researchers and policymakers. Examples of potential applications include research to understand the influence of biomass burning on indoor air quality and the outdoor atmospheric environment, and the development of effective management decisions to relieve the environmental burden and reduce health risk.

Since the early research conducted by Crutzen et al. (1979), a series of efforts have been made to develop a biomass burning emission inventory, especially in developed countries (Reddy and Venkataraman, 2002; Ito and Penner, 2004; van der Werf et al., 2006; Nelson et al., 2012; Shon, 2015). Compared with the developed countries, research by Chinese scientists on this issue started relatively late. The initial studies on biomass burning emission inventory across China (Streets et al., 2001; Tian et al., 2002; Streets et al., 2003; Cao et al., 2005) or in certain regions (Zheng et al., 2009; Huang et al., 2011) were developed mainly based on EFs developed for foreign nations (Turn et al., 1997; Andreae and Merlet, 2001; U.S. EPA, 2002) because of the lack of local measurements in China. However, this approach could introduce relative great uncertainty in emission estimates because of the differences in crop types and the combustion conditions between China and other countries.

In recent years, various research activities have focused on the emission characteristics of biomass burning in China, including local EF and chemical species profile tests. Li et al. (2007b) and Li et al. (2009) conducted field measurements to determine the EF for several of the main household biofuels in Beijing, Chongqing, Henan and Shandong. Li et al. (2007c) determined the EF for wheat and maize burning in field and Cao et al. (2008) measured EFs for the ~~indoor-domestic~~ burning of rice straw, wheat straw, corn ~~strawstovers~~ and cotton stalks. Zhang et al. (2008) measured CO₂, CO, NO, NO₂, NO_x and PM EFs of rice, wheat and corn straw and Wang et al. (2009) launched a study on characteristics of gaseous

pollutants from biofuel stoves in China. More recently, Zhang et al. (2013b) carried out experiments on EFs for ~~open-in-field~~ burning of sugar cane leaves and rice straw in southeast China. Ni et al. (2015) conducted laboratory burn tests to determine the EFs of wheat straw, rice straw and corn stalks, considering the impacts of the fuel moisture content.

Based on the local EFs, emission inventories that focused on certain provinces (Li et al., 2015; He et al., 2015) or city group regions (He et al., 2011; Fu et al., 2013) were developed. In our previous study, we reported an emission inventory with high resolution in the Beijing–Tianjin–Hebei region of China (Zhou et al., 2015). To produce a national emission inventory, several studies of biomass burning have been carried out without distinguishing the detailed crop straws (Lu et al., 2011; Yan et al., 2006; Tian et al., 2011). Moreover, there are several studies that have focused on certain pollutants (Huang et al., 2012d; Chen et al., 2013; Zhang et al., 2013a; Kang et al., 2016; Li et al., 2016), and certain crop straws (Zhang et al., 2008; Hong, et al., 2016; Sun, et al., 2016). In recent years, the comprehensive biomass emission inventory is limited. Most of recent studies are concentrated upon biomass open burning, including the multi-year trend analysis on certain or multiple pollutants (Wang and Zhang, 2008; Song et al., 2009; ~~Huang et al., 2012c~~; Shi et al., 2014; Shon, 2015; Xu et al., 2016; Zhang et al., 2016). Few studies have covered recent firewood burning (see next paragraph for details regarding the reason for this). In addition to the EF, detailed activity data are also important for a reliable emission inventory, such as straw domestic or in-field crop residue burning ratios, which are not currently publicly available. Gao et al. (2002) produced a study on the percentage of straw used as fuel and for direct incineration in 2000. Wang et al. (2008) investigated the percentage of in-field crop residue ~~crop open~~ burning in 2006 of six regions in China, which were divided according to the similarities of agriculture, climate, economy and region. Tian et al. (2011) estimated the proportion of crop straw domestic burning and in-field crop residue ~~open~~ burning in 2007 for seven and three regions of China, respectively. Thus, there is limited information about the ratio of straw used as fuel ~~and to that crop residue burning~~ in the field that reflects the status of China in recent years for different provinces. ~~Moreover, because of the lack of firewood consumption in the energy statistical yearbook after 2007, there are few reports containing a comprehensive biomass burning emission inventory for China.~~ Moreover, because of the lack of firewood consumption record in the China Energy Statistical Yearbook (NBSC, 2009-2015), few studies have developed a comprehensive biomass burning emission inventory in China in recent years. China Energy Statistical Yearbook provides official information on the energy construction, production and consumption, including the detailed firewood consumption in various regions. However, the firewood consumption data is no longer contained in the NBSC (2009-2015) since 2008, as a result, there are few literature containing a comprehensive biomass burning emission inventory for China.

Consequently, we have identified several weaknesses in the current biomass burning emission inventories. First, not all biomass burning sources have been included in recent years, especially since 2008, because of the lack of firewood consumption data in the various statistical yearbooks (e.g. China Energy Statistical Yearbook, China statistical yearbook, China rural statistical yearbook). Second, the source-specific EFs used in emission estimation need to be updated based on the systematic combing of local tests in the latest research. Third, the proportion of crop straw

domestic burning and in-field crop residue burning, which could reflect the recent conditions of different provinces in China needs to be investigated.
Fourth, the current biomass burning emission inventory for China is generally at province resolution because detailed activity data cannot be directly
obtained from the various statistical yearbooks in China.~~First, not all biomass burning sources have been included in recent years, especially after~~
~~2007, because of the lack of firewood consumption data in the statistical yearbook. Second, the source-specific EFs used in emission estimation~~
~~need to be updated based on the systematic combing of local tests in the latest research. Third, the proportion of crop straw domestic burning and~~
~~open burning, which could reflect the recent situation of different provinces in China needs to be investigated. Fourth, the current biomass burning~~
~~emission inventory for China is generally at province resolution because detailed activity data cannot be directly obtained from the yearbook.~~

Activity data at coarse resolution are likely to be associated with greater uncertainty in grid emissions generated according to source-based gridded spatial surrogates (e.g., population) using GIS technology (Zheng et al., 2014). As a result, it is of great importance to develop an integrated and model-ready biomass burning emission inventory with high spatial and temporal resolution.

In this study, a comprehensive biomass burning emission inventory including crop straw domestic combustion and in-field crop residue burning, firewood and livestock excrement combustion, forest and grassland fire was developed for the Chinese mainland (excluding Hong Kong, Macao, and Taiwan) in 2012, based on detailed activity data and satellite burned area data. In addition, we attempt to take full account of the source-specific EFs measured in China. A range of important information for emissions estimation (e.g., province-specific straw domestic combustion/in-field crop residue burning ratio, detailed firewood combustion quantities and uneven temporal distribution coefficient) were obtained from a field investigation, systematic combing of latest research and regression analysis of statistical data. A 1-km resolution emission inventory was generated using GIS software. The gaseous and particulate pollutants examined in this research included SO₂, NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, NH₃, CO, EC, OC, CO₂, CH₄ and Hg, covering the major precursors of complex pollution, greenhouse gases and heavy metals released from biomass burning. The detailed emission inventory given by this paper could provide valuable information to support the further biomass burning pollution research and the development of a targeted control strategy of all regions across the Chinese mainland.

The remainder of this paper is structured as follows. Section 2 describes the methodology including the emission estimation method, the selection and handling of activity data and corresponding parameters, determination of EFs, spatial and temporal allocation, ~~and~~ speciation of PM_{2.5} and ~~VOENMVOC~~s. Section 3.1 describes the total emission in China, and the contribution of various biomass burning sources and crop straws. Section 3.2 describes the emission from different regions, ~~and~~ contributions of different biomass sources and crop straws of each province. Spatial and temporal distribution of biomass burning emissions is discussed in Secs. 3.3 and 3.4, respectively. Section 3.5 presents the emissions of PM_{2.5} and ~~VOENMVOC~~ species. Uncertainty in biomass burning emission estimates is described in Section 3.6. The comparison between this study and other studies appears in Section 3.7. Section 4 summarizes the conclusions.

2 Methodology

2.1 General description

The biomass burning considered in this study is mainly divided into two categories, domestic combustion and open burning. Domestic combustion mainly involves crop straw, firewood and livestock excrement (mainly used in pastoral and semi-pastoral areas) burning. Open burning includes in-field crop residue burning, forest and grassland fire. Details of the source classifications are shown in Table 1. ~~The biomass burning considered in this study is mainly divided into two categories including domestic combustion and in field burning. Domestic combustion mainly involves crop straw, firewood and livestock excrement (mainly used in pastoral and semi-pastoral areas). In field burning includes seasonal crop residue waste burning, grassland and forest fires. Details of the source classifications are shown in Table 1.~~

A bottom-up approach was used to develop the biomass burning emission inventory for all districts or counties. The annual biomass burning emissions (E_i) were calculated using Eq. (1) as follows:

$$E_i = \sum (A_i \times EF_{i,j}) \times 1000, \quad (1)$$

where subscripts i and j represent the type of pollutant and biomass burning source; E is the annual typical pollutant emission (Mg/yr); A is annual amount of dry biomass burned (Mg/yr), for which the detailed calculation method is shown in Sec. 2.2; and EF is the emission factor (g/kg), for which a detailed description is presented in Sec. 2.3.

2.2 Activity data

2.2.1 Straw burning

The burning mass of straw ~~indoor-domestic burning~~ and ~~outdoor-in-field crop residue~~ burning can be calculated using Eq. (2) as follows:

$$A_{i,k} = P_{i,k} \times N_k \times R_{i,k} \times D_k \times CE_k, \quad (2)$$

where subscripts i and k represent region (district or county) and crop type, respectively; $A_{i,k}$ is the annual burning mass of crop straw (Mg/yr); $P_{i,k}$ is the amount of crop-specific yields per year (Mg/yr); N_k is the residue-to-production ratio of each straw type (ton/ton); $R_{i,k}$ is percentage of crop straw burned as fuel or in field burning; D_k is dry matter fraction of each straw type; and CE_k is the combustion efficiency of each straw type.

There are currently no statistics on the amount of each crop yield at the county resolution ($P_{i,k}$) in ~~the statistical various~~ yearbooks in China. Therefore, in this study, we conducted a correlation analysis between grain yield and crop yield at prefecture resolution, and found a good correlation ($R = 0.747$, detailed analysis is provided in the Supplement, Fig. S1). The grain yield at prefecture resolution was summarized from China Statistical

Yearbook in 2012 (NBSC, 2013b). The crop yield at prefecture resolution was summarized from statistical yearbooks edited by National Bureau of Statistics in 2012 for each province. Next, the $P_{i,k}$ was calculated based on the various types of crop yield at prefecture resolution and grain yield at county resolution, which was summarized from a range of statistical yearbooks in 2012 for each province (including the city yearbooks which are publically available for some cities), NBSC (2013a) and NBSC (2013b). The total straw amount of China in 2012 calculated in this study is 832.5 Tg, which is similar to NDRC (2014) (817.4 Tg). Next, the $P_{i,k}$ at county level was calculated based on the various types of crop yield at prefecture resolution and grain yield at county resolution. Grain yield at county resolution was summarized from a range of statistical yearbooks edited by National Bureau of Statistics in 2012 for each province and city, NBSC (2013a) and NBSC (2013b). The total straw amount of China in 2012 calculated in this study is 832.5 Tg, which is similar to the data of Chinese governmental annual statistical reports about the straw utilization and burning (NDRC, 2014; the amount of straw can be collected is 817.4 Tg). The map at prefecture and county resolution is shown in Fig. S2 in the Supplement.

The variable $R_{i,k}$ is important for biomass burning emission estimation, and the information that can representing the recent status in China needs to be updated because of the continued economic development and the gradual implementation of national control policies for straw residue open in-field crop residue burning. In this study, we conducted a detailed investigation of recent literature to collect the percentage of crop straw burned as domestic fuel and burned as waste for each province. For some provinces where the current reporting is limited (e.g., Heilongjiang, Zhejiang, Guangdong, Inner Mongolia, and Hebei), a questionnaire survey was launched. Details of the questionnaire survey are presented in the Supplement (S3). The percentage of crop straw indoor-domestic burning and outdoor-in-field crop residue burning for each province is summarized in Table 2. According to our estimation, the amount of straw indoor-domestic and outdoor-in-field burning for China in 2012 was 0.26 billion tons-Mg and 0.19 billion tons-Mg, respectively, which is similar to other recently published results for 2012 (0.26 billion tons-Mg indoor-domestic burning, Tian et al., 2014) and 2009 (0.215 billion tons-Mg outdoor-in-field crop residue burning, MA, 2011).

The N_k , D_k and CE_k values were obtained according to the literature collection. Detailed parameters used in this study are summarized in Table 3.

2.2.2 Firewood

Firewood consumption is recorded as non-commodity energy in the China energy statistical yearbook. However, detailed firewood consumption has not been publicly available since 2008⁷. For more recent years, we obtained the total firewood consumption for China in 2012 and for each province in 2010 (Tian et al., 2014; IEA, 2012). However, these data could not support the development of an emission inventory at high resolution. There are several detailed statistics available in the yearbook, such as the rural population, gross agricultural output and timber yield, which are likely to have a relationship with the firewood consumption. Therefore, we produced a correlation analysis between the three statistics and the

firewood consumption of each province for different years in which the firewood consumption data were available at province resolution, as shown in Fig. 1. The best correlation relationship was found between rural population and firewood consumption. The correlation coefficient for the different years ranged from 0.66 to 0.82, therefore, we choose rural population as the surrogate to calculate the detailed firewood consumption. The firewood consumption at county resolution was obtained based on the rural population at county resolution and the total firewood consumption reported by Tian et al. (2014) and IEA (2012). China's rural population, gross agriculture output and timber yield of each province come from NBSC (1999-2008a). Firewood consumption comes from NBSC (1999-2008b).

2.2.3 ~~Biomass burning of forest/grassland fires~~ Forest and grassland burning

The burning mass of forest/grassland can be calculated from the annual mass of forest/grassland burned (Mg/yr) as Eq. (3):~~The burning mass of forest/grassland can be calculated using Eq. (3) as follows:~~

$$A = (\sum_{j=1}^{10} BA_{x,j} \times FL_{x,j} \times CF_j) \times 10^{-6}, \quad (3)$$

where subscripts j, and x represent the land cover type, and location, respectively, $BA_{x,j}$ is the burned area (m^2) of land cover type j at x, $FL_{x,j}$ is the biomass fuel loading (the aboveground biomass density in this study; g/m^2) of land cover type j at x, and CF_j is the combustion factor (the fraction of burned aboveground biomass) of land cover type j.

Burned area data for 2012 were derived from the moderate-resolution imaging spectroradiometer (MODIS) direct broadcast burned area product (MCD64A1; <http://modis-fire.umd.edu>). This product employs an automated algorithm for mapping MODIS post-fire burned areas, and deriving the approximate burn date within each burn cell combined with surface reflectance, land cover products, and daily active fires. The MCD64A1 product has a primary spatial resolution of 500 m. The daily burned areas could be obtained from the product.

Earlier research on the estimation of FL values for forest and grassland typically employed an averaged value of aboveground biomass density. However, these values do not well reflect the spatial variations of FL for each vegetation type. In this study, numerous local FL were collected for each province and vegetation type. The type of vegetation burned in each pixel was determined by the 1 km resolution MODIS Land Cover product produced by Ran et al. (2010). We considered 10 vegetation types as forest and grassland (i.e., evergreen needleleaf forest, evergreen broadleaf forest, deciduous needleleaf forest, deciduous broadleaf forest, mixed forest, closed shrublands, open shrublands, woody savannas, savannas, and grassland). The values of FL employed in this study are listed in Table 4. As for CF, it has usually been set as a constant in previous literature. In our paper, CF values were collected for each vegetation type, and the CF in each pixel was determined by the MODIS Land Cover product and the CF of typical vegetation. The CF of forest, closed shrublands, open shrublands, woody savannas, and grassland were set as 0.25, 0.5, 0.85, 0.4, and 0.95, respectively (Michel et al., 2005; Kasischke et al., 2000; Hurst et al., 1994).

$$A = AR \times B \times \eta, \quad (3)$$

where A is the annual burning amount of forest/grassland (ton/a); AR is the damaged area of grassland or forest fire per year (hm^2/a); B represents the dry biomass density of grassland or forest (ton/hm^2); and η is the combustion efficiency of grassland or forest.

The damaged area of forest fire and grassland fire could be obtained from NBSC (2013c) and NBSC (2013d), respectively. The specific locations at which fire occurred can be determined according to Moderate Resolution Imaging Spectroradiometer (MODIS) fire point data; the B and η of forest is different from the climatic zones, which is shown in Table 4; B for grassland is 1800 kg/ha (Lu et al., 2011), and η for grassland is 80.0% according to the 'Guide for the Preparation of Atmospheric Pollutant Emission Inventories for Biomass Burning' (EPD, 2014).

2.2.4 Livestock manure

The mass of biomass burned by animal waste was calculated using Eq. (4) as follows:

$$A = S \times Y \times C \times R, \quad (4)$$

where, A is the annual discharge of livestock manure burned (tonMg/yra); S represents the amount of each livestock type in pastoral and semi-pastoral land at the end of the year (head/yra); Y is a single livestock annual fecal output per year (tonMg/head); C represents livestock manure dry matter fraction; and R is the proportion of total livestock manure directly combusted.

The S values were taken from the [China governmental annual statistical reports, including EOCAIY \(2013\)](#) and [NBSC \(2013c\)](#). The Y values were related to the large animals only. Among these, single cattle annual manure output was 10 tons-Mg and single horse annual manure output was 7.3 tons-Mg (Li and Zhao, 2008). The livestock annual manure output of other animals was set at 8 tonsMg , according to [Tian et al. \(2011\)](#). The C value was set as 18% ([Tian et al., 2011](#)) and R was 20% ([Li, 2007a](#); [Liu and Shen, 2007](#)). Since not all regions use livestock manure in biomass burning, we consider only the pastoral and semi-pastoral areas including Tibet, Inner Mongolia, Gansu, Xinjiang, Qinghai province in this study ([Tian et al., 2011](#)).

2.3 Determination of EFs

In order to ensure the accuracy of the emission inventory as much as possible, it is important to choose the appropriate EF. The EFs used in this study were mainly based on localized measurements. When selecting the EFs, we applied the following principles: first, for a certain type of biomass source or crop type, we prioritized the use of localized measured EFs from the literature. Second, for the biomass sources or crop types which lacked localized measurements, we prioritized results from developing foreign countries similar to our country above those of developed countries. Third, when localized measured data of a certain crop type were missing, the average value of the mainstream literature in the foreign country was

used as an estimate. After extensive literature research on EFs, the resultant EFs for ~~indoor-domestic~~ and ~~outdoor-open~~ biomass burning for each chemical species and each source are summarized in Tables 5 and 6, respectively.

2.4 Spatial distribution

In order to obtain the detailed spatial distribution characteristics of biomass emission, and to provide grid based data for the air quality model simulation, the biomass burning inventory in this study ~~is~~ assigned into 1×1 km grid cells based on the source-specific surrogate. We applied GIS software as the main tool to produce the spatial distribution. In this paper, the approaches used to determine spatial distribution varied between biomass sources; thus, we selected different methods of spatial allocation according to the homologous source characteristics. ~~The regions in which open biomass burning occurred can be located according to the MODIS fire counts data (van der Werf et al., 2006; Liu et al., 2015). Detailed description about the MODIS fire data are shown in Supplement (S4). The outdoor biomass sources (e.g. outdoor straw burning, forest fire, and grassland fire) were treated as point sources that were located based on MODIS fire data, and land use data provided by Ran et al. (2010). Farmland fire point, forest fire point and grassland fire point are the spatial surrogates of outdoor straw burning, forest fire and grassland fire, respectively. The regions in which in-field crop residue burning occurred can be located according to the MODIS fire counts data (MOD14/MYD14) (van der Werf et al., 2006; Huang et al., 2012e). Farmland fire point is the spatial surrogates of in-field crop residue burning, land use data (MODIS Land cover) is provided by Ran et al. (2010). Detailed description about the MODIS fire counts data (MOD14/MYD14) are shown in Supplement (S4). As for forest and grassland fire, the emission of forest and grassland fire are estimated in 500 resolution, it can be resampled into 1km grid using GIS software.~~ The emissions of straw, firewood, and livestock excrement combustion were treated as area sources and the spatial surrogates used to distribute these biomass sources were population density of different land use types (e.g. rural population density, grassland population density) (Zheng et al., 2009; Huang et al., 2012c). The population density of different land use types is according to the land use data provided by Ran et al. (2010) and 1 km grid population distribution data provided by Fu et al. (2014). Detailed calculation method and equation of gridded emission are presented in Supplement (S4).

2.5 Temporal distribution

According to the temporal resolution of MODIS fire counts data (MOD14/MYD14), ~~the monthly/daily emission of outdoor straw burning, forest fire and grassland fire emission can be estimated based on the number of typical fire points the monthly/daily emission of in-field crop residue burning can be estimated based on the number of typical fire points, the monthly/daily emission of forest and grassland fire emission can be calculated by the Julday emission of forest and grassland fire.~~ For ~~indoor-domestic~~ biomass source, the monthly uneven coefficient was mainly

derived from our survey questionnaire. Details of the questionnaire survey are presented in the Supplement (S3). The daily ~~indoor-domestic~~ emission is equally allocated from the monthly emission.

2.6 Speciation of ~~VOC~~NMVOCs and PM_{2.5}

The detailed species emission of ~~VOC~~NMVOCs and PM_{2.5} is necessary information of model simulation for different chemical mechanism selection (e.g., CB05). The speciation of ~~VOC~~NMVOCs and PM_{2.5} is the main research object of the chemical composition of the atmospheric emission source, which has received extensive attention by domestic scholars in recent years (Song et al., 2007; Li et al., 2007c; Liu et al., 2008).

In this study, the species emission was mainly estimated based on the total emission, and ~~VOC~~NMVOC and PM_{2.5} source profiles (mass fraction) of biomass sources collected from literature review. In terms of the data selection, we prioritized domestic measurement with the species as much as possible. Therefore, the ~~VOC~~NMVOCs source profile mainly refers to data from Liu et al. (2008) and Akagi et al. (2011), including ~~94~~ species covering alkane, alkene, alkyne, ~~benzene-series-compounds~~aromatic and so on; the PM_{2.5} source profile data is cited from the work of Li et al. (2007c) and Watson et al. (2001), including 36 species, such as element, ion and so on.

3 Results and discussion

3.1 Total emissions in China

3.1.1 Contributions by biomass burning sources

The annual emissions of biomass burning in mainland China are presented in Table 7; The total annual emissions of SO₂, NO_x, PM₁₀, PM_{2.5}, ~~VOC~~NMVOC, NH₃, CO, EC, OC, CO₂, CH₄ and Hg for Chinese mainland in 2012 are ~~336.8 Gg, 990.7 Gg, 3728.3 Gg, 3526.7 Gg, 3474.2 Gg, 401.2 Gg, 34380.4 Gg, 369.7 Gg, 1189.5 Gg, 675299.0 Gg, 2092.4 Gg and 4.12 Mg~~332.8 Gg, 972.5 Gg, 3676.0 Gg, 3479.4 Gg, 3429.6 Gg, 395.8 Gg, 33987.9 Gg, 367.1 Gg, 1151.7 Gg, 665989.0 Gg, 2076.5 Gg and 3.65 Mg, respectively. The contribution of different sources to the total emissions ~~of various pollutantsto various chemical species~~ is shown in Fig. 2. It shows that the straw ~~indoor-domestic burning, outdoor-in-field crop residue~~ burning and firewood combustion are the dominant biomass burning sources with the total contribution ranging from ~~95.986.02%~~ to ~~99.497.58%~~ for various species. However, the largest contributing sources to different species are not similar. Compared with other sources, straw ~~indoor-domestic~~ burning contributed most to SO₂, CO, CH₄ and ~~Hg~~NMVOC, accounting for ~~58.557.8%, 58.758.1%, 53.653.2%~~ and ~~42.249.2%~~ of total emissions, respectively. Straw ~~indoor-domestic~~ burning has a direct impact on residents and the prolonged exposure under high ~~indoor-domestic~~ biomass burning emission (e.g., SO₂, CO, CH₄ and Hg) can cause many adverse health effects (e.g. acute respiratory infections and chronic bronchitis) (Emily and Martin, 2008). The contribution of firewood to each species cannot be neglected, especially for EC (~~51.651.3%~~) and NH₃

(41.841.2%). According to the localized measurement of EF by Li et al. (2009), the average EC EF for firewood (1.49 g/kg) is 3.5 times of crop residue (0.43 g/kg). EF of firewood NH₃ is larger than the average of various straws. This results in a large contribution by firewood for these two species. The contribution of straw ~~indoor-domestic burning~~ and ~~outdoor-in-field crop residue~~ burning to NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, Hg, OC and CO₂ is nearly equal. Straw burning has an important influence on indoor air quality and ~~the~~ outdoor atmospheric environment.

In addition to the sources mentioned above, the contribution of livestock excrement burning, forest and grass~~land~~ fires is relatively small. It is mainly due to the small amount of biomass fuel consumption. The biomass fuel consumption of these three biomass sources are 10614Gg, 6647Gg and 505 Gg, respectively, which is significantly lower than that of straw domestic combustion (201582 Gg), in-field crop residue burning (147178 Gg) and firewood combustion (127250 Gg). The contribution of livestock excrement burning to PM₁₀, PM_{2.5}, NH₃, EC, OC, CO₂ and CH₄ is 2.55%, 2.51%, 3.49%, 1.53%, 2.03%, 1.69% and 2.12%, respectively. The contribution of forest and grass fires to biomass burning emissions to most chemical species in China is negligible (0.19–0.66%), except for the contribution of forest fire to Hg emissions (2.65%). The contribution of livestock excrement burning to PM₁₀, PM_{2.5}, NH₃, EC, OC, CO₂ and CH₄ is 2.52%, 2.47%, 3.44%, 1.52%, 1.96%, 1.67% and 2.10%, respectively. The contribution of forest and grassland fires to biomass burning emissions to most chemical species in China is small (0.9–3.7%), except for the contribution of forest fire to Hg emissions (14.0%).

3.1.2 Contributions by various crop straw

As mentioned in Section 3.1.1, straw burning is the important biomass burning source with considerable influence on the chemical species that most strongly impact the air quality, climate change and human health. Furthermore, the major crop straw type contribution was analysed. Figure 3 shows the contributions of 12 different types of crop straw ~~indoor-domestic burning~~ and ~~outdoor-in-field crop residue~~ burning to total straws burning emissions for various species to various chemical species in 2012 from the perspective of the ~~whole country-mainland China~~. Figure 3c indicates that corn, rice and wheat straw are the major crops straw burned as fuel or as waste in China. The contribution is more than 80% to the total straw burned emissions of all pollutants studied in this paper. Corn, rice and wheat are the three major food crops in China with large planting area (the output of these three kinds of grain accounts for 70% of the total grain output in China, NBSC, 2013c), resulting in a large amount of straw production. Among the various crops, corn straw burning has ~~the greatest~~large contribution to all of the chemical species except for CH₄. Rice straw ~~is major contributor~~ has the largest contribution to CO₂, ~~VOENMVOC~~, CH₄ and NH₃ emissions, accounting for 32.90%, 32.43%, 31.61% and 30.12%, respectively; wheat straw has a ~~a~~ considerable contribution to Hg, SO₂ and OC emissions, accounting for 29.46%, 26.47% and 25.91%, respectively. Compared with the three kinds of crop mentioned above, the total contribution of soybean, cotton, sugar cane, potato, peanut and rape to the various chemical species is relatively small, accounting for 8.1–19.2% of the total emissions s for all pollutants; the contribution of sesame,

sugar beet and hemp burning to various chemical species emission is negligible, never exceeding 0.5%. In addition, Fig. 3a and Fig. 3b indicate that for most of the chemical species, the contribution of ~~corn straw outdoor in-field~~ corn residue burning is larger than that of ~~indoor domestic~~ burning, except for SO₂, EC and CO₂. Contrary to that for corn straw, emissions of all chemical species (except for SO₂, NO_x and EC) from wheat straw ~~indoor domestic~~ burning is greater than those from ~~outdoor in-field~~ crop residue burning. For rice straw, the contribution of ~~outdoor in-field~~ crop residue burning to NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, EC and OC emissions is larger than ~~indoor domestic~~ burning.

3.2 Emissions from different regions

3.2.1 Total emissions for different provinces

The total biomass burning emissions in ~~the~~ 31 provinces in 2012 are presented in Table 7. These results indicate that Heilongjiang, Shandong, Henan, Hubei, Anhui, Sichuan, Jilin, Inner Mongolia, Hunan and Jiangsu province are the major contributors, with the total emission contributions of ~~various pollutant emissions between~~ ranging from ~~5453~~ to ~~and 6665~~ % for ~~various a species~~ pollutants. The province with most contribution to total emission of NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, NH₃, OC, CH₄, Hg and CO₂ is Heilongjiang; while Shandong province has the highest emission of SO₂, CO and, ~~EC and Hg~~. It could be attributed to different types of biomass consumption in each province due to geographical location, climate conditions and population density. Detailed discussion about the contribution by biomass source and crop straw type of different regions is shown below.

3.2.2 Contributions by biomass sources of each province

The emission of detailed biomass sources of each province is presented in Fig. 4. The provinces with major contribution to total pollutants emissions for each biomass source are various. Straw burning emissions mainly distributed in Shandong, Henan, Heilongjiang, Hebei, Anhui, Sichuan, Jilin and Hunan province. The total contribution of these provinces to various pollutants is more than 58%. It is due to the large amount of cultivated land in the north plain region as cultivated land in this region prioritizes economic crops that produce rich straw resources. Several regions in which firewood produce a large emission are Hunan, Yunnan, Hubei, Hebei, Sichuan, Guangdong, Shaanxi, Liaoning and Jiangxi province. More than 54% firewood combustion emission is contributed by these provinces. These areas are mainly distributed in the south of China, a mountainous region in which the forest cover is higher than 30% (NBSC, 2013c). Livestock excrement combustion emissions mainly distributed in Tibet, Inner Mongolia, Gansu, Xinjiang, and Qinghai province, since only pastoral and semi-pastoral areas burn livestock manure as fuel in China. Emissions from forest and grassland fires are mainly distributed in Tibet, Yunnan, ~~Zhejiang~~ Heilongjiang, Xinjiang, Inner Mongolia and Sichuan province. This is owing to the high vegetation cover and climatic conditions in these areas.

The contribution of biomass sources to total emissions in each province is also distinct. Straw burning has a large contribution to various pollutants emissions in Heilongjiang (79–~~95~~~~97~~%), Ningxia (~~84~~~~87~~–98%), Shandong (~~73~~~~74~~–95%), Jilin (~~73~~~~74~~–95%), Henan (~~60~~~~61~~–~~94~~~~93~~%), Anhui (~~50~~~~51~~–91%) and ~~Sichuan-Shanxi~~ (~~57~~~~61~~–~~89~~~~90~~%) province. The economic income of the rural areas in these provinces is relatively low. A large number of crop residues ~~is are~~ consumed as main non-commodity energy. In addition, firewood resources ~~is are~~ scarce in these areas and as a result, the usage of straw is very high. Figure 4 also indicates that, for most provinces (e.g. Beijing, Tianjin, and Hebei), the contribution of the ~~indoor domestic~~ straw burning is greater than ~~outdoor straw in-field crop residue~~ burning. This is mainly attributable to the gradual response to the prohibition of burning straw and the introduction of straw resource utilization measures. The emission contribution of ~~straw burned in-field crop residue burnings~~ is higher than that of straw domestic burning in Hebei, Heilongjiang and Anhui province. It suggests that the prohibition of burning straw measures in these provinces still needs to be strengthened. Several regions in which firewood produce a large component of total emissions of various pollutants are Beijing (~~56~~~~47~~–~~76~~~~90~~%), Guangdong (31–83%), Yunnan (~~37~~~~31~~–~~84~~~~79~~%), Fujian (~~29~~~~30~~–~~80~~~~81~~%), Hainan (26–~~64~~~~77~~%) and Guizhou (27–74%) province. The straw amounts in the rural areas of these provinces are relative low. Firewood is the mainly non-commodity energy used by rural people. It is worth noting that though the biomass fuel consumption in Beijing is small, ~~the firewood is main biofuel due to the server restriction of straw open burning.~~ Compared to straw burning emission contribution (9%–~~45~~~~41~~%), firewood emission (~~55~~~~47~~%–90%) represents a large proportion of the total biomass burning in Beijing. ~~It is mainly due to the server restriction of in-field crop residue burning.~~ Firewood gradually replaces straw as the main non-commodity biomass energy source in suburban Beijing in recent years (Wang, 2010; Liu, 2012). In addition, Tibet and Inner Mongolia are the major provinces where livestock excrement produces a large component of total pollutant emissions. Less crop straw and little firewood is used as a fuel source and thus fierce has a large contribution to total biomass emissions in these provinces. Forest and grassland fires have a small contribution to pollutant emissions in each province. The contribution of Hg emission by forest fire in ~~Inner-Mongolia, Sichuan, Yunnan, Qinghai, Tibet and Xinjiang~~ ~~Jiangsu, Zhejiang, Fujian, Hunan, Yunnan, Tibet and Qinghai~~ province is considerable (exceeding 10%), which mainly due to the high EF of Hg for forest fire.

3.2.3 Contributions from different crop straws of each province

As the largest biomass source, crop straw burning represents a major contribution to the total emissions from biomass burning. The 12 different types of straw burning emission of each province are further analysed in Fig. 5. The corn straw burning emission is concentrated in Heilongjiang, Shandong, Inner Mongolia, Hebei, Henan, Shanxi and Sichuan province, with the total contribution more than 72%. Wheat crop straw emissions mainly distributed in Henan, Shandong, Anhui, Hebei, Jiangsu, Sichuan, Shaanxi, Hubei and Shanxi province. More than 89% wheat- crop straw combustion emission is contributed by these provinces. Rice crop straw combustion emissions mainly distributed in Heilongjiang, Hunan, Jiangsu, Sichuan, Anhui, Hubei, Guangxi, Guangdong and Zhejiang province, with the total contribution more than 71%. The water condition, light and

heat are better for the cultivation of rice in the South. Low temperature, long sunshine duration, and the large temperature difference between day and night are suitable for wheat growing in the North. In addition, soybean, cotton, sugar cane, potato, peanut and rape straw have a small contribution to the various chemical species, and these straw are mainly distributed in Heilongjiang, Xinjiang, Guangxi, Sichuan, Henan and Sichuan province, respectively.

3.2.4 Emissions intensity at county resolution

At county resolution, we found that the spatial distributions of emissions for various chemical species are similar, taking PM_{2.5} as an example to analyse the emission intensity (e.g., per unit area, per capita) at county resolution. Figure 6a shows the county-level geographic distribution of PM_{2.5} emissions in 2836 counties. The numbers of counties within different emission ranges were shown in Fig. 6d. The spatial diversity of various counties emission is obvious. There are ~~403~~ 406 counties without biomass burning, because they are mainly distributed in the urban areas of developed cities, such as the Dongcheng and Xicheng districts in Beijing, the Jing'an district in Shanghai. The total emission of ~~the 33.432.3%~~ the total districts and counties (~~948917~~ 948917) in China were less than 0.25 Gg. The cumulative frequency analysis result indicated that the emission in most of the counties (i.e., more than 90%) were less than ~~3.24.0~~ 3.24.0 Gg, including the regions with low crop yield or scarce population. The emission of ~~the 31.130.9%~~ the 31.130.9% of the total districts and counties (~~883875~~ 883875) were more than the average emission across all counties (~~1.2271.245~~ 1.2271.245 Gg). The two largest emission (approximately 16 Gg) appeared in Longjiang and Wuchang which are major grain-producing counties in Heilongjiang province.

Figure 6b shows the PM_{2.5} emissions intensities per unit area. The most of high values (more than 3 Mg km⁻² ~~yr~~ yr⁻¹) mainly appeared in the north and central region of China (e.g., Hebei, Jiangsu, Shandong, Anhui, Jiangxi, Hunan), where the land is relatively flat and give priority to agricultural activity, with a substantial amount of crop straw from a relatively small area. The most counties with lower intensity concentrated in Tibet, Qinghai and Xinjiang province. In addition, it could be found that some rural counties in Heilongjiang, Jilin, and Liaoning provinces show substantial emissions, but relative lower intensity (e.g., Nenjiang in Heilongjiang, ~~7~~ 7 Dunhua in Jilin, Chaoyang in Liaoning) due to the large area of these counties.

PM_{2.5} emissions intensities per capita is illustrated in Fig. 6c. Because of the diversity of population density and biomass energy utilization, the emissions intensities per capita among various counties presents obvious difference. The counties with emission intensity more than ~~20.10~~ 20.10 kg per ¹ ~~yr~~ yr⁻¹ are mainly distributed in Heilongjiang, Jilin, Tibet and Sichuan province. The high emission intensity in northeast China are mainly attributed to the large amount of biomass burning emissions from straw and firewood burning. The high emission intensity in southwest China mainly because these regions are less economically developed (depending on non-commercial energy as straw, firewood) and prone to forest and grassland fire burning. Besides, and population in there are relatively ~~low~~ small. The ~~most~~ counties with lower emissions intensities per capita concentrated in Henan, Guangdong, and Shanxi provinces, attributed to the large amount of people there.

3.3 Spatial distribution of biomass burning emissions

As chemical species showed a similar distribution, PM_{2.5} ~~was~~ taken as an example to discuss ~~the~~ the grid emission distribution. Figure 7 shows the 1 × 1 km grid distribution. It illustrates that high biomass emissions are distributed in Henan, Heilongjiang, Shandong, Anhui, Hebei and Sichuan provinces; ~~the these~~ areas with ~~higher high~~ emission are mainly scattered in ~~major agricultural region of China's~~ northeast to central-south ~~China's~~ ~~major agricultural region~~, showing a zonal distribution. The biomass burning emissions are concentrated in the regions with greater agricultural and rural activity, and lower economic income. These regions characterized by dense population, ~~abundant~~ cultivated areas and tree resources. Low emissions are mainly distributed in the part of southwest, northwest regions and downtown areas of the majority of urban areas. The scarce populations and crop yields in part of southwest and northwest areas, and lower agricultural activity in downtown areas result in lower emissions. ~~Specially, some urban areas in the north China Plain are surrounded by suburban and rural areas, the main fuel used in these urban areas is commodity energy. Besides, there is no agricultural activity in the field. Therefore, little biomass burning emission produced by these areas. In addition, it should be noted that grid distribution result allocated from emission at county resolution could reflect the fact that there is no grid emission within several urban regions in eastern China. However, The~~ error will be brought in grid emissions if they are allocated from the emission inventory at coarse preliminary resolution (e.g., provincial or prefectural resolution before spatial allocation) based on the gridded surrogates (e.g., rural population). Consequently, gridded emissions, which were obtained through spatial allocation from emission inventory at county resolution, could better represent the actual situation.

3.4 Temporal variation in biomass burning emission

Figure 8 shows the 12 species emissions in each month, indicating that there are different ~~monthly variations~~ in the chemical species emissions ~~in different months~~. ~~The chemical species showing large monthly variations were SO₂, NO_x, PM₁₀, OC, VOC and PM_{2.5}. This is because the main contribution of these species emission sources is from straw outdoor burning. The outdoor burning straw mainly occurs in the harvest season and thus shows the obvious monthly variation features. The chemical species showing large monthly variation were SO₂, NO_x, PM₁₀, OC, NMVOC and PM_{2.5}. This is because in-field crop residue burning is the main source for these species. Besides, the in-field burning of crop residue mainly in the harvest season and thus shows the obvious monthly variation features.~~ The sources of NH₃, CO and EC emissions are dominated by straw and firewood ~~indoor domestic~~ burning and the contributions of these two kinds of source to the total emissions of these species are ~~74.0873.1%~~, ~~76.875.9%~~, and ~~87.5486.9%~~, respectively. The temporal distribution of these two sources was more uniform ~~compared to in-field crop residue burning~~ at the monthly scale, and thus monthly emissions of these three chemical species showed less temporal distinction. Despite the temporal variations of some pollutants at the monthly scale, the overall trends of emissions to ~~most each~~ species show a certain similarity: ~~April, June and October are the top four months with peak higher emissions and the contribution of these three months to the totals for various pollutant are 8.7–~~

11.2%, 8.4–12.7% and 9.0–12.8%, respectively, due to the ~~crop straw outdoor burning of the main crop in the harvest season.~~ While as for EC, the emission in February, January, October and December are relatively higher due to the biomass domestic burning in heating season. April, May, June and October are the top four months with higher emissions, due to the in-field crop residue burning. While as for EC, the emission in February, January, October and December are relatively higher due to the biomass domestic burning in heating season.

Burning activity mainly occurs in the harvest season (crop residue burning) or crop sowing season (clearing the cultivated land and increasing the soil fertility for the next sowing) ~~and it varies by burning habit in different regions.~~ In addition, the sowing and harvest seasons vary in different regions because of climate conditions. Because of the differences in ~~burning activity elimate conditions and climate conditions and sowing practices in various region each region,~~ monthly emission features vary regionally and to consider this, we divided China into seven areas, again taking PM_{2.5} as an example to analyse the pollutant emission characteristics (Fig. 9). Regions located in south China (including Fujian, Guangdong, Hainan and Guangxi provinces) and southwest China (including Chongqing, Sichuan, Guizhou, Yunnan and Tibet provinces) have climates that are highly suited to arable agriculture because of the sufficient heat and abundant rainfall. As indicated by Fig. 9, the south regions have relatively small peaks of PM_{2.5} emissions in February, April and August, these periods are consistent with local sowing and harvest times in south region. As a result of the climate ~~differences discrepancies,~~ crops in these areas are sown earlier than in northern areas. February and April are the sowing season of beans, the harvest season of the first-round and second-round crop (e.g., rice), respectively (CAAS, 1984). For the southwest region, the emission peaks are mainly distributed in February, May and August, which differ from south regions due to the inclusion of May, ~~which owing to the burning of is the harvest season of rapeseed straw and large emission of forest fire.-~~

For the central region (including Henan, Hubei and Hunan provinces), the main crops are winter wheat and summer corn, and the harvest season of these two crops are the end of May and the end of September (MOA, 2000), ~~respectively.-~~ The peak emissions in the east region (including Shanghai, Jiangsu, Zhejiang, Anhui and Jiangxi provinces) are mainly distributed from May to July, ~~while where~~ May, June and July are the harvest seasons of rapeseed, wheat and rice in east region, respectively. The northern plains of China (including Beijing, Tianjin, Hebei, Shanxi, Inner Mongolia and Shandong provinces), include the largest agricultural area in the country, accounting for 34% of the rural population, 27% of the farmland and 35% of the harvest crops (NBSC, 2013c). ~~These is~~ region differs from the eastern and central parts firstly in the usage of firewood, since here firewood is also used as heating energy and therefore the consumption of firewood in winter is greater than in summer. In addition, for the ~~in-field burning of outdoor straw crop residue burning,~~ northern winter wheat and corn are mainly harvested in June and October, respectively, and April and May are the sowing seasons of spring rice and soybeans. Northeast region (including Liaoning, Jilin and Heilongjiang provinces) shows high value in October, April and November. The high value in April was a result of burning activity. The peak in October was mainly due to the harvesting of corn and November is the harvest season for rice. In the northwest region (including Shaanxi, Gansu, Qinghai, Ningxia and Xinjiang provinces), the peaks in March–April and October ~~are is~~ due to burning activities for next sowing and corn harvesting, respectively.

Furthermore, the daily PM_{2.5} emissions are estimated according to the monthly emissions and the biomass sources daily non-uniformity coefficient, and are shown in the Supplement (Fig. S3). It could be found that the main emission peaks ~~s appearedis~~ in early April, early June and the whole month of December. This is due to (1) burning activities for the next sowing in the south, southwest, and northeast regions; (2) the harvest season of winter wheat in the central, east, and north regions; and (3) the harvest season of corn in the central, northeast, northwest regions.

3.5 Emissions of PM_{2.5} and ~~VOCNMVOC~~ species

The total PM_{2.5} emission of biomass burning emission in this study is ~~34793527~~ Gg. According to our calculation based on the method described in Sec. 2.6., OC is the largest contributor of PM_{2.5} accounting for ~~33.433.7~~ % of total emission. Cl⁻, EC, K⁺, NH₄⁺, K, and SO₄²⁻ are also the major species of PM_{2.5}, and the contribution of these species is ~~48.2946.63~~%. ~~AdditionallyBesides~~, there are several species have less emission (e.g. Al, Si, Mg). Detailed PM_{2.5} components emissions are presented in Supplement (Fig. S4).

~~The total VOC emission is 3429.6 Gg in this study. The alkenes are the major contributor of biomass burning VOC emissions. The contribution of alkenes to the total VOC emission is approximately 39%, more than it of alkane (25%), benzene series compounds (18.4%), alkynes (11.1%) and halocarbons (6.5%). Among the 91 species, ethylene, acetylene, propylene and 1 butylene are the major species of alkenes and alkynes, with the total contribution accounting for 29.7%. Methyl chloride, ethane, n propane, isobutene, and n butane are the main species of alkanes and halocarbons, with the total contribution accounting for 17.5%. Benzene, toluene, mp xylene and styrene are the major species of benzene series compounds, with the total contribution of 12.1%. Several species mentioned above are key for the formation of secondary air pollution, such as ethylene, propylene, toluene, mp xylene and halocarbons (Huang et al., 2011). It illustrates that the biomass burning emission control is very necessary for the air quality improvement. Detailed VOC species emission is shown in the Supplement (Fig. S5).~~

~~The total NMVOC emission is 3474 Gg in this study. The alkenes are the major contributor of biomass burning NMVOC emissions. The contribution of alkenes to the total NMVOC emission is approximately 34%, more than that of alkane (28%), aromatics (24%), alkynes (13%) and others (1%). Among these species, ethylene, acetylene, propylene and 1-butylene are the major species of alkenes and alkynes, with the total contribution accounting for 40.1%. Ethane, n-propane, n-butane, and n-dodecane are the main species of alkanes, with the total contribution accounting for 14.0%. Benzene, toluene, styrene, mp-xylene and ethyl benzene are the major species of aromatics, with the total contribution of 16.6%. Several species mentioned above are key for the formation of secondary air pollution, such as ethylene, propylene, toluene, mp-xylene and ethyl benzene (Huang et al., 2011). It illustrates that the biomass burning emission control is urgently needed for the air quality improvement. Detailed NMVOC species emission is shown in the Supplement (Fig. S5).~~

3.6 Uncertainties in biomass burning emission estimates

The Monte Carlo method is used to analyse the uncertainty of this emission inventory, which was used in uncertainties estimation for many inventories studies (e.g., Streets et al., 2003; Zhao et al., 2011; Zhao et al., 2012). Activity data (Zheng et al., 2009) and EFs (Zhao et al., 2011) are assumed to be normal distributions. The coefficients of variation (CV, the standard deviation divided by the mean) of activity data and emission factors were obtained from literature review. CV of activity data for firewood and crop straw burning were set as 20% (Zhao et al., 2011; Ni et al., 2015). As the data source of activity data for livestock excrement is same as the crop straw burning (i.e., government statistic data), CV is also set as 20%. MCD64A1 burned data products has been shown to be reliable in big fires (Giglio et al., 2013), and the CV of burned area of forest and grassland fire is from the reported standard deviation (Giglio et al., 2010). The biomass fuel loadings (Saatchi et al., 2011; Shi et al., 2015) and combustion factor (van der Werf et al., 2010) of forest and grassland fire were within a CV of approximately 50%. The CV of EF for each pollutant for each biomass burning type is shown in the supplement S8 and S9. The Monte Carlo method is used to analyse the uncertainty of this emission inventory, which was used in uncertainties estimation for many inventories studies (e.g., Streets et al., 2003; Zhao et al., 2011; Zhao et al., 2012). Activity data are assumed to be uniform distributions (Zheng et al., 2009) and EFs are assumed to be lognormal distributions (Zhao et al., 2011). And then we ran 10000-20000 Monte Carlo simulations to estimate the range of emissions with a 95% confidence interval. Uncertainty ranges of different pollutants in emission estimation are in Table 8. The uncertainty of NH_3 , SO_2 , EC, PM_{10} , and OC are large compared to other chemical species. The total uncertainty for emissions of these species are (-78%, 76%), (-76%, 75%), (-103%, 37%), (-37%, 80%), and (-78%, 24%), respectively. NH_3 , EC, and SO_2 exist the highest uncertainties in livestock excrement combustion. The parameters used in activity data estimation of livestock excrement exist large uncertainties, such as the proportion of total livestock manure directly combusted. Besides, the localized measurements of EF for livestock manure are limited, resulting in the large uncertainty of emission. In addition to livestock excrement source, straw burning is another large uncertainty contributor to NH_3 , EC, and SO_2 due to the parameter used in the estimation of straw burning amount (e.g., percentage of crop straw burned as fuel or in-field burning). The PM_{10} uncertainty is attributed in straw burning with the uncertainty ranging from -46.65% to 101.94%. It is mainly due to the limited localized measurements of EF. The uncertainty of OC is mainly attributed to firewood burning (-179.82% to 60.75%). Because the firewood consumption could not be directly obtained from yearbook after 2007, the data used in this study is estimated by statistical regression. This process may bring uncertainty to some extent. From the perspective of source, the uncertainty of forest fire (ranging from -624% to 631% for all pollutants) is the highest, following by grassland burning (ranging from -378% to 290% for all pollutants), livestock excrement (ranging from -300% to 295% for all pollutants), and firewood combustion (ranging from -189% to 188% for all pollutants). The uncertainty of crop straw (ranging from -114% to 114% for all pollutants) is the smallest. Uncertainty ranges of different pollutants in emission estimation are in Table 8. The total uncertainty of SO_2 , NH_3 and EC are large compared to other chemical species. The total uncertainty for emissions of these species are (-54%, 54%), (-49%, 48%) and (-61%, 61%), respectively. NH_3 , EC and SO_2 exist the highest uncertainties in

livestock excrement combustion, forest and grassland fire. The emission factors used in emission estimation of livestock excrement exist large uncertainties, which is mainly due to lack of localized measurements of EF. The large uncertainty of forest and grassland fire emission due to the uncertainty of biomass fuel loadings and combustion factor used in the estimation. Though the uncertainty exists in this study, compared with the limited research of national and comprehensive emission with uncertainty analysis (Table 8), our emission inventory is relatively reliable due to the selection of localized and specific crop EFs.

3.7 Comparison with other studies

In this paper, the national biomass burning emission inventory published after 2000 has been compared with this study (Fig. 10). It could be found that the relatively high difference (range from -80% to ~~426366~~% for various species) occur between our estimation and earlier studies (e.g., published paper before 2006) due to the economic development and EF localization. Compared with recent studies, the SO₂, NO_x, PM_{2.5}, EC, and OC emissions of our estimation are close to those derived from Lu et al. (2011), with the difference ranging from ~~-3234~~% to ~~+615~~%. While the PM₁₀, ~~VOENMVOC~~, CH₄ and NH₃ emission in this study is lower than Lu et al. (2011). The EFs of PM₁₀, ~~VOENMVOC~~, CH₄ and NH₃ for various crop types used in this study is generally lower than the EF without ~~disgusting-specific~~ crop types in Lu et al. (2011). The SO₂, NO_x, CH₄ and CO₂ emissions in this study are close to those in Tian et al. (2011), with the difference ranging from ~~-4849~~% to ~~4240~~%. The difference of CO emission is relatively high. The major emission difference of the straw ~~indoor-domestic~~ burning, ~~straw outdoor-in-field crop residue~~ burning, and firewood combustion between our paper and Tian's et al. (2011) research are -78%, -17%, and -122%. The reason is also the selection of EF. Our localized EF for crop and firewood is lower than EFs in Tian et al. (2011). In addition, for NH₃ emission, compared with the earlier studies, our estimation is close to that derived from recent research (Kang et al., 2016). The difference is less than 17%. For Hg emission, our estimation is lower than Huang et al. (2012d), but is close to Chen et al. (2013). The EF of Hg is classified by stems and leaves (40 ng/g and 100 ng/g for firewood; 35 ng/g and 319 ng/g for crop residues) in Huang et al. (2012d), which is higher than the localized EF classified by specific crop (mean EF is 6.08 ng/g) and firewood (7.2 ng/g).

4 Conclusions

In this study, a comprehensive biomass burning emission inventory with high spatial and temporal resolution was developed for mainland China in 2012, based on the county-level activity data, satellite data and updated source-specific EFs. The emission involves crop straw domestic combustion and in field burning, firewood and livestock excrement combustion, forest and grassland fires. The total annual emissions of SO₂, NO_x, PM₁₀, PM_{2.5}, ~~VOENMVOC~~, NH₃, CO, EC, OC, CO₂, CH₄ and Hg are 336.8 Gg, 990.7 Gg, 3728.3 Gg, 3526.7 Gg, 3474.2 Gg, 401.2 Gg, 34380.4 Gg, 369.7 Gg,

1189.5 Gg, 675299.0 Gg, 2092.4 Gg and 4.12 Mg, 332.8 Gg, 972.5 Gg, 3676.0 Gg, 3479.4 Gg, 3429.6 Gg, 395.8 Gg, 33987.9 Gg, 367.1 Gg, 1151.7 Gg, 665989.0 Gg, 2076.5 Gg, and 3.65 Mg, respectively.

The straw ~~indoor-domestic burning, and outdoor-in-field crop residue~~ burning and firewood combustion are the major biomass burning sources, while the largest contributing source to various pollutants is different. Straw ~~indoor-domestic~~ burning is the major source of SO₂, CO, CH₄ and ~~Hg~~ NMVOC emission, while firewood contributes most to EC and NH₃ emission. ~~Corn, rice and wheat straw are the major crop types, with the total contribution exceeding 80% for each pollutant. Corn straw burning has the greatest contribution to EC, NO_x and SO₂ emissions; rice straw burning is dominant contributor to CO₂, VOC, CH₄ and NH₃ emissions. In terms of crop straw burning, corn, rice and wheat straw are the major crop types, with the total contribution exceeding 80% for each pollutant of straw burned emissions. Corn straw burning has the greatest contribution to EC, NO_x and SO₂ emissions; rice straw burning has higher contribution to CO₂, NMVOC, CH₄ and NH₃ emissions; wheat straw burning has a considerable contribution to Hg and OC.~~ Straw burning emissions are concentrated in agricultural provinces. Firewood burning emissions are mainly distributed in southern regions of China, where the tree resource is abundant. The corn and wheat straw burning emission are mainly distributed in the northern China, while the rice straw burning emission is concentrated in the southern China. Gridded emissions result show that high emission is concentrated in northeast and central-south region of China with more agricultural and rural activity. It also illustrates that gridded emissions, which were obtained through spatial allocation from emission inventory at county resolution instead of province or prefecture resolution, could better reflect the actual situation. Monthly distributions reveal the higher emissions in April, ~~September~~ May, June and October due to the burning activity before sowing and harvesting of main crops. Regional differences of temporal distribution are attributed in the diversity of main planted crop and the climate conditions in each region. OC, Cl⁻, EC, K⁺, NH₄⁺, K, and SO₄²⁻ are the major PM_{2.5} species, with the total contribution of 80%. Several species with higher contribution to ~~VOC~~ NMVOCs (e.g., ethylene, propylene, toluene, mp-xylene and ~~ethyl benzene~~ ~~halocarbons~~) are key species for the formation of secondary air pollution. The comparison with other studies presents that the emission inventory in this study is relatively reliable. The detailed emission inventory given by this paper could provide detailed information to support the further biomass burning pollution research and the development of a targeted control strategy of all regions across the Chinese mainland.

EF and speciation of chemical species are the key parameter in the emission estimation. More localized EF of different biomass fuel types ~~indoor and outdoor burning~~ within diverse burning conditions, ~~more detained~~ PM_{2.5} and ~~VOC~~ NMVOC source profiles that contain as much components as possible still needs to expand in the future. In addition, the higher temporal resolution (e.g. hourly resolution) satellite data are necessary to provide hourly emission information for the numerical simulation of biomass burning pollution research and effective control.

Acknowledgements. The MODIS Thermal Anomalies/Fire products were provided by Land Process Distributed Active Archive Center (LPDAAC). The China Land Cover product was provided by National Science & Technology Infrastructure of China, National Earth System Science Data Sharing Infrastructure. The 1

km population distribution dataset was provided by National Science & Technology Infrastructure of China, National Earth System Science Data Sharing Infrastructure. This study was supported by the Natural Sciences Foundation of China (No. 51408014), the National Science and Technology Support Project of China (No. 2014BAC23B02 & 2014BAC23B04) and the Graduate Student Science and Technology Fund (ykj-2015-12315). The authors are grateful to the anonymous reviewers for their insightful comments.

References

- [Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, doi:10.5194/acp-11-4039-2011, 2011.](#)
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15, 955–966, doi: 10.1029/2000gb001382, 2001.
- [Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part 1, The nature and sources of cloud-active aerosols, *Earth Sci. Rev.*, 89, 13–41, doi:10.1016/j.earscirev.2008.03.001, 2008.](#)
- Bao, J. C., Yu, J. H., Feng, Z., Chen, B. H., Lei, C. and Yang, J.: Situation of Distribution and Utilization of Crop Straw Resources in Seven Western Provinces, *Chinese Journal of Applied Ecology*, 25, 181–187, 2014 (in Chinese).
- Bi, Y. Y.: Study on Straw Resources Evaluation and Utilization in China, Ph.D. thesis, Chinese Academy of Agriculture Sciences, China, Beijing, 2010 (in Chinese).
- Cao, G. L., Wang, F. C. and Wang, Y. Q.: Emission inventory of TSP, PM₁₀ and PM_{2.5} emission from biomass burning in China, *Journal of Process Engineering*, 700–704, 2004 (in Chinese).
- Cao, G. L., Zhang, X. Y., Wang, D. and Deng, F. C.: Inventory of Atmospheric Pollutants Discharged from Biomass Burning in China Continent, *China Environmental Science*, 25, 389–393, 2005 (in Chinese).
- Cao, G. L., Zhang, X. Y., Gong, S. L. and Zheng, F. C.: Investigation on emission factors of particulate matter and gaseous pollutants from crop residue burning, *Journal of Environmental Sciences*, 20, 50–55, doi: 10.1016/s1001-0742(08)60007-8, 2008 (in Chinese).
- Chang, C., Tian, J. Q., Feng, D. C., Jin, X. X., Xia, S. Y., Zhang, Y., Lu, X. Q., Lv, D. M. and Yu, G. J.: Straw Resources and Problems and Measures of Straw Returning to Field in Liaoning Province, *Liaoning Agricultural Sciences*, 6, 71–73, 2012 (in Chinese).
- Chang, D. and Song, Y.: Estimates of biomass burning emissions in tropical Asia based on satellite-derived data, *Atmos. Chem. Phys.*, 10, 2335–2351, 2010.

Chen, C., Wang, H. H., Zhang, W., Hu, D., Chen, L. and Wang, X. J.: High-resolution inventory of mercury emissions from biomass burning in China for 2000–2010 and a projection for 2020, *Journal of Geophysical Research–Atmospheres*, 118, 12248–12256, doi: 10.1002/2013jd019734, 2013.

Cheng, Z., Wang, S., Fu, X., Watson, J. G., Jiang, J., Fu, Q., Chen, C., Xu, B., Yu, J., Chow, J.C. and Hao, J.: Impact of biomass burning on haze pollution in the Yangtze River delta, China: a case study in summer 2011, *Atmos. Chem. Phys.*, 14, 4573–4585, doi: 10.5194/acp-14-4573-2014, 2014.

Chinese Academy of Agricultural Sciences (CAAS): *Agricultural Regionalization of China*, China Agriculture Press, Beijing, 275 pp, 1984 (In Chinese).

Christian, T., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P. J., Hao, W. M., Saharjo, B. H. and Ward, D. E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African, and other fuels, *J. Geophys. Res.*, 108(D23), 4719, 2003.

Crutzen, P. J., Heidt, L. E., Krasnec, J. P., Pollock, W. H. and Seiler, W.: Biomass Burning as a Source of Atmospheric Gases CO, H₂, N₂O, NO, CH₃CL and COS, *Nature*, 282, 253–256, doi: 10.1038/282253a0, 1979.

Emily, P. G. and Martin, D.: *Health Effects of Indoor Air Pollution from Biomass Cooking Stove*, Public Health Perspective, 2008.

EOCAIY: *China Animal Industry Yearbook 2013*, China Agriculture Press, Beijing, 2013 (in Chinese).

~~EOCAY: China Agriculture Yearbook 2013, China Agriculture Press, Beijing, 2013 (in Chinese).~~

EPD: Guide for compiling atmospheric pollutant emission inventory for biomass burning, Environmental Protection Department, 2014, available at http://www.zhb.gov.cn/gkml/hbb/bgg/201501/t20150107_293955.htm, (in Chinese).

Fang, F., Li, X., Shi, Z. L., Wang, F., Chang, Z. Z., Zhang, S., Shun, R. H., Bao, Z. and Qiu, L.: Analysis on Distribution and Use Structure of Crop Straw Resources in Huang–Huai–Hai Plain of China, *Transactions of the CSAE*, 31, 228–234, 2015 (in Chinese).

Fang, J.Y., Liu, G.H. and Xu, S.L.: Biomass and net production of forest vegetation in China. *Acta. Eco.Sin.*, 16, 497-508, 1996 (in Chinese).

Fang, J. Y., Wang, G. G., Liu, G. H. and Xu, S. L.: Forest biomass of China: An estimate based on the biomass-volume relationship, *Ecol. Appl.*, 8, 1084–1091, 1998.

Fernandez, A., Davis, S. B., Wendt, J. O. L., Cenni, R., Young, R. S. and Witten, M. L.: Public health – Particulate emission from biomass combustion, *Nature*, 409, 998–998, doi: 10.1038/35059169, 2001.

[Friedli, H. R., Radke, L. F., Prescott, R., Hobbs, P. V., and Sinha, P., Mercury emissions from the August 2001 wildfires in Washington State and an agricultural waste fire in Oregon and atmospheric mercury budget estimates, *Global Biogeochem. Cycle*, 17\(2\), 1039, doi:10.1029/2002GB001972, 2003.](#)

Fu, J. Y., Jiang, D. and Huang, Y. H.: China 1 km grid population distribution data set, Publish system for global change science research data, 2014 (in Chinese).

Fu, X., Wang, S. X., Zhao, B., Xing, J., Cheng, Z., Liu, H. and Hao, J. M.: Emission inventory of primary pollutants and chemical speciation in 2010 for the Yangtze River Delta region, China, *Atmos. Environ.*, 70, 39–50, doi: 10.1016/j.atmosenv.2012.12.034, 2013.

Fullerton, D.G., Bruce, N. and Gordon, S. B.: Indoor air pollution from biomass fuel smoke is a major health concern in the developing world, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102, 843–851, doi: 10.1016/j.trstmh.2008.05.028, 2008.

Gao, L. W., Ma, L., Zhang, W. F., Wang, F. H., Ma, W. Q. and Zhang, F. S.: Estimation of Nutrient Resource Quantity of Crop Straw and Its Utilization Situation in China, *Transactions of the CSAE*, 25, 173–179, 2009 (in Chinese).

Gao, X. Z., Ma, W. Q., Ma, C. B., Zhang, F. S. and Wang, Y. H.: Analysis on the Current Status of Utilization of Crop Straw in China, *Journal of Huazhong Agricultural University*, 21, 242–247, 2002 (in Chinese).

[Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.](#)

[Giglio, L., Randerson, J. T. and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database \(GFED4\), *J. Geophys. Res., Biogeosci.*, 118, 317–328, 2013.](#)

Han, L. J., Yan, Q. J., Liu, X. Y. and Hu, J. Y.: Straw Resources and Their Utilization in China, *Transactions of the CSAE*, 18, 87–91, 2002 (in Chinese).

He, M., Wang, X. R., Han, L., Feng, X. Q. and Mao, X.: Emission Inventory of Crop Residues Field Burning and Its Temporal and Spatial Distribution in Sichuan Province, *Environmental Science*, 36, 1208–1216, 2015 (in Chinese).

He, M., Zheng, J. Y., Yin, S. S. and Zhang, Y. Y.: Trends, temporal and spatial characteristics, and uncertainties in biomass burning emissions in the Pearl River Delta, China, *Atmos. Environ.*, 45, 4051–4059, 2011.

Hou, X. Q., Zhang, H. T. and Yang, J. X.: Study in Crop Straw Comprehensive Utilization Mode of Xinjiang, Master thesis, Xinjiang Agriculture University, China, Xinjiang, 2012 (in Chinese).

[Hu, H. F., Wang, Z. H., Liu, G. H., et al. Vegetation carbon storage of major shrublands in China. *J. Pla. Eco.*, 30, 539-544, 2006 \(in Chinese\).](#)

Huang, C., Chen, C. H., Li, L., Cheng, Z., Wang, H. L., Huang, H. Y., Streets, D. G., Wang, Y. J., Zhang, G. F. and Chen, Y. R.: Emission inventory of anthropogenic air pollutants and VOC species in the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 11, 4105–4120, doi: 10.5194/acp-11-4105-2011, 2011.

Huang, G. B.: Evaluation and Trend Analysis of Crop Straw Comprehensive Utilization in Fujian Province, *Research*, 66–67, 2012a (in Chinese).

Huang, X., Song, Y., Li, M. M., Li, J. F. and Zhu, T.: Harvest season, high polluted season in East China, *Environ. Res. Lett.*, 7, doi: 10.1088/1748-9326/7/4/044033, 2012b.

Huang, X., Song, Y., Li, M. M., Li, J. F., Huo, Q., Cai, X. H., Zhu, T., Hu, M. and Zhang, H. S.: A high-resolution ammonia emission inventory in China, *Global Biogeochem. Cycles*, 26, doi: 10.1029/2011gb004161, 2012c.

Huang, X., Li, M. M., Hans, R. F., Song, Y., Di, C. and Lei, Z.: Mercury Emissions from Biomass Burning in China, *Environ. Sci. Technol.*, 45, 9442–9448, 2012d.

Huang, X., Li, M. M., Li, J. F. and Song, Y.: A high-resolution emission inventory of crop burning in fields in China based on MODIS Thermal Anomalies/Fire products, *Atmos. Environ.*, 50, 9-15, 2012e.

Hurst, D. F., Griffith, D. W. T. and Cook, G. D.: Trace gas emissions from biomass burning in tropical Australian savannas, *J. Geophys. Res.*, 99, 16441–16456, 1994.

International Energy Agency (IEA), IEA Statistics 2012, IEA Publication, 2012.

Intergovernmental Panel on Climate Change (IPCC): Climate Change 2007, [available at http://www.ipcc.ch](http://www.ipcc.ch), 2011.

Ito, A. and Penner, J. E.: Global estimates of biomass burning emissions based on satellite imagery for the year 2000, *Journal of Geophysical Research–Atmospheres*, 109, doi: 10.1029/2003jd004423, 2004.

Jiang, R. and Bell, M.: The Characterization and Health Effects of Indoor Particulate Matter Pollution from Biomass Burning in Northeastern China, *Epidemiol.*, 19, 89–90, 2008.

Kanabkaew, T. and Nguyen, T. K. O.: Development of Spatial and Temporal Emission Inventory for Crop Residue Field Burning, *Environ. Model. Assess.*, 16, 453–464, doi: 10.1007/s10666-010-9244-0, 2011.

Kang, Y. N., Liu, M. X., Song, Y., Huang, X., Yao, H., Cai, X. H., Zhang, H. S., Kang, L., Liu, X. J., Yan, X. Y., He, H., Zhang, Q., Shao, M. and Zhu, T.: High-resolution ammonia emissions inventories in China from 1980 to 2012, *Atmos. Chem. Phys.*, 16, 2043–2058, doi: 10.5194/acp-16-2043-2016, 2016.

Kasischke, E. S., Stocks, B. J., O'Neill, K., French, N. H. F. and Bourgeau-Chavez, L. L.: Direct effect of fire on the boreal forest carbon budget, in Biomass Burning and Its Inter-Relationships With the Climate System, 51–68, Dordrecht, Norwell, Mass, 2000.

Kaufman, Y. J. and Fraser, R. S.: The effect of smoke particles on clouds and climate forcing, *Science*, 277, 1636–1639, doi: 10.1126/science.277.5332.1636, 1997.

5 Koppmann, R., von Czapiewski, K. and Reid, J. S.: A review of biomass burning emissions, part I: gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds, *Atmos. Chem. Phys. Discuss.*, 5, 10455–10516, doi:10.5194/acpd-5-10455-2005, 2005.

Langmann, B., Duncan, B., Textor, C., Trentmann, J. and van der Werf, G. R.: Vegetation fire emissions and their impact on air pollution and climate, *Atmos. Environ.*, 43, 107–116, doi: 10.1016/j.atmosenv.2008.09.047, 2009.

10 Levine, J. S.: Global biomass burning: A case study of the gaseous and particulate emissions released to the atmosphere during the 1997 fires in Kalimantan and Sumatra, Indonesia, in Biomass Burning and Its Inter-Relationships with the Climate System, 15–31, Dordrecht, Norwell, Mass, 2002.

Li, Q., Jiang, J. K., Cai, S. Y., Zhou, W., Wang, S. X., Duan, L. and Hao, J. M.: Gaseous Ammonia Emissions from Coal and Biomass Combustion in Household Stoves with Different Combustion Efficiencies, *Environ. Sci. Technol.*, 3, 98–103, 2016.

15 Li, Q. J.: Study on the Comprehensive Utilization of Straw in Guangxi, Master thesis, College of Agriculture, China, Guangxi, 2013a (in Chinese).

Li, G. J.: Status of Anda Area Livestock Manure Handling and Effective Utilization, *Veterinary Ophthalmology*, 12, 25–17, 2007a (in Chinese).

20 Li, G. J. and Zhao, R. H.: The effective utilization and treatment of livestock excrement in Anda area, *Contemporary Animal Husbandry*, 3, 43–45, 2008 (in Chinese).

Li, J. F., Song, Y., Li, M. M. and Huang, X.: Estimating Air Pollutants Emissions from Open Burning of Crop Residues in Jiangnan Plain, *Acta Scientiarum Naturalium Universitatis Pekinensis*, 51, 647–656, 2015 (in Chinese).

Li, M., He, W. Y., Li, Z. Y. and Zheng, R.: Study on the Distribution Characteristics and Utilization of Crop Straws in Chongqing City, *South China Agriculture*, 7, 32–34, 2013b (in Chinese).

25 Li, X. H., Duan, L., Wang, S. X., Duan, J. C., Guo, X. M., Yi, H. H., Hu, J. N., Li, C. and Hao, J. M.: Emission characteristics of particulate matter from rural household biofuel combustion in China, *Energy & Fuels*, 21, 845–851, doi: 10.1021/ef060150g, 2007b.

- Li, X. H., Wang, S. X., Duan, L., Hao, J., Li, C., Chen, Y. S. and Yang, L.: Particulate and trace gas emissions from open burning of wheat straw and corn stover in China, *Environ. Sci. Technol.*, 41, 6052–6058, doi: 10.1021/es0705137 2007c.
- Li, X. H., Wang, S. X., Duan, L., Hao, J. M. and Nie, Y. F.: Carbonaceous Aerosol Emissions from Household Biofuel Combustion in China, *Environ. Sci. Technol.*, 43, 6076–6081, doi: 10.1021/es803330j; 2009.
- 5 Liu, G. and Shen, L.: Quantitative assessment of biomass energy and its geographical distribution in China, *Journal of Natural Resources*, 22(1), 9–19, 2007 (in Chinese).
- Liu, H. Y., Zhou, J. G., Zhou, P., Xiao, H. A. and Wu, J. S.: Analyze Regional Characteristic and Influencing Factors of Different Crop Straw Treatments in South Central of China, *Quaternary Sciences*, 34, 848–855, 2014 (in Chinese).
- Liu, M. X., Song, Y., Yao, H., Kang, Y. N., Li, M. M. and Huang, X.: Estimating emissions from agricultural fires in the North China Plain based on MODIS fire radiative power, *Atmos. Environ.*, 112, 326–334, 2015.
- 10 Liu, P., Na, W., Wang, Q. L., Wang, X. M., Zhang, W. D. and Wang, X. F.: Analysis on Evaluation and Energy Utilization of Main Crop Stalk Resource in Jilin Province, *Journal of Jilin Agricultural Sciences*, 35, 58–64, 2010 (in Chinese).
- Liu, Y., Shao, M., Fu, L. L., Lu, S. H., Zeng, L. M. and Tang, D. G.: Source profiles of volatile organic compounds (~~VO~~ENMVOCs) measured in China: Part I, *Atmos. Environ.*, 42, 6247–6260, doi: 10.1016/j.atmosenv.2008.01.070, 2008.
- 15 Liu, Z. R.: Analysis of Beijing rural life energy consumption and study on the policies, Master thesis, Southwest University, China, Chongqing, 2012 (in Chinese).
- Lu, B., Kong, S. F., Han, B., Wang, X. Y. and Bai, Z. P.: Inventory of Atmospheric Pollutants Discharged from Biomass Burning in China Continent in 2007, *China Environmental Science*, 31, 186–194, 2011, http://manu36.magtech.com.cn/Jweb_zghjcx/CN/Y2011/V31/I2/186, (in Chinese).
- 20 MA: Investigation and Evaluation Report on Crop Straw Resources in China, Ministry of Agriculture, 2011, available at <http://www.kjs.moa.gov.cn/>, (in Chinese).
- McMeeking, G. R.: The optical, chemical, and physical properties of aerosols and gases emitted by the laboratory combustion of wildland fuels, Ph.D. Dissertation, Department of Atmospheric Sciences, Colorado State University, 109–113, Fall 2008.
- MEP: Measures of Prohibiting Straw Burning and Comprehensive Utilization Management, Ministry of Environmental Protection of the People’s Republic of China, 1999 (in Chinese).
- 25 Michel, C., Liousse, C., Gre igoire, J.M., Tansey, K., Carmichael, G.R. and Woo, J.H.: Biomass burning emission inventory from burnt area data given by the SPOT-VEGETATION system in the frame of TRACE-P and ACE-Asia campaigns, *J. Geophys. Res.*, 110, 2005.

- Ministry of Agriculture of the People's Republic of China (MOA), Database of Farming Season, Ministry of Agriculture of the People's Republic of China, [available at http://www.zzys.gov.cn/](http://www.zzys.gov.cn/), 2000 (in Chinese).
- National Agricultural Technology Extension Service Center (NATESC): China Organic Fertilizer Resources, China Agriculture Press, Beijing, 1999 (in Chinese).
- 5 National Bureau of Statistics of China (NBSC): China Statistical Yearbook 1999–2008, China Statistics Press, Beijing, 1999–2008a, [available at http://www.stats.gov.cn/tjsj/ndsj/](http://www.stats.gov.cn/tjsj/ndsj/), (in Chinese).
- National Bureau of Statistics of China (NBSC): China Energy Statistical Yearbook 1999–2008, China Statistics Press, Beijing, 1999–2008b, [available at http://www.stats.gov.cn/tjsj/tjcbw/](http://www.stats.gov.cn/tjsj/tjcbw/), (in Chinese).
- ~~National Bureau of Statistics of China (NBSC): China Energy Statistical Yearbook 2009–2015, China Statistics Press, Beijing, 2009–2015, [available at http://www.stats.gov.cn/tjsj/tjcbw/](http://www.stats.gov.cn/tjsj/tjcbw/), (in Chinese).~~
- 10 National Bureau of Statistics of China (NBSC): China County Statistical Yearbook 2013, China Statistics Press, Beijing, 2013a (in Chinese).
- National Bureau of Statistics of China (NBSC): China Statistical Yearbook for Regional Economy 2013, China Statistics Press, Beijing, 2013b, [available at http://www.stats.gov.cn/tjsj/tjcbw/](http://www.stats.gov.cn/tjsj/tjcbw/), (in Chinese).
- National Bureau of Statistics of China (NBSC): China Statistical Yearbook 2013, China Statistics Press, Beijing, 2013c, [available at http://www.stats.gov.cn/tjsj/ndsj/](http://www.stats.gov.cn/tjsj/ndsj/), (in Chinese).
- 15 ~~National Bureau of Statistics of China (NBSC): China Statistical Yearbook on Environment 2013, China Statistics Press, Beijing, 2013d (in Chinese).~~
- NDRC: National utilization and burning of straw in 2012, National Development and Reform Commission, 2014 (in Chinese).
- Nelson, P. F., Morrison, A. L., Malfroy, H. J., Cope, M., Lee, S., Hibberd, M. L., Meyer, C. P. and McGregor, J.: Atmospheric mercury emissions in Australia from anthropogenic, natural and recycled sources, Atmos. Environ., 62, 291-302, 2012.
- 20 Ni, H. Y., Han, Y. M., Cao, J. J., Chen, L. W. A., Tian, J., Wang, X. L., Chow, J. C., Watson, J. G., Wang, Q. Y., Wang, P., Li, H. and Huang, R. J.: Emission characteristics of carbonaceous particles and trace gases from open burning of crop residues in China, Atmos. Environ., 123, 399–406, doi: 10.1016/j.atmosenv.2015.05.007, 2015.
- Penner, J. E., Dickinson, R. E. and Oneill, C. A.: Effects of Aerosol from Biomass Burning on the Global Radiation Budget, Science, 256, 1432–1434, doi: 10.1126/science.256.5062.1432, 1992.
- 25 ~~[Pu, S. L., Fang, J. Y. and He, J. S.: Spatial distribution of grassland biomass in China, Acta. Phyt. Sci., 28, 491-498, 2004 \(in Chinese\).](#)~~

- Qin, D. D. and Ge, L.: Study on the comprehensive utilization of straw in Anhui Province, Modern Agricultural Technology, 8, 264–265, 2012 (in Chinese).
- Ran, Y. H., Li, X. and Lu, L.: Evaluation of four remote sensing based land cover products over China, Int. J. Remote Sens., 31, 391–401, [available at http://www.tandfonline.com/doi/abs/10.1080/01431160902893451](http://www.tandfonline.com/doi/abs/10.1080/01431160902893451), 2010.
- 5 Reddy, M. S. and Venkataraman, C.: Inventory of aerosol and sulphur dioxide emissions from India. Part II – biomass combustion. Atmos. Environ. 36, 699–712, doi: 10.1016/s1352-2310(01)00464-2, 2002.
- [Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M. and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, Proc. Natl. Acad. Sci. U. S. A., 108, 9899–9904, 2011.](#)
- 10 ~~Shekar, R. M. and Chandra, V.: Inventory of Aerosol and Sulphur Dioxide Emissions from India. Part II: Biomass Combustion, Atmos. Environ., 36, 699–712, 2002.~~
- Shi, Y. S. and Yamaguchi, Y.: A high-resolution and multi-year emissions inventory for biomass burning in Southeast Asia during 2001–2010, Atmos. Environ., 98, 8–16, doi: 10.1016/j.atmosenv.2014.08.050, 2014.
- [Shi, Y. S., Matsunaga, T. and Yamaguchi, Y.: High-Resolution Mapping of Biomass Burning Emissions in Three Tropical Regions, Environ. Sci. Technol., 49, 10806–10814, 2015.](#)
- 15 Shon, Z. H.: Long-term variations in PM_{2.5} emission from open biomass burning in Northeast Asia derived from satellite-derived data for 2000–2013, Atmos. Environ., 107, 342–350, doi: 10.1016/j.atmosenv.2015.02.038, 2015.
- Song, Y., Liu, B., Miao, W., Chang, D. and Zhang, Y.: Spatiotemporal variation in ~~nonagricultural~~non-agricultural open fire emissions in China from 2000 to 2007, Global Biogeochem. Cycles, 23, GB2008, doi: 10.1029/2008GB003344, 2009.
- 20 Song, Y., Shao, M. and Liu, Y.: Source apportionment of ambient volatile organic compounds in Beijing, Environ. Sci. Technol., 41, 4348–4353, 2007.
- Streets, D. G., Gupta, S., Waldhoff, S. T., Wang, M. Q., Bond, T. C. and Bo, Y. Y.: Black carbon emissions in China. Atmos. Environ., 35, 4281–4296, doi: 10.1016/S1352-2310(01)00179-0, 2001.
- Streets, D. G., Yarber, K. F., Woo, J. -H. and Carmichael, G. R.: Biomass burning in Asia: Annual and seasonal estimates and atmospheric emissions, Global Biogeochem. Cycles, 17, 1099, doi: 10.1029/2003GB002040, 2003.
- 25 Streets, D. G., Hao, J. M., Wu, Y., Jiang, J. K., Chan, M., Tian, H. Z. and Feng, X. B.: Anthropogenic mercury emissions in China, Atmos. Environ., 39, 7789–7806, doi: 10.1016/j.atmosenv.2005.08.029, 2005.

- Sun, J. F., Peng, H. Y., Chen, J. M., Wang, X. M., Wei, M., Li, W. J., Yang, L. X., Zhang, Q. Z., Wang, W. X. and Mellouki, A.: An estimation of CO₂ emission via agricultural crop residue open field burning in China from 1996 to 2013, *J. Cleaner Prod.*, 112, 2625–2631, doi: 10.1016/j.jclepro.2015.09.112, 2016.
- Tang, X. B., Huang, C., Lou, S. R., Qiao, L. P., Wang, H. L., Zhou, M., Chen, M. H., Chen, C. H., Wang, Q., Li, G. L., Li, L., Huang, H. Y. and Zhang, G. F.: Emission Factors and PM Chemical Composition Study of Biomass Burning in the Yangtze River Delta Region, *Environmental Science*, 35, 1623–1632, 2014 (in Chinese).
- Tian, H. Z., Hao, J. M., Lu, Y. Q. and Zhou, Z. P.: Evaluation of SO₂ and NO_x Emissions Resulted from Biomass Fuels Utilization in China, *Acta Scientiae Circumstantiae*, 22, 204–208, 2002 (in Chinese).
- Tian, H. Z., Zhao, D. and Wang, Y.: Emission Inventories of Atmospheric Pollutants Discharged from Biomass Burning in China, *Acta Scientiae Circumstantiae*, 31, 349–357, 2011 (in Chinese).
- ~~Tian, X. R., Shu, L. F. and Wang, M. Y.: Direct Carbon Emissions from Chinese Forest Fires, 1991–2000, *Fire Safety Science*, 12, 6–10, 2003 (in Chinese).~~
- Tian, Y. S.: Current Development Situation and Trend of China's Rural Energy in 2013, *Energy of China*, 36, 10–14, 2014, [available at http://www.zgln.chinajournal.net.cn/WKB3/WebPublication/paperDigest.aspx?paperID=8703e983-c6a0-447d-adff-60e4b8ca5c51](http://www.zgln.chinajournal.net.cn/WKB3/WebPublication/paperDigest.aspx?paperID=8703e983-c6a0-447d-adff-60e4b8ca5c51), (in Chinese).
- Turn, S. Q., Jenkins, B. M., Chow, J. C., Pritchett, L. C., Campbell, D., Cahill, T. and Whalen, S. A.: Elemental characterization of particulate matter emitted from biomass burning: Wind tunnel derived source profiles for herbaceous and wood fuels, *Journal of Geophysical Research–Atmospheres*, 102, 3683–3699, doi: 10.1029/96jd02979, 1997.
- U.S. Environmental Protection Agency (U.S. EPA): Compilation of air pollutant emission Factors, AP-42, 5th ed., Washington, DC, 2002.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S. and Arellano, A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, 2006.
- ~~van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.~~
- Wang, J. J.: The Statistical Analysis of Beijings Rural Energy Consumption, Master thesis, Hebei University, China, Hebei, 2010 (in Chinese).

- Wang, S. X. and Zhang, C. Y.: Spatial and Temporal Distribution of Air Pollutant Emissions from Open Burning of Crop Residues in China, Sciencepaper Online, 3, 329–333, 2008 (in Chinese).
- Wang, S. X., Wei, W., Du, L., Li, G. H. and Hao, J. M.: Characteristics of gaseous pollutants from biofuel–stoves in rural China, Atmos. Environ., 43, 4148–4154, doi: 10.1016/j.atmosenv.2009.05.040, 2009.
- 5 Wang, Y. and Zhao, Q. R.: Current situation and measures of Jiangsu straw utilization, Jiangsu Agricultural Economy, 31, 66–67, 2011 (in Chinese).
- Watson, J. G., Chow, J. C. and Houck, J. E.: PM_{2.5} chemical source profiles for vehicle exhaust, vegetative burning, geological material, and coal burning in Northwestern Colorado during 1995, Chemosphere, 43, 1141–1151, doi: 10.1016/s0045-6535(00)00171-5, 2001.
- Wei, W., Wang, S. X., Chatani, S., Klimont, Z., Cofala, J. and Hao, J. M.: Emission and speciation of non–methane volatile organic compounds from anthropogenic sources in China, Atmos. Environ., 42, 4976–4988, doi: 10.1016/j.atmosenv.2008.02.044, 2008.
- 10 Xu, P., Liao, Y. J., Lin, Y. H., Zhao, C. X., Yan, C. H., Cao, M. N., Wang, G. S., and Luan, S. J.: High-resolution inventory of ammonia emissions from agricultural fertilizer in China from 1978 to 2008, Atmos. Chem. Phys., 16, 1207–1218, doi:10.5194/acp-16-1207-2016, 2016.
- ~~Xu, P., Zhang, Y. S., Gong, W.W., Hou, X. K., Kroeze, C., Gao, W. and Luan, S. J.: An inventory of the emission of ammonia from agricultural fertilizer application in China for 2010 and its high-resolution spatial distribution, Atmos. Environ. 115, 141–148, doi: 10.1016/j.atmosenv.2015.05.020, 2015.~~
- 15 ~~Xu, P., Zhang, Y. S., Gong, W.W., Hou, X. K., Kroeze, C., Gao, W. and Luan, S. J.: An inventory of the emission of ammonia from agricultural fertilizer application in China for 2010 and its high-resolution spatial distribution, Atmos. Environ. 115, 141–148, doi: 10.1016/j.atmosenv.2015.05.020, 2015.~~
- Yan, W. L., Liu, D. Y., Sun Y., Wei, J. S. and Pu, M. J.: Analysis of the Sustained Fog and Haze Event Resulting from Crop–Burning Residue in Jiangsu Province, Climatic and Environmental Research, 19, 237–247, 2014 (in Chinese).
- Yan, X. Y., Ohara, T. and Akimoto, H.: Bottom–up estimate of biomass burning in mainland China, Atmos. Environ., 40, 5262–5273, doi: 10.1016/j.atmosenv.2006.04.040, 2006.
- 20 Zha, S. P., Zhang, S. Q., Cheng, T. T., Chen, J. M., Huang, G. H., Li, X. and Wang, Q. F.: Agricultural Fires and Their Potential Impacts on Regional Air Quality over China, Aerosol and Air Quality Research, 13, 992–1001, doi: 10.4209/aaqr.2012.10.0277, 2013.
- Zhang, F. C. and Zhu, Z. H.: Harvest Index of Crops in China, Scientia Agricultura Sinica, 23, 83–87, 1990 (in Chinese).
- Zhang, H. F., Ye, X. N., Cheng, T. T., Chen, J. M., Yang, X., Wang, L. and Zhang, R. Y.: A laboratory study of agricultural crop residue combustion in China: Emission factors and emission inventory, Atmos. Environ. 42, 8432–8441, doi: 10.1016/j.atmosenv.2008.08.015, 2008.
- 25

- Zhang, L. B., Liu, Y. Q. and Hao, L.: Contributions of open crop straw burning emissions to PM_{2.5} concentrations in China, *Environ. Res. Lett.*, 11, doi:10.1088/1748-9326/11/1/014014, 2016.
- Zhang, J., Smith, K. R., Ma, Y., Ye, S., Jiang, F., Qi, W., Liu, P., Khalil, M. A. K., Rasmussen, R.A. and Thorneloe, S.A.: Greenhouse gases and other airborne pollutants from household stoves in China: a database for emission factors, *Atmos. Environ.* 34, 4537–4549, doi: 10.1016/s1352-2310(99)00450-1, 2000.
- Zhang, Q., Klimont, Z., Streets, D. G., Huo, H. and He, K. B.: Emission model of anthropogenic sources and estimation of emission inventory in 2001, China, 16, 223–231, *Progress in Natural Science*, 2006 (in Chinese).
- Zhang, W., Wei, W., Hu, D., Zhu, Y. and Wane, X.J.: Emission of Speciated Mercury from Residential Biomass Fuel Combustion in China. *Energy & Fuels*, 27, 6792–6800, doi: 10.1021/ef401564r, 2013a.
- Zhang, Y. R., Li, Y., Jiang, T. M. and Zhang, W. A.: Distribution and Comprehensive Utilization of Straw Resources of Main Crops in Guizhou, *Guizhou Agricultural Sciences*, 3, 262–267, 2015 (in Chinese).
- Zhang, Y. S., Shao, M., Lin, Y., Luan, S. J., Mao, N., Chen, W. T. and Wang, M.: Emission inventory of carbonaceous pollutants from biomass burning in the Pearl River Delta Region, China, *Atmos. Environ.*, 76, 189–199, doi: 10.1016/j.atmosenv.2012.05.055, 2013b.
- Zhao, B., Wang, P., Ma, J. Z., Zhu, S., Pozzer, A. and Li, W.: A high-resolution emission inventory of primary pollutants for the Huabei region, China, *Atmos. Chem. Phys.*, 12, 481–501, doi: 10.5194/acp-12-481-2012, 2012.
- Zhao, C.: Utilization Status of Crop Straws and Sustainable Development in Beijing, *Journal of Agriculture*, 5, 42–46, 2015 (in Chinese).
- Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B. and Hao, J.: Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China, *Atmos. Chem. Phys.*, 11, 2295–2308, doi: 10.5194/acp-11-2295-2011, 2011.
- Zheng, B., Huo, H., Zhang, Q., Yao, Z. L., Wang, X. T., Yang, X. F., Liu, H. and He, K. B.: High-resolution mapping of vehicle emissions in China in 2008, *Atmos. Chem. Phys.*, 14, 9787–9805, doi: 10.5194/acp-14-9787-2014, 2014.
- Zheng, J. Y., Zhang, L. J., Che, W. W., Zheng, Z. Y. and Yin, S. S.: A highly resolved temporal and spatial air pollutant emission inventory for the Pearl River Delta region, China and its uncertainty assessment, *Atmos. Environ.*, 43, 5112–5122, doi: 10.1016/j.atmosenv.2009.04.060, 2009.
- Zhou, Y., Cheng, S.Y., Lang, J. L., Chen, D. S., Zhao, B. B., Liu, C., Xu, R. and Li, T. T.: A comprehensive ammonia emission inventory with high-resolution and its evaluation in the Beijing–Tianjin–Hebei (BTH) region, China. *Atmos. Environ.*, 106, 305–317, doi: 10.1016/j.atmosenv.2015.01.069, 2015.

Zong, Z., Wang, X., Tian, C., Chen, Y., Qu, L., Ji, L., Zhi, G., Li, J. and Zhang, G.: Biomass burning contribution to regional PM_{2.5} during winter in the North China, Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2016-97, in review, 2016.

5

10

15

Table Captions List:

Table 1. The classification of biomass burning emission source.

Table 2. ~~Indoor-Straw domestic~~ and ~~outdoor-in-field straw-crop residue~~ burning proportions of each province.

Table 3. Residue-to-production ratio (N_k), dry matter fraction (D_k) and combustion efficiency (CE_k) used in this study.

Table 4. ~~Forest and grassland biomass fuel loadings in each province.~~~~Forest biomass and combustion efficiency of each province.~~

Table 5. Emission factors used in the estimation of ~~indoor-domestic~~ biomass burning emissions.

Table 6. Emission factors used in the estimation of ~~outdoor-open~~ biomass burning emissions.

Table 7. Biomass burning emission inventory in the 31 provinces or municipalities of China in 2012.

Table 8. Uncertainty ranges of different pollutants in emission estimates (min, max). (Unit for emission estimate: Gg)

Figure Captions List:

Figure1. Regression analysis between firewood consumption at province resolution and (1) rural population, (2) gross agricultural output, and (3) timber yield, respectively.

Figure 2. Contribution of different source to the total biomass burning emissions in China, 2012.

Figure 3. Contributions of 12 crop straw types to total straws burning emissions for various species.

~~Contributions of 12 crop straws burning to total pollutant emissions for various species.~~

Figure 4. Contributions of different biomass sources to the emission in each province (Gg).

Figure 5. Contributions of different crop straw types to the emission in each province (Gg).

Figure 6. Biomass emission inventory at county resolution and intensity ($PM_{2.5}$).

Figure 7. Gridded distribution of $PM_{2.5}$ annual emissions.

Figure 8. Monthly variation of different biomass sources emission for each chemical species.

Figure 9. Monthly variation of different biomass sources emission for $PM_{2.5}$ emissions in different regions.

Figure 10. Comparison of the emissions inventory derived by this study with the emissions estimated by previous research.

Table 1. The classification of biomass burning emission source.

I	II	III
<u>Domestic combustion</u>	<u>Firewood</u>	<u>Firewood</u>
	<u>Livestock excrement</u>	<u>Cattles</u>
		<u>Horses</u>
		<u>Donkeys</u>
		<u>Mules</u>
		<u>Camels</u>
<u>Open burning</u>	<u>Crop straw</u>	<u>Corn, wheat, cotton, sugar cane, potato, peanut, rapeseed, sesame, sugar beet, hemp, rice, soybean</u>
	<u>In-field crop residue</u>	<u>Corn, wheat, cotton, sugar cane, potato, peanut, rapeseed, sesame, sugar beet, hemp, rice, soybean</u>
	<u>Forest</u>	<u>Evergreen Needleleaf Forest</u>
		<u>Evergreen Broadleaf Forest</u>
		<u>Deciduous Needleleaf Forest</u>
		<u>Deciduous Broadleaf Forest</u>
		<u>Mixed Forest</u>
		<u>Closed Shrublands</u>
		<u>Open Shrublands</u>
		<u>Evergreen Needleleaf Forest</u>
	<u>Grassland</u>	<u>Woody Savannas</u>
		<u>Savannas</u>
		<u>Grasslands</u>

I	II	III
<u>Domestic combustion</u>	<u>Firewood</u>	<u>Firewood</u>
	<u>Livestock excrement</u>	<u>Cattles</u>
		<u>Horses</u>

		Donkeys
		Mules
		Camels
	Crop straw	Corn, wheat, cotton, sugar cane, potato, peanut, rapeseed, sesame, sugar beet, hemp, rice, soybean
In-field burning	Crop straw	Corn, wheat, cotton, sugar cane, potato, peanut, rapeseed, sesame, sugar beet, hemp, rice, soybean
	Grassland fire	Grassland fire
		Tropics
		South subtropical zone
		Central subtropical zone
		North subtropical zone
	Forest fire	Warm temperate zone
		Temperate zone
		Cold temperate zone
		Tibetan region

Table 2. ~~Indoor Straw domestic~~ and ~~outdoor-in-field straw-crop residue~~ burning proportions of each province.

Province	Straw indoor <u>domestic</u> burning ratio	<u>In-field</u> crop residue <u>outdoor</u> -burning burning ratio	Province	Straw indoor <u>domestic</u> burning ratio	<u>In-field</u> crop residue <u>burning ratio</u> outdoor -burning ratio
Beijing	0.0923 ^a	0.096 ^b	Hubei	0.283 ^j	0.197 ^o
Tianjin	0.42 ^a	0.165 [*]	Hunan	0.4 ^c	0.2 ^c
Hebei	0.35 [*]	0.165 [*]	Guangdong	0.17 [*]	0.197 [*]
Shanxi	0.45 ^c	0.2 ^c	Guangxi	0.2226 ^k	0.2273 ^k
Inner Mongolia	0.338 [*]	0.246 [*]	Hainan	0.45 ^c	0.2 ^c
Liaoning	0.396 ^e	0.2 ^c	Chongqing	0.4922 ^l	0.1211 ^l
Jilin	0.3 ^c	0.259 ^f	Sichuan	0.45 ^c	0.2 ^c
Heilongjiang	0.26 [*]	0.5 [*]	Guizhou	0.35 ^m	0.2 ^c
Shanghai	0.2 ^c	0.148 [*]	Yunnan	0.2 ^c	0.1 [*]
Jiangsu	0.3 ^g	0.225 ^g	Xizang	0.338 ^d	0.148 ^d
Zhejiang	0.3 [*]	0.3 [*]	Shaanxi	0.338 ^d	0.159 ^o
Anhui	0.29 ^h	0.319 [*]	Gansu	0.338 ^d	0.159 ^o
Fujian	0.3 ^c	0.188 ⁱ	Qinghai	0.338 ^d	0.159 ^o
Jiangxi	0.23 [*]	0.2 ^c	Ningxia	0.338 ^d	0.159 ^o
Shandong	0.45 ^c	0.2 ^c	Xinjiang	0.143 ⁿ	0.137 ⁿ
Henan	0.3 ^c	0.2 ^c			

^a Fang et al. (2015). ^b Zhao et al. (2015). ^c Tian et al. (2011). ^d Bao et al. (2014). ^e Chang et al. (2012). ^f Liu et al. (2010). ^g Wang and Zhao (2011). ^h Qin and Ge (2012). ⁱ Huang (2012a). ^j Liu et al. (2014). ^k Li et al. (2013a). ^l Li et al. (2013b). ^m Zhang et al. (2015). ⁿ Hou et al. (2013). ^o EPD (2014).

^{*} The result from our questionnaire.

Table 3. Residue-to-production ratio (N_k), dry matter fraction (D_k) and combustion efficiency (CE_k) of crop straw used in this study.

Crops	N_k	D_k	CE_k
Corn	1.269 ^a	0.87	0.92
Wheat	1.3 ^b	0.89	0.92
Cotton	3 ^b	0.83	0.9
Sugar cane	0.3 ^c	0.45	0.68
Potato	0.5 ^d	0.45	0.68
Peanut	1.5 ^b	0.94	0.82
Rapeseed	1.5 ^d	0.83	0.9
Sesame	2.2 ^d	0.83	0.9
Sugar beet	0.1 ^b	0.45	0.9
Hemp	1.7 ^e	0.83	0.9
Rice	1.323 ^a	0.89	0.93
Soybean	1.6 ^d	0.91	0.68

Crops	N_k	D_k^f	CE_k^f
Corn	1.269 ^a	0.87 ^f	0.92 ^f
Wheat	1.3 ^b	0.89 ^f	0.92 ^f
Cotton	3 ^b	0.83 ^f	0.9 ^f
Sugar cane	0.3 ^c	0.45 ^f	0.68 ^f
Potato	0.5 ^d	0.45 ^f	0.68 ^f
Peanut	1.5 ^b	0.94 ^f	0.82 ^f
Rapeseed	1.5 ^d	0.83 ^f	0.9 ^f
Sesame	2.2 ^d	0.83 ^f	0.9 ^f
Sugar beet	0.1 ^b	0.45 ^f	0.9 ^f
Hemp	1.7 ^e	0.83 ^f	0.9 ^f
Rice	1.323 ^a	0.89 ^f	0.93 ^f
Soybean	1.6 ^d	0.91 ^f	0.68 ^f

^a Zhang et al. (1990). ^b Bi et al. (2010). ^c Han et al. (2002). ^d NATESC (1999). ^e Gao et al. (2009). ^f He et al. (2015).

Table 4. Forest and grassland biomass fuel loadings in each province, biomass and combustion efficiency of each province.

Province	Climatic zone	Forest biomass ^a	Combustion efficiency ^a
Hainan	Tropics	348	0.2
Guangdong	South-subtropical zone	178	0.2
Guangxi		178	0.2
Yunnan	South-subtropical zone, Central-subtropical zone	138	0.2
Fujian	Central-subtropical zone	143	0.2
Jiangxi		143	0.2
Chongqing		143	0.2
Zhejiang	Central-subtropical zone, North subtropical zone	121	0.2
Guizhou		121	0.2
Sichuan	Central-subtropical zone, Tibetan region	118	0.2
Shanghai	North-subtropical zone	98	0.2
Anhui		98	0.2
Hubei		98	0.2
Hunan		98	0.2
Jiangsu	North-subtropical zone, Warm temperate zone	77	0.2
Beijing	Warm temperate zone	55	0.1
Tianjin		55	0.1
Hebei		55	0.1
Shanxi		55	0.1
Shandong		55	0.1
Henan		55	0.1
Shaanxi		55	0.1
Xinjiang		106	0.1
Inner Mongolia	Temperate zone	157	0.1
Liaoning		157	0.1
Jilin		157	0.1
Heilongjiang		157	0.1
Gansu		157	0.1
Ningxia		157	0.1
Tibet	Tibetan region	121	0.2

Province	Biomass fuel loadings (g/m ²)				
	Needleleaf Forest ^{a,d}	Broadleaf Forest ^{a,e}	Mixed Forest ^{a,f}	Shrublands ^{a,g}	Grassland ^{a,h}
Heilongjiang	8140	7610	7875	1387	180
Jilin	9340	10710	10025	1387	140
Liaoning	2620	8250	5435	1387	160
Inner Mongolia	8140	4470	6305	1387	90
Gansu	8900	6630	7765	1500	90
Ningxia	6910	6280	6595	1386	50
Qinghai	8800	5430	7115	1545	110

<u>Shaanxi</u>	<u>3730</u>	<u>7550</u>	<u>5640</u>	<u>1442</u>	<u>100</u>
<u>Xinjiang</u>	<u>14410</u>	<u>3060</u>	<u>8735</u>	<u>1387</u>	<u>70</u>
<u>Tibet</u>	<u>13990</u>	<u>6490</u>	<u>10240</u>	<u>2007</u>	<u>60</u>
<u>Beijing</u>	<u>1560</u>	<u>6750</u>	<u>4155</u>	<u>1387</u>	<u>170</u>
<u>Hebei</u>	<u>2480</u>	<u>5150</u>	<u>3815</u>	<u>1388</u>	<u>150</u>
<u>Henan</u>	<u>1550</u>	<u>5560</u>	<u>3555</u>	<u>1388</u>	<u>140</u>
<u>Shandong</u>	<u>1280</u>	<u>5660</u>	<u>3470</u>	<u>1387</u>	<u>130</u>
<u>Shanxi</u>	<u>3640</u>	<u>4790</u>	<u>4215</u>	<u>1387</u>	<u>130</u>
<u>Tianjin</u>	<u>0</u>	<u>6760</u>	<u>3380</u>	<u>1378</u>	<u>160</u>
<u>Anhui</u>	<u>1690</u>	<u>10360</u>	<u>6025</u>	<u>2447</u>	<u>140</u>
<u>Hubei</u>	<u>1680</u>	<u>8060</u>	<u>4870</u>	<u>1573</u>	<u>160</u>
<u>Hunan</u>	<u>2080</u>	<u>10650</u>	<u>6365</u>	<u>3471</u>	<u>150</u>
<u>Jiangxi</u>	<u>1820</u>	<u>9370</u>	<u>5595</u>	<u>3699</u>	<u>140</u>
<u>Fujian</u>	<u>2910</u>	<u>9700</u>	<u>6305</u>	<u>3773</u>	<u>160</u>
<u>Guangdong</u>	<u>2060</u>	<u>8970</u>	<u>5515</u>	<u>3702</u>	<u>140</u>
<u>Hainan</u>	<u>4810</u>	<u>9220</u>	<u>7015</u>	<u>3739</u>	<u>150</u>
<u>Jiangsu</u>	<u>2630</u>	<u>5530</u>	<u>4080</u>	<u>1371</u>	<u>120</u>
<u>Shanghai</u>	<u>3060</u>	<u>9250</u>	<u>6155</u>	<u>1371</u>	<u>110</u>
<u>Zhejiang</u>	<u>1710</u>	<u>10500</u>	<u>6105</u>	<u>3682</u>	<u>160</u>
<u>Chongqing</u>	<u>8090</u>	<u>9900</u>	<u>8995</u>	<u>3010</u>	<u>170</u>
<u>Guangxi</u>	<u>2110</u>	<u>9280</u>	<u>5695</u>	<u>3142</u>	<u>150</u>
<u>Guizhou</u>	<u>2210</u>	<u>11410</u>	<u>6810</u>	<u>3431</u>	<u>150</u>
<u>Sichuan</u>	<u>8090</u>	<u>9900</u>	<u>8995</u>	<u>3006</u>	<u>170</u>
<u>Yunnan</u>	<u>5760</u>	<u>14510</u>	<u>10135</u>	<u>3534</u>	<u>150</u>

* ~~Tian et al. (2003)~~.^a Fang et al.(1996,1998). ^b Hu et al. (2006). ^c Pu et al. (2004). And all the biomass here calculated using the aboveground biomass density.

^d Needleleaf forest including needleleaf deciduous forest and needleleaf evergreen forest. ^e Broadleaved forest including broadleaved deciduous forest and broadleaved evergreen forest. ^f The biomass of mixed forest is the mean of needleleaf forest and broadleaved forest. ^g Shrublands including Closed Shrublands and Open Shrublands. ^h Grassland including

Woody Savannas, Savannas and Grasslands.

Table 5. Emission factors used in the estimation of ~~indoor~~ domestic biomass burning emissions.

Material	SO ₂	NO ₂	PM ₁₀	PM _{2.5}	NMVOC	NH ₃	CO	EC	OC	CO ₂	CH ₄	Hg	
	g/kg*											ng/g*	
Domestic burning	Corn	1.33 ^h	1.86 ^{a,h,h}	7.39 ^h	6.87 ^h	7.34 ^h	0.68 ^h	82.37 ^{a,b,h,i,m}	0.95 ^a	2.25 ^a	1491 ^{b,i}	3.91 ^{b,m}	7.94 ^{n,o}
	Wheat	1.2 ^{a,h}	1.19 ^{a,b,h,i}	8.86 ^b	8.24 ^h	9.37 ^h	0.37 ^h	136.46 ^{a,b,h,i,m}	0.42 ^a	3.46 ^f	1246.7 ^{b,i}	8.3 ^b	11.09 ^{n,o}
	Cotton	0.53 ^{d,k,e,g,h}	2.49 ^a	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.82 ^a	1.83 ^a	963.42 ^b	6.08 ^b	3.12 ^{n,o}
	Sugar cane	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Potato	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Peanut	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	4.82 ^{n,o}
	Rape	1.36 ^h	1.65 ^f	13.73 ^h	12.77 ^h	7.97 ^h	0.52 ^h	133.5 ^f	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Sesame	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Sugar beet	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Hemp	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	121.7 ^{b,h}	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	6.5 ^{n,o}
	Rice	0.48 ^h	1.92 ^{a,h,i}	6.88 ^h	6.4 ^h	8.4 ^h	0.52 ^h	79.7 ^{a,b,f,h}	0.49 ^a	2.01 ^a	1147.4 ^{a,b,i}	4.8 ^b	5.56 ^{n,o}
	Soybean	0.53 ^{d,k,e,g,h}	1.12 ^{d,e,f,g}	7.69 ^b	7.15 ^h	8.82 ^{b,j}	1.3 ^{b,e}	80.7 ^f	0.51 ^{d,e,g,c}	2.21 ^{d,e,c}	963.42 ^b	6.08 ^b	4.48 ^{n,o}
	Feces	0.28 ^h	0.58 ^h	8.84 ^h	7.15 ^h	3.13 ^h	1.3 ^h	19.8 ^h	0.53 ^g	2.2 [*]	1060 ^g	4.14 ^g	-
	Firewood	0.4 ^{e,g}	1.49 ^{a,f,h}	5.66 ^{b,i}	5.22 ^{h,d}	3.13 ^j	1.3 ^e	48.25 ^{b,f,h}	1.49 ^c	1.14 ^c	1445.2 ^{b,m}	2.48 ^{b,m}	7.2 ^{n,o}

Note: Lowercase letters indicate the data source.

5 Sources are from the following: ^aCao et al. (2008). ^bWang et al. (2009). ^cLi et al. (2009). ^dReddy and Venkataraman (2002). ^eAndreae and Merlet (2001). ^fTang et al. (2014). ^gTian et al. (2011). ^hEPD (2014). ⁱCao et al. (2004). ^jWei et al. (2008). ^kTurn et al. (1997). ^lZhang et al. (2008). ^mZhang et al. (2000). ⁿChen et al. (2013). ^oZhang et al. (2013a).

* The unit of emission factor.

Table 6. Emission factors used in the estimation of open outdoor-biomass burning emissions.

	Material	SO ₂	NO _x	PM ₁₀	PM _{2.5}	NM VOC	NH ₃	CO	EC	OC	CO ₂	CH ₄	Hg
		g/kg											
		ng/g											
Open burning	Corn	0.44 ^{a,b,c}	4.3 ^{a,b,c}	11.95 ^c	11.7 ^{b,c}	10 ^b	0.68 ^{b,c}	53 ^{a,c,h}	0.3 ^{b,h}	4.35 ^{b,h}	1350 ^{b,h}	4.4 ^b	7.94 ^o
	Wheat	0.85 ^{a,b,c}	3.3 ^{a,b,c}	7.73 ^c	7.58 ^c	7.5 ^{b,c}	0.37 ^b	55.8 ^{a,b,c,d}	0.37 ^{b,h}	3.9 ^{b,h}	1390 ^{b,h}	3.4 ^b	11.09 ^o
	Cotton	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	1.3 ⁿ	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	3.12 ^o
	Cane	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	11.02 ^d	1 ⁿ	40.08 ^d	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	6.5 ^o
	Potato	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	6.5 ^o
	Peanut	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	4.82 ^o
	Rape	0.53 ^{c,f,h,g}	1.12 ^g	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	34.3 ^g	0.23 ^g	1.08 ^g	1410 ^b	3.9 ^b	6.5 ^o
	Sesame	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	6.5 ^o
	Beet	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	6.5 ^o
	Hemp	0.53 ^{c,f,h,g}	3.16 ^{c,f,h,g}	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	1.3 ⁿ	66.1 ^{b,c,i}	0.42 ^b	3.3 ^b	1410 ^b	3.9 ^b	6.5 ^o
	Rice	0.53 ^c	1.42 ^{c,g}	5.78 ^c	5.73 ^{c,g,h}	7.25 ^{c,d}	0.53 ^{b,c}	46.03 ^{d,g,h}	0.16 ^{g,h}	2.03 ^{g,h}	1393 ^b	3.9 ^b	5.56 ^o
	Soybean	0.53 ^{c,f,h,g}	1.08 ^g	6.93 ^c	6.79 ^c	9.5 ^{c,f,i}	0.53 ^{b,c}	32.3 ^g	0.13 ^g	1.05 ^g	1410 ^b	3.9 ^b	4.48 ^o
	Evergreen Needleleaf Forest	1 ^p	1.8 ^p	13.1 ^s	12.7 ^r	28 ^r	3.5 ^r	118 ^r	0.2 ^r	7.8 ^t	1514 ^r	6 ^r	113 ^{u,o}
	Evergreen Broadleaf Forest	0.45 ^r	2.6 ^r	12.8 ^r	10.2 ^r	24 ^r	0.76 ^r	92 ^r	0.5 ^r	4.7 ^r	1643 ^r	5.1 ^r	113 ^{u,o}
	Deciduous Needleleaf Forest	1 ^p	3 ^p	13.1 ^s	12.7 ^r	28 ^r	3.5 ^r	118 ^r	0.2 ^r	7.8 ^t	1514 ^r	6 ^r	113 ^{u,o}
	Deciduous Broadleaf Forest	1 ^p	1.3 ^r	12.8 ^r	12.3 ^r	11 ^r	1.5 ^r	102 ^r	0.6 ^r	9.2 ^r	1630 ^r	5 ^r	113 ^{u,o}
	Mixed Forest	1 ^p	1.3 ^r	12.8 ^r	12.3 ^r	14 ^r	1.5 ^r	102 ^r	0.6 ^r	9.2 ^r	1630 ^r	5 ^r	113 ^{u,o}
	Closed Shrublands	0.68 ^r	3.9 ^r	8.5 ^r	7.9 ^r	4.8 ^r	1.2 ^r	68 ^r	0.5 ^r	6.6 ^r	1716 ^r	2.6 ^r	80 ^{v,o}
	Open Shrublands	0.68 ^r	3.9 ^r	8.5 ^r	7.9 ^r	4.8 ^r	1.2 ^r	68 ^r	0.5 ^r	6.6 ^r	1716 ^r	2.6 ^r	80 ^{v,o}
	Woody Savannas	0.68 ^r	3.9 ^r	8.5 ^r	7.9 ^r	4.8 ^r	1.2 ^r	68 ^r	0.5 ^r	6.6 ^r	1716 ^r	2.6 ^r	80 ^{v,o}
	Savannas	0.68 ^r	2.8 ^r	9.9 ^r	6.3 ^r	9.3 ^r	0.5 ^r	59 ^r	0.4 ^r	2.6 ^r	1692 ^r	1.5 ^r	80 ^{v,o}
	Grasslands	0.68 ^r	2.8 ^r	9.9 ^r	6.3 ^r	9.3 ^r	0.5 ^r	59 ^r	0.4 ^r	2.6 ^r	1692 ^r	1.5 ^r	80 ^{v,o}

Note: Lowercase letters indicate the data source.

Sources are from the following: ^a Li et al. (2015). ^b Li et al. (2007^c). ^c EPD (2014). ^d Zhang et al. (2013b). ^e Tian et al. (2011). ^f Wang and Zhang (2008). ^g Tang et al. (2014). ^h Ni et al. (2015). ⁱ Streets et al. (2003). ^j Andreae and Merlet (2001). ^k Chang ~~et al.~~ and Song (2010). ^l Christian et al. (2003). ^m Kanabkaew and Nguyen (2011). ⁿ Chen et al. (2013). ^o Zhang et al. (2013a). ^p Andreae and Rosenfeld (2008). ^r Akagi et al. (2011). ^s Song et al. (2009). ^t McMeekin et al. (2008). ^u Friedli et al. (2003). ^v Streets et al. (2005).

*The unit of emission factor.

Table 7. Biomass burning emission inventory in the 31 provinces or municipalities of China in 2012.

Province	SO ₂	NO _x	PM ₁₀	PM _{2.5}	NMVOC	NH ₃	CO	EC	OC	CO ₂	CH ₄	Hg
	unit:Gg											unit:Mg
Beijing	0.5	1.9	6.9	6.5	4.8	1.2	58	1.3	1.9	1507	3.1	0.01
Tianjin	1.2	3.1	11.6	10.9	10.4	1.3	116	1.4	3.7	2136	6.6	0.01
Hebei	21.4	52.8	200.2	188.6	178.7	21.4	2023	22.7	65.9	36308	115.4	0.22
Shanxi	9.4	22.9	83.5	78.8	74.9	8.1	777	8.7	27.5	14668	43.9	0.09
Inner-Mongolia	16.1	45.2	217.5	204.9	154.5	24.1	1309	16.8	65.3	32278	103.6	0.16
Liaoning	13.8	40.7	144.3	136.1	128.2	17.4	1277	18.0	42.5	27369	72.4	0.14
Jilin	16.5	54.3	179.6	171.5	165.6	15.7	1395	14.7	58.3	29529	84.7	0.16
Heilongjiang	30.0	117.5	397.4	383.3	395.4	32.5	2878	22.8	132.1	65619	200.7	0.36
Shanghai	0.3	0.8	3.1	3.0	3.6	0.2	33	0.2	1.1	566	2.1	0.00
Jiangsu	14.7	39.7	154.5	146.8	167.0	12.4	1614	10.2	52.6	27527	102.2	0.16
Zhejiang	3.8	12.4	48.2	45.7	48.2	6.1	451	5.5	13.6	9986	28.6	0.05
Anhui	19.7	56.5	210.1	199.9	209.9	19.7	2046	17.6	71.4	38539	127.7	0.23
Fujian	3.0	10.7	40.6	38.1	36.4	6.3	387	6.4	10.5	8905	22.9	0.04
Jiangxi	8.0	27.4	105.0	99.0	102.7	14.3	998	13.4	28.6	22445	62.0	0.10
Shandong	34.7	77.9	304.3	287.4	296.6	25.2	3318	24.9	108.9	50493	192.2	0.33
Henan	33.1	82.1	313.0	296.5	301.3	26.6	3294	26.0	112.7	52896	194.5	0.35
Hubei	13.2	40.7	158.2	149.0	147.6	19.9	1530	19.1	44.5	31167	91.5	0.16
Hunan	15.6	51.6	199.2	187.4	198.0	24.1	1949	22.8	54.0	39478	118.5	0.19
Guangdong	5.9	21.1	78.9	74.2	70.0	12.8	726	12.5	21.0	17619	43.4	0.08
Guangxi	8.1	29.1	105.6	100.0	108.0	15.2	1001	12.6	31.8	21300	61.5	0.11
Hainan	1.4	5.0	19.1	17.9	17.7	3.0	191	2.9	5.1	4032	11.0	0.02
Chongqing	5.5	15.6	61.3	57.4	58.7	7.4	619	7.3	17.0	11564	35.6	0.06
Sichuan	19.3	53.0	212.1	199.6	206.4	22.1	2115	21.0	63.2	38192	125.1	0.23
Guizhou	6.4	19.5	74.7	70.1	62.9	10.4	679	10.8	19.8	14944	38.9	0.08
Yunnan	8.8	27.6	108.7	101.4	90.7	17.2	972	17.4	31.8	22370	54.0	0.20
Tibet	3.0	15.4	40.6	37.6	24.8	5.4	305	2.4	26.5	7554	14.8	0.30
Shaanxi	8.5	22.2	86.5	81.1	71.6	11.1	832	12.0	25.9	16701	47.0	0.10
Gansu	6.2	15.9	66.6	62.5	52.6	8.1	579	7.8	20.1	11814	35.2	0.07
Qinghai	0.9	2.3	10.5	9.8	7.2	1.2	84	0.9	3.6	1683	5.2	0.02
Ningxia	1.6	4.1	14.9	14.1	14.8	1.2	144	1.2	5.1	2515	8.5	0.02
Xinjiang	6.2	21.8	71.5	67.5	64.9	9.8	682	8.5	23.7	13596	39.5	0.08

<u>Total</u>	<u>336.8</u>	<u>990.7</u>	<u>3728.3</u>	<u>3526.7</u>	<u>3474.2</u>	<u>401.2</u>	<u>34380</u>	<u>369.7</u>	<u>1189.5</u>	<u>675299</u>	<u>2092.4</u>	<u>4.12</u>
--------------	--------------	--------------	---------------	---------------	---------------	--------------	--------------	--------------	---------------	---------------	---------------	-------------

5

10

15

Table 8. Uncertainty ranges of different pollutants in emission estimates (min, max). (Unit for emission estimate: Gg)

Species	Emission estimate	Uncertainty ranges *	Previous study
			Street et al., 2003
<u>SO₂</u>	<u>337</u>	<u>(-54%, 54%)</u>	<u>(-245%, 245%)</u>
<u>NO_x</u>	<u>991</u>	<u>(-37%, 37%)</u>	<u>(-220%, 220%)</u>
<u>PM₁₀</u>	<u>3728</u>	<u>(-7%, 6%)</u>	
<u>PM_{2.5}</u>	<u>3527</u>	<u>(-13%, 1%)</u>	
<u>NMVOC</u>	<u>3474</u>	<u>(-9%, 9%)</u>	<u>(-210%, 210%)</u>
<u>NH₃</u>	<u>401</u>	<u>(-49%, 48%)</u>	<u>(-240%, 240%)</u>
<u>CO</u>	<u>34380</u>	<u>(-4%, 4%)</u>	<u>(-250%, 250%)</u>
<u>EC</u>	<u>370</u>	<u>(-61%, 61%)</u>	<u>(-430%, 430%)</u>
<u>OC</u>	<u>1190</u>	<u>(-20%, 19%)</u>	<u>(-420%, 420%)</u>
<u>CO₂</u>	<u>675299</u>	<u>(-3%, 3%)</u>	
<u>CH₄</u>	<u>2092</u>	<u>(-9%, 9%)</u>	<u>(-195%, 195%)</u>
<u>Hg</u>	<u>0.00412</u>	<u>(-31%, 32%)</u>	

* 95% confidence interval.

Table 8. Uncertainty ranges of different pollutants in emission estimates (min, max). (Unit for emission estimate: Gg)

Species	Emission estimate	Uncertainty ranges [±]	Previous study
			Streets et al. (2003)
SO ₂	333	(=76%, 75%)	(=245%, 245%)
NO _x	972	(=62%, 24%)	(=220%, 220%)
PM ₁₀	3676	(=37%, 80%)	
PM _{2.5}	3479	(=54%, 34%)	
VOC	3430	(=55%, 21%)	(=210%, 210%)
NH ₃	396	(=78%, 76%)	(=240%, 240%)
CO	33988	(=41%, 47%)	(=250%, 250%)
EC	367	(=103%, 37%)	(=430%, 430%)
OC	1152	(=78%, 24%)	(=420%, 420%)
CO ₂	665989	(=38%, 43%)	
CH ₄	2076	(=55%, 12%)	(=195%, 195%)
Hg	0.00365	(=26%, 51%)	

[±] 95% Confidence Interval.




Figure 1. Regression analysis between firewood consumption at province resolution and (1) rural population, (2) gross agricultural output, and (3) timber yield, respectively.

Note: It is referred by circles, crosses and triangles, respectively. The regression equation of each figure is provided in the top, middle and bottom, respectively.




Figure 2. Contribution of different source to the total biomass burning emissions in China, 2012.




Figure 3. Contributions of 12 crop straw types burning to total pollutant straws burning emissions for various species.




Figure 4. Contributions of different biomass sources to the emission in each province (Gg).

Note: The numbers 1–12 represent the species $\text{SO}_2 \times 10$, NO_x , VOC , $\text{NH}_3 \times 10$, $\text{EC} \times 10$, OC , $\text{CO} / 10$, PM_{10} , $\text{PM}_{2.5}$, $\text{CO}_2 / 100$, CH_4 and $\text{Hg} \times 1000000$, respectively.






Figure 5. Contributions of different crop straw types to the emission in each province (Gg).


Note: The numbers 1–12 represent the species $\text{SO}_2 \times 10$, NO_x , VOCNMVOC , $\text{NH}_3 \times 10$, $\text{EC} \times 10$, OC , $\text{CO}/10$, PM_{10} , $\text{PM}_{2.5}$, $\text{CO}_2/100$, CH_4 and $\text{Hg} \times 1000000$, respectively.




(a) The annual emissions of PM_{2.5} in each county



(b) PM_{2.5} emissions intensities per unit area



(c) PM_{2.5} emissions intensities per capita



(d) The number of annual PM_{2.5} emissions within different range of county resolution

Figure 6. Biomass emission inventory at county resolution and intensity (PM_{2.5}).





Figure 7. Gridded distribution of PM_{2.5} annual emissions.



5 Figure 8. Monthly variation of different biomass sources emission for each chemical species.




Figure 9. Monthly variation of different biomass sources emission for PM_{2.5} emissions in different regions.




Figure 10. Comparison of the emissions inventory derived by this study with the emissions estimated by previous research.