

***Interactive comment on* “Sensitivities of modelled water vapour in the lower stratosphere: temperature uncertainty, effects of horizontal transport and small-scale mixing” *by* Liubov Poshyvailo et al.**

Liubov Poshyvailo et al.

l.poshyvailo@fz-juelich.de

Received and published: 23 March 2018

We thank the referee for the detailed review and for very helpful comments. We give a point-by-point reply below, where the reviewer comments are repeated in italics. The positions of the corrected sentences in the revised version are noted in the brackets, and the revised text is also given in the quotation marks point-by-point below.

[Printer-friendly version](#)

[Discussion paper](#)



General remarks

This discussion paper addresses the impact of two important yet poorly understood processes controlling water vapour in the UTLS: horizontal transport and small-scale mixing. The CLaMS model is well suited to study this problem. The results help to constrain parametrizations of these processes in climate models, and thus the scientific merit of this paper is significant... This paper is suitable for publication in ACP after consideration of the specific comments and suggestions provided in detail below.

Thank you very much for this positive comment. In the revised version all comments have been taken into account, particularly we improved the discussion part, presentation of figures and formulation of the statements throughout the text related to the remarks of the Reviewer #1.

Major Specific Comments

1. *...I strongly suggest that the study focus on the impacts of horizontal transport and small-scale mixing only...*

Thank you for this suggestion. However, Referee #2 pointed out that the discussion of the reanalysis datasets/tropopause temperature is also important and interesting and should not be removed, as *it consolidates some previous results and, in my opinion, helps to put the significance of the other results presented here into context.* So, we prefer to leave the discussion about the reanalysis TTL temperatures and the impact on LS H₂O. Also we added some references regarding TTL processes and their impact on LS H₂O distribution (p2, L22).

"...Related to the mean upward transport, the TTL includes the region of very low temperatures around the cold-point tropopause, where the moist tropospheric air is freeze-dried to stratospheric values (Brewer, 1949). Thus, the tropical cold-point temperatures control the amount of H₂O, which enters the stratosphere

(e.g., Wang et al., 2015; Kim and Alexander, 2015). The dehydration occurs as a result of the slow upward and large-scale horizontal motion of air in this region (Holton and Gettelman, 2001), where the nucleation and sedimentation of ice crystals take place, which in essence is a microphysical process controlled by TTL temperatures. The freezing is sensitive not just to large-scale TTL temperatures, but also to microphysical processes controlling the ice crystal number densities, particle size distribution, and fall speed. Several studies focused on the modelling of the detailed cloud microphysical processes (e.g., Jensen and Pfister, 2004; Jensen et al., 2005,2012). Other recent papers have examined the effect of cloud microphysical processes on the humidity of the TTL and stratosphere using cloud models of varying complexity..."

2. *The discussion of the effects of the transport barriers (p12-14) could be improved.*

a *...how you are able to discern the direction of transport from these experiments...*

We agree that this point was not discussed appropriately enough. Here we changed the order of the description, from Fig.6a to Fig.6c (p12, L13 – p14, L22), and we added an explanation regarding Fig.6b.

BAR-15 represents suppressed horizontal transport from the subtropics into the tropics and vice versa. Similarly, in-mixing of mid- and high-latitude air and transport from the subtropics to extratropics is presented with BAR-35.

"...Similarly, in-mixing of mid- and high-latitude air (see BAR-35) has a very small impact on tropical mean H₂O, in agreement with the findings of Ploeger et al. (2012). In contrast, transport from the subtropics into the tropics has a strong effect. Suppressing such transport by applying a barrier at 15° N/S (BAR-15) changes the PDF substantially, as evident from the difference between the simulation BAR-15 and the reference cases. The isolation of the tropics due to the lack of horizontal transport in the BAR-15 simulation (all the way from the surface to 600 K) between equator and subtropics

Printer-friendly version

Discussion paper



(both ways) causes dry air at the equator. Thus, with the barrier at 15° N/S the fraction of dry air at the equatorial region increases. The comparison of BAR-15 with BAR-35 shows that transport from the subtropical region into the tropics increases H₂O..."

b *Also, the relationship to the AoA results seems inconsistent at times...*

We agree, that the discussion was to sketchy. So we, added more explanation regarding it (p15, L1).

"...The pure transport effects of horizontal exchange between tropics and mid-latitudes are evident from mean age of air (AoA), the mean transit time for air through the stratosphere for the different model experiments with horizontal transport barriers. Figure 7 shows CLaMS calculations of the AoA for the reference case (Fig. 7a), simulation with transport barriers in the subtropics at 35° N/S (Fig. 7b) and the absolute difference between them (Fig. 7c). These horizontal transport barriers at 35° N/ effectively isolate the tropical pipe from the in-mixing of older stratospheric air from mid-latitudes... A similar result has been recently found by Garny et al. (2014). Furthermore, Garny et al. (2014) presents a nice explanation of the recirculation process, describing recirculation as a process when an air parcel enters the tropical stratosphere and travels along the residual circulation to the extratropics, where it can be mixed back into the tropics, and thus recirculates along the residual circulation again. In this way, the age of air of the parcels increases steadily while performing multiple circuits through the stratosphere..."

c *Please also clarify what you mean on p14, L1-2: "The fact that his drying occurs only with..."*

With both, BAR-15 and BAR-35, recirculation is suppressed, as the region of the extratropics is isolated from the tropics with the barriers. This suppresses the recirculation of extratropical air into the tropics (p15, L28).

"...The fact that this drying occurs only with transport barriers at 15° N/S and

Printer-friendly version

Discussion paper



not with barriers at 35° N/S, shows that it is not related to the suppression of recirculation of aged air from mid-latitudes, which has been affected by methane oxidation. In fact, processes in the subtropics (e.g., monsoon circulations) have a strong effect in moistening the global stratosphere, and suppressing these processes in BAR-15 causes drying. ..."

- d *The second paragraph of p14 states that horizontal transport exports dry (moist) air out of the tropics in the winter (summer). This is followed by a statement that "the entire annual cycle of LS H₂O in the NH extratropics is related to horizontal transport out of the subtropics". There is some disconnect between these two statements that need to be clarified.*

Thank you for pointing this unclear formulation out. We changed the formulation of the statement (p16, L5). Also we added some changes to the Fig.9, we think this will add more clarity to the explanation.

"...Therefore, during winter, horizontal transport exports dry air out of the tropics into the NH, and, during summer, moist air. Consequently, the entire annual cycle of the H₂O in the NH extratropical LS is related to horizontal transport from low latitudes, as argued by Ploeger et al. (2013). The boreal summer maxima are related to monsoonal circulations and transport out of the tropics, along the eastern and western flanks (Randel and Jensen, 2013)..."

3. *Please reconcile the seemingly contradicting results between Fig. 6c vs. Figs. 8acd and Figs. 9acd. Fig. 6c suggests that the lower stratosphere in the NH extratropics is wetter with BAR-15 than with BAR-35, but the opposite seems to be true in Figs. 8 and 9.*

It is not straightforward to compare Fig. 6c with Figs. 8acd and Figs. 9acd. This is due to the fact that Figs. 8,9 only show the mean value for 2011 while Fig. 6 shows the distribution of all H₂O values. Therefore, it is only possible to compare the mean value of H₂O from the distributions in Fig. 6 (as presented by the

[Printer-friendly version](#)[Discussion paper](#)



vertical lines) with Figs. 8acd, 9acd. These values are around 5 ppmv for both BAR-15 and BAR-35, and agree well between all figures so that no contradiction occurs.

4. *p18, L3 - p19, L8: The descriptions and explanations provided in this section are confusing. For example, although it states (on p18, L3-4) that a clear response to mixing is found in the lower stratosphere below 430K (i.e., moistening with increased small-scale mixing), I see significant drying in the UTLS \hat{A} ij350K with the addition of weak small-scale mixing in Figs. 12b and f.*

Thank you for pointing this potential for confusion out. Our sensitivity studies are suitable only for the stratosphere, as the simple H₂O parametrization in CLaMS is adequate only above the cold-point tropopause. Also we added a discussion of this issue, and reformulated some part of the text (p6, L12; p.19, L.9). In Figs. 12,13 (now it is 13,14) we also show the tropopause for better illustration of the separation of stratosphere from the troposphere.

"...Note, that the CLaMS H₂O calculation gives meaningful results only above the tropopause due to the simple parametrization of ice microphysics and not including a convection parametrization. In the stratosphere, however, CLaMS H₂O has been shown to agree well with the observations (e.g., Ploeger et al., 2013)..."

"...A clear response to mixing is found for the LS (below \approx 430 K), which is moistened with increasing small-scale mixing. In the following, we consider total water above the tropical tropopause as an indicator of changes in transport because it is not affected by chemistry (here methane oxidation). As the moistening in the LS below 430 K is also evident in total water, but not in methane and mean age, it is largely related to enhanced diffusive cross-tropopause transport of moist air..."

Also, on p18, L15, it states that an analogous (drying) signal is found in methane (Fig. 13b), but the figure shows moistening?

This paragraph discusses the region above 430K, where the air becomes younger and contains more CH₄ (in SH/NH in the middle and upper stratosphere).

"...A related signal (above 430K in the middle and upper stratosphere) is evident in methane (Fig. 14b) and mean age (Fig. 14f), but not in total water (Fig. 13f)...."

5. *In the description about small-scale mixing in the CLaMS model, the authors cite the Riese et al. (2012) study. Please include a more detailed explanation of how realistic this mixing parametrization might be, and how it relates to observations.*

Thanks for this suggestion to improve the explanation and discussion of CLaMS mixing. We had included already references to some other papers (Konopka 2004, 2005). In the revised version we extended the explanation of CLaMS mixing and in particular added a discussion about how realistic the mixing parametrization in CLaMS (p.5, L27) is.

"...A validation of the CLaMS mixing scheme was presented by Konopka et al. (2005a) in comparison to CRISTA-1 observations. Importantly, the CLaMS mixing parametrization affects both vertical and horizontal diffusivity. Horizontal diffusivity is largely associated with deformation in the horizontal flow. The vertical mixing is mainly related to the vertical shear (Konopka et al., 2004, 2005b)..."

I am assuming that the small-scale mixing refers to both horizontal and vertical diffusion.

Thank you for this comment. And yes, it refers to both horizontal and vertical mixing. This is pointed out more clearly in the revised version (see the answer above).

A recent study by Podglajen et al. (2017) quantified the magnitude of vertical diffusion in the TTL using aircraft observations and found the average diffusivity to be about 0.1 m²s⁻¹ with a strong vertical gradient. Ueyama et al. (2015)

[Printer-friendly version](#)[Discussion paper](#)

found that the “an increase in vertical diffusivity coefficient by 2 orders of magnitude (from 0.001 to 0.1 m²s⁻¹) moistens near the cold point tropopause by approximately 0.5 ppmv”. How do your results based on your small-scale mixing parameterization compare to these results?

Thank you for this comment. We added information about the connection between small-scale mixing and vertical diffusivity in the revised manuscript. Therefore, we estimated the vertical diffusivity for the different CLaMS sensitivity simulations and provide the numbers in a new Fig. 19 (p.25-27).

"...In addition, we estimated the vertical diffusivity coefficient for the TTL for the different model simulations. The result suggests a non-linear response of H₂O to the small-scale mixing in CLaMS (details are considered in Sect. 4)..."

"...Although, it is clear qualitatively that a decreasing critical Lyapunov exponent enhances mixing, it is also desirable, at least for comparison purposes, to quantify this effect. Because of the similarity between the mixing procedure in CLaMS and physical diffusion, the vertical mixing intensity can be quantified by computing the induced vertical diffusivity K_z (in m²/s) (Konopka et al., 2007)... Finally, it should be noted that the mixing in CLaMS induces both vertical and horizontal diffusion. However, given the larger vertical gradients of H₂O compared to horizontal gradients in the UTLS, the impact of small-scale horizontal diffusion is assumed to be much smaller than the impact of vertical diffusion, especially in the tropics..."

...p6 L15, why are there two numbers for the Lyapunov coefficients that lead to good agreement between observations and simulations?

Thank you for this remark. We added some detailed explanation to the text (p7, L5). And the brief explanation is that these two values, 1.2 and 1.5, describe well the stratospheric behaviour when compared to the observations, depending on whether the model is run in a 2D or 3D set-up (1.2 for 2D, 1.5 for 3D).

[Printer-friendly version](#)[Discussion paper](#)

"...provide good agreement between observations and simulation results, as found from comparison of CLaMS simulations with observations from infrared limb-sounding from the research aircraft Geophysica (Khosrawi et al., 2005). In particular, using the value of 1.2 day^{-1} gives a better agreement with observations in the 2D version of CLaMS (Konopka et al., 2003). Furthermore, Konopka et al. (2004, 2005b) showed that the value of $\lambda_c = 1.5 \text{ day}^{-1}$ (corresponding to the critical deformation of $\gamma_c = 1.5$) for the chosen horizontal resolution and time step here, turns out to be optimal for the 3D version of CLaMS. Even for such a small difference in the small-scale mixing strengths, annual mean H_2O concentrations in the extratropical LS differs by about 10-15% (Riese et al., 2012; McKenna et al., 2002a). In our study we use a value of $\lambda_c = 1.5 \text{ day}^{-1}$ for the reference run, 2.0 day^{-1} to represent weak mixing, and 1.0 day^{-1} for modelling strong mixing to cover the range of realistic small-scale mixing strength..."

6. *Some of the terminology needs to be better described. For example, please define what you mean by stratospheric H_2O (e.g., p14, L1: "global stratosphere becomes substantially drier (up to about 1 ppmv)". Is this a vertically integrated quantity averaged from 90S to 90N?*

Thank you for these remarks. This explanation refers to the comparison between Figs.8ad. Hence, the statement concerned the maximum local difference. These maximum differences of H_2O in the stratosphere can reach up to 1 ppmv. We improved the explanation (p.15, L.27).

"...Without such transport from the subtropics (Fig. 8a, 8d), the stratosphere becomes substantially drier (maximal differences in the stratosphere are up to about 1 ppmv)..."

Also, the term "subtropics" is used rather liberally. Sometimes it refers to barriers at 15N/S and other times it refers to the barriers at 35N/S, which can be very confusing. Please clarify your definition and be consistent throughout the text.

[Printer-friendly version](#)[Discussion paper](#)

We improved the explanation and clarified the use of the BAR-terminology in the revised version (p6, L31). In particular, we explain how both barriers suppress the impact of the subtropics, depending on which transport is considered.

"...The two types of barriers, BAR-15 and BAR-35, are located at the edge of the subtropics. BAR-15 is located at the equatorward edge and BAR-35 at the poleward edge of the subtropics. So, both of them inhibit the transport from the subtropics. BAR-15 suppresses horizontal transport from the subtropics into the tropics, and BAR-35 suppresses transport from the subtropics to the extratropics...."

Along the same line, the barrier is sometimes defined at a specific latitude (e.g., "at 15S/S", p14 L2) and other times within a range of latitudes (e.g., "between 10 and 30 S/N", p13 L23).

Thank you for this remark regarding the terminology. The barriers are always 10 degrees in width, centred at 0, 15, 35 degrees. We tried to remove all ambiguous formulations throughout the paper (p6, L29).

"...The transport barriers are defined in the model and centred at the given latitude. Their thickness is 10° in latitude (to inhibit diffusive mixing transport), and the barriers extend from the ground to 600 K potential temperature..."

- 7. The Introduction lacks a discussion on the role of convection in lower stratospheric (or TTL) H₂O. It is briefly mentioned in the Discussion Section as a possible explanation for the biases in JRA-55 simulations, but convection is an important feature that needs to be discussed early on. The role of convectively-injected ice is discussed in the Introduction, but its relative importance on stratospheric humidity is still a topic of debate.*

Thank you for this comment. We agree that convection is an important process controlling stratospheric water vapour. We included a discussion of convection in the introduction in the revised manuscript (p3, L3).

[Printer-friendly version](#)[Discussion paper](#)

"...Sublimation of ice, injected by deep convection, has also been argued to be an important factor for the H₂O budget of the tropical LS (e.g., Avery et al., 2017; Jensen and Pfister, 2004). Convection affects the transport of water and ice and influences the temperatures over the convective region, in turn affecting dehydration (e.g., Fueglistaler et al., 2009). The predominant impact of convection has been shown to moisten the TTL by up to 0.7 ppmv at 100 hPa level and even more below this level (e.g., Ueyama et al., 2014, 2015). Similarly, Schoeberl et al. (2014) argued that an increase of convection will increase stratospheric H₂O and tropical cirrus around the cold-point tropopause. At higher levels in the TTL, however, the moistening effect of convection appears very weak (e.g., Schiller et al., 2009)..."

8. *Figures 6 and 10: The vertical lines are very difficult to see because of the size of the plots and the overlapping lines (e.g., black solid line in panel c?). Vertically stacked panels like Fig. 17 are much easier to see.*

Thank you for this comment. We changed the Figures 6 and 10 (now it is Fig.11) as proposed by the Reviewer.

Minor Specific Comments

1. *It would be helpful to provide the approximate magnitudes ($\hat{\Delta}l_{ij}$ ppmv) of the effects of horizontal mixing and small-scale mixing on stratospheric H₂O in the Abstract and Conclusions (e.g., in the sentence at the top of p22, "Furthermore, our results suggest that the NH subtropics are a critical source region of moisture for the global stratosphere, . . ."). It is worth pointing out that the impacts of these processes are of same order of magnitude as the impact of temperature difference of $\hat{\Delta}l_{ij}2K$.*

Thank you for the comment. We changed the text according to the Reviewer's suggestions in the Abstract and the Conclusions (p1, L14).

"...Comparison of tropical entry H₂O from the sensitivity 15° N/S barrier simulation and the reference case shows differences of up to around 1 ppmv... For the sensitivity simulation with varied mixing strength differences in tropical entry H₂O between the weak and strong mixing cases amount to about 1 ppmv, with small-scale mixing enhancing H₂O in the LS...."

2. *Dessler et al. (2013) paper may not be the most appropriate reference for the first sentence of the Introduction, unless you're specifically emphasizing the feedback effects of stratospheric water vapor. In addition to the papers by Forster and Shine, I would consider the Shindell (2001) paper instead.*

We changed the reference in the revised version according to the Reviewer's suggestion (p2, L1).

"...Stratospheric water vapour (H₂O) is a crucial factor for global radiation, as it cools the stratosphere and warms the troposphere (e.g., Forster and Shine, 1999, 2002; Shindell, 2001; Nowack et al., 2015)..."

3. *p 2, L13-14: In your discussion of the rapid transport from the tropics to mid latitudes above the subtropical jets, are you distinguishing between transports in the "tropically controlled transition region" vs. the shallow branch of the Brewer-Dobson circulation, as in Rosenlof et al. (1997)?*

Thank you for this comment regarding clarification of the terminology. We do not distinguish between transports in the "tropically controlled transition region" vs. the shallow branch of the Brewer-Dobson circulation. In our opinion both processes are strongly related, and the transport in the tropically controlled transition region may even belong to the shallow branch of the Brewer-Dobson circulation.

4. *p2, L15-19: You may want to clarify that the dehydration occurs due to the nucleation and sedimentation of ice crystals, which in essence is a microphysical process controlled by TTL temperatures. Hardiman et al. (2015), which is a study*

[Printer-friendly version](#)[Discussion paper](#)

based on global climate models that have difficulty simulating cloud microphysical processes, is not the most appropriate reference here. I would consider citing Jensen et al. (2004,2005, 2012) studies on the modeling of detailed cloud microphysical processes. Several papers have examined the effect of cloud microphysical processes on humidity of the TTL and stratosphere using cloud models of varying complexity (e.g., Schoeberl et al., 2014; Ueyama et al., 2015).

Thank you for these clarifications. We changed the text and referencing according to the suggestions (p2, L22).

"...Related to the mean upward transport, the TTL includes the region of very low temperatures around the cold-point tropopause, where the moist tropospheric air is freeze-dried to stratospheric values (Brewer, 1949). Thus, the tropical cold-point temperatures control the amount of H₂O, which enters the stratosphere (e.g., Wang et al., 2015; Kim and Alexander, 2015). The dehydration occurs as a result of the slow upward and large-scale horizontal motion of air in this region (Holton and Gettelman, 2001), where the nucleation and sedimentation of ice crystals take place, which in essence is a microphysical process controlled by TTL temperatures. The freezing is sensitive not just to large-scale TTL temperatures, but also to microphysical processes controlling the ice crystal number densities, particle size distribution, and fall speed. Several studies focused on the modelling of the detailed cloud microphysical processes (e.g., Jensen and Pfister, 2004; Jensen et al., 2005, 2012). Other recent papers have examined the effect of cloud microphysical processes on the humidity of the TTL and stratosphere using cloud models of varying complexity (e.g., Ueyama et al., 2015; Schoeberl et al., 2014)..."

5. *p3, L3-6: I see now that the discussion of temperature and freeze-drying process appears here. I would move this ("The tropical stratospheric entry H₂O mixing ratios ... temperature and vertical velocity fields.") after the paragraph on TTL transport processes in the previous page.*

[Printer-friendly version](#)[Discussion paper](#)

We moved it to the paragraph above (p2, L31).

"...The tropical stratospheric entry H₂O mixing ratios can be well simulated by the advection through the large-scale temperature field and instantaneous freezing, often described as the "advection-condensation" paradigm (Pierrehumbert and Rocca, 1998; Fueglistaler and Haynes, 2005)..."

6. *p3, L8-10: What exactly are you referring to with respect to the relationship between summer max tropical H₂O and the monsoons? Mixing processes? Deep convection in the monsoon regions? Relatively high tropopause heights?*

This is indeed a good question. The summertime monsoon systems very likely involve all three mentioned processes. The relative strength of these processes, however, is an open question and a topic of current research. Based on our model experiments, we can not estimate which processes dominates in the real atmosphere. We, therefore, rewrote this sentence (p3, L18).

"...The summer maximum of tropical H₂O mixing ratios has been argued to be also related, to some degree, to the subtropical monsoon circulations like the Asian monsoon. However, the strength of this effect and the detailed processes involved (e.g., deep convection, large-scale upwelling) is a matter of debate..."

7. *p 3, L18-24: The discussion of TWV here seems a bit out of place. Could you better tie this to the overall discussion about processes that affect lower stratospheric H₂O?*

Thank you for this suggestion. We shifted this sentence to the paragraph above (p3, L10).

"...Above the TTL, H₂O behaves mainly as a tracer, and the tape recorder signal imprinted at the cold-point tropopause ascends deep into the tropical stratosphere (Mote et al., 1996). At higher altitudes in the stratosphere, methane oxidation results in a chemical source for stratospheric H₂O (e.g., LeTexier et

[Printer-friendly version](#)[Discussion paper](#)

al., 1988; Rohs et al., 2006). As a net result of this oxidation process each methane molecule is converted into approximately two H₂O molecules. Hence, the total water vapour (TWV), $TWV = 2CH_4 + H_2O$, is unchanged by transport in the stratosphere and can be regarded approximately constant (e.g., Dessler et al., 1994; Mote et al., 1998; Randel et al., 1998). Therefore, the sum $2CH_4 + H_2O$ is an important value to indicate the amount of water entering the stratosphere (e.g., Kampfer, 2013). The annual cycle of TTL temperatures (minimum in boreal winter, maximum in summer) is imprinted on H₂O mixing ratios entering the stratosphere, forming the so-called “tape recorder” signal (Mote et al., 1995, 1996)…”

8. *p5, L20: In the sentence, “Thus, we use the fifth year. . .for our further analysis”, make it clear that you are referring to the analysis of the sensitivity simulations.*

We rewrote this sentence.

“...Thus, we use the fifth year of the perpetuum simulation for our further analysis...”

9. *Table 2: What is “PR”? Would a graph be better for illustrating your point?*

Thank you for this remark concerning the terminology. PR-files are the files after each year of perpetuum run. We agree that this terminology is not helpful for the readership and that also giving the differences between all the spin-up years provides not too much insight. The necessary information is simply that after the fourth year the maximum changes are below 1%. This information is now given in the revised manuscript and all other unnecessary information is removed (p6, L19).

“...After the fourth year, the maximum relative change of H₂O mixing ratios between further years of the simulation is very small with the defined resolution and the time step (maximum year to year changes are below 1.0%). Thus, we use the fifth year of the perpetuum simulation for our further analysis...”

Printer-friendly version

Discussion paper



10. *p7, L1: Is "10-15%" an annual mean value?*

Yes, it is an annual mean differences for 2003 (Riese et al. 2012).

"...Even for such a small difference in the small-scale mixing strengths, annual mean H₂O concentrations in the extratropical LS differs by about 10-15% (Riese et al., 2012; McKenna et al., 2002a)..."

11. *Figure 2: What is the reason for showing the annual cycle at 400K? What about the 380K level (Figs. 5 and 16 are based on this level)? A brief statement of the sensitivity of your results to the different potential temperatures in the lower stratosphere may be useful. Also, is the MLS averaging kernel applied to the model data for a more accurate comparison to MLS data?*

Thank you for this questions. We preferred to consider the H₂O distribution at 400K, as the 380K surface may be well located below the tropopause in certain regions *e.g. Asian monsoon) and hence mixes stratospheric and tropospheric characteristic. However, for model and reanalysis intercomparison the level 380K is often used. To have better comparability to these studies, we changed Fig.16 380K now.

For MLS data the averaging kernel was not applied. As shown by Ploeger et al. (2013) the MLS averaging kernel has only a weak effect on H₂O at middle and low latitudes in the lower stratosphere, while having a strong effect at high latitudes. As the main focus of this paper is an intercomparison of different model simulations, we prefer to show the full model resolution without smearing out structures with the averaging kernel (p.11, L5).

"... Oscillations in MLS H₂O at high latitudes are a known effect of the broad averaging kernel (Ploeger et al., 2013). At low latitudes the effects of the MLS averaging kernel on H₂O are much smaller and we do not apply it to the model data here, in order not to smear out the structure in the simulated H₂O..."

Printer-friendly version

Discussion paper



12. *p8, L9-10: "The impact of horizontal transport. . .mixing strength covered." Effects look very similar to me. I would suggest removing this statement to avoid confusion.*

We removed this statement.

13. *Figure 5: There are four temperature contours in panels a and b, but only three contour levels are mentioned in the caption. Devise a better way to label these temperature contours?*

Thank you for this remark. We now added the temperature values in Fig.5 to clarify the plot. The small contours in the Figs.5ab, without the labels belong to 193 K (what should be obvious from the plot and labelling). Also we changed the caption for Fig.5.

14. *p11, L7: Please be more specific about what you mean by "agree slightly better" (quantify).*

We changed the text in the revised version (p12, L5).

"...Overall, regarding the global H₂O distributions and maps in the LS, CLaMS modelling results with ERA-Interim are drier when compared to JRA-55, resulting from lower TTL temperatures in ERA-Interim. The agreement between CLaMS based on ERA-Interim and JRA-55 with the observations strongly depends on the considered region and season. And it is not possible to conclude from our analysis which reanalysis results in simulated H₂O in the best agreement with the observations..."

15. *p11, L14: I think you mean to say "... show that the interhemispheric transport is rather unimportant..."*

It is changed in the revised version.

Printer-friendly version

Discussion paper



"...the insignificant difference between the reference (REF) and an equatorial transport barrier (BAR-0) simulations shows that the interhemispheric transport is rather unimportant for tropical mean H₂O mixing ratios..."

16. *p13, L11: Where is the "double peak structure"?*

We added a short explanation to the text.

"...In the tropics, the age distribution in Fig. 7b shows a weak double peak structure up to about 500 K, indicating that the subtropics are regions of particularly fast transport, likely related to subtropical processes like monsoon circulations..."

17. *Figures 12 and 13: Why are these two figures separated? It makes more sense to combine the two and make a 4 x 4 panel figure: TWV, H₂O, 2CH₄, AoA. Does the sum of Fig. 12a and Fig. 13b equal Fig. 12e?*

Thank you for the remark. We had the two figures combined in a first draft version. But, we think that it is confusing to have a 4 x 4 panel Figure. So, we decided to separate it in two plots. In our opinion, it makes sense to separate it in this way (H₂O + TWV; and transport tracers CH₄ + AoA).

18. *Figure 15: The lines in these figures are very difficult to see.*

Thank you for pointing that out. We changed the representation of Figure 15 (now it is Fig. 16).

19. *p20, top: It appears that during boreal summer, weak mixing moistens the Asian monsoon region while very strong mixing moistens the N American monsoon region. Do you have an explanation for this interesting result?*

This is indeed an interesting result. However, we have no good explanation for this hitherto. We will study his point in the future. Additionally, now we added the discussion about this issue into the text (p22, L12).

Printer-friendly version

Discussion paper



"...During boreal winter the SH subtropical jet substantially moistens with increasing mixing, during boreal summer the NH jet moistens. In particular, the moist anomaly of the Asian and American monsoons during boreal summer is affected by small-scale mixing. Without mixing only a weak anomaly occurs in the Asian monsoon, while the moist anomaly in the American monsoon is absent. With increased mixing the Asian monsoon moist anomaly first increases (MIX-weak and REF cases). When mixing becomes very strong (MIX-strong) this behaviour changes and the entire jet region becomes strongly moistened such that the Asian monsoon moist anomaly relative to the entire jet region weakens... Overall, small-scale mixing in the CLaMS simulations and related diffusive upward moisture moisture transport seem crucial for the development of Asian and American monsoon moisture anomalies, and in particular for the American monsoon (where no anomaly occurs without including small-scale mixing)..."

20. *p20, L30: Where do you get "380K"? The explanation you provide is certainly plausible, but do these biases occur in mid and high latitudes where the H₂O biases are observed?*

Thank you for pointing this out. Fig.17 (now it is Fig.18) is done for mid-and high latitudes (we forgot to mention it in the caption), we changed the description in the revised version accordingly. The related discussion in the revised version is improved (p25, L13).

"...This behaviour is the clearest at around 370 K (Fig. 18), but it is also visible at levels below and above (not shown). The different shape of the JRA-55 PDFs, with the peak at much higher mixing ratios, suggests that high H₂O mixing ratios are deposited in the extratropical LS, from potential temperature levels of about 350 K up to at least about 400 K, potentially related to the convective scheme in the reanalysis...."

21. *Figure 17: It would be helpful to add MLS data to these plots.*

[Printer-friendly version](#)[Discussion paper](#)

Thank you for the suggestion. But, we would not like to overload Fig.17 with too many lines. A comparison between CLaMS and MLS data has been made already at an earlier stage in the paper. Here we want just to discuss and compare the reanalysis with CLaMS model results (which were based on the temperature and winds taken from the same reanalysis, ERA-Interim and JRA-55). The aim is to emphasize that the JRA-55 own reanalysis H₂O products have too high values and are different from CLaMS H₂O, which is actually consistent with JRA-55 tropopause temperatures (as it is based on a rather simple dehydration parametrization).

Technical Corrections

Thank you for these detailed corrections. We changed everything regarding to your suggestions.

1. *p6, L15: “. . .our study, . . .”*

Corrected in the revised version.

2. *p8, L7: “. . . tropical (10S-10N) entry of H₂O . . .”*

Corrected in the revised version.

3. *p8, L14: Rewrite this sentence “. . . reaches the values of $\hat{L}_{ij}0.8$ ppmv” as “. . . drying of $\hat{L}_{ij}0.8$ ppmv in (month)”.*

Corrected in the revised version.

4. *p10, L17: “. . . LS H₂O between MLS and CLaMS simulations . . .”*

Corrected in the revised version.

5. *p11, L13; p12, L1: I'm slightly confused by the use of the term "vanishing difference". Are the differences decreasing over time?*

Corrected in the revised version.

6. *p 12, L3: ". . . barrier at 15 N/S changes the PDF . . ."*

Corrected in the revised version.

7. *p12, L5: ". . . (blue line in Fig 6b), . . ."*

Corrected in the revised version.

8. *p12, L7: Rearrange the sentence: "equatorial transport barrier has no effect and cross-equatorial transport is unimportant"*

Corrected in the revised version.

9. *p14, L9: Replace with ". . .very weak drying effect on. . ."*

Corrected in the revised version.

10. *p16, L5: I'm not sure what you mean by "Therefore"*

We removed that sentence, and reformulated the previous statement. Corrected in the revised version.

11. *p16, L12: "H2O (a-d) and total water (e-h)"*

Corrected in the revised version.

12. *p20, L18: "On the contrary, . . ." is a very long sentence.*

Corrected in the revised version.