

Dear editor and referees,

We would like to thank the referees for the valuable comments. There is a major concern that both referees share and we would like to address that in the first place, and then answer separately for each question.

This major concern refers to the calibration process, and not including the 2-way transmission (T_p). From this, it derives the concern of performing the calibration during the event if the aerosol load is high. We must say that the calibration process and obtaining backscatter coefficient profiles are two separate processes but thanks to the referees comments we realized that not including the T_p , the process is less intuitive and harder to explain. We recalculated everything including the T_p calculated with the AOD and values for the calibration factor for each site are provided. The entire methodology section has changed. Each process is described in a subsection (calibration on section 3.1 and inversion and validation on section 3.2 and 3.2.1). The distinction between CL^* and CL is no longer necessary, so equations has been modified accordingly. We hope that this new structure provides a better understanding of the methodology applied.

We revised the manuscript according to the rest of suggestions and we replied to all the concern raised by the referees. Next paragraphs show comments and questions from referees (in red) and our response (in black).

Concerning Referee #1:

The authors state that the desert dust event was unusual. In what sense? Why it was unusual? Please add some comments.

Valenzuela et al., (2012) presented a classification of dust events affecting southern Spain during the period 2005-2010. They established 3 different typical patterns for the dust plumes corresponding with different synoptic scenarios. According to their results, only 22% of the desert dust events followed the same pattern described on Section 4.1.

The intensity of the event is also unusual for that time of the year. For this pattern, Valenzuela et al. (2012) reported a maximum AOD in Granada AERONET station of 0.98, being the pattern with the lower intensity of the three pattern (maximum AOD for the other pattern classifications were 1.6 and 1.4). During the dust event presented in this study, a maximum AOD of 1.77 was measured at Granada AERONET station (see new Fig. 6) which is significantly higher than the value reported by Valenzuela et al. (2012).

Also, there are two recent publications on the same event (Sorribas et al., 2017; Titos et al., 2017) that demonstrates the extraordinariness of the event. In particular, Titos et al. (2017) showed that the PM10 concentrations were very high, with 90% of stations exceeding the PM10 daily limit of 50 ug/m^3 (especially in southern Spain where the average PM10 was above 150 ug m^3). Also, at Montsec, the aerosol light extinction coefficient (from nephelometer + maap) at ground level was the highest measured in this station (Titos et al., 2017).

These references are now included in the reference list and on Section 4 we have included the following information:

“Sorribas et al. (2017) studied the same event and compared it with meteorological parameters, aerosol properties and ozone from historical data sets on a site in southern Spain. They concluded that the event was exceptional because of its unusual intensity, its impact on surface measurements and the month of occurrence. In addition, Titos et al. (2017) also analyzed this event using 250 air quality monitoring stations over Spain to investigate the impact and temporal evolution of the event on surface PM10 levels. They also investigated aerosol optical properties, including attenuated backscatter from ceilometer during the event at Montsec station (one of the station included in ICENET). They concluded that the impact on surface PM10 was exceptional and highlighted the complexity of the event.”

And later on Section 4.1:

“For dust events following a similar pattern that the one described here during the period 2005-2010, Valenzuela et al., (2012) reported a maximum AOD of 0.98 at Granada, which is significantly lower than the maximum measured at Granada station during this event.”

* P4 line 15: I doubt that the telescope field-of-view is 1.8mrad. Please verify and provide the correct value. 1.8mrad is the FoV of so-called x-ceilometers manufactured by Jenoptik, now Lufft.

This was already changed on the previous revision of the manuscript:

“The laser beam divergence is less than 0.3 mrad and the laser backscattered signal is collected on a telescope with a field of view of 0.45 mrad”

* CL and CL* are not really constant. They change over time. There was a discussion within E-Profile and ToProf about the vocabulary. I suggest using calibration value instead of calibration constant.

Calibration value, factor or parameter is used instead of constant. In any case, CL* is no longer used in the new version.

* I dont understand why the authors focus on CL which is systematically biased by ignoring the two-way transmission of particles when their method includes the use of sun photometer measurements in order to estimate the correct lidar ratio. I would understand it in case that there are no co-located sun photometer measurements which is not the case for ICENET. The description of the method should be adapted. Either there are information about AOD and hence CL* could be derived or AOD is not available.

As explained before, the entire methodology section has changed. The two-way transmission is now included.

* p7 lines 3-11: it is missing a description about the treatment of lower 250 or 300m where the overlap correction is very large and often unreliable.

A constant value is considered:

“The first 300 m of the profile are assigned to the value at 300 m to avoid very large overlap correction.”

* the authors highlight that the best calibration values are obtained for low AOD. But then they derive calibration values during a desert dust episode with high AOD. This is very strange for me. For me it would make more sense deriving calibration values before and after the desert dust event at times with low AOD and interpolate them for the desert dust measurements. This would even enable comparisons for ceilometer and lidar during night-time.

This is completely different with the new methodology section. Whenever the AOD matches the integral of a backscattering coefficient profile multiplied by the lidar ratio, a CL value is provided. The CL of a site would be an average value for a long-time series of calibrations. The aim is not to obtain calibrations during the event, but in order to obtain inverted backscattering coefficient profiles, at the reference height, the ratio between RCS and molecular backscatter (Eq. 3) need to match.

* The ceilometers were measuring 5 days. The calibration was only performed on 11 half-hour intervals. It would be good if the method description includes a few sentences how the calibration values for each individual profile were derived. Example: Figure 7 shows that there is a calibration value for each profile.

This has changed with the new methodology section. The calibration factor for each site is an average of a long time series of calibrations. The values for each site are provided on the new Table 2.

* the unit of center of mass is missing. It should be m or km (e.g. p8 lines 23/24, but also on other parts of the manuscript **Minor critical comments**)

The relative difference between the center of mass of each profile is provided, not the center of mass. The center of mass is shown on Fig. 9b (units are meters).

* I found 5 times “Iberian Ceilometer Network (ICENET)”. It is sufficient to explain the abbreviation one time

This has been corrected. Only abstract and introduction indicate both name and acronym.

* P4 line 17: the authors wrote “the overlap is 90% complete at 555 m agl”. The manufacturer provides an individual overlap function for each single ceilometer. I strongly doubt that all 5 ceilometers have the same overlap. I suggest providing a range.

A range is provided in the new manuscript:

“the overlap is 90% complete between 555 and 885 m agl”

* P4 line 35 “Raman-shifted channels” this formulation is a bit sloppy. The channels are not Raman-shifted. **Re-phrase!**

‘Raman channel’ is used instead.

* p5 line 11-12: please add the emitted laser energy P0 in the details of the calibration value.

The emitted laser energy per pulse of the ceilometers is provided on Section 2:

“The energy per pulse is 8.4 μJ with a repetition frequency in the range of 5 – 7 kHz”

* p5 line 23: the rcs signal is also normalized to the number of laser shots which vary from profile-to-profile

The manuscript has been modified:

“the range corrected signal ($RCS(z) = P(z) \cdot z^2$), using an overlap function determined by the manufacturer and corrected for the number of laser shots”

* p5 line 23/24: “and also, the overlap function of the instrument is factory determined.” Although it is correct that the overlap function is determined by the manufacturer, I guess the authors wanted to point out that the overlap correction is already included in the rcs.

That is correct:

“the range corrected signal ($RCS(z) = P(z) \cdot z^2$), using an overlap function determined by the manufacturer and corrected for the number of laser shots”

* p5 line 28: “a region in the ceilometer profile”. Here is missing the depth of the region. On page 7 the authors wrote about 990m. It should be mentioned already here.

We changed the manuscript accordingly:

“we select regions of 990 m with a difference in gradients below 1%.”

* p6 line 9: “low aerosol load, where the transmittance due to particles is close to 1”. Please provide an AOD value what you consider as low. For instance the transmittance is 0.9 for an AOD of 0.1. Is this considered low? Or it is still too high?

With the new structure of the methodology, we think this is no longer an issue.

* p6 line 25 ff “continuously calculating“ Do you really mean continuously or you mean repeatedly? And there are missing some details. I doubt that CL is calculated for every single profile. Do you use a temporal average for improving signal-to-noise ratio?

The section including those lines has changed completely. In any case, references to “continuously” has been changed. Also, with the new description of the calibration process it is clear that profiles are averaged for one hour.

* p7 line 3/4 “In this study 30-min profiles are used for comparison with lidar profiles.” Is the calculation of the calibration value based on 30min too or a longer average is used?

The calibration is performed on one hour averaged profiles. In order to compare with lidar, profiles for the inversion are averaged for 30 minutes.

* p7 line 17: “has to be close to the CL” what is close? Maybe the author could write something like “has to be close to Cl within x%”

This has changed with the new methodology. A range of mean±std of the calibration factor is used.

* p11 line 21: “total attenuated backscatter” what is the difference between attenuated backscatter and total attenuated backscatter?

They are the same, and manuscript has been changed to include ‘total’ in all cases. Total makes reference to particle + molecular.

* table 5: it is obvious when the site, the ceilometer and the sun photometer have the same geographical coordinates, that ceilometer and sun photometer are co-located. Hence it is not needed to write: co-located

Co-located has been removed from the table.

* Figure caption 1: Raman with capital R

Changed.

* Figure 2: middle image: unit of center of mass difference is missing. it’s m or km?

It is a relative difference between the center of mass of both profiles. It has no units.

* Figure caption 3: Monte Carlo instead of Montecarlo

Changed.

* Figure 7: it’s attenuated backscatter or total attenuated backscatter?

It is total attenuated backscatter. We changed the figure caption.

* Figure 9: I suggest adding the sun photometer values in the top panel. As the lidar ratio was derived, it is straight forward to derive the integrated backscatter from sun photometer AOD.

AOD and Ångström exponent are included in the new Figure 6 on section 4.1.

Concerning Referee #2:

1) The title does not clearly reflect the content of the paper. The input of sun photometer data is essential for the proposed method, but the title does not provide any hint that the described method is not usable for standalone ceilometer instruments. Further, the manuscript provides no proof, why the described dust outbreak event is extraordinary.

We modified the title to include a reference to sun-photometer data and we have removed the ‘extraordinary’ from the title:

“Near real time processing of a ceilometer network assisted with sun-photometer data: monitoring a dust outbreak over the Iberian Peninsula”

2) What makes the described event extraordinary? Is it the meteorological situation, intensity, duration or something else? If the event was extraordinary in a certain aspect, this statement should be proved by comparison with typical events. If the extraordinariness of the event cannot be corroborated, the title should be adopted.

Valenzuela et al., (2012) presented a classification of dust events affecting southern Spain during the period 2005-2010. They established 3 different typical patterns for the dust plumes corresponding with different synoptic scenarios. According to their results, only 22% of the desert dust events followed the same pattern described on Section 4.1.

The intensity of the event is also unusual for that time of the year. For this pattern, Valenzuela et al. (2012) reported a maximum AOD in Granada AERONET station of 0.98, being the pattern with the lower intensity of the three pattern (maximum AOD for the other pattern classifications were 1.6 and 1.4). During the dust event presented in this study, a maximum AOD of 1.77 was measured at Granada AERONET station (see new Fig. 6) which is significantly higher than the value reported by Valenzuela et al. (2012).

Also, there are two recent publications on the same event (Sorribas et al., 2017; Titos et al., 2017) that demonstrates the extraordinariness of the event. In particular, Titos et al. (2017) showed that the PM10 concentrations were very high, with 90% of stations exceeding the PM10 daily limit of 50 $\mu\text{g}/\text{m}^3$ (especially in southern Spain where the average PM10 was above 150 $\mu\text{g}/\text{m}^3$). Also, at Montsec, the aerosol light extinction coefficient (from nephelometer + maap) at ground level was the highest measured in this station (Titos et al., 2017).

These references are now included in the reference list and on Section 4 we have included the following information:

“Sorribas et al. (2017) studied the same event and compared it with meteorological parameters, aerosol properties and ozone from historical data sets on a site in southern Spain. They concluded that the event was exceptional because of its unusual intensity, its impact on surface measurements and the month of occurrence. In addition, Titos et al. (2017) also analyzed this event using 250 air quality monitoring stations over Spain to investigate the impact and temporal evolution of the event on surface PM10 levels. They also investigated aerosol optical properties, including attenuated backscatter from ceilometer during the event at Montsec station (one of the station included in ICENET). They concluded that the impact on surface PM10 was exceptional and highlighted the complexity of the event.”

And later on Section 4.1:

“For dust events following a similar pattern that the one described here during the period 2005-2010, Valenzuela et al., (2012) reported a maximum AOD of 0.98 at Granada,

which is significantly lower than the maximum measured at Granada station during this event.”

3) Why is the transmittance due to particles neglected? The authors describe how sun photometer data are used to constrain the Klett-Fernald inversions. Thus, all relevant information for the calculation of T_p is available. Why is it not used?

We agree with the reviewer. This has changed in the new methodology section.

4) The authors shall provide an estimation of the difference between CL^* and CL for different particle optical depths.

With the new approach, there is no CL^* .

5) It would be nice to have an example plot for illustrating the calibration method which is described at page 7. It would be even better to show two examples, one of a clear day and one of a polluted day.

With the new methodology section we hope that it is clear enough and that there is no need to include an additional plot.

6) The manuscript suffers from a major internal conflict. First, it is introduced that the calibration parameter CL can be retrieved only on days with low aerosol load. But in the next part, a period with very high aerosol load is used for demonstration and validation of the method. All CL values which are derived during the dust period will lead to a systematic bias of profiles if they are applied to measurements in clean conditions. The retrieved CL values are systematically too small due to neglecting T_p . To overcome this problem, the authors should derive and present a longer time series of CL values, with clean periods before and after the dust event. According to the theory, only the maximum values of CL in this time series (clean periods) are ok. Those maximum values before and after the dust event should be used for the retrieval of the attenuated backscatter profiles during the event. The lidar profiles can be used for the validation of these ('clean') CL values. But, it makes no sense to derive CL values during the dust event (even if constrained with lidar profiles) without taken into account T_p . All attempts to constrain CL values measured during the dust event should be removed from the manuscript. This includes Figures 2 and 3, and all text below equation 7 and the begin of section 4. Besides, the use of the correlation coefficient and center of mass as measures of goodness of the calibration seems to be of little help. Even if the calibration value is wrong, both profiles should have the same shape (and therefore the same C_{mass} and high R) because they are measured under the same atmospheric conditions.

The calibration and validation of inversion are independent processes and with the new structure of Section 3 (Methodology) we hope there is no doubt about the calibration process (not during the event) and the validation of the inversion during the event.

Figure 2 is one of the most important figures in the manuscript. Especially Figure 2a (modified to match the new methodology section) where it is shown that, at the reference height (Z_{ref}) which is a key parameter for the Klett-Fernald inversion, the ratio between the RCS and molecular backscatter (technically the same as CL in Eq. 3 in the manuscript) has to be similar to a long-term calibration factor calculated with the calibration method proposed. In particular, inversions that present at Z_{ref} this ratio between \pm the standard

deviation of the mean calibration factor have smaller difference in NMB with the lidar profiles (considered the truth) so the long term calibration factor can be used to determine automatically if an inversion is good or not. The center of mass does not vary significantly with good or bad calibrations, which is good because highlights that the center of mass can be obtained directly from the total attenuated backscatter and no inversion is needed. However, the R does vary even the atmospheric conditions are the same (both instruments are measuring the same). The way the inversion algorithm distributes the backscatter coefficient with range depends on Z_{ref} , and if Z_{ref} is not correct, the inverted profiles differ from the lidar profile. On Fig. 2c better R values correspond with the values within the mean $CL_{\pm std}$, and the worst case have a R of about 0.5 (a rejected profile).

7) The link between the two parts of the manuscript (methodology and results) is weak.

With the methodology section this link between sections is stronger.

8) The description of the meteorological situation during the event is quite lengthy while some interesting measurement data are not provided, e.g. ,depolarization profiles from the lidar, time series of optical depth and Angstrom exponent, fine-to-coarse mode fraction etc. from the photometer network.

The intention of this manuscript is to describe the capabilities of the network of ceilometers for the monitoring of singular events such as the dust event described. Thus, the description of the event is aimed to show the path of the dust plumes reaching the Iberian Peninsula but we do not intend to give a full characterization of the event.

In any case, as indicated in previous questions we added some references and extra information about the event on Section 4:

“Sorribas et al. (2017) studied the same event and compared it with meteorological parameters, aerosol properties and ozone from historical data sets on a site in southern Spain. They concluded that the event was exceptional because of its unusual intensity, its impact on surface measurements and the month of occurrence. In addition, Titos et al. (2017) also analyzed this event using 250 air quality monitoring stations over Spain to investigate the impact and temporal evolution of the event on surface PM10 levels. They also investigated aerosol optical properties, including attenuated backscatter from ceilometer during the event at Montsec station (one of the stations included in ICENET). They concluded that the impact on surface PM10 was exceptional and highlighted the complexity of the event.”

And later on Section 4.1:

“For dust events following a similar pattern to the one described here during the period 2005-2010, Valenzuela et al., (2012) reported a maximum AOD of 0.98 at Granada, which is significantly lower than the maximum measured at Granada station during this event.”

9) In general, the description of the event would be more useful if the authors could provide references to other studies about dust over the Iberian Peninsula. How often do events like the described dust outbreak occur at the Iberian Peninsula? What are typical

pathways and meteorological conditions? What are typical optical properties (e.g. optical depth) of the dust? What are typical geometrical properties of the dust layers? What makes this event special compared to others?

Valenzuela et al., (2012) presented a classification of dust events affecting southern Spain during the period 2005-2010. They established 3 different typical patterns for the dust plumes corresponding with different synoptic scenarios. According to their results, only 22% of the desert dust events followed the same pattern described on Section 4.1.

The intensity of the event is also unusual for that time of the year. For this pattern, Valenzuela et al. (2012) reported a maximum AOD in Granada AERONET station of 0.98, being the pattern with the lower intensity of the three pattern (maximum AOD for the other pattern classifications were 1.6 and 1.4). During the dust event presented in this study, a maximum AOD of 1.77 was measured at Granada AERONET station (see new Fig. 6) which is significantly higher than the value reported by Valenzuela et al. (2012).

On Section 4.1 we added the following information:

“For dust events following a similar pattern that the one described here during the period 2005-2010, Valenzuela et al., (2012) reported a maximum AOD of 0.98 at Granada, which is significantly lower than the maximum measured at Granada station during this event.”

10) abstract, the last sentence “... quantitative optical aerosol characterization with ceilometers...” is misleading. Ceilometers can be used for the quantification of the aerosol layering or aerosol load in terms of particle backscatter coefficients, but they cannot be used for characterization. ‘Characterization’ is often used as synonym for ‘detection of aerosol type’ or ‘retrieval of intensive optical properties (like lidar ratio or Angstrom exponent)’. Ceilometers cannot provide this kind of information.

The term ‘characterization’ could be used on very different contexts depending on the instrumentation available, but we understand the referee concern. We used the term ‘monitoring’ in the title and the last sentence of the abstract was changed to:

“Results reveal that it is possible to obtain quantitative optical aerosol properties (particle backscatter coefficient) and discriminate the quality of these retrievals with ceilometers over large areas”

11) p2, l22: It is strange to write about the inadequate quality of satellite products in a paper about aerosol profiles from ceilometers. Certainly, products from ceilometers are very useful, but usually they do not provide profiles of higher quality than space-borne lidars.

We do not question the quality of satellite products and we never intended to suggest that ceilometers are better than satellite products. It is a fact that the temporal and spatial resolution of satellites is, sometimes, inadequate for specific applications. Also, validation of satellite retrievals often requires ground measurements. Those ground measurements do not need to be, and we do not intend them to be, coming from ceilometer. Thus, to avoid any doubt, we modified the sentence:

“The main disadvantage of measurements from space-borne platforms is the low temporal resolution, since the measurements are limited to the satellite passes over a region”

12) p2, l28: There are more relevant references, e.g. Flentje et al. 2010.

We included the reference.

13) introduction: The order of sentences in the last two paragraphs seems to be somewhat randomly. Maybe due to copy-and-paste? An outline to the structure of the manuscript is missing.

We modified the last part of the introduction and included an outline.

14) equation 1: CL* is not a constant in a strict sense. It changes on long time scales, e.g. due to laser aging or window contamination. Better to use ‘parameter’ instead of ‘constant’.

All references to constant has been removed and ‘value’, ‘factor’ or ‘parameter’ is used instead.

15) p7, l29: how can negative CL values be explained? NMB is usually calculated as mean value of the bias profile ($b_{\text{ceil}}(z) - b_{\text{lidar}}(z)$) / $b_{\text{lidar}}(z)$

The reference height is calculated by Rayleigh fit automatically and, due to signal-to-noise ratio, sometimes the window selected to check the slope has, on average, a negative value. This is definitely a bad region for calibration, and therefore is rejected. The problem is when this reference height is automatically selected for the inversion algorithm. If this is the best Zref obtained, the inversion is not going to be valid. It is important to note, one of the findings here is that the calibration factor at Zref need to be around the value of the long-term calibration factor. This is what makes the automatic and unsupervised near-real time inversion possible.

16) Figure 5: This kind of trajectory plot seems to be less informative than the traditional plots (with a map projection and a time-altitude plot below). To which times of observation do the red dots refer to?

This figure is aimed to show the height intervals where African dust may have had an impact at two of the ceilometer network sites. The plots are in the form height vs latitude of back-trajectories, which is not the most common format but it is the simplest and clearer way to show what is intended. Please note that latitude and height are the relevant variables. The southern trajectories arrive from Africa (this is clear for Granada but also holds for Badajoz). The map projection would show the same information but it would be needed to use a color code for the changing heights along each trajectory, which makes the plots not so clear in the end. Time-latitude plots do not give the information we need to show.

The red dot in a trajectory corresponds to the air parcel’s position at 12UTC. When this position is located to the north of the dust activation areas, then no dust is carried along with the air parcel even if that parcel came from northern Africa. This corresponds to the

height intervals depicted in green. Figure caption already indicated the meaning of the red dots (circles), but following the suggestion of the Reviewer we have explicitly added the corresponding time.

17) Figure 6: Is it really necessary to show this figure?

With the detailed description of section 4.2, we agree that the figure is not necessary. We removed it from the manuscript.

18) Figure 7: The labels of the color bar are unreadable. Time axes of the different stations have different tick scales.

We modified the Figure accordingly.

19) Figure 8: Where does the uncertainty of 15% comes from? Please, indicate the measurement times of the individual profiles by vertical lines in figure 7.

It comes from Fig. 2a. As we move away from the mean CL, the NMB increases and this value is used as an estimation of the uncertainty. We also added the vertical lines on Fig 7.

20) Figure 9: Please, add time series of AOD and columnar mean lidar-ratio (whenever Integrated bsc and AOD are available). Also time series of Angstroem exponents and fine-to-corase mode fractions would be interesting.

AOD and Ångström exponent is included in the new Figure 6 is section 4.1.

21) The readability of the text could be improved by splitting some long sentences into shorter ones, e.g.

- * p1, 123+24
- * p1, 125+26
- * p3, 11+2
- * p3, 118-20
- * p5, 128-31

We looked at the manuscript carefully and modified some long sentences.

22) p1, 125: it is not clear to which method the term "this method" refers.

It is said at the beginning of the paragraph:

“In this work we describe a method that uses aerosol optical depth (AOD) measurements from the AERONET network that it is applied for the calibration and automated quality assurance of inversion of ceilometer profiles”

23) p1, 126: the date of the event is described several times in the manuscript as ... on 20 February and lasted until 24 February... -> it would be better to write "between... and .." or "... lasted from .. to..."

We modified that throughout the manuscript

24) p1, l21 + p2 l1: what refers "their" to?

The sentences has been modified:

“The aerosol direct effects depend on the optical properties and spatial and vertical distribution of the aerosol in the atmosphere. In spite of the recent advances on instrumentation that has improved the ability of characterizing key aerosol properties and increase the spatial resolution, the associated uncertainties are still considered to be one of the majors in climate forcing (Boucher et al., 2013).”

25) p2, l6: The terms "in-situ" and "surface measurements" are often used for ground based remote sensing instruments like lidars, in contrast to space-borne instruments. Maybe a term like "measurements of aerosol properties at ground level" could be used instead.

We changed the term to ‘ground level aerosol measurements’.

26) p2, l7: when speaking about a covered area, it would be better to use "Europe" instead of "European Union".

Changed.

27) p2, l10: (and elsewhere in the paper): vertical resolved -> vertically resolved

Changed

28) p2, l31: make complicate -> hinder?

Changed

29) p3, l18-20: the calibration is used to validate the inversion?

Changed

30) p4, l2: comparable -> calibrated?

Changed

31) p4, l4: constraint -> constrain

Changed

32) p5, l16: not overlap -> no overlap

Changed

33) p5, 128: what means "computing the Rayleigh fit"? -> ... particle free regions are determined by comparing the gradient... If the difference is below 1%, we can assume particle free conditions...

We changed the sentence:

“The Rayleigh fit, compares the gradient with altitude (the slope) of both profiles, and looks for a region in the ceilometer profile that has the same trend than the expected molecular profile”

34) p7, 112: ... has the influence of ... -> ...is influenced by...

Changed

35) p10, 126 ... northern African... -> ... northern Africa...

Changed

References:

Sorribas, M., Adame, J.A., Andrews, E. and Yela, Y.: An anomalous African dust event and its impact on aerosol radiative forcing on the Southwest Atlantic coast of Europe in February 2016, *Science of the Total Environment*. 583, 269–279, doi: 10.1016/j.scitotenv.2017.01.064

Titos, G., Ealo, M., Pandolfi, M., Pérez, N., Sola, Y., Sicard, M., Comerón, A., Querol, X., and Alastuey, A.: Spatiotemporal evolution of a severe winter dust event in the western Mediterranean: Aerosol optical and physical properties. *Journal of Geophysical Research: Atmospheres*, 122, 4052–4069, doi:10.1002/2016JD026252

Valenzuela, A., Olmo, F.J., Lyamani, H., Antón, M., Quirantes, A. and Alados-Arboledas, L.: Classification of aerosol radiative properties during African desert dust intrusions over southeastern Spain by sector origins and cluster analysis. *Journal of Geophysical Research*, 117, D06214, doi:10.1029/2011JD016885, 2012.

Near real time processing of ceilometer network assisted with sun-photometer data: monitoring a dust outbreak over the Iberian Peninsula

Alberto Cazorla^{1,2}, Juan Andrés Casquero-Vera^{1,2}, Roberto Román^{1,2}, Juan Luis Guerrero-Rascado^{1,2},
5 Carlos Toledano³, Victoria E. Cachorro³, José Antonio G. Orza⁴, María Luisa Cancillo^{5,6}, Antonio
Serrano^{5,6}, Gloria Titos⁷, Marco Pandolfi⁷, Andres Alastuey⁷, Natalie Hanrieder⁸, Lucas Alados-
Arboledas^{1,2}

¹Andalusian Institute for Earth System Research, IISTA-CEAMA, University of Granada, Junta de Andalucía, Granada, Spain

²Department of Applied Physics, University of Granada, Granada, Spain

10 ³Grupo de Óptica Atmosférica (GOA), Universidad de Valladolid, Valladolid, Spain

⁴SCOLab, Física Aplicada, Universidad Miguel Hernández, Elche, Spain

⁵Department of Physics, University of Extremadura, Badajoz, Spain

⁶Institute of Water Research, Climate Change and Sustainability, IACYS. University of Extremadura, Spain

⁷Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona, Spain

15 ⁸German Aerospace Center (DLR), Institute of Solar Research, Plataforma Solar de Almería, Almería, Spain

Correspondence to: Alberto Cazorla (cazorla@ugr.es)

Abstract. The interest on the use of ceilometers for optical aerosol characterization has increased in the last few years. They
operate continuously almost unattended and are also much less expensive than lidars, hence they can be distributed in dense
networks over large areas. However, due to the low signal-to-noise-ratio it is not always possible to obtain particle backscatter
20 coefficient profiles, and the vast amount of data generated requires an automated and unsupervised method that ensures the
quality of the profiles inversions.

In this work we describe a method that uses aerosol optical depth (AOD) measurements from the AERONET network that it
is applied for the calibration and automated quality assurance of inversion of ceilometer profiles. The method is compared
with independent inversions obtained by co-located multiwavelength lidar measurements. A difference smaller than 15% in
25 backscatter is found between both instruments. This method is continuously and automatically applied to the Iberian
Ceilometer Network (ICENET) and a case example during an unusually intense dust outbreak affecting the Iberian Peninsula
between 20 - 24 February 2016 is shown. Results reveal that it is possible to obtain quantitative optical aerosol properties
(particle backscatter coefficient) and discriminate the quality of these retrievals with ceilometers over large areas. This
information has a great potential for alert systems and model assimilation and evaluation.

30 1 Introduction

Atmospheric aerosol is one of the main responsible factors of climate radiative forcing through multiple processes including
aerosol-radiation and aerosol-cloud interactions (IPCC, 2014). The aerosol direct effects depend on the optical properties and

spatial and vertical distribution of the aerosol in the atmosphere. In spite of the recent advances on instrumentation that has improved the ability of characterizing key aerosol properties and increase the spatial resolution, the associated uncertainties are still considered to be one of the majors in climate forcing (Boucher et al., 2013).

In this sense, the implementation of observational networks is crucial for spatial characterization of aerosol properties. Ground level aerosol measurement networks represent key tools in the study of aerosol radiative forcing. These observational networks provide surface measurements distributed over large areas, e.g. the Global Atmospheric Watch, GAW (GAW, 2011) and ACTRIS (www.actris.eu) for Europe. On the other hand, one of the recognized instruments for the retrieval of column-integrated aerosol properties is the robotic sun and sky photometer that is used in the global Aerosol Robotic NETwork (AERONET; Holben et al. 1998; Dubovik et al. 2006). Lidar systems are well known active remote sensing instruments for the vertically resolved characterization of aerosol optical and microphysical properties (Winker et al., 2003). GAW Atmospheric Lidar Observation Network (GALION) has emerged as an initiative of the GAW aerosol program (GAW, 2008). Its main objective is to provide the vertical component of the aerosol distribution through advanced laser remote sensing in a network of ground-based stations. Among other networks, GALION includes the European Aerosol Research Lidar Network (EARLINET) that provides vertical aerosol profile observations over Europe based on 27 instruments in 16 countries (Pappalardo et al. 2014), the Micro Pulse Lidar Network, MPLNET (Welton et al. 2001), and the Latin American Lidar Network, LALINET (Guerrero-Rascado et al. 2016).

In order to obtain a larger spatial coverage than ground-based networks, in the last few years some space missions have been promoted focusing on aerosol measurements from satellites, e.g. the Lidar in Space Technology Experiment, LITE (McCormick, 1997) and the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation, CALIPSO (Winker et al. 2003). The main disadvantage of measurements from space-borne platforms is the low temporal resolution, since the measurements are limited to the satellite passes over a region.

The usefulness of vertically resolved aerosol characterization has been proven by monitoring dust outbreaks (e.g. Guerrero-Rascado et al, 2008; 2009; Cordoba-Jabonero et al. 2011; Bravo-Aranda et al. 2015; Preissler et al. 2011; 2013; Granados-Muñoz et al. 2016; Valenzuela et al. 2017), biomass burning plumes (e.g. Alados-Arboledas et al. 2011; Ortiz-Amezcuca et al. 2016), or the volcanic ash plume from the Eyjafjallajökull eruption on April 2010 (Navas-Guzmán et al., 2013; Pappalardo et al. 2013; Sicard et al., 2012). Precisely, this singular event caused aviation problems and drew the attention to the use of ceilometers for vertically resolved aerosol characterization (Flentje et al., 2010).

The complexity of lidar systems requires staff to be trained in their operation and the analysis procedures are not fully automated in many stations. In this sense, continuous operation of lidar systems is not feasible for most stations. In addition, economic and operational costs hinder the implementation of dense lidar networks. On the other hand, ceilometers are one-wavelength (near infrared) lidars with simple technical specifications (eye-safe low pulse energy and high pulse repetition frequencies) allowing for unattended and continuous operation. Originally designed for cloud base determination, their performance has been improved in the last few years. It has been shown their capabilities for determining planetary boundary layer (e.g., Wiegner et al., 2006; Münkler et al., 2007; Haeffelin et al., 2011; Pandolfi et al., 2013), detection and forecast of

fogs (Haefelin et al., 2016), and recent efforts have been conducted to quantify the aerosol optical information that can be derived from ceilometers (Frey et al., 2010; Heese et al., 2010, Wiegner et al., 2014).

The main advantage of the use of ceilometers for aerosol characterization is, on the one hand, the automatic and much simpler operation compared to lidars and, on the other hand, the possibility of installing them distributed over large areas.

5 Meteorological services such as those in Germany, France, the Netherlands or United Kingdom are deploying ceilometers networks to cover their national territories with the objective of reaching a spatial density of nearly one device every 100km (e.g. Haij and Klein-Baltink 2007; Flentje et al. 2010). Due to a dense number of instruments and continuous measurements, operative networks need an automated processing and a protocol that ensures the quality of the data.

10 In this sense, two programs in Europe are dealing with the use of automated lidars and ceilometers for aerosol and cloud properties characterization. The COST Action ES1303 TOPROF (TOwards operational ground based PROFiling with ceilometers, doppler lidars and microwave radiometers for improving weather forecasts) aims in one of its working groups at better characterizing the parameters that can be derived from ceilometer measurements and related uncertainties. At the same time, E-PROFILE, a program of EUMETNET (EUropean METeorological services NETwork), focuses on the harmonization of ceilometer measurements and data provision across Europe.

15 In this study we present the implementation of procedures to manage a regional ceilometer network for aerosol characterization over the Iberian Peninsula, the Iberian Ceilometer Network (ICENET). An automatic calibration procedure is applied to the ceilometers and this calibration is used to validate the elastic inversion automatically applied to the profiles. This method uses additional aerosol optical depth (AOD) information during the calibration for the quality assurance of the data.

20 All processes can be performed unattended and in near real-time with the objective of obtaining reliable vertically resolved aerosol optical properties. This information is especially useful on strong events, such as mineral dust outbreaks, volcanic plumes, severe biomass burning episodes or contamination episodes. Thus, the aerosol information obtained can be potentially used as an alert system for aviation or weather services, or feed models for assimilation and validation in near-real time.

25 The capabilities of this distributed network are explored by characterizing an unusually intense dust outbreak affecting the Iberian Peninsula on 20 -24 February 2016 and a multi-wavelength (MW) Raman lidar is used to validate the retrievals from ceilometers.

30 Next section describes the Iberian Ceilometer Network and related instrumentation used in this study. Section 3 presents the methodology, including the calibration of ceilometers (Section 3.1) and the use of the calibration for inversion validation, and a validation with an independent lidar system (Sections 3.2 and 3.2.1 respectively). The results are presented on Section 4, with a description of the dust event (Sections 4.1 and 4.2) and retrievals from ceilometers (Section 4.3). Finally, conclusions are presented on Section 5.

2 Instrumentation: the Iberian Ceilometer Network

An initiative of the Atmospheric Physics Group of the University of Granada has been the coordination of a network of ceilometers (ICENET) combined with sun photometers for the characterization of atmospheric aerosol with the objective of obtaining reliable vertically resolved aerosol optical properties in near real-time. On a first stage, the goal is obtaining the total attenuated backscatter for all ceilometers in the network, i.e. to obtain **calibrated** output from ceilometers and, on a second stage, applying an inversion algorithm to the ceilometer profiles in order to obtain the particle backscatter coefficient. All sites of this new network have a co-located AERONET CE318 sun-sky photometer (Cimel Electronique) that is used to constrain the ceilometers calibration and inversion retrievals. In addition, the high performance lidar system MULHACEN located at the EARLINET Granada station is used as an independent validation of the inversions. This nested approach combining high performance systems like those operated in EARLINET and the distributed ceilometer plus sun photometer is an example of synergy among active and passive remote sensing observations in ACTRIS research infrastructure (www.actris.eu).

Figure 1 shows a map of the ceilometer distribution over the Iberian Peninsula, and Table 1 presents the characteristics of each site.

All sites operate a Jenoptik (now Lufft) CHM15k-Nimbus ceilometer and have a co-located AERONET sun photometer, except Montsec station (MSA) that has the photometer 770 m above the ceilometer and at a horizontal distance of 2 km approximately (Titos et al, 2017). The ceilometer at Murcia (UMH) was not operative during the outbreak studied in this work. The CHM15k is a ceilometer that operates with a pulsed Nd:YAG laser emitting at 1064 nm. The energy per pulse is 8.4 μJ with a repetition frequency in the range of 5 – 7 kHz. The laser beam divergence is less than 0.3 mrad and the laser backscattered signal is collected on a telescope with a field of view of 0.45 mrad. The signal is detected by an avalanche photo diode in photon counting mode. Complete overlap of the telescope and the laser beam is found about 1500 m above the instrument (Heese et al., 2010). According to the overlap function provided by the manufacturer, the overlap is 90% complete **between 555 and 885 m agl**. The vertical resolution used is 15 m and the maximum height recorded is 15360 m agl. Ceilometers at Granada (UGR), Tabernas (PSA) and Valladolid (UVA) operate at a temporal resolution of 15 s while ceilometers at Montsec (MSA) and Badajoz (UEX) operate at a temporal resolution of 1 min.

The process of calibration for ceilometers described on the next section is assisted with AOD data from co-located AERONET stations. All sun photometers near the ceilometers belong to the Iberian network for aerosol measurements (RIMA), a regional network federated to AERONET. This means that all instruments are routinely calibrated following the same protocol and the data are quality controlled. The sun-photometer provides solar extinction measurements at 340, 380, 440, 675, 870, 936 and 1020 nm, allowing for computing the AOD at these wavelengths (except 936 nm). The AOD uncertainty ranges from ± 0.01 in the infrared-visible to ± 0.02 in the ultraviolet channels (Holben et al., 1998). For comparison with the ceilometers the AOD is extrapolated to 1064 nm by the Ångström law (Ångström, 1964) using the AOD measurements at 870 and 1020 nm. Level 1.5 AERONET data, which are automatically cloud-screened and delivered in near real time, are used in this analysis.

At UGR station a multi-wavelength Raman lidar system (MULHACEN) is used for validation of the ceilometer inversions. The upgraded LR331-D400 (Raymetrics Inc.) operated at IISTA-CEAMA (Andalusian Institute for Earth System Research) is part of EARLINET since April 2005. This lidar system is a ground-based, six wavelength system with a pulsed Nd:YAG laser. The emitted wavelengths are 355, 532 and 1064 nm with output energies per pulse of 60, 65 and 110 mJ, respectively. It has elastic backscatter channels at 355, 532 and 1064 nm and Raman channels at 387 (from N₂), 408 (from H₂O) and 607 nm (from N₂). Full overlap is reached around 1220 m agl although the overlap is complete at 90% between 520 and 820 m agl (Navas-Guzmán et al., 2011). Appropriate overlap corrections are derived following the procedure of Wandinger et al. (2002).

3 Methodology

Elastic lidars and ceilometers principle of measurement is the same and retrieving optical properties in both systems follow the lidar equation (the dependency with the wavelength has been omitted for simplicity since it is always the same in ceilometers):

$$P(z) = C_L \cdot \frac{O(z)}{z^2} \beta(z) \cdot T^2(z). \quad (1)$$

where $P(z)$ is the backscattered power received in the telescope from a distance z , C_L is a parameter that depends on the geometry and characteristics of the instrument and universal constants, and the term z^2 accounts for the acceptance solid angle of the receiver optics with the distance to the laser. The backscattered signal collected by the telescope depends on the overlap between the laser beam and the telescope field of view, and the degree of overlap is quantified by $O(z)$, ranging from 0, if there is no overlap, to 1, if overlap is complete. $\beta(z)$ is the atmospheric backscatter coefficient and $T(z)$ estimates the atmospheric transmittance of the laser signal (squared due to travel back and forth). Also, both properties can be split into contributions of particles and molecules ($\beta(z) = \beta_m(z) + \beta_p(z)$; $T(z) = T_m(z) \cdot T_p(z)$) (Fernald, 1984).

On Eq. (1) the only properties depending on the medium are $\beta(z)$ and $T(z)$. Thus, the atmospheric attenuated backscatter is defined as:

$$\beta_{att}(z) = \beta(z) \cdot T^2(z) \quad (2)$$

3.1 Ceilometer calibration

The ceilometers used in this study provides as output the range corrected signal ($RCS(z) = P(z) \cdot z^2$), using an overlap function determined by the manufacturer and corrected for the number of laser shots. Therefore, the only parameter that needs to be addressed is C_L .

Wiegner et al. (2014) describes a method to find the C_L^{\blacksquare} parameter in ceilometers, commonly referred as ceilometer calibration. This method compares the RCS from the ceilometer in a particle-free region with the molecular attenuated backscatter that can be calculated using Rayleigh theory. The Rayleigh fit compares the gradient with altitude (the slope) of both profiles, and looks for a region in the ceilometer profile that has the same trend than the expected molecular profile. In this study, we select regions of 990 m with a difference in gradients below 1%. Thus, in that region or reference height (z_{ref}), C_L^{\blacksquare} can be calculated:

$$C_L^{\blacksquare}(z_{ref}) = \frac{RCS(z_{ref})}{\beta_m(z_{ref}) \cdot T_m^2(z_{ref}) \cdot T_p^2(z_{ref})} \quad (3)$$

At this reference height, the backscattering is only due to molecules. The transmittance due to molecules (T_m) can be easily determined from Rayleigh theory but the transmittance due to particles (T_p) is unknown. However, if a col-located sun-photometer is available, T_p can be calculated:

$$T_p^2(z_{ref}) = e^{-2 \cdot AOD} \quad (4)$$

using the AOD at 1064 nm.

When trying to automate the calibration process, the main problem is that z_{ref} must be a particle free region and, due to the low signal to noise ratio, finding z_{ref} is not always possible. In some cases, the region might be a non-particle free region that, on average, follows the molecular trend (they have a similar gradient). Also, we might find several regions that meet the criteria, but it is complicated to discriminate automatically which one is the most appropriate one. A way to ensure that the z_{ref} selected is a molecular region is by applying Klett-Fernald inversion algorithm (Klett, 1981, 1985; Fernald et al., 1972; Fernald, 1984) as follow.

First, we need to determine the z_{ref} . Thus, after finding z_{ref} , the Klett-Fernald inversion is applied. Heese et al. (2010) and Wiegner et al. (2012; 2014) showed the capabilities of ceilometers applying this inversion algorithm studying a few cases. Using the AOD measurements, the lidar ratio (Lr) of the inversion can be adjusted. This can be done matching the integral of the particle extinction coefficient profile (i.e. particle backscatter coefficient profile multiplied by the Lr) with the AOD. Wiegner et al. (2012) applied this procedure to a Jenoptik CHM15kx ceilometer obtaining reasonable values for the Lr.

In summary, the calibration process follows the next steps:

1. First, temporal averaging of the profiles is performed (hourly averages are used for the calibration). The first 300 m of the profile are assigned to the value at 300 m to avoid large overlap correction.
2. Second, for each profile a set of potential z_{ref} is obtained by comparing the profiles of the RCS and β_m , which is obtained from a standard atmosphere profile scaled to ground temperature and pressure. The slopes are calculated over a 990 m window. All regions with slope differences below 1% are selected.

3. For each z_{ref} , and Lr from 20 to 80 sr, Klett-Fernald inversion is applied and the resulting profile for the backscattering coefficient is integrated and multiplied by the Lr and compared to the AOD. The pair z_{ref} and Lr that minimizes the difference between the integral of the particle extinction coefficient profile (i.e. particle backscatter coefficient profile multiplied by the Lr) with the AOD is selected.

5 4. Finally, C_L^{\blacksquare} is calculated using Eq. (3) if the minimum difference calculated on step 3 is below 10%.

C_L^{\blacksquare} calculated with this method uses Eq. (4) to calculate T_p^2 . In the case of MSA the sun-photometer is 770 m above the ceilometer and the T_p^2 calculated would not be representative of the entire column. However, MSA is a remote mountain site and the effect of accounting for T_p^2 values obtained using the AOD measured from 770 m above the ceilometer can be considered as negligible.

10 Thus, the total attenuated backscatter can be calculated applying the following equation:

$$\beta_{att}(z) = \frac{RCS(z)}{C_L(z_{ref})} \quad (5)$$

Calibration values can be used individually or averaged over a period of time. Table 2 shows a mean calibration factor (\pm standard deviation) calculated using this method for all sites in ICENET for the period 1 May 2014 to 1 May 2016.

3.2 Ceilometer inversion

Total attenuated backscatter obtained applying the calibration factor allows the comparison between ceilometers since the signal is corrected for instrument characteristics. Also, a long-time series of the calibration allows determining possible problems or degradation of the systems. However, the total attenuated backscatter is influenced by transmission, so in order to be able to monitor and compare singular events at multiple sites, the backscattering coefficient is more appropriate. Section 3.1 showed that it is possible to apply Klett-Fernald inversion to ceilometer data, but the challenge is to determine automatically, without human supervision, if the inversion is correct or not.

A common step between the calibration proposed on Section 3.1 and Klett-Fernald inversion is finding a reference height (Z_{ref}). At the Z_{ref} selected for the inversion with Rayleigh fit, applying Eq. (3), we can obtain a value that has to be close to the C_L^{\blacksquare} calculated for the instrument on a longer period of time. If a simultaneous AOD measurement is available, the calibration process itself provides the inverted backscattering coefficient profile (steps 3 and 4 of the calibration process) and the inversion can be marked as valid or invalid based on the value of the C_L^{\blacksquare} compared to a long term C_L^{\blacksquare} value. If no simultaneous AOD measurement is available (e.g during nigh time or partially cloudy skies), an approximation of the AOD needs to be used in order to apply Eq. (4). In this case, an interpolated value or an averaged value can be used.

30 The next section quantifies the differences between these backscattering coefficient inversions and the inversions calculated independently with a multi-wavelength Raman lidar.

3.2.1 Lidar – ceilometer comparison

During the dust outbreak affecting the Iberian Peninsula between 20 – 24 February 2016, the multi-wavelength lidar operated on 22 February between 7:30 to 14:00 UTC and on 23 February between 8:00 and 13:30 UTC. Elastic inversions using Klett – Fernald were applied to 30-min average profiles at 1064 nm using a fixed lidar ratio of 50 sr. Thus, a total of 24 particle backscatter coefficient profiles were obtained. Coherence of the inversion at 1064 nm was checked against Klett – Fernald and Raman methods at 355 and 532 nm. The resolution of the multi-wavelength lidar (7.5 meters) has been downscaled to 15 meters for the comparison with the ceilometer.

The ceilometer elastic inversion, using Klett – Fernald method, was also applied to 30-min average profiles for the same period; a total of 15 profiles were successfully inverted (a reference height was found automatically). The calibration factor at the reference height was calculated using the average AOD for the entire dust event. If negative C_L^{\square} are discarded, a total of 11 profiles are comparable with lidar inversions.

Each one of the derived C_L^{\square} values at the reference height selected for the inversion are compared with long-term C_L^{\square} calculated for the Granada station ceilometer over a long period of time (see Table 2) classifying the situation according to statistical parameters measuring the agreement between the ceilometer and lidar retrievals. The normalized mean bias (NMB) in particle backscatter of the ceilometer and lidar profile is calculated following Eq. (6). The center of mass of the profiles is calculated with Eq. (7), and also the relative difference between ceilometer and lidar center of mass. Finally, the coefficient of correlation (R) of the profiles is determined.

$$NMB = \frac{\overline{\beta_{ceil}} - \overline{\beta_{lidar}}}{\overline{\beta_{lidar}}} \quad (6)$$

$$C_{mass} = \frac{\int_{z_{min}}^{z_{max}} z \cdot \beta(z) dz}{\int_{z_{min}}^{z_{max}} \beta(z) dz} \quad (7)$$

In Eq. (6), $\overline{\beta_{ceil}}$ and $\overline{\beta_{lidar}}$ are the mean particle backscatter coefficient from ceilometer and lidar respectively for the entire retrieved profile, and β in Eq. (7) may refer to ceilometer or lidar particle backscatter coefficient depending on the case.

Figure 2 shows for the 11 comparable profiles, the retrieved calibration factors at the reference height on the ceilometer profile versus the NMB (top panel), center of mass relative differences (middle panel) and R (bottom panel). It is evident that ceilometer profiles with a calibration factor closer to the mean calibration factor have inversions closer to the lidar inversions.

Figure 2 also shows that, for the statistics NMB and R, the difference between the calibration factor and the mean calibration factor are related, and the farther the profile calibration factor is from the mean value, the worse the mentioned statistics. Thus, it seems feasible to determine the quality of the profiles by selecting an appropriate threshold for this difference. Considering a maximum discrepancy between the particular calibration factor and the long term calibration factor equal to one standard deviation of the mean value of the calibration factor (dotted lines on Fig. 2), we obtain four profiles that have a NMB smaller

than 15%; the center of mass of the profiles is practically the same, with a relative difference smaller than 2% and finally, R of the profiles are above 0.92.

A sequence of ceilometer and lidar particle backscatter profiles from 23 February 2016 is shown on Fig. 3. The first ceilometer profile (marked in blue) has a calibration factor of $2.57 \cdot 10^{11}$ ($\text{m}^3 \text{sr}$) and hence is rejected according to the threshold described above. For this case, the NMB of the ceilometer and lidar profiles is -0.31, the center of mass relative difference is -0.06 and the R is 0.84. The other four ceilometer profiles (marked in red) have calibration factors within the standard deviation of the mean calibration factor. The profiles on 23 February at 9:00 and 9:30 UTC correspond with a decoupled dust layer. Those profiles have a NMB of -0.08 and 0.1 respectively, the center of mass relative difference is -0.01 and -0.03 respectively, and R is 0.95 and 0.97 respectively. The profiles on 23 February at 12:00 and 12:30 UTC show that the previous dust layer is mixed with the boundary layer. In these cases, profiles have a NMB of 0.14 and -0.12 respectively, the center of mass relative difference is 0.006 and -0.01 respectively, and R is 0.99 and 0.93 respectively.

4. Results

The capabilities of the ceilometer network for aerosol optical properties characterization and the near real-time processing have been tested with the analysis of the African dust outbreak that affected the Iberian Peninsula on 20 February 2016 and persisted until 24 February 2016.

Sorribas et al. (2017) studied the same event and compared it with meteorological parameters, aerosol properties and ozone from historical data sets on a site in southern Spain. They concluded that the event was exceptional because of its unusual intensity, its impact on surface measurements and the month of occurrence. In addition, Titos et al. (2017) also analyzed this event using 250 air quality monitoring stations over Spain to investigate the impact and temporal evolution of the event on surface PM_{10} levels. They also investigated aerosol optical properties, including attenuated backscatter from ceilometer during the event at Montsec station (one of the station included in ICENET). They concluded that the impact on surface PM_{10} was exceptional and highlighted the complexity of the event.

Next section provides a detailed description of the dust mobilization and arrival of the plumes to the Iberian Peninsula.

4.1 Description of the dust episode

The evolution of the dust outbreak is illustrated in Fig. 4 where a sequence of the false-color RGB dust images from MSG-SEVIRI is shown. This product makes use of three thermal satellite channels to contrast the brightness temperature signal between surface, cloud and dust (Lensky and Rosenfeld, 2008), in a color scheme in which dust appears in magenta. The presence of dust plumes over the High Plateau located between the Saharan and Tell Atlas in Algeria at 12:00 UTC on 20 February 2016 is shown in Fig. 4a. Dust migrated NW and passed over the Alboran Sea from the Algerian-Moroccan border around 14:00 UTC, reaching the southern Iberian Peninsula at 18:00 UTC (Fig. 4b) and continued moving north-westward (Fig. 4c). A second dust plume migrated northwards on 21 February 2016 at 16:00 UTC (Fig. 4d).

SYNOP meteorological observations and aerodrome routine (METAR) and special (SPECI) reports in Northern Africa recorded a strong reduction in horizontal visibility (MOR, Meteorological Optical Range), down to 2 km, between 7:00 and 8:00 UTC (20 February) at distant locations situated at the edges of the Great Western Erg in Algeria. At the eastern part of the Western Erg and in the Great Eastern Erg, visibility lowered to less than 5 km between 9 and 11:30 UTC. In good
5 accordance with the satellite images, at the Saharan Atlas and the High Plateau area, with heights over 1000 m, visibility less than 2 km were recorded at 10:00 UTC at Mecheria, while at the other stations in that area values went down to 2-3 km at 12:00 UTC. High relative humidity and clouds were found in the western most sites, also in agreement with the satellite images. It is remarkable that no significant visibility reduction was reported at the north-faced downslope areas of the Tell Atlas and in the Rif mountains, close or at the coast of Algeria and Morocco. This indicates that dust was uplifted before passing over
10 the northern slope of the Atlas range and the North African coast. Correspondingly, no station in the southern Iberian Peninsula reported a reduction in visibility when the dust plume reached Spain.

The entrance of dust-laden air masses above the ground level in the Iberian Peninsula is confirmed by a back-trajectory analysis performed with HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory) model (Draxler and Hess, 1998; Stein et al, 2015) using ERA-Interim data of 0.5° resolution. The trajectory analysis provides an estimate of the range of heights at
15 which the dust-laden air masses passed over the study sites. This is illustrated in Fig. 5 for two stations: on the south where dust reached first the Iberian Peninsula (Granada); and in the westernmost station, on the cyclonic track of the dust (Badajoz). The height vs latitude plots show that the dust plumes reaching Granada on February 20, 18:00 UTC and Badajoz on February 21, 00:00 UTC, arrived at mid-levels in the lower troposphere after being uplifted in the southern slope of the Saharan Atlas from heights between 500 and 1200 m above the ground in that area (black trajectory on Fig. 5). Those trajectories end at
20 around 3250 and 2500 m asl in Granada and Badajoz respectively. Figure 5 also shows that the trajectories reaching Granada at about the height of 2250 m asl passed over Africa before the dust mobilization took place while the ones below had no African origin. Finally, the trajectory reaching Granada at 4250 m asl followed the mid-upper level tropospheric circulations. In Badajoz, results are analogous: at 1000 m asl and below trajectories arrived from the north, between 1250 m asl and 2000 m asl they arrived from Africa but the air parcels were located north of the area where dust was observed in the morning of
25 February 20. The air parcels reaching Badajoz between 2250 m asl and 3250 m asl were previously located in the area where dust was being observed, while at upper levels trajectories followed the mid-upper circulation pattern.

In terms of aerosol load, Fig. 6 shows the time series of AOD at 1064 nm (top panel) and Ångström exponent between 440
and 880 nm (bottom panel) for the entire period of the event for all sites in ICENET. We observe that the increase in AOD and
decrease of Ångström exponent correspond with the arrival of the plumes at each site. The strongest part of the event occurs
30 on 21-22 February and the second plume is observed clearly for Granada station on 22 February. For dust events following a similar pattern that the one described here during the period 2005-2010, Valenzuela et al., (2012) reported a maximum AOD of 0.98 at Granada, which is significantly lower than the maximum measured at Granada station during this event.

4.2 Synoptic scenario and context

Several atmospheric features at mid-upper levels were relevant for this episode as they promoted instability near the surface and induced dust transport in the lower free troposphere:

(1) the amplification and break up of a Rossby wave in the eastern Atlantic resulted in a trough that became isolated as a cut-off low over the Atlas Mountains on 19 February. A shallow cyclone was then originated leeward of the Atlas. From early 21 February, the cut-off low displaced off the Moroccan coast and centered southwest of St. Vincent Cape. On 22 February it decayed bringing the Iberian Peninsula under the influence of the Azores and North African subtropical highs, with dominant zonal flows. (2) An upper-level anticyclone over a wide area centered over Niger-Chad, which during the episode intensified and extended northwards to the western Mediterranean. This high-pressure influenced circulation at mid-upper levels in combination with the cut-off low. (3) Moisture flux at mid tropospheric levels, which entered from the central Atlantic into the African continent below 20°N and was transported to northern Africa (at 400-550 hPa according to the radiosoundings in the area) between the upper-level trough and the high-pressure system. The tropical air masses are well appreciated in the satellite imagery as an elongated cloud band moving north and eastward, and so are the convective clouds formed ahead of the band. The tropical-extratropical interaction between the advected tropical moisture and the upper-level trough located over the Atlas range is linked to convective precipitation in northwestern Africa, see Knippertz (2003) and references therein. Divergence at upper levels (250 hPa) and low-level (850 hPa) convergence are found over the area where the gust front mobilized the dust on 20 February. The interaction with the Ahaggar Mountains in southern Algeria possibly enhanced convection and low-level instability. Convective precipitation was registered at several locations of eastern Spain when the cloud band passed over the area in the second half of the episode. From 22 February onwards the cloud band and local convective situations were gradually displaced to the Mediterranean, as zonal flows began to dominate.

At low levels, the low-pressure formed in the lee of the Atlas moved to the SW of the St. Vincent Cape on 21 February following the upper level instability. The low was then intensified and influenced northern Africa and most of the Iberian Peninsula. In addition, high pressures over the western Mediterranean were formed when the Rossby wave train progressed to the east and retreated poleward. Then, the North African high, which was previously located over Libya at 850 hPa, extended to Tunisia and Algeria and was gradually intensified in connection with the northward extension of the high-pressure at upper levels, which arrived (along with the cloud band on its western flank) to the western Mediterranean Basin.

The advection of dust-laden air masses to the Iberian Peninsula was driven by both the low located to the SW of the Iberian Peninsula and the North African high. The presence of these two synoptic systems corresponds to one of the typical synoptic situations leading to dust transport over the Iberian Peninsula (Rodriguez et al., 2003; Escudero et al., 2005). During the episode, however, two distinct strong plumes were transported from northern Africa to the Iberian Peninsula in consecutive days and showed a different evolution. Dust mobilized by the gust front on 20 February south of the Saharan Atlas and north of the Ahaggar migrated west and northward to the Iberian Peninsula, as showed in the satellite images, forming a curve-shaped plume over Iberia due to the cyclonic shear imposed by the low. The second strong dust plume was mobilized and

transported northwards on 21 February on the western side of the North African high, driven by the intensification of this high pressure system, which was the dominating feature in the second half of the episode. In this second case, dust was advected mostly below the cloud band and affected the eastern part of the Iberian Peninsula as well as most of the western Mediterranean basin.

- 5 The low pressure system weakened on early 22 February and the region was increasingly dominated by the Azores and the North African highs. As a consequence, zonal flow swept the first dust plume along northern Spain from west to east in subsequent days. The second dust plume, which was moving northward along eastern Iberia, was also displaced to the Mediterranean. The study region was then under high pressures and the event ended.

4.3 Ceilometer data analysis

- 10 The vertical structure of the dust event described above has been monitored with the ICENET. Firstly, by obtaining the total attenuated backscatter profiles using the calibration factors from Table (2) and, secondly, by applying the inversion and obtaining particle backscatter profiles. In addition, the integral of the backscatter profiles multiplied by the lidar ratio is used to estimate the AOD during the event and the center of mass of the backscatter profiles is considered as an indicator of the presence of a decoupled aerosol layer (a dust plume in this case) or the entrainment of the aerosol layer into the boundary layer. All these products were calculated in near real-time and serve as an example of the promising capabilities for real-time
15 characterization of singular events with a network of distributed ceilometers.

- Figure 7 shows time series of total attenuated backscatter profiles, i.e. calibrated profiles, for the five ceilometers used in this study. From top to bottom the series correspond to Granada, Tabernas, Badajoz, Valladolid and Montsec stations respectively. Tabernas station is covered by clouds during most of the event and Montsec station is also affected by clouds during part of
20 the event.

- Dust arrives first at the stations in Granada and Tabernas (on 20 February at 18:00 UTC). As the dust plume moves northwestward we observe the dust plume in Badajoz (on 21 February at 00:00 UTC), and Valladolid (on 21 February at 06:00 UTC). At Montsec, the dust plume is detected on 21 February at 12:00 UTC). The second plume brings the cloud band and this is visible at Tabernas station around 12:00 UTC on February 21 and a bit later at 21:00 UTC on 21 February at Montsec
25 station. Finally, the displacement of the dust from west to east at the end of the event, when the cut-off low weakens appears as a dust plume at Valladolid on 22 February at 15:00 UTC, at Badajoz station on 22 February at 21:00 UTC, and at Granada station on 23 February at 06:00 UTC. Tabernas and Montsec are influenced by the second dust plume and the cloud band and this is not as clearly visible as on the other stations. Another feature that is observed in Fig. 7 is that the dust plumes, specially the first one, are entrained into the boundary layer rapidly.

- 30 After applying the inversion, a quantitative comparison of stations is possible as shown in Fig. 8 for different stages of the dust outbreak (the specific times are shown on the vertical lines in Fig. 7). The beginning of the outbreak, when the first plume arrives at the different stations is shown in Fig. 8a. The center of mass of the dust plumes is about 3000 m asl for all stations. Additionally, for Granada and Badajoz, we observe that the height of the peak in particle backscatter coefficient is in

accordance with the backward trajectory analysis shown on Section 4.1. The arrival of the second plume is shown in Fig. 8b for all sites on 22 February at 06:00 UTC. At this stage, we observe that Granada and Tabernas stations (which are only 100 km apart) show very different behavior in particle backscatter and also in the height of the dust plume. Finally, Fig. 8c shows the final part of the outbreak when dust is mobilized eastwards to the Mediterranean Sea. In this case, dust is below 2000 m asl for Granada and Tabernas, whereas for the rest of the stations it is still observed above 3000 m asl. In general, the particles backscatter coefficient profiles indicate a stronger intensity of the event in this stage of the event, after the second plume arrives specially for Granada and Tabernas.

For the entire dust outbreak period and all stations the integral of the backscatter profiles is shown in Fig. 9a. This parameter allows identifying the beginning of the dust event for each station. Thus, it is observed and increase of the integral of the backscatter in Granada around 20 February at 19:30 UTC, in Badajoz is detected around 21 February at 5:30 UTC, in Valladolid at 16:30 UTC and in Montsec at 17:00 UTC. Due to clouds, this increase of the integral of the backscatter is not observed in Tabernas. The influence of the dust load after the first plume masks the arrival of the second plume, but the dust mobilization towards the Mediterranean sea is observed again at Badajoz (around 22 February at 20:00 UTC) and in Montsec at 23:00 UTC. The change of the integral of backscatter to larger values is in coincidence with the starting time observed in the total attenuated backscatter temporal series and it is in accordance with the satellite observations and back trajectory analysis. Additionally, the center of mass of the particle backscatter coefficient profiles is used to monitor the evolution of the profile region with more predominance of aerosol particles. Thus, in Fig. 9b for Granada before the event, the center of mass is about 1500 m asl, and when the dust arrives the center of mass is elevated to 2500 m. After 9 hours the center of mass is about the same as before the event, indicating that, possibly, the dust plume is no longer decoupled, and it is entrained into the boundary layer. A similar behavior is observed for Badajoz, Valladolid and Montsec stations. Again, the second plume is not observed in changes of the center of mass but the mobilization of dust towards the Mediterranean Sea is observed as an increase of the center of mass of the profiles for Badajoz, Valladolid and Tabernas.

5 Conclusions

The use of ceilometers for the characterization of optical aerosol properties is possible but, due to the weak signal, it is important to screen out profiles in order to ensure the quality of the inversion. In addition, due to the vast amount of data, it is important to perform all these operations in an automated, unsupervised way and, preferably, in near real-time. The methodology proposed uses ancillary data from sun photometer in order to constraint the calibration of the ceilometers. The time series of this calibration is used to determine the quality of the inversions selecting those that present, at the reference height, a ratio of the backscattering signal to molecular attenuated backscatter within the mean calibration factor \pm the standard deviation. A comparison with independent lidar measurements indicates that this method allows the automatic discrimination of the quality of the inversions with ceilometers. During this comparison a difference smaller than 15% in backscatter coefficient is observed. Thus, it is feasible to routinely provide particle backscatter coefficient profiles with ceilometers.

The inverted profiles obtained with ceilometers could be used for elevated aerosol layer alert by setting a threshold on the particle backscatter coefficient values of the profile and are potentially useful for model assimilation and evaluation since all the processing is automated and in near-real time.

This method has been applied to a group of ceilometers (ICENET) and tested during a dust outbreak reaching Spain on 20 February 2016 until 24 February 2016. This dust event affected all ICENET stations with two distinct plumes reaching the Iberian Peninsula following different paths and a final stage where zonal flows swept the dust towards the Mediterranean Sea. This scheme of dust mobilization is unusual for this season of the year and the intensity, spatial coverage and duration of the event makes it perfect as a test for monitoring purposes with the ceilometer network. The calibration of the ceilometers allows a qualitative monitoring of the event while the inversions provide quantitative information. Thus, ceilometers can complement 10 lidar stations that, in principle would operate intermittently and with less spatial density. It is worth noting that it has been observed differences on profiles 100 km apart. This reinforces the need of providing vertical profiles of aerosol optical properties with a dense spatial resolution.

Parameters extracted from the particle backscatter coefficient profiles such as the integral or the center of mass can also give a quantitative idea of the presence of an elevated aerosol layer. These parameters are expected to increase with an elevated 15 aerosol layer and the second one can be used as a rough indicator for the deposition velocity of an elevated aerosol layer by comparing a time series of these values.

Acknowledgments

This work was supported by the Spanish Ministry of Economy and Competitiveness through projects CGL2012-39623-C02-01, CGL2013-45410-R, CGL2014-56255-C2-1-R, CMT2015-66742-R, CGL2015-70741-R, CGL2015-73250-JIN and 20 CGL2016-81092-R, by the Andalusia Regional Government through project P12-RNM-2409, by the Castilla y Leon Regional Government through project VA100U14, by the Junta de Extremadura (Ayuda a Grupos de Investigación GR15137) and by the European Union's Horizon 2020 research and innovation program through project ACTRIS-2 (grant agreement No 654109). The authors thankfully acknowledge the FEDER program for the instrumentation used in this work. This work was also partially funded by the University of Granada through the contract "Plan Propio. Programa 9. Convocatoria 2013". Marco 25 Pandolfi is funded by a Ramón y Cajal Fellowship (RYC-2013-14036) awarded by the Spanish Ministry of Economy and Competitiveness. The authors would like to acknowledge the valuable contribution of the discussions in the working group meetings organized by COST Action ES1303 (TOPROF).

References

- Alados-Arboledas, L., Müller, D., Guerrero-Rascado, J. L., Navas-Guzmán, F., Pérez-Ramírez, D. and Olmo, F. J.: Optical and microphysical properties of fresh biomass burning aerosol retrieved by Raman lidar, and star-and sun-photometry, *Geophys. Res. Lett.*, 38, L01807, doi:10.1029/2010GL045999, 2011.
- 5 Ångström, A.: The parameters of atmospheric turbidity. *Tellus* 16, 64–75, 1964.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens B. and Zhang, X.Y.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V.
- 10 Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 2013.
- Bravo-Aranda, J., Titos, G., Granados-Muñoz, M., Guerrero-Rascado, J., Navas-Guzmán, F., Valenzuela, A., Lyamani, H., Olmo, F., Andrey, J. and Alados-Arboledas, L.: Study of mineral dust entrainment in the planetary boundary layer by lidar depolarization technique. *Tellus B*, 67. doi:http://dx.doi.org/10.3402/tellusb.v67.26180, 2015.
- Córdoba-Jabonero, C., Sorribas, M., Guerrero-Rascado, J. L., Adame, J. A., Hernández, Y., Lyamani, H., Cachorro, V., Gil,
- 15 M., Alados-Arboledas, L., Cuevas, E., and de la Morena, B.: Synergetic monitoring of Saharan dust plumes and potential impact on surface: a case study of dust transport from Canary Islands to Iberian Peninsula, *Atmos. Chem. Phys.*, 11, 3067-3091, doi:10.5194/acp-11-3067-2011, 2011.
- Draxler, R.R. and Hess, G.D.: An Overview of the HYSPLIT_4 Modelling System for Trajectories, Dispersion, and Deposition. *Australian Meteorological Magazine*, 47 (1998) 295-308, 1998.
- 20 Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B.N., Mishchenko, M., Yang, P., Eck, T.F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W.J., Leon, J.-F., Sorokin, M. and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust. *J. Geophys. Res.*, 111, D11208, doi:10.1029/2005JD006619. 2006.
- Escudero, M., Castillo, S., Querol, X., Avila, A., Alarcón, M., Viana, M. M., Alastuey, A., Cuevas, E. and Rodríguez, S.: Wet and dry African dust episodes over eastern Spain. *Journal of Geophysical Research: Atmospheres*, 110(D18), 2005.
- Fernald, F. G., Herman, B. M., and Reagan, J. A.: Determination of Aerosol Height Distributions by Lidar, *Journal of Applied Meteorology*, 11, 482-489, 1972.
- Fernald, F. G.: Analysis of atmospheric lidar observations: some comments, *Appl. Optics*, 23, 652–653, 1984.
- Flentje, H., Heese, B., Reichardt, J., and Thomas, W.: Aerosol profiling using the ceilometer network of the German
- 30 Meteorological Service, *Atmos. Meas. Tech. Discuss.*, 3, 3643-3673, doi:10.5194/amtd-3-3643-2010, 2010.
- Frey, S., Poenitz, K., Teschke, G., and Wille, H.: Detection of aerosol layers with ceilometers and the recognition of the mixed layer depth, *ISARS*, 2010. 3646, 3647, 2010.

- GAW report 178. Plan for the implementation of the GAW Aerosol Lidar Observation Network GALION (Hamburg, Germany, 27-29 March 2007) (WMO TD No. 1443), 52 pgs, November 2008.
- GAW report 197. World Meteorological Organization Global Atmospheric Watch. Addendum for the period 2012-2015 to the WMO Global Atmospheric Watch (GAW) Strategic Plan 2008 – 2015, 63 pgs, May 2011.
- 5 Granados-Muñoz, M. J., Bravo-Aranda, J. A., Baumgardner, D., Guerrero-Rascado, J. L., Pérez-Ramírez, D., Navas-Guzmán, F., Veselovskii, I., Lyamani, H., Valenzuela, A., Olmo, F. J., Titos, G., Andrey, J., Chaikovskiy, A., Dubovik, O., Gil-Ojeda, M., and Alados-Arboledas, L.: A comparative study of aerosol microphysical properties retrieved from ground-based remote sensing and aircraft in situ measurements during a Saharan dust event, *Atmos. Meas. Tech.*, 9, 1113-1133, doi:10.5194/amt-9-1113-2016, 2016.
- 10 Guerrero-Rascado, J.L., Ruiz, B., Alados-Arboledas, L.: Multi-spectral Lidar characterization of the vertical structure of Saharan dust aerosol over southern Spain. *Atmospheric Environment*, 42,11, 2668–2681, 2008.
- Guerrero-Rascado, J. L., Olmo, F. J., Avilés-Rodríguez, I., Navas-Guzmán, F., Pérez-Ramírez, D., Lyamani, H., and Alados Arboledas, L.: Extreme Saharan dust event over the southern Iberian Peninsula in september 2007: active and passive remote sensing from surface and satellite, *Atmos. Chem. Phys.*, 9, 8453-8469, doi:10.5194/acp-9-8453-2009, 2009.
- 15 Guerrero-Rascado, J.L., Landulfo, E., Antuña, J.C., Barbosa, H. M. J., Barja, B., Bastidas, A. E., Bedoya, A. E., da Costa, R., Estevan, R., Forno, R. N., Gouveia, D. A., Jimenez, C., Larroza, E. G., Lopes, F. J. S., Montilla-Rosero, E., Moreira, G. A., Nakaema, W. M., Nisperuza, D., Alegria, D., Manera, M., Otero, L., Papandrea, S., Pallota, J. V., Pawelko, E., Quel, E. J., Ristori, P., Rodrigues, P. F., Salvador, J., Sanchez, M. and Silva, A.: Latin American Lidar Network (LALINET): diagnosis on network instrumentation, *Journal of Atmospheric and Solar-Terrestrial Physics*, 138-139, 112–120.
- 20 doi:10.1016/j.jastp.2016.01.001. 2016.
- Haefelin, M., et al. (2011), Evaluation of mixing-height retrievals from automatic profiling lidars and ceilometers in view of future integrated networks in Europe, *Boundary Layer Meteorol.*, 143, 49–75, doi:10.1007/s10546-011-9643-z.
- Haefelin, M., Angelini, F., Morille, Y., Martucci, G., Frey, S., Gobbi, G. P., Lolli, S., O’Dowd, C. D., Sauvage, L., Xueref-Rémy, I., Wastine, B. and Feist, D. G.: Evaluation of Mixing-Height Retrievals from Automatic Profiling Lidars and Ceilometers in View of Future Integrated Networks in Europe. *Boundary-Layer Meteorol.* 143: 49. doi:10.1007/s10546-011-9643-z, 2012.
- 25 Haefelin, M., Laffineur, Q., Bravo-Aranda, J.-A., Drouin, M.-A., Casquero-Vera, J.-A., Dupont, J.-C., and De Backer, H.: Radiation fog formation alerts using attenuated backscatter power from automatic lidars and ceilometers, *Atmos. Meas. Tech.*, 9, 5347-5365, doi:10.5194/amt-9-5347-2016, 2016.
- 30 Haij, M. de, Wauben, W. and Klein-Baltink, H.: Continuous mixing layer height determination using the LD-40 ceilometer: a feasibility study. KNMI Scientific Report WR 2007-01, 98 pp, 2007.
- Heese, B., Flentje, H., Althausen, D., Ansmann, A., and Frey, S.: Ceilometer lidar comparison: backscatter coefficient retrieval and signal-to-noise ratio determination, *Atmos Meas Tech*, 3, 1763-1770, 2010.

- Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET - A federated instrument network and data archive for aerosol characterization, *Remote Sens. Env.*, 66, 1-16, 1998.
- IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp. 2014.
- 5 Klett, J. D.: Stable analytic inversion solution for processing lidar returns, *Appl. Optics*, 20, 211–220, 1981.
- Klett, J. D.: Lidar inversion with variable backscatter/extinction ratios, *Appl. Optics*, 24, 1638–1643, 1985.
- Knippertz, P.: Tropical-extratropical interactions causing precipitation in Northwest Africa: Statistical analysis and seasonal variations. *Mon. Wea. Rev.*, 131, 3069–3076. doi:10.1175/1520-0493, 2003.
- 10 Lensky, I. M. and Rosenfeld, D.: Clouds-Aerosols-Precipitation Satellite Analysis Tool (CAPSAT), *Atmos. Chem. Phys.*, 8, 6739-6753, doi:10.5194/acp-8-6739-2008, 2008.
- Mattis, I., D'Amico, G., Baars, H., Amodeo, A., Madonna, F., and Iarlori, M.: EARLINET Single Calculus Chain – technical – Part 2: Calculation of optical products, *Atmos. Meas. Tech.*, 9, 3009-3029, doi:10.5194/amt-9-3009-2016, 2016.
- 15 McCormick M.P.: The Flight of the Lidar In-Space Technology Experiment (LITE). In: Ansmann A., Neuber R., Rairoux P., Wandinger U. (eds) *Advances in Atmospheric Remote Sensing with Lidar*. Springer, Berlin, Heidelberg, 1997.
- Münel C, Eresmaa N, Räsänen J, Karppinen A.: Retrieval of mixing height and dust concentration with lidar ceilometer. *Boundary-Layer Meteorol* 124: 117–128, 2007.
- Navas-Guzmán, F., Guerrero-Rascado, J. L., and Alados-Arboledas, L.: Retrieval of the lidar overlap function using Raman signals, *Óptica Pura y Aplicada*, 44(1), 71-75, 2011.
- 20 Navas-Guzmán, F., Müller, D., Bravo-Aranda, J. A., Guerrero-Rascado, J. L., Granados-Muñoz, M. J., Pérez-Ramírez, D., Olmo, F. J. and Alados-Arboledas, L.: Eruption of the Eyjafjallajökull Volcano in spring 2010: Multiwavelength Raman lidar measurements of sulphate particles in the lower troposphere, *J. Geophys. Res. Atmos.*, 118, 1804–1813, doi:10.1002/jgrd.50116, 2013.
- 25 Ortiz-Amezcuca, P., Guerrero-Rascado, J. L., Granados-Muñoz, M. J., Benavent-Oltra, J. A., Böckmann, C., Samaras, S., Stachlewska, I. S., Janicka, Ł., Baars, H., Bohlmann, S., and Alados-Arboledas, L.: Microphysical characterization of long-range transported biomass burning particles from North America at three EARLINET stations, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-946, in review, 2016.
- Pandolfi, M., Martucci, G., Querol, X., Alastuey, A., Wilsenack, F., Frey, S., O'Dowd, C. D., and Dall'Osto, M.: Continuous atmospheric boundary layer observations in the coastal urban area of Barcelona during SAPUSS, *Atmos. Chem. Phys.*, 13, 4983-4996, doi:10.5194/acp-13-4983-2013, 2013
- 30 Pappalardo, G., Amodeo, A., Pandolfi, M., Wandinger, U., Ansmann, A., Bösenberg, J., Matthias, V., Amiridis, V., De Tomasi, F., Frioud, M., Larlori, M., Komguem, L., Papayannis, A., Rocadenbosch, F. and Wang X.: Aerosol lidar intercomparison in

- the framework of the EARLINET project. 3. Raman lidar algorithm for aerosol extinction, backscatter, and lidar ratio. *Appl Opt.* 43(28):5370-85, doi: 10.1364/AO.43.005370 2004.
- Pappalardo, G., Mona, L., D'Amico, G., Wandinger, U., Adam, M., Amodeo, A., Ansmann, A., Apituley, A., Alados Arboledas, L., Balis, D., Boselli, A., Bravo-Aranda, J. A., Chaikovskiy, A., Comeron, A., Cuesta, J., De Tomasi, F.,
5 Freudenthaler, V., Gausa, M., Giannakaki, E., Giehl, H., Giunta, A., Grigorov, I., Groß, S., Haefelin, M., Hiebsch, A., Iarlori, M., Lange, D., Linné, H., Madonna, F., Mattis, I., Mamouri, R.-E., McAuliffe, M. A. P., Mitev, V., Molero, F., Navas-Guzman, F., Nicolae, D., Papayannis, A., Perrone, M. R., Pietras, C., Pietruczuk, A., Pisani, G., Preißler, J., Pujadas, M., Rizi, V., Ruth, A. A., Schmidt, J., Schnell, F., Seifert, P., Serikov, I., Sicard, M., Simeonov, V., Spinelli, N., Stebel, K., Tesche, M., Trickl, T., Wang, X., Wagner, F., Wiegner, M., and Wilson, K. M.: Four-dimensional distribution of the 2010 Eyjafjallajökull volcanic
10 cloud over Europe observed by EARLINET, *Atmos. Chem. Phys.*, 13, 4429-4450, doi:10.5194/acp-13-4429-2013, 2013.
- Pappalardo, G., Amodeo, A., Apituley, A., Comeron, A., Freudenthaler, V., Linne, H., Ansmann, A., Bosenberg, J., D'Amico, G., Mattis, I., Mona, L., Wandinger, U., Amiridis, V., Alados-Arboledas, L., Nicolae, D., and Wiegner, M.: EARLINET: towards an advanced sustainable European aerosol lidar network, *Atmos Meas Tech*, 7, 2389-2409, 2014.
- Preißler, J., Wagner, F., Pereira, S. N. and Guerrero-Rascado, J. L.: Multi-instrumental observation of an exceptionally strong
15 Saharan dust outbreak over Portugal, *J. Geophys. Res.*, 116, D24204, doi:10.1029/2011JD016527, 2011.
- Preißler, J., Wagner, F., Guerrero-Rascado, J. L. and Silva, A. M.: Two years of free-tropospheric aerosol layers observed over Portugal by lidar, *J. Geophys. Res. Atmos.*, 118, 3676–3686, doi:10.1002/jgrd.50350., 2013.
- Rodríguez, S., Querol, X., Alastuey, A. Viana, M.M. and Mantilla, E.: Events Affecting Levels and Seasonal Evolution of Airborne Particulate Matter Concentrations in the Western Mediterranean. *Environmental Science & Technology* 37 (2), 216-
20 222, DOI: 10.1021/es020106p, 2003.
- Sicard, M., Guerrero-Rascado, J. L., Navas-Guzmán, F., Preißler, J., Molero, F., Tomás, S., Bravo-Aranda, J. A., Comerón, A., Rocadenbosch, F., Wagner, F., Pujadas, M., and Alados-Arboledas, L.: Monitoring of the Eyjafjallajökull volcanic aerosol plume over the Iberian Peninsula by means of four EARLINET lidar stations, *Atmos. Chem. Phys.*, 12, 3115-3130, doi:10.5194/acp-12-3115-2012, 2012.
- 25 **Sorribas, M., Adame, J.A., Andrews, E. and Yela, Y.: An anomalous African dust event and its impact on aerosol radiative forcing on the Southwest Atlantic coast of Europe in February 2016. *Science of the Total Environment*, 583, 269–279, doi: 10.1016/j.scitotenv.2017.01.064, 2017.**
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System. *Bull. Amer. Meteor. Soc.*, 96, 2059-2077, doi: 10.1175/BAMS-D-14-00110.1,
30 2015.
- Titos, G., Ealo, M., Pandolfi, M., Pérez, N., Sola, Y., Sicard, M., Comerón, A., Querol, X., and Alastuey, A.: Spatiotemporal evolution of a severe winter dust event in the western Mediterranean: Aerosol optical and physical properties. *Journal of Geophysical Research: Atmospheres*, 122, 4052–4069, doi:10.1002/2016JD026252, 2017.**

Valenzuela, A., Olmo, F.J., Lyamani, H., Antón, M., Quirantes, A. and Alados-Arboledas, L.: Classification of aerosol radiative properties during African desert dust intrusions over southeastern Spain by sector origins and cluster analysis. *Journal of Geophysical Research*, 117, D06214, doi:10.1029/2011JD016885, 2012.

5 Valenzuela, A., Costa, M.J., Guerrero-Rascado, J.L., Bortoli, D., Olmo, F.J.: Solar and thermal radiative effects during the 2011 extreme desert dust episode over Portugal. *Atmospheric Environment*, 148, 16–29, 2017.

Wandinger, U., Müller, D., Böckmann, C., Althausen, D., Matthias, V., Bösenberg, J., Weiß, V., Fiebig, M., Wendisch, M., Stohl, A. and Ansmann, A.: Optical and microphysical characterization of biomass-burning and industrial-pollution aerosols from multiwavelength lidar and aircraft measurements, *J. Geophys. Res.*, 107(D21), doi:10.1029/2000JD000202, 2002

10 Welton, E. J., Campbell, J. R., Spinhirne, J. D. and Scott, V. S.: Global monitoring of clouds and aerosols using a network of micro-pulse lidar systems, in *Lidar Remote Sensing for Industry and Environmental Monitoring*, U. N. Singh, T. Itabe, N. Sugimoto, (eds.), *Proc. SPIE*, 4153, 151-158, 2001.

Wiegner, M., Emeis, S., Freudenthaler, V., Heese, B., Junkermann, W., Munkel, C., Schafer, K., Seefeldner, M., and Vogt, S.: Mixing layer height over Munich, Germany: Variability and comparisons of different methodologies, *J. Geophys. Res.*, 111, D13201, doi:10.1029/2005JD006593, 2006.

15 Wiegner, M. and Geiss, A.: Aerosol profiling with the Jenoptik ceilometer CHM15kx, *Atmos Meas Tech*, 5, 1953-1964, 2012.

Wiegner, M., Madonna, F., Biniotoglou, I., Forkel, R., Gasteiger, J., Geiss, A., Pappalardo, G., Schafer, K., and Thomas, W.: What is the benefit of ceilometers for aerosol remote sensing? An answer from EARLINET, *Atmos Meas Tech*, 7, 1979-1997, 2014.

20 Winker, D. M., Pelon, J. R., and McCormick, M. P.: CALIPSO mission: spaceborne lidar for observation of aerosols and clouds. *Proc. SPIE* 4893, *Lidar Remote Sensing for Industry and Environment Monitoring III*, (21 March 2003); doi: 10.1117/12.466539, 2003, 1-11. 2003.




Figure 1. Map of the Iberian Peninsula showing the location of the ceilometers. In Granada station (circled in red) a co-located multiwavelength Raman lidar is also available.




Figure 2. Ceilometer Calibration Factor (C_L) vs. normalized mean bias (NMB) on top panel, relative difference in center of mass on middle panel and coefficient of correlation (R) on bottom panel. Horizontal line indicates the mean C_L for the dust event period and dashed lines indicates the 33% around this mean value.

5




Figure 3. Lidar and ceilometer particle backscatter profiles for five cases on 23 February 2016. The first case (marked in blue) is a rejected ceilometer profile and the other four cases (marked in red), are cases with a ceilometer calibration factor within the 33% of the median calibration factor. Errors on lidar profiles have been estimated with a **Monte Carlo** technique (Pappalardo et al., 2004; Mattis et al., 2016) and are in the order of 10^{-9} ($\text{m}^{-1} \text{sr}^{-1}$), therefore they are not shown.




Figure 4. False-color RGB dust image from MSG-SEVIRI showing different stages of the dust outbreak (dust appears magenta).





Figure 5. Evolution of the air parcels reaching Granada on February 20, 18:00 UTC (left) and Badajoz on February 21, 00:00 UTC (right) at different heights. Back-trajectories were calculated from the ground level to 5000 m asl, at every 250 m. Lines in grey indicate trajectories arriving at the lowest levels, with no African history; in green are trajectories that passed over the southern slope of the Saharan Atlas before the observed dust mobilization; in black are the trajectories followed by the parcels residing at the times and area where dust was observed; the trajectories residing at higher levels are depicted in blue. One representative trajectory is shown for each evolution and the altitude interval is shown in the same color as the representative trajectory. The brown line corresponds to the ground level for the trajectories more associated to the dust advection (thick black lines). The location of the air parcels around the time of observation (12 UTC) of the dust plumes is shown as a red circle.

5

10






Figure 6. Sun-photometer time series representing the AOD at 1064 nm (top panel) and Ångström exponent between 440 and 800 nm (bottom panel) for all sites in ICENET during the dust event.






Figure 7. Ceilometer time series of total attenuated backscatter representing the evolution of the dust outbreak between 20 and 24 of February 2016 (the color scale is logarithmic). Red vertical lines indicate the time of the profiles in Fig. 8: First line of each site indicates the times for Fig. 8a, second line for Fig. 8b, and third line for Fig. 8c.



5

Figure 8. Particle backscatter coefficient profiles for all stations at the beginning (a), middle (b) and final stage (c) of the outbreak (note that x axis has different scale and the profiles start at ground level). Shaded area represents the 15% uncertainty.






Figure 9. Time series of the integral of the particle backscatter coefficient for all stations (top panel) and time series of the center of mass of the backscatter profiles for all stations (bottom panel).

5 **Table 1.** Description of the Iberian ceilometer network sites.

Site (code)	Managed by	Location (Lat°/Lon°)	Height (m asl)	Additional Instruments
Granada (UGR)	Atmospheric Physics Group. University of Granada.	37.16°N/3.58°W	680	CIMEL CE 318 Multi-Wavelength lidar
PlataformaSolar de Almería-Tabernas (PSA)	Institute of Solar Research. German Aerospace Center.	37.09°N/2.36°W	500	CIMEL CE 318
Badajoz (UEX)	AIRE Group. University of Extremadura.	38.88°N/7.01°W	199	CIMEL CE 318
Valladolid (UVA)	Atmospheric Optics Group, University of Valladolid.	41.66°N/4.71°W	705	CIMEL CE 318
Montsec (MSA)	Institute of Environmental Assessment and Water Research. Spanish Research Council.	42.02°N/0.74°E	800	CIMEL CE 318 (42.05°N/0.73°E/1570m amsl)
Murcia (UMH)	Statistical and Computational Physics	39.98°N/1.13°W	69	CIMEL CE 318

	Lab, Miguel Hernández University.			
--	-----------------------------------	--	--	--

Table 2. Mean calibration factors for ceilometers in ICENET for the period 1 May 2014 to 1 May 2016.

Site code	CL (m³ sr)
UGR	$(1.6 \pm 0.4) \cdot 10^{11}$
PSA	$(3.3 \pm 0.7) \cdot 10^{11}$
UEX	$(3.7 \pm 0.9) \cdot 10^{11}$
UVA	$(3.8 \pm 0.5) \cdot 10^{11}$
MSA	$(3.1 \pm 0.7) \cdot 10^{11}$
UMH	$(4.4 \pm 0.9) \cdot 10^{11}$