

1 **Long-term series and trends in surface solar radiation at Athens,**
2 **Greece**

3

4 Stelios Kazadzis^{1,2}, Dimitra Founda², Basil E. Psiloglou², Harry Kambezidis², Nickolaos
5 Mihalopoulos^{2,3}, Arturo Sanchez-Lorenzo^{4,5}, Charikleia Meleti⁶, Panagiotis I. Raptis^{1,2}, Fragiskos
6 Pierros², Pierre Nabat⁷

7

8 [1] {Physikalisch-Meteorologisches Observatorium Davos, World Radiation Center (PMOD/WRC)
9 Dorfstrasse 33, CH-7260 Davos Dorf, Switzerland}

10 [2] {Institute of Environmental Research and Sustainable Development, National Observatory of Athens,
11 Greece}

12 [3] {Department of Chemistry, Univ. of Crete, Heraklion, Crete}

13 [4] {Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza,
14 Spain}

15 [5] {Department of Physics, University of Extremadura, Badajoz, Spain}

16 [6] {Physics Department, Aristotle University of Thessaloniki, Greece}

17 [7] {CNRM UMR 3589, Météo-France/CNRS, Toulouse, France}

18 Corresponding author: S. Kazadzis, kazadzis@noa.gr

1 **Abstract**

2 We present a long-term series of solar surface radiation (SSR) from the city of Athens, Greece. SSR
3 measurements were performed from 1954 to 2012, and before that (1900-1953) sunshine duration
4 (SD) records have been used in order to reconstruct monthly SSR. Analysis of the whole dataset
5 (1900-2012) mainly showed very small (0.02%) changes in SSR from 1900 to 1953, including a
6 maximum decrease of 2.9% per decade in SSR during the 1910 to 1940 period, assuming a linear
7 change. For the dimming period 1955-1980, a -2% change per decade has been observed that
8 matches various European long-term SSR measurement related studies. This percentage at Athens
9 is in the lower limit, compared to other studies in the Mediterranean area. For the brightening period
10 1980-2012 we have calculated a +1.5% change per decade, which is also in the lower limit of the
11 reported positive changes in SSR around Europe. Comparing the 30-year periods 1954-1983 and
12 1983-2012, we have found a difference of 4.5%. However, measurements of the first 30 year period
13 are associated with higher uncertainties than those of the second period, especially when looking at
14 year to year changes. The difference between the two periods was observed for all seasons except
15 winter. Analyzing SSR calculations of all sky and clear sky (cloudless) conditions/days, we report
16 that most of the observed changes in SSR after 1954 can be attributed partly to cloudiness and
17 mostly to aerosol load changes.

18

1 **1 Introduction**

2 In the past decades surface solar radiation (SSR) and the transmission of the atmosphere have been
3 of increasing interest because of the related impacts on climate. Most of the energy in the Earth-
4 atmosphere system is introduced by solar radiation. It provides heating, which creates pressure
5 gradients and ultimately wind and triggers water, carbon and oxygen cycles through evaporation
6 and photosynthesis. These processes define the climatological conditions, and changes of incoming
7 solar radiation rapidly affect the energy balance (Wild et al., 2015). Interest in solar radiation
8 changes has also been raised after the development of solar energy applications, which are
9 continuously growing in number over the recent years. Changes in SSR have been recorded over the
10 last century and can be caused either by natural events such as volcanic eruptions or human-related
11 activities, mainly in polluted regions (Wild, 2016). At longer scales (thousands of years) changes in
12 SSR might have been caused by changes in the Earth's orbit and Sun solar output (Lean, 1997;
13 Ohmura, 2006).

14 Systematic continuous measurements of SSR were established in the middle of the 20th century at
15 selected meteorological observatories. Solar variations have been investigated in several studies
16 using ground based SSR measurements from various monitoring networks worldwide (e.g.,
17 Ohmura, 2009) and also by satellite-derived estimations (e.g. Kambezidis et al., 2010). Overall,
18 most of these studies (Gilgen et al., 1998; Noris and Wild, 2009; Wild, 2009 and 2016 and
19 references therein) have reported a worldwide decrease of solar incoming radiation in the period
20 1960-1985 (known as dimming period), followed by an increase (brightening period) thereafter.
21 These changes were reported to be higher in more polluted and urban areas but have also been
22 recorded in isolated regions such as the Arctic (Stanhill, 1995) and Antarctica (Stanhill and Cohen
23 1997). Other recent studies have investigated the effect of urbanization on global brightening and
24 dimming, and found no marked differences among urban and rural SSR time series (Tanaka et al.,
25 2016) and Imamovic et al. (2016). Changes in atmospheric transmission due to variations in
26 cloudiness and aerosol concentration are the main factors to be investigated in order to determine
27 the possible causes of such trends in SSR (Wild, 2009). However, due to the aerosol-cloud
28 interactions and the aerosol indirect effect on SSR, the two factors (clouds and aerosols) are not
29 completely mutually exclusive in explaining SSR changes.

30 The cloud and aerosol radiative effects on solar radiation variations over the past decades have been
31 investigated by numerous studies during the last years. The inter-annual variations in cloudiness is
32 crucial for studying SSR time series, but its decadal variability is not always connected with the
33 widespread dimming and brightening effects (Wang et al., 2012; Wild, 2016). Aerosols play a

1 significant role in incoming radiation, by scattering and absorbing light and by acting as cloud-
2 condensation nuclei. Over the 20-year dimming phase (from 1960 to 1980) and the 15-year
3 brightening phase (from 1990 to 2005), it was found that the aerosol effects (direct and indirect)
4 played the most important role in SSR variation (Dudok de Wit et al., 2015). Concerning Central
5 Europe, Ruckstuhl et al. (2008) suggested that the brightening phase under cloud-free conditions is
6 in line with decreasing anthropogenic aerosol emissions (Streets et al., 2006). Nabat et al. (2013)
7 using a blending of remote sensing and model products showed that a decreasing Aerosol Optical
8 Depth (AOD) trend of 0.05 per decade in Europe for the period of their study (1979-2009). In
9 addition, Nabat et al. (2014) reported that anthropogenic aerosol decline in Europe from 1980 to
10 2012 statistically explains $81 \pm 16\%$ of the observed brightening. Overall, changes in anthropogenic
11 aerosol emissions are now considered as the major cause of brightening and dimming effects (Wild,
12 2016). The gaseous and particulate air pollutants may reduce solar radiation by up to 40% during air
13 pollution episodes (Jauregui and Luyando, 1999). Aerosol related attenuation is much larger during
14 forest fires, dust events and volcanic eruptions. Vautard et al. (2009) have also reported a decline of
15 the frequency of low-visibility conditions such as fog, mist and haze in Europe over the past 30
16 years, suggesting a significant contribution of air-quality improvements

17 Long-term series of SSR measurements are essential for such studies. One of the main constraints in
18 studying SSR temporal changes is the small number of sites with reliable long-term records, even
19 over areas with a relatively high density of stations such as Europe, Japan or the USA. In Europe for
20 example, there are currently less than 80 stations with more than 40-years homogeneous data
21 (Sanchez-Lorenzo et al., 2015), with very few of them operating over Southern Europe. Recently, a
22 high-quality dataset of SSR has been set up over Italy (Manara et al., 2016), but there is still lack of
23 high quality long-term trends in other countries around the Mediterranean Basin.

24 In addition, even more sporadic measurements are available before the 1950s (Stanhill and Ahiman,
25 2016); the few studies of them have pointed out an SSR increase in the first decades of the 20th
26 century and a maximum around 1950 (Ohmura, 2006 and 2009). This topic is still controversial due
27 to the few long-term series available (Antón et al., 2014). Recently, there have been efforts to
28 reconstruct SSR series in periods with no direct measurements available, using other variables such
29 as sunshine duration (SD), which is available in a large number of sites since the late 19th century
30 (e.g., Stanhill and Cohen, 2005, for USA; Sanchez-Lorenzo and Wild 2012, for Switzerland;
31 Matuszko 2014, for Poland). For example, Sanchez-Lorenzo and Wild (2012) used data from 17
32 stations in Switzerland, considered SD as a proxy and successfully reconstructed SSR time series
33 since the late 19th century. They calculated that the variability in SSR monthly anomalies can be

1 explained by SD anomalies in a range of 76%-96%, and a monthly root mean squared error of 4.2
2 W m^{-2} between recorded and estimated SSR for all-sky conditions and of 5.5 W m^{-2} for clear-sky
3 conditions. Other studies have tried to use pan evaporation as a proxy of SSR, for the first half of
4 the 20th century (Stanhill and Möller, 2008). Kambezidis et al. (2016) used monthly re-analysis
5 datasets from the Modern Era Retrospective-Analysis for Research and Applications (MERRA) and
6 calculated shortwave radiation trends over the period 1979-2012 for the Mediterranean basin. They
7 reported an increase in MERRA of $+0.36 \text{ W m}^{-2}$ per decade, with higher rates over the western
8 Mediterranean ($+0.82 \text{ W m}^{-2}$ per decade).

9 A few studies discuss the brightening/dimming effect in the southeastern Mediterranean. Zerefos et
10 al. (2009) have studied the Ultraviolet A (UVA) changes for the area of Thessaloniki (Greece) from
11 1984 to 2008. They calculated a 5% positive trend per decade linked to a negative trend in aerosol
12 optical depth (AOD) for the area due to air pollution abatement strategies. The variability in
13 shortwave downward solar irradiance received at Earth's surface over Thessaloniki, Greece, for the
14 period 1993-2011 (Bais et al., 2013), showed an upward trend in SSR after 1990 ($+0.33\%$ per
15 year). They also reported signs of a slowdown in the upward trend in SSR during the beginning of
16 the 2000s. Founda et al., (2014) have studied the SD long-term variability over Athens area. They
17 reported a 7% decline in the annual SD from 1951-1982 and a 3% increase from 1983-2011 under
18 all sky conditions. Under near clear sky conditions, these percentages are -7% and + 9% for the
19 dimming and brightening periods respectively. Similarly, Founda et al. (2016a) analyzed long-term
20 SD and total cloud cover time series over 15 sites in Greece (the oldest one beginning in 1897).
21 They have shown an increase in SD almost at all stations since the mid-1980s, which in certain
22 areas of Southeastern Greece amounts to an increase of 20 h per year. This increase is not
23 accompanied with synchronous decrease in total cloud cover, possibly evidencing to decreasing
24 aerosol loads, despite the fact that their impact on SD should be lower than on SSR (Sanchez-
25 Romero et al., 2014). Yildirim et al. (2014) have analyzed 41 years of SD measurements in 36
26 stations in Turkey. They reported a decreasing trend (between 1970 to about 1990) at most of the
27 stations. After 1990, they observed either zero trend variation or a reduction in the decreasing rate
28 of SD at most of the locations. They concluded that the decreasing period might be attributed to
29 human-induced air pollution. Founda et al. (2016b) have investigated the visibility trends over
30 Athens area from 1931 to 2013. They reported a deterioration in the visibility up to 2004 and a
31 slight recovery afterwards, negatively/positively correlated with relative humidity/wind speed and
32 positively correlated with AOD from 2000 to 2013. Finally, Alexandri et al., 2017 studied the
33 spatio-temporal variability in SSR over the eastern Mediterranean for the 1983–2013 period, using
34 the Satellite Application Facility on Climate Monitoring Solar surfAce RADIation Heliosat satellite-

1 based product (SAF). They reported a positive (brightening) and statistically significant SSR trend
2 at the 95% confidence level ($0.2 \pm 0.05 \text{ W m}^{-2} \text{ year}^{-1}$ or $(0.1 \pm 0.02\% \text{ year}^{-1})$) being almost the same
3 over land and sea.

4 In this work, measurements of SSR, recorded over 60 years at the center of Athens, are presented.
5 In addition, with the use of the SD measurements that are conducted in Athens since 1900, we could
6 reconstruct the time series of SSR during the first half of the 20th century. These time series (1900-
7 2012) are the oldest, uninterrupted and high quality SSR time series in the SE Mediterranean and
8 one of the oldest in Europe, providing unique information about the variations and trends in the area
9 for the past decades. Time-series of SSR over Athens are presented to try answering questions such
10 as:

11 Are the dimming–brightening patterns observed in Europe over the past century also observed, at
12 the same extent, over the eastern Mediterranean?

13 Is SSR variability during the first decades of the 20th century in Athens in line with the trends
14 reported at other locations over this period?

15 Can we verify that anthropogenic aerosols play the most important role on the brightening/dimming
16 observed SSR after 1950, in agreement with results from other European regions?

17

18 **2 Data and Methodology**

19 **2.1 DDR data collection and analysis**

20 The SSR data used in this study cover the period from December 1953 to December 2012 and were
21 measured by a series of pyranometers that are mentioned in Table 1. These instruments have been
22 operating continuously at the Actinometric Station of the National Observatory of Athens
23 (ASNOA) (Hill of Pnyx, Thissio), that is located near the center of Athens, Greece (38.00° N ,
24 23.73° E , 110 m above mean sea level). Table 1 presents the instruments and the period of
25 operation, as well as the maximum error on the integrated daily values. References mentioned in
26 Table 1 describe the exact type of errors and uncertainties related to the sensors. In the period 1953-
27 1986, the maximum daily error was about 5%, and 2% afterwards. The spectral response of the
28 sensors is in the range of 285-2800 nm; since 1986 a first-class Eppley PSP pyranometer (WMO,
29 1983) is operating at ASNOA. Since 1992, frequent calibrations (every two years) have been
30 performed by the NOAA's Laboratory of Meteorological Device Calibration (LMDC, 2016) in order
31 to ensure the high quality of measurements. LMDC follows the standard calibration procedure for
32 thermopile pyranometers (ISO 9847, 1992), with exposure to real sunlight conditions and

1 comparison with a standard thermopile pyranometer (Secondary Standard). LMDC's reference
 2 pyranometer, Kipp & Zonen CM21, is regularly calibrated in PMOD/WRC, Davos, Switzerland.

3

4 Table 1: History of SSR instruments used at ASNOA. SSR measurements refer to the total solar
 5 radiation on a horizontal surface.

	Instrument	Period	Class	Maximum error (daily integral)	Reference	Class	Comments	Resolution
1	Solarigraph GOREZYNSKI	1953-1959	2nd	5%	Coulson (1975)	B	One instrument being used	1 hour
2	Eppley 180° pyranometer (No. 3604)	1960-1966	2nd	5%	Coulson (1975), Drummond (1965)	B	Manual measurements archiving with mvoltmeter	1 hour
3	Eppley 180° pyranometer (No. 3604) coupled with a Leeds-Northup recorder, Speedomax, type G	1966-1968	2nd	5%	Coulson (1975), Drummond (1965)	B	Same instrument as #2 with Speedomax recorder	1 hour
4	Eppley 180° pyranometer (No. 3034) coupled with a Leeds-Northup recorder, Speedomax, type G	1968-1973	2nd	5%	Coulson (1975), Drummond (1965)	B	New instrument, same recorder	1 hour
5	Eppley pyranometer, type 8-48 and type 8-48A coupled with a Leeds-Northup recorder, Speedomax, type G	1974-1986	2nd	3-5%	Hulstrom (1989)	B	Type 8-48 and type 8-48A instruments were measuring alternatively for three years each	1 h
6	Eppley Precision Spectral Pyranometer (PSP)	1986-now	1st	1-2%	Hulstrom (1989)	A	Regular recalibrations. Coupled with a, A/D recorder (Campbell Scientific Ltd.) Datalogger, type CR-21X at the beginning until 2003, a CR10X until 2012	1 min

6

7 SSR data are processed using a set of quality-control (QC) tests in order to ensure the quality of the
 8 data set. The QC procedures include rejection of:

- 9
- Measurements for solar elevation angles less than 5 degrees;

1 • SSR values equal to or less than 5 W m^{-2} , during sunrise and sunset, due to the
2 pyranometers' sensitivity;

3 • SSR values greater than 120% of the seasonally corrected solar constant.

4 After the initiation of diffuse horizontal radiation measurements at ASNOA in 1991, the following
5 quality criteria were added for rejection:

6 • diffuse horizontal values greater than the corresponding SSR ones;

7 • diffuse horizontal values greater than 80% of the seasonally correct solar constant;

8 • direct-beam solar component exceeding the extraterrestrial solar irradiance.


9 Also, both total and diffuse horizontal measurements are corrected for the night-time dark-signal
10 offset of the pyranometers.

11 Mean daily SSR values were calculated from the data set of this study (December 1953 – December
12 2012); only months with more than 20 days of measurements were considered in the analysis. Over
13 the 60 years of measurements, only three months (January and February of 1998 and March of
14 2012) did not fulfill this criterion.

15 When trying to use such long term series, it is evident that the data quality differs as instrument
16 performance have been improved, quality assurance and quality control procedures have been
17 standardized, and finally the information flow on the day to day instrument performance issues is
18 much more complete in the recent time. At ASNOA, (after 1986 only) the instruments were
19 calibrated or checked with a reference instrument on a yearly basis to identify changes in the
20 calibration and drifts. As reported, the addition of diffuse irradiance measuring instruments
21 provided the opportunity to improve also minute based measurement quality. Before 1986, the
22 instruments reported in table 1 have been used. According to the log books there has always been a
23 certain overlap when changing from one instrument to another. Reports mention that there were
24 instrument drifts that have been corrected with no further information from 1953 to 1970.
25 Instrument overlaps after 1986 were used to eliminate possible instrument related offsets. However,
26 instrument differences (e.g. thermal offset of PSP instrument compared with 8-48 pyranometer,
27 Vignola et al., 2016) could theoretically have an effect in the order of $1\text{-}2 \text{ W m}^{-2}$ on the series
28 continuation. In this case the subtraction of the night time dark signal (more specific the mean of the
29 previous and next night signal was subtracted for a specific day) reduces at least in half the
30 problem. However, in order to answer to the reviewer question, the remaining offset was not
31 considered in our analysis, as partly it have been tackled through the overlapping
32 measurements/homogenization procedures. In addition, the inclusion of diffuse radiation in the

1 quality assurance tests after 1991 could cause a major improvement on the newest data compared
2 with the old ones. However, this recent improvement in quality control cannot be linked with a
3 systematic impact on SSR measurements and changes compared to the past, other than higher
4 uncertainty on the integrated (monthly, yearly) SSR values. For the 1953-1986 time series there is a
5 number of publications that have been using the SSR-NOA time series. More specific: Macris,
6 (1959), has used the 1954-1956 SSR measurements to identify the relationship between SSR and
7 sunshine duration. Katsoulis and Papachristopoulos, (1978), have used the SSR data from 1960 to
8 1976 in order to calculate SSR statistics for daily, seasonal and yearly solar radiation levels at
9 Athens, Greece. Notaridou and Lalas, (1979), have used the 1954-1976 SSR data in order to verify
10 an empirical formula on global net radiation over Greece. Flocas, (1980) has used the 1961-1975
11 SSR time series to compare them with sunshine duration data for the same period. Kouremenos et
12 al., (1985) have used the SSR data from 1955-1980 in order to correlate their changes with various
13 atmospheric parameters. Zabara, (1986) has used the 1965-1980 time series to verify a monthly
14 solar radiation calculation method. Katsoulis and Leontaris, (1981), have used the 1960-1977 data
15 to verify tools describing the solar radiation distribution over Greece. Finally, the percentages of
16 errors reported in table 1 are not directly linked with possible instrument drifts that can impact the
17 SSR time series analysis. So results of measurements before 1986 have to be used with caution and
18 accompanied by a report on the different level of uncertainties of the past and recent data.

19 Figure 1 shows the intra-annual variability of SSR at ASNOA based on the measurements from all
20 instruments during the period 1953-2012. Daily SSR at Athens ranges between approximately 6 to
21 27 MJ m⁻² during the year. Mean and standard deviations were calculated using the 60 year record
22 for each day.



1

2 **Figure 1.** Average intra-annual variability of Surface Solar Radiation (SSR) at Actinometric Station of the
 3 National Observatory of Athens (red), along with the inter-annual variability for a given day (± 1 standard
 4 deviation, blue), calculated over the period 1953-2012.
 5

6 The results of figure 1 show the average yearly pattern of SSR at ASNOA. The day to day
 7 variability that is shown as “noise” in the plotted blue line comes from the 60 year averaging of
 8 each day and is mostly related with the amount of cloudiness for each of the averaged days.
 9 Minimum and maximum SSRs at solstices, compared to a cloudless sky aerosol free model, are also
 10 related with the highest probability of the presence of clouds during winter months. For the
 11 calculation of each of the daily averages the available data points vary from 55 to the possible
 12 maximum of 60.

13 Concerning the data availability for SSR and SD data, SSR monthly means calculated here have
 14 been retrieved from daily calculated SSRs. Over the 59 years (708 months) of SSR data, 98% of the
 15 months had none or one day missing, 3 months had from 10-20 missing days and 2 months from
 16 20-30 missing days. For SD, 1931-1940 monthly data used have been taken from the NOA
 17 measurement annals. From 1940 on, hourly measurements have been used in order to derive daily
 18 and monthly measurements. The SD time series have no gaps with only six missing days during
 19 December 1944 (Founda et al., 2014).

20

21 **2.2 Sunshine duration data**

22 In addition, collocated measurements of SD recorded at ASNOA have been used. According to
 23 WMO (2010), the SD during a given period is defined as the sum of the sub-periods for which the

1 direct solar irradiance exceeds 120 W m^{-2} . In Athens, SD has been recorded using classical
2 Campbell-Stokes heliographs (since 1894) and been replaced by electronic instrumentation in 1998
3 (EKO, MS-091 analog SD sensor). Monthly SD values since January 1900 have been used in this
4 study. A more analytical study of these time series can be found in Founda et al. (2014).

5 Complementary to this study, cloud-cover observations from the Hellenic National Meteorological
6 Service (HNMS) from 1954 have also been used. These observations are recorded at a site 7 km
7 away from ASNOA. All cloud observations at HNMS are conducted every 3 hours and are
8 expressed in octas.

9 **2.3 Aerosol Optical Depth (AOD)**

10 In order to examine the AOD impact on SSR, we have used the longest satellite based AOD series
11 available for the area. This is the AOD time series from Advanced Very High Resolution
12 Radiometer (AVHRR). AOD retrievals at 630 nm over global oceans at spatial resolution of $0.1^\circ \times$
13 0.1° and one overpass per day have been used. Data used were downloaded from NOAA Climate
14 Data Record (CDR) version 2 of aerosol optical thickness (Zho and Chan, 2014), and cover the
15 period from August 1981 to December 2009. AVHRR AOD embodies a large variety of
16 uncertainties, including radiance calibration, systematic changes in single scattering albedo and
17 ocean reflectance (Mishchenko et al, 2007). Current dataset radiances have been recalibrated using
18 more accurate MODIS data (Chan et al, 2013). We used daily data at the region around Athens
19 (longitude: 37.5° - 38.2° N, latitude: 23.2° - 24.4° E) which includes 50 active available (ocean) grid-
20 points. The above region was selected based on data availability on each grid, within 50 km from
21 ASNOA.

22 To complement the analysis on the evolution of aerosols, the recent climatology developed by
23 Nabat et al., (2013) has been considered over the period 1979-2012. This product provides monthly
24 averages of AOD at 550 nm over the Mediterranean region at 50 km resolution. It is based on a
25 combination of satellite-derived (MODIS instrument) and model-simulated products (MACC
26 reanalysis and RegCM-4 simulations), which have been selected among many available datasets,
27 from an evaluation against ground-based measurements of the AERONET network. Thus this
28 climatology is able to give the best possible atmospheric aerosol content over the period 1979-2012.
29 For the present work, the AOD time series over the grid cell of the ASNOA (38.00° N, 23.73° E)
30 has been extracted and is referred to as the ChArMEx data thereafter.

31

32 **2.4 Clear-sky SSR**

1 For the determination of the clear sky (defined here as the cloudless) days, we have used both the
2 cloud octas and SD data. Daily observations have been used for this analysis. We have defined as a
3 clear sky day each day that fulfills the following criteria:

- 4 - the mean daily cloudiness (in octas) should be less than 1.5, and
- 5 - the total daily SD should be higher than 90% of its theoretical (astronomical) value.

6 The procedure for calculating a single mean cloud octa value for each day was the following:

7 We have first excluded night-time cloud observations; then, we have weighted each observation
8 based on the hour of the observation. Weights have been calculated based on the solar radiation
9 contribution of the specific time slot and day of the month, compared with the daily clear sky SSR
10 integral, of the particular day and month.

11


12 2.5 Reconstruction of SSR from SD


13 We have used the 1900-2012 SD time series in order to extend our SSR time series back to 1900.

14 There are different methods that are used in order to estimate SSR values from SD. In this work we
15 have tried two methods based on the linear regression between SSR and SD (Sanchez-Lorenzo and
16 Wild, 2012). For all-sky conditions the monthly anomalies, obtained as differences from the 1983-
17 2012 mean, of SSR and SD have been calculated. Then for each month a linear regression has been
18 used to estimate the relationship between the SSR and the SD:

$$19 \text{SSR} = a * \text{SD} + b \quad (1)$$

20 Figure 2 shows four out of the total twelve regressions together with its' statistics.





1 Figure 2. Linear regression of SSR and SD anomalies for March, June, October and December,
 2 1983-2012.

3 Table 2. Monthly regression statistics of SSR vs SD anomalies (see eq. 1).

month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
a	22.47	34.99	39.53	46.65	57.88	44.67	51.87	46.23	34.28	28.02	27.32	23.10
R	0.842	0.895	0.887	0.840	0.799	0.757	0.773	0.772	0.812	0.888	0.916	0.842

4

5 The correlation coefficients (R) they vary from 0.75 to 0.91. This implies that the SD monthly
 6 anomalies explain between 56% and 83% of the variability of the SSR monthly anomalies. It has
 7 to be noted that coefficient b is less than 10^{-3} for all months.

8 The second method to derive a relationship between SD and SSR, was based on the broadly
 9 accepted formula of Ångström (1924):

10

$$11 \quad \text{SSR}/\text{SSR}_{\max} = c + d * (\text{SD}/\text{SD}_{\max}) \quad (2)$$

12

13 where SSR_{\max} and SD_{\max} refer to the theoretical extra-terrestrial value of SSR and the astronomical
 14 value of SD, respectively, while c and d are constants usually defined monthly. This formula can
 15 only be used in large data sets as a statistical approach. That is because for different cloud heights,
 16 thicknesses and positioning, the constants can show a large variability (Angell, 1990). The 1983-
 17 2012 period was chosen for determining the SSR vs. SD relationship as SSR measurements have
 18 lower uncertainties compared with the 1953-2012 period. We thus calculated $c=0.237$ and $d=0.458$
 19 and an R^2 equal to 0.81.

20 We also followed the same procedure to calculate the coefficients of the Ångström formula
 21 separately for each month and for each season during the control period 1983-2012. For individual
 22 months, calculated $\text{SSR}/\text{SSR}_{\max}$ vs $\text{SDU}/\text{SDU}_{\max}$ coefficients of determination ranged from 0.5
 23 to 0.65 for winter months, 0.32 to 0.67 for spring months, 0.47 to 0.53 for autumn months and 0.1
 24 to 0.38 for summer months. So coefficients of determination using the monthly based data were

1 much lower than the first reported method. The low coefficients for the summer period are related
 2 with the small range of values of SDU/SDU max and SSR/SSRmax that are related with the
 3 absence of clouds.

4 We have used both the monthly regression coefficients from the first method and the yearly based
 5 Ångström formulas in order to investigate the impact of the different methods on the SSR
 6 reconstruction. Results of the reconstructed SSR yearly values from 1900-1953 showed maximum
 7 differences of 1% in the calculated SSR percent anomalies, while for monthly values the higher
 8 difference was 2%. In order to avoid the use of theoretical normalization values such as SDUmax
 9 and SSRmax needed for the second method, we have reconstructed the SSR time series based on
 10 the monthly based results of the first method as proposed in Sanched-Lorenzo and Wild, 2012.

11


12 **3 Results**


13 **3.1 Long-term variations and trends (1900-2012)**

14 Based on methods described in Section 2, we have reconstructed monthly SSR from 1900 to
 15 1953. Using the full dataset of reconstructed (1900-1953) and measured SSR (1954-2012)
 16 we have calculated the mean monthly SSR values and used them for de-seasonalising the
 17 results shown in Figure 3. The de-seasonalizing was determined by: a. calculating the
 18 average SSR (SSR_{mi}) for each month (i) out of the 12 months of the given year, for all 1983-
 19 2012 years, b. calculating the changes in % in SSR ($SSR\%(i,y)$) for each month (i) of each
 20 year (y) as:

$$21 \quad SSR\%(i,y) = \frac{SSR_{iy} - SSR_{mi}}{SSR_{mi}} * 100 \quad (3)$$

22





1
 2 **Figure 3.** Full time series of de-seasonalised SSR percent changes (using the 1900-2012 monthly
 3 averages). Upper panel: different colors represent seasonal analysis, lower panel: black bold line
 4 represents the annual series, and light blue line the mean monthly values

5
 6 According to Figure 3, the month-to-month variation (shown with light grey line) can reach more
 7 than 30% in comparison with the mean monthly average of the whole data set. Annual means show
 8 a 10%-12% (peak to peak) decrease in SSR from 1910 to late 1930's and then an increase of 12%
 9 from 1940 to early 1950's. The simulated SSR results follow the observed decline of SD reported in
 10 Founda et al., (2014), where a decrease from 1910 to 1940's is shown.

11 Subsequently, there is a decrease during the late 1950's and then a positive change of the order of
 12 20% till today with an episode in the early 1990's that shows low SSR values. Measured SSR in
 13 1991-1993 period differs by 5% compared with the one in 1990. The latest can be linked with the
 14 Pinatubo volcanic eruption and its known effect in the SSR (e.g. Zerefos et al., 2012).

15 Analytical linear trends of each of the sub-periods and for every season are presented in Table 2. It
 16 has to be noted that the trend determination and its statistical significance do not take into account
 17 measurement or SSR reconstruction related uncertainties, which are different for the different
 18 periods.

19 **Table 2.** Annual and seasonal SSR trends in percent per decade over the period 1900-2012 and
 20 different sub-periods. Percentages in parenthesis show the limits of the 95% confidence bounds.


Season	1900-2012	1900-1952	1953-1982	1983-2012
Winter	-0.11 (±0.47)	-0.90 (±1.46)	-6.43 (±3.83)	+0.52 (±3.26)
Spring	+0.54 (±0.37)	+0.38 (±1.12)	-0.60 (±3.10)	+2.77 (±3.10)

Summer	+0.59 (± 0.21)	+0.28 (± 0.48)	-1.14 (± 2.90)	+1.38 (± 2.55)
Autumn	+0.21 (± 0.44)	+0.11 (± 0.97)	-1.28 (± 3.42)	-1.50 (± 1.83)
Year	+0.39 (± 0.22)	+0.04 (± 0.71)	-2.33 (± 2.28)	+0.80 (± 1.96)

1

2 Looking at the 1900-2012 period the seasonal and annual linear trends in SSR are less than 1% per
3 decade. A positive change of 0.39% per decade has been calculated from annual values. For the
4 whole data set, all seasons show positive trends, except winter. For the periods with simulated SSR
5 values (1900-1952), even smaller trends have been detected for spring and summer. The measuring
6 period of 1954-2012 has been split into two sub-periods of 1954-1982 and 1983-2012. The first
7 sub-period shows a negative annual change of -2.33% per decade in SSR, which is also reflected in
8 all seasons with predominant changes during winter (-6.43% per decade). The second sub-period
9 shows a positive trend of +0.80% per decade with the highest ones in spring (+2.77% per decade)
10 and summer (+1.38% per decade) and negative in autumn (-1.50% per decade). Looking at the
11 trend significance described by the 95% confidence bounds, we can see significant positive trends
12 for 1900-2012 (yearly, summer and spring) and significant negative trends for yearly analysis and
13 winter of 1953-1982.

14 In order to have a better understanding of the SSR changes over the 113-year period (1900-2012),
15 we have calculated the decadal SSR trends for different time-windows (15 to 40 years). Figure 5
16 shows the results of this analysis.



17

18 **Figure 4.** Trends in SSR (% per decade) calculated for different sliding time windows. The value of
19 the trend has been calculated at the central year of each time window.

20

1 For the first two decades of the 20th century there appears a decrease in SSR, in line with other
2 long-term SD series as recently shown by Stanhill and Ahiman (2016). Then, in all calculations an
3 increase is shown from mid-1930's to late 1940's, in line with the early brightening effect pointed
4 out by other authors (Ohmura, 2009; Sanchez-Lorenzo et al., 2008). It should be reminded that this
5 period is based on estimations of SSR from SD measurements, which thus include additional
6 uncertainties. Nevertheless, early dimming and brightening periods have been reported in Stanhill
7 and Ahiman (2016). The results can be partly supported by trends in anthropogenic black carbon
8 (McConnell et al., 2007; Lamarque et al., 2010) and biomass-burning (Lamarque et al., 2010)
9 emissions in Europe. The dimming period from 1950's to 1970's can be observed in all time
10 windows with a brightening effect after late 1970s.


11 The 40-year and 30-year time windows in the analysis presented in Figure 5 show the maximum
12 rate of increase in early 1940's (resulting in an increase of 2% per decade and 3% per decade,
13 respectively). Then a maximum rate of decrease is observed in early-mid 1960's, followed by a
14 positive rate of increase after 1990's. Shorter time windows (15 years) are also interesting as they
15 are able to capture the Pinatubo effect in early 1990's.

16


17 **3.2 Variations and trends in SSR for the 1954-2012 measurement period**

18 In order to further analyze the whole 59-yr SSR data set of this study, we have divided it in two 30-
19 yr climatological sub-periods: 1954-1983, and 1983-2012 (the common year is meant to have equal
20 (30 full years) duration for both periods). Investigating a possible seasonal dependence, the relative
21 difference in SRR for every month from its mean monthly value over the whole measurement
22 (1954-2012) period was calculated.

23 Figure 5 shows the mean daily insolation for each month for the two sub-periods and the whole 59-
24 year period. Examining the monthly average differences between the two periods, we observe that
25 for spring and summer months these are of the order of 6%. In addition, for all months SSR
26 differences of the 1983-2012 period compared to the 1954-1983 period are positive with an
27 exception of November (-1.9%) and December (-1.2%). In general, the second measurement period
28 shows a 3% to 8% larger monthly SSR than the first measurement period.



1
 2 **Figure 5.** Intra-annual variability of monthly mean daily SSR over the sub-periods of 1954-1983
 3 (blue line) and 1983-2012 (red line) and the entire period of 1954-2012 (black line). The green line
 4 (right axis) represents the monthly relative difference between the two 30-year sub-periods (recent
 5 minus older period).



6
 7 **Figure 6.** Percent change per decade for different time scales and different months using 1954 as
 8 the starting year (the last 30 year period is magnified with changes presented as percent per
 9 decade).

10

11 We have also calculated decadal trends in time windows of 15 to 30 years for the entire SSR
 12 measurement period (see Figure 6), only for the 1954 to 2012 period, Figure 6 shows the SSR
 13 change per decade for the months of January, April, July, October and yearly (all months). The
 14 figure is showing a trend analysis for the entire data set with time windows from 10 to 59 years,
 15 where each time window starts from 1953. For all months SSR changes become positive for time

1 windows of 35 years and higher (1953-1988 time window and any larger window starting from
2 1953). Negative trends calculated from 1954 to any given year up to 1989 are mainly due to the
3 large negative changes during the winter period. Especially during the 1954-1974 period, winter
4 SSR changes show a 18% per decade decrease. Linear trends in SSR from 1954-2012 showed a
5 positive trend of the order of 1% per decade, while individual months vary from 0.5% per decade to
6 1.5% per decade. Mostly positive trends are detected using any time window centered after 1975.
7 Larger trends are calculated for time windows centered at 1975 to 1980 and after 2000 (in the order
8 of 5% per decade using the 15-year time window). For the period 1954 to 1970 mainly negative
9 trends are shown.

10


11 **4. Comparison between all-sky and clear-sky SSR records variation**

12 We have used the 59-year data set (1954-2012) in order to quantify the factors controlling the SSR
13 variations in Athens, Greece, focusing mainly on two known dominant factors, clouds and aerosol
14 load.

15 **4.1 The role of clouds**

16 Figure 7 shows the 1954-2012 time series of yearly anomalies based on daily SSR, together with
17 yearly total cloud coverage in weighted octas. The yearly de-seasonalised SSR values for all-sky
18 conditions show a drop of ~14% from 1960 to 1970 and then a continuous increase excluding the
19 Pinatubo period in the early 1990's. Most pronounced positive changes can be seen during the last
20 15 years with a change of the order of about 15%.

21



22

1 **Figure 7.** De-seasonalised yearly percent deviations from mean for SSR (black line) and cloud
2 octas. Grey lines are related with measurements possible uncertainties/drifts.

3 Going back to the measurement uncertainties for the 1954-1983 period where a number of
4 instruments have been used in order to build the presented time series; we have tried to investigate
5 possible instrument drifts and their effect on the calculated long term trends. In order to indirectly
6 try to tackle this issue we included in figure 7 a shaded area representing a possible (one direction)
7 “uncertainty” based on reconstructing the 1954-1983 series using: the 1984-2012 measured SSR
8 data and the sunshine duration data for 1954-1983. The reconstruction has been performed in the
9 same way as the 1900-1953 one. The one direction “uncertainty”, points out possible drifts and
10 instrument exchange related uncertainties. However, that does not mean that we believe more on the
11 reconstructed through sunshine duration 1954-1983 series than the actual SSR measurements. If this
12 was the case, we would have decided to present a 1983-2012 high quality measuring period and a
13 1900-1983 reconstructed one. There are various of such papers published quite recently (small
14 measuring period compared with the reconstructed one: Garcia et al., 2014; [1992-2013
15 measurements reconstructed back to 1933] and Anton et al., 2017 [1887-1950 using radiative
16 transfer modelling]) while in our case we would like to try use the best way possible the historical
17 SSR measurements of NOAA during the 1954-83 period.

18 Using the 1984-2012 measurements and the 1900-1983 reconstruction data set we have
19 recalculated all trends presented in figure 4 and table 2. Differences for the 15 year window
20 differences on the calculated trends outside the 54-83 period are less than 1%, with maximum
21 differences at the late 60’s 1-3%. For the 30 year window maximum differences are in the order of
22 1-2%, while for the 40 year window, maximum differences are less than 1%.

23 This particular exercise cannot be defined as an uncertainty assessment on the 1954-83
24 measurements, as reconstructed data cannot be used as a reference. Moreover, SSR is much more
25 sensitive than SD to aerosol optical depth change. So, in locations where the number of cloudless
26 days is relatively high SD reconstruction tends to “smooth” the SSR variability, however the
27 opposite can be said in cases with constant cloudiness.

28 Figure 8 shows the correlation between annual mean SSR and cloud cover. From the best-fit linear
29 regression line it is deduced that a -1.54 MJ m^{-2} (or -9.6%) change in mean daily insolation
30 accounts for a change of 1 octa in cloud cover.




Figure 8. Correlation between annual means of daily insolation and cloud cover over the period 1954-2012. The straight line represents the best-fit regression line to the data points. The year 1953 has not been included in the analysis since it does not include measurements for all months.


However, the great scatter of the data points and the low correlation of the two parameters in Figure 8 ($R^2=0.229$) indicate that the cloud cover can only partly explain the changes in SSR. In addition, there is no significant change in cloudiness over the 59 year period for Athens, Greece. Calculating linear changes of cloudiness from data shown Figure 8, shows a non-significant change of -0.4% per decade which can practically have a limited effect on SSR changes during the examined period. Nevertheless, it is worth mentioning that different cloud properties like cloud optical thickness and cloud phase, not described by the measurements of cloud cover, can influence SSR.

4.2 Clear sky records

In order to minimize the cloud influence and investigate the possible role of direct aerosol effects on Athens SSR series, we had to select clear-sky (or cloudless) days. We have used daily SSR measurements from 1954 to 2012 and we have separated the cloudless days according to the criteria mentioned in Section 2.2.

For considering the SSR seasonality, we have calculated a five-degree polynomial derived from the maximum daily SSR (for all years of the data set), as a function of the day of the year (Figure 9). Afterwards we have calculated the ratio of the daily SSR to the SSR calculated by this function. Seasonal and yearly means of this ratio have been estimated and have been used to describe cloudless-sky SSR percentage changes on a seasonal and yearly basis. This approach has been chosen since averaging a random set of cloudless days, within each month during the 59-year


1 period, could cause solar elevation-related (due to the change of maximum solar elevation within
2 each month) discrepancies, when calculating the monthly average SSR. It can be emphasized that
3 the clear sky selection criterion could possibly eliminate a few cases with very high aerosol optical
4 depth.



5
6 **Figure 9.** Clear-sky SSR measurements (red dots) and all-sky SSR measurements (blue dots)
7 derived with the cloud octa (cloudiness<1.5) and sunshine duration (SD>0.9) related criteria. The
8 black line represents the polynomial fit to the daily SSR_{max} values.
9

10 Using the clear sky conditions seasonal and yearly averages of SSR have been calculated. The use
11 of seasonal instead of monthly SSR has been introduced in order to improve the averaging SSR-
12 related statistics, since the average number of cloudless days (per year) can be relatively low
13 especially during the winter months. For all cases the ratios of the mean daily cloudless SSR to the
14 SSR_{max} derived from the daily best-fit curve in Figure 9 has been calculated and deviations of this
15 ratio from its 59-yr mean have been calculated for each year.

16



1
2 **Figure 10.** Changes in yearly mean SSR to relative to the 1954-2012 average for cloudless sky (in
3 %; black), AVHRR AOD series (red) and ChArMEx AOD climatology (blue; Nabat et al., 2013)
4 for Athens area is shown in the right axis.

5
6 Figure 10 shows that most of the SSR variation observed for the measuring period has to be
7 explained by other factors than changes in cloudiness (see figure 7 for variations due to cloudiness).
8 Different seasons with the exception of wintertime show similar patterns to the year-to-year
9 variability. Individual seasonal calculated SSR variability do not exceed by more than $\pm 5\%$, the
10 SSR variability of all sky data, with the exception of the winter season. Comparing clear sky and all
11 sky yearly mean SSR, we find a high correlation ($R^2 = 0.71$), which can be explained as a
12 combination of: aerosol changes driving the SSR changes and by the number of clear sky days
13 during the year. There is a decrease of more than 15% in the clear sky SSR from the start of the
14 series to the end of 1960's. A decline after 1983 could possibly be related with El Chichon volcanic
15 eruption.

16 All sky SSR measurements and AOD from AVHRR have been used in order to find the AOD effect
17 in all sky data. For yearly AOD and SSR averages from 1981 to 2009 we have calculated a
18 correlation coefficient of -0.55 with a rate of SSR reduction per 0.1 units of AOD equal with -3.8%.
19 For monthly based comparisons, all months revealed a correlation coefficient of -0.2 with a rate of -
20 1.5% per 0.1 AOD with better results for summer and Autumn months (-0.30, -2.2%/0.1 AOD and -
21 0.30, -1.5%/0.1 AOD)


22 The Pinatubo-related drop of 6% from the early 1990's to the mid-1993 can also be seen in both
23 cloudless and all-sky datasets and also to the increase in AOD in the AVHRR dataset (Figure 10).

1 Since, the ~6% drop from 1990 to 1991-1993 is shown for all seasons, we can argue that it
2 describes the effect of the eruption on SSR data for the Athens station. However, as shown in figure
3 7 cloudiness for 1991 is also high, while is much lower for 1992 and 1993. Combined with the
4 stratospheric AOD figure, it seems that 1991 related decrease is also related with cloud increase
5 while 92 and 93 one with the Pinatubo related aerosol effect.

6 Concerning the stratospheric AOD in Athens the ChArMEX AOD dataset revealed two main peaks
7 of 0.12 for 1983 and 0.09 for 1992 due to El Chichon and Pinatubo eruptions, respectively, while
8 stratospheric AOD after 1995 is lower than 0.01. These two peaks are possibly associated with
9 decreases in SSR as measured at ASNOA.

10 Finally, the ~13% change from 1995 to 2012 shown for all skies (Fig. 7) and clear skies (Fig. 10) is
11 accompanied with a drop of ~25% in AOD measured by AVHRR. The year to year variations of
12 clear sky SSR series and the AVHRR-related AOD show an anti-correlation with $R=-0.78$ ($N=29$),
13 verifying the hypothesis that SSR clear sky changes are associated with aerosol load changes, at
14 least within the common AVHRR/measurement period (1982-2009).

15 Similar to the AVHRR data the ChArMEX 4-D aerosol climatology is shown in figure 10, providing
16 similar conclusions, namely the AOD negative trend of 0.03 or 14% per decade from 1979 to 2012.
17 Differences between the AVHRR and ChaArMEX data can be explained in part by the different
18 AOD wavelengths presented here (630 vs 550 nm) and also by a general negative bias of AVHRR
19 over the Mediterranean compared to AERONET (Nabat et al., 2014). The smooth decline in the
20 ChArMEX AOD data is due to the method used to build this product and uses the trend and not the
21 interannual variability which is not included in the global model that was used.



22
23

Figure 11. Ratio of cloudless vs all days, per season and yearly

1

2 In addition to figure 10 we have included figure 11 showing the ratio of cloudless days to all
 3 available days for each season and for each year. Figure 11 shows a minimum (less than 30%
 4 during a year) of the number of cloudless days from mid-1970's to early 1990's. It is mostly linked
 5 with the decrease of cloudless days during summer months. The figure provides a hint on the SSR
 6 relative changes observed during this period, but it cannot directly interpret year to year SSR
 7 changes as they depend also on cloud fraction and properties for cloudy days. In addition, it can
 8 only partly be linked with fig. 10 as aerosol effects on cloudless sky calculated SSRs depend mostly
 9 on AOD levels and not on the number of days included in the calculations.

10 Differences in the ratio of cloudless days shown in figure 12 and in the almost constant cloud octa
 11 variability shown in figure 7 is partly attributed to the different definition of a cloudless day that is
 12 based on the cloud radiative effect for fig. 12 and on observation of cloud percentage in the sky for
 13 fig. 7. However, this can also be an indication of changes in cloud properties (e.g. change in
 14 optically thin clouds that could have small radiation effect but are marked as cloudy conditions
 15 from the observer).

16 In Table 3 we have calculated the linear trends for the 1953-2012 period and for both clear sky and
 17 all sky measurements and the 1953-1982 and 1983-2012 sub-periods for clear sky measurements.
 18 Results shows comparable changes per decade (2% for the clear sky and 1.5% for the all sky cases).
 19 Seasonal analysis show that clear sky trends for summer, autumn and winter months are higher than
 20 the ones derived for all skies. Such differences are linked with the seasonal variability and long-
 21 term changes in cloudiness for the specific seasons.

22 Table 3: Clear sky and all sky data trends comparison for the whole 1953-2012 period and the two
 23 30-yr sub-periods (% per decade). Percentages in parenthesis show the limits of the 95% confidence
 24 bounds.

Season	Clear sky 1953-2012	All skies 1953-2012	Clear sky 1953-1982	Clear sky 1983-2012
Winter	0.91 (±2.31)	-6.43 (±3.83)	-7.01 (±3.16)	0.55 (±2.41)
Spring	1.22 (±1.12)	-0.60 (±3.10)	-0.92 (±1.11)	2.62 (±1.97)
Summer	2.03 (±0.78)	-1.14 (±2.90)	-0.36(±0.83)	1.31(±0.81)
Autumn	2.74 (±1.37)	-1.28 (±3.42)	-1.03(±1.84)	-1.48 (±1.73)
Year	2.17 (±1.21)	-2.33 (±2.28)	-1.44(±2.35)	1.94 (±2.08)

25

1 Clear sky results for the 1953-2012 period show significant positive changes in SSR for all seasons
2 except winter. Looking individually at the 1953-1982 and 1983–2012 periods we have calculated
3 significant negative trends only for the winter over the first and for summer and spring over the
4 second.

5 The effect of various parameters on SSR has been discussed by Kambezidis et al. (2016) in their
6 study about the global dimming/brightening effect over the Mediterranean in the period 1979-2012.
7 They show that the influence of parameters related to the atmospheric transparency, like water
8 vapor, aerosols and trace gases, as well as changes in the surface albedo on SSR have been larger in
9 the southern parts of the Mediterranean, over the Balkan countries and central Turkey. This
10 outcome is in agreement with the conclusion of the present study that other factors than cloudiness
11 play significant role in the SSR variations.

12 A comparison of the SSR results in Athens with visibility observations since 1931 (Founda et al.,
13 2016) did not show any correlation among SSR and horizontal visibility. For the first part of the
14 common dataset (1930-1959) the visibility decline is accompanied with a SSR increase. However
15 from 1950 till today visibility shows a monotonical decrease. The steep visibility decrease from
16 1931 till early 90's is not accompanied by a relative SSR decrease excluding individual sub-periods.

17 However, simulated SSR is driven purely by changes in sunshine duration, in this case the SD
18 variability in founda et al., 2014 is almost stable after 1950 so SD can not be also linked with the
19 visibility reported decrease. Studying the literature for similar cases, similar conclusions have been
20 drawn by Liepert and Kukla (1997) showing an SSR decrease over 30 years of measurements
21 accompanied by a visibility increase and no significant changes in the cloud cover conditions, in
22 Germany. This Athens SSR vs visibility relationship can be partly explained by the fact that SSR
23 and visibility have different response on cloud conditions, water vapor and rainfall, and also by the
24 fact that visibility is affected by aerosols only in the first few hundred meters above the surface,
25 while SSR is affected by the columnar AOD, which in the case of Athens can be significantly
26 different due to aerosol long-range transport in altitude (e.g. Saharan dust; Léon et al., 1999).

27

28 **5 Conclusions**

29 Surface solar radiation (SSR) at National Observatory of Athens, in the center of the city, is
30 presented using a unique dataset covering a period of 59 years (1954-2012). Sunshine duration (SD)
31 records for another 54 years have been used as a proxy to reconstruct SSR time series for the period
32 from 1900 to 2012.

1 The data accuracy of such historic radiation dataset is more difficult to be assessed especially going
2 back to the 50' and 60's where instruments, operational procedures and quality control were not at
3 the same level as in the recent 30 years. Quality assessment procedures in the presented time series
4 have been applied with criteria based on instrument characteristics and the availability of additional
5 collocated measurements. Year to year fluctuations of the measured SSR in addition to the reversal
6 of the downward tendencies at the ASNOA site adds credibility to the measured variations. That is
7 because a typical radiometer behavior is to lose sensitivity with time indicating spurious downward,
8 but not upward trends. The more recent (after 1986) SSR measurements can be characterized as
9 high-quality radiation data with known accuracy. Considering the measurements from 1954 to 1970
10 there has been sporadic reports mentioning the homogenization and calibration procedures, while
11 for 1970 to 1986 there is more information on the instrument quality control.

12 Reporting of the results from the 1954-1986 period should be accompanied with the fact that the
13 uncertainties of the measurements of this period are linked with higher uncertainties than after
14 1986. For the reconstruction of the 1900-1953 series, only the 1983-2012 SSR and SD
15 measurements were used in order not to link possible instrument uncertainties to the extrapolated
16 period. However, reconstruction of the 1900-1983 time series using the 1984-2012 dataset leads to
17 small differences in the determination of the long term trends, especially for more than 20 year
18 running average windows.

19 De-seasonalized SSR data analysis from 1900 to 2012 showed high month to month variability that
20 could reach up to 25%, mainly related with monthly cloudiness variations. During the period 1910-
21 mid-1930s where only few datasets have reported worldwide SSR results, we observe a -2.9% per
22 decade or a total of -8.7 % decrease in SSR, assuming linear changes in SSR during this period.
23 This early dimming was followed by a +5% per decade increase from 1930 to the 1950s. Similar
24 results have been found at Washington DC and at Potsdam, Germany (Stanhill and Achiman, 2016).
25 They have reported an early brightening at both locations in the 1930's. For the SSR measurement
26 period of 1953 to 1980, European related studies presented in Wild (2009) showed a -1% down to -
27 7% change per decade in SSR measurements over various European sites (dimming period). For the
28 Mediterranean region, Manara et al. (2016) showed a decrease of the order of -2% to -4% per
29 decade in Italy. We are reporting a change in SSR of -2% per decade in Athens. Finally, for the
30 brightening (1990-2012) phase again Wild et al. (2009) reported a +1.6% up to +4.7% per decade
31 positive change in SSR while we have calculated a +1.5% per decade, which is lower than the +(3-
32 6)% per decade reported in Manara et al. (2016) for Italy. A summary of the above findings can be
33 seen in table 4.

1
2 Table 4: Summary of per cent SSR changes per decade for various locations
3

Period	Location	Trend % per decade	Reference
1893-2012	Potsdam, Germany	0.71	Stanhill and Achiman, 2014
1900-2012	Athens, Greece	0.40 (± 0.26)	This work
1959-1988	Europe	-2.0	Ohmura and Lang, 1989
1971-1986	Europe	-2.3	Norris and Wild, 2007
1959-1985	Italy	-6.4(± 1.1) / -4.4(± 0.8)	Manara et al, 2016
1953-1982	Athens, Greece	-2.33(± 2.28)	This work
1985-2005	Europe	2.5	Wild, 2009
1990-2012	Italy	6.0 (± 1.1) / 7.7 (± 1.1)	Manara et al, 2016
1986-2013	Athens, Greece	0.80 (± 1.96)	This work

4
5 The decadal variations of SSR measured since 1954 at Athens, Greece, originate from the
6 alterations in the atmosphere's transparency (namely by clouds and aerosols). Using an analysis of
7 SSR calculations of all sky and clear sky (cloudless) days we end up that since cloud cover changes
8 during the 59 period were very small, most of the observed decadal changes can be related with
9 changes in the aerosol load of the area. An additional hint in support of this conclusion is the high
10 correlation of clear sky and all sky yearly SSR. We also found an anti-correlation between either
11 clear sky and all sky SSR measurements and AOD time series from AVHRR (1981-2009) or
12 ChArMEx (1979-2012). Looking at linear trends over the 59 year period, clear sky changes per
13 decade were 2% while it was 1.5% for all sky conditions. The most pronounced changes have been
14 calculated for summer and autumn seasons (2% and 2.7% respectively).

15 16 **Acknowledgements**

17 The authors wish to thank all the past and present NOA staff members who carefully collected and archived the long-
18 term data used in this study. This study contributes to the Chemistry-Aerosol Mediterranean Experiment (ChArMEx)
19 Work package 6 on trends. The work was partly funded by the Greek national project "Aristotelis", work package 1:
20 "Study of long term variations of Solar Radiation in the region of Athens". A. S. L. was supported by postdoctorial
21 fellowships (JCI-2012-12508 and RYC-2016-20784) and a project (CGL2014-55976-R) funded by the Spanish
22 Ministry of Economy, Industry and Competitiveness. We would like to thank the anonymous reviewers, Dr. Tanaka and
23 the editor Dr. Dulac for their efforts on substantially improving this work.

1 **References**

2 Alexandri, G., Georgoulas, A. K., Meleti , C., Balis, D., Kourtidis, K. A., Sanchez-Lorenzo, A.,
3 Trentmann, J. und Zanis, P.: A high resolution satellite view of surface solar radiation over the
4 climatically sensitive region of Eastern Mediterranean, *Atmosph. Res.*, 188, 107–121,
5 doi:10.1016/j.atmosres.2016.12.015, 2017.

6

7 Angell, J.K.: Variation in United States cloudiness and sunshine duration between 1950 and the
8 drought year of 1988, *Climate*, 3, 296-308, 1990.

9

10 Ångström, A.: Solar and terrestrial radiation. Report to the international commission for solar
11 research on actinometric investigations of solar and atmospheric radiation, *Q. J. R. Met. Soc.*, 50,
12 121-126, 1924.

13

14 Anton M., Vaquero J.M., Aparicio A.J.P.: The controversial early brightening in the first half of
15 20th century: A contribution from pyrliometer measurements in Madrid (Spain), *Global and
16 Planetary Change*, 115, 71-75, 10.1016/j.gloplacha.2014.01.013, 2014.

17

18 Anton M., R. Roman, A. Sanchez-Lorenzo, J. Calbo, J.M. Vaquero, Variability analysis of the
19 reconstructed daily global solar radiation under all-sky and cloud-free conditions in Madrid during
20 the period 1887–1950, *Atmos. Res.*, 191, 94-100, doi:10.1016/j.atmosres.2017.03.013, 2017

21

22 Bais, A. F., Drosoglou, T., Meleti, C., Tourpali, K., and Kouremeti, N.: Changes in surface
23 shortwave solar irradiance from 1993 to 2011 at Thessaloniki (Greece), *Int. J. Climatol.*, 33, 2871–
24 2876, doi:10.1002/joc.3636, 2013.

25

26 Chan, P. K., Zhao, X. P., and Heidinger, A. K.: Long-term aerosol climate data record derived from
27 operational AVHRR satellite observations, dataset *Papers in Geosciences*, 140791,
28 doi:10.7167/2013/140791, 2013.

29

1 Coulson, K.L.: Solar and Terrestrial Radiation: Methods and Measurements, Academic Press, New
2 York, 1975.
3

4 Drummond, A.J. and Roche, J.J.: Corrections to be applied to measurements made with Eppley (and
5 other) spectral radiometers when used with Schott colored glass filters, *J. App. Meteor.*, 4, pp.741-
6 744, doi:10.1175/1520-0450(1065)004<0741:<TBATM>2.0.CO,-2, 1965.
7

8 Dudok de Wit, T., Ermolli, I., Haberreiter, M., Kambezidis, H., Lam, M., Liliensten, J., Matthes, K.,
9 Mironova, I., Schmidt, H., Seppälä, A., Tanskanen, E., Tourpali, K. & Yair, Y. (Eds.): *Earth's
10 Climate Response to a Changing Sun*, Les Ulis Cedex: EDP Sciences , doi:10.1051/978-2-7598-
11 1733-7, 2015.
12

13 Flocas A., Estimation and prediction of global solar radiation over Greece, *Sol. Energy*, 24, 63-70,
14 doi: 10.1016/0038-092X(80)90021-3, 1980.
15

16 Founda, D., Kalimeris A., Pierros F.: Multi annual variability and climatic signal analysis of sun-
17 shine duration at a large urban area of Mediterranean (Athens). *Urban Climate*, 10, 815-830,
18 doi:10.1016/j.uclim.2014.09.008, 2014.
19

20 Founda, D., Kazadzis, S., Mihalopoulos, N., Gerasopoulos, E., Lianou, M. and Raptis, P.I.: Long-
21 term visibility variation in Athens (1931–2013): a proxy for local and regional atmospheric aerosol
22 loads. *Atm. Chem. Phys.*, 16, 11219-11236, doi:10.5194/acp-16-11219-2016, 2016.
23

24 Founda, D., Pierros, F. and Sarantopoulos, A.: Evidence of Dimming/Brightening over Greece from
25 long-term observations of Sunshine Duration and Cloud Cover, *Perspectives in Atmospheric
26 Sciences*, pp. 753-759, Springer, ISBN 978-3-319-35094-3, 2016.
27

28 García, R. D., Cuevas, E., García, O. E., Cachorro, V. E., Pallé, P., Bustos, J. J., Romero-Campos,
29 P. M., and de Frutos, A. M.: Reconstruction of global solar radiation time series from 1933 to 2013
30 at the Izaña Atmospheric Observatory, *Atmos. Meas. Tech.*, 7, 3139-3150, doi:10.5194/amt-7-
31 3139-2014, 2014.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Gilgen, H., Wild, M., and Ohmura, A., Means and trends of shortwave irradiance at the surface estimated from global energy balance archive data. *J. of Climate*, 11, 2042-2061, 1998.

Hulstrom, R. L. (Ed.): *Solar Resources, Solar Heat Technologies: Fundamentals and Applications 2*, The MIT Press, Cambridge, 1989.

Imamovic A., Tanaka K., Folini D., Wild M. Global dimming and urbanization: did stronger negative SSR trends collocate with regions of population growth? *Atm. Chem. Phys.*, 16, 2719-2725, doi:10.5194/acp-16-2719-2016, 2016.

ISO 9847: *Solar energy - Calibration of field pyranometers by comparison to a reference pyranometer*, International Organization for Standardization, 1992,

Jauregui, E., and E. Luyando.: Global radiation attenuation by air pollution and its effects on the thermal climate in Mexico City, *International Journal of Climatology* 19, no. 6: 683-694, 1999.

Kambezidis, H., Demetriou D., Kaskaoutis, D., Nastos, P.: Solar dimming/brightening in the Mediterranean EGU General Assembly 2010, held 2-7 May, 2010 in Vienna, Austria, p.10023, 2010.

Kambezidis, H., Kaskaoutis, D., Kalliampakos, G., Rashki, A. and Wild, M.: The solar dimming/brightening effect over the Mediterranean Basin in the period 1979 - 2012, *J. Atm. Sol.Terr, Phys.*, doi:10.1016/j.jastp.2016.10.006, 2016.

Katsoulis, B. and Leontaris, S.: The distribution over Greece of global solar radiation on a horizontal surface, *Agr. Methodol.*, 23, 217-229, doi: 10.1016/0002-1571(81)90106-0, 1981.

1 Katsoulis, B. and Papachristopoulos, E.: Analysis of solar radiation measurements at Athens
2 Observatory and estimates of solar radiation in Greece, *Sol. Energy*, 21, 217-226, doi:
3 10.1016/0038-092x(78)90024-5, 1978.

4

5 Kouremenos, D., Antonopoulos, K. and Domazakis, E.: Solar radiation correlations for Athens, *Sol.*
6 *Energy*, 35, 259-269, doi: 10.1016/0038-092x(85)90105-7, 1985

7

8 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C.,
9 Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J.,
10 Cooper, O. R., Kainuma, M., Mahowald, N., Mc-Connell, J. R., Naik, V., Riahi, K., and van
11 Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of
12 reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039,
13 doi:10.5194/acp-10-7017-2010, 2010.

14

15 Lean, J.: The Sun’s variable duration and its relevance for earth, *Annual Review of Astronomy and*
16 *Astrophysics* 35.1 33-67, 1997.

17

18 Léon, J.F., Chazette, P., and Dulac, F.: Retrieval and monitoring of aerosol optical thickness over an
19 urban area by spaceborne and ground-based remote sensing, *Appl. Opt.*, 38, 6918-6926, 1999.

20

21 Liepert B. and Kukla G., Decline in global solar radiation with increased horizontal visibility in
22 Germany between 1964 and 1990, *J. Climate*, 10, 2391-2401, doi: 10.1175/1520-
23 0442(1997)010<2391:DISGRW>2.0.Co;2, 1997.

24

25 LMDC,: Laboratory of Meteorological Device Calibration, <http://www.meteo.noa.gr/lmdc.html>, B.
26 Psiloglou, Scientific Responsible, personal contact, Nov. 2016.

27

28 Macris G.J., Solar energy and sunshine hours in Athens, Greece, *Mon. Wea. Rev.*, 87, 29-32, 1959

29

1 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J.
2 R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th Century industrial black carbon
3 emissions altered Arctic climate forcing, *Science*, 317, 5843, doi:10.1126/science.1144856, 2007.
4

5 Manara, V., Brunetti, M., Celozzi, A., Maugeri, M., Sanchez-Lorenzo, A., and Wild, M.: Detection
6 of dimming/brightening in Italy from homogenized all-sky and clear-sky surface solar radiation
7 records and underlying causes (1959–2013), *Atmos. Chem. Phys.*, 16, 11145-11161,
8 doi:10.5194/acp-16-11145-2016, 2016.
9

10 Matuszko, D.: Long-term variability in solar radiation in Krakow based on measurements of
11 sunshine duration, *Int. J. Climatol.*, 34, 228 – 234, doi: 10:1002/joc.3681, 2014.
12

13 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J.
14 R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th Century industrial black carbon
15 emissions altered Arctic climate forcing, *Science*, 317, 1381-1384, doi:10.1126/science.1144856,
16 2007.
17

18 Mishchenko, M. I., Geogdzhayev, I. V., Rossow, W. B., Cairns, B., Carlson, B. E., Lacis, A. A.,
19 Liu, L., and Travis, L. D.: Long-term satellite record reveals likely recent aerosol trend, *Science*,
20 315, 1543, doi:10.1126/science.1136709, 2007.
21

22 Nabat, P., Somot, S., Mallet, M., Chiapello, I., Morcrette, J.J., Solmon, F., Szopa, S., Dulac, F.,
23 Collins, W., Ghan, S. and Horowitz, L.W.,: A 4-D climatology (1979-2009) of the monthly
24 tropospheric aerosol optical depth distribution over the Mediterranean region from a comparative
25 evaluation and blending of remote sensing and model products, *Atmos. Meas. Tech.*, 6, 1287-1314,
26 doi:10.5194/amt-6-1287-2013.
27

28 Nabat, P., Somot, S., Mallet, M., Sanchez-Lorenzo, A. and Wild, M.: Contribution of anthropogenic
29 sulfate aerosols to the changing Euro-Mediterranean climate since 1980, *Geoph. Res. Lett.*, 41,
30 5605-5611, doi: 10.1002/2014GL060798, 2014.
31

1 Norris, J. R., and Wild M., Trends in aerosol radiative effects over Europe inferred from observed
2 cloud cover, solar “dimming,” and solar “brightening”, *J. Geophys. Res.*, 112, D08214,
3 doi:10.1029/2006JD007794, 2007.
4

5 Norris, J. R., and M. Wild : Trends in aerosol radiative effects over China and Japan inferred from
6 observed cloud cover, solar “dimming,” and solar “brightening”, *J. Geophys. Res.*, 114, D00D15,
7 doi:10.1029/2008JD011378, 2009.
8

9 Notaridou V. and D. Lalas, The distribution of global and net radiation over Greece, *Sol. Energy* 22,
10 504-514, doi: 10.1016/0038-092X(79)90022-7 1978.
11

12 Ohmura, A., Observed long-term variations of solar irradiances at the Earth’s surface, *Space Sci.*
13 *Rev.*, 125, 111–128, doi:10.1007/s11214- 006-9050-9, 2006.
14

15 Ohmura, A.: Observed decadal variations in surface solar radiation and their causes, *J. Geophys.*
16 *Res.*, 114, D00D05, doi:10.1029/2008JD011290, 2009.
17

18 Ohmura, A. and H. Lang: Secular variation of global radiation over Europe, in *Current Problems in*
19 *Atmospheric Radiation*, edited by J. Lenoble and J. F. Geleyn, pp. 98–301, Deepak, Hampton, Va. ,
20 1989.
21

22 Ruckstuhl C, Philipona R, Behrens K, Collaud Coen M, Durr B, Heimo. A, Matzler C, Nyeki S,
23 Ohmura A, Vuilleumier L, Weller M, Wehrli C, Zelenka A.: Aerosol and cloud effects on solar
24 brightening and the recent rapid warming, *Geophys. Res. Lett.*, 35, L12708, doi:
25 10.1029/2008gl034228, 2008.
26

27 Sanchez-Lorenzo, A., Calbó J, Martin-Vide J: Spatial and temporal trends in sunshine duration over
28 Western Europe (1938-2004), *J. of Climate* 21, 6089-6098, doi: 10.1175/2008JCLI2442.1, 2008.
29

30 Sanchez-Lorenzo, A. and Wild, M.: Decadal variations in estimated surface solar radiation over
31 Switzerland since the late 19th century, *Atmos. Chem. Phys.*, 12, 8635-8644,
<https://doi.org/10.5194/acp-12-8635-2012>, 2012.

1
2
3
4
5

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Sanchez-Lorenzo, A., Wild M., Brunetti M., Gujjarro J. A., Hakuba, M. Z. , Calbó, J. , Mystakidis, S. and Bartok, S.: Reassessment and update of long-term trends in downward surface shortwave radiation over Europe (1939–2012), *J. Geophys. Res. Atmos.*, 120, 9555–9569, doi:10.1002/2015JD023321, 2015.

Sanchez-Romero, A., A. Sanchez-Lorenzo, J. Calbó, J. A. González, and C. Azorin-Molina : The signal of aerosol-induced changes in sunshine duration records: A review of the evidence, *J. Geophys. Res. Atmos.*, 119, 4657–4673, doi:10.1002/2013JD021393, 2014.

Stanhill, G. and Ahiman, O.: Early global radiation measurements: a review. *Int. J. Climatol.*, 37, 1665-1671, doi:10.1002/joc.4826, 2017.

Stanhill, G.: Global irradiance, air pollution and temperature changes in the Arctic, *Philos. Trans. R. Soc. A*, 352, 247 –258, doi:10.1098/rsta.1995.0068, 1995.

Stanhill, G., and Cohen S.: Recent changes in solar irradiance in Antarctica, *J. Clim.*, 10, 2078–2086, doi: 10.1175/1520-0442(1997)010<2078:RCISII>2.0.CO;2, 1997.

Stanhill, G. and Cohen, S.: Solar radiation changes in the United States during the twentieth century: Evidence from sunshine duration measurements. *J. of Climate*, 18, 1503-1512, doi: 10.1175/JCLI3354.1, 2005.

Stanhill, G. and Möller, M.: Evaporative climate change in the British Isles. *International Journal of Climatology*, 28(9), pp.1127-1137, 2008.

Stanhill, G., and O. Ahiman: Radiative forcing and temperature change at Potsdam between 1893 and 2012, *J. Geophys. Res. Atmos.*, 119, 9376–9385, doi:10.1002/2014JD021877, 2014.

1 Streets, D. G., Y. Wu, and M. Chin, Two-decadal aerosol trends as a likely explanation of the
2 global dimming/brightening transition, *Geophys. Res. Lett.*, 33, L15806,
3 doi:10.1029/2006GL026471. 2006.
4

5 Tanaka, K., Ohmura, A., Folini, D., Wild, M., and Ohkawara, N.: Is global dimming and
6 brightening in Japan limited to urban areas?, *Atmos. Chem. Phys.*, 16, 13969-14001,
7 <https://doi.org/10.5194/acp-16-13969-2016>, 2016.
8

9 Vautard, R., and P. Yiou, Control of recent European surface climate change by atmospheric flow,
10 *Geophys. Res. Lett.*, 36, L22702, doi:10.1029/2009GL040480, 2009.
11

12 Vignola F., Michalsky J., and Stoffel T., *Solar and Infrared Radiation Measurements*, ISBN
13 9781439851906, CRC Press, 2012.
14

15 Wang, K. C., Dickinson, R. E., Wild, M., and Liang, S.: Atmospheric impacts on climatic
16 variability of surface incident solar radiation, *Atmos. Chem. Phys.*, 12, 9581-9592,
17 doi:10.5194/acp-12-9581-2012, 2012.
18

19 Wild M.: Global dimming and brightening: A review, *J. of Geoph. Res.*, 114, D00D16, DOI:
20 10.1029/2008JD011470, 2009.
21

22 Wild M, Folini D, Hakuba MZ, Schär C, Seneviratne SI, Kato S, Rutan D, Ammann C, Wood EF,
23 König-Langlo G.: The energy balance over land and oceans: an assessment based on direct
24 observations and CMIP5 climate models, *ClimDyn* 44:3393–3429. doi:10.1007/s00382-014-2430-
25 z, 2015.
26

27 Wild, M.: Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for
28 global warming. *WIREs Clim. Change*, 7, 91–107, doi:10.1002/wcc.372, 2016
29

1 WMO: “Measurement of radiation”, in Guide to Meteorological Instrument and Observing
2 Practices, Chapter 9, fifth ed., WMO-No. 8, 1983.
3
4 WMO: Scientific Assessment of Ozone Depletion: 2010, report 52, World Meteorological
5 Organization (WMO), Global Ozone Research and Monitoring Project, Geneva, Switzerland;
6 National Oceanic and Atmospheric Administration (NOAA), Washington, DC, USA; National
7 Aeronautics and Space Administration (NASA), Washington, DC, USA; United Nations
8 Environment Program (UNEP), Nairobi, Kenya; and the European Commission, Research
9 Directorate General, Brussels, Belgium, 2010.
10
11 U. Yildirim, I. O. Yilmaz, and B. G. Akinoglu, “Trend analysis of 41 years of sunshine duration
12 data for Turkey,” Turkish Journal of Eng. Env. Sci., 37, 286–305, doi: 10:3906/muh-1301-11, 2013.
13
14 Zabara K., Estimation of the global solar radiation in Greece, Sol. & Wind Tech., 3, 267-272, 1986.
15
16 Zerefos, C.S., K. Eleftheratos, C. Meleti, S. Kazadzis, A. Romanou, C. Ichoku, G. Tselioudis, and
17 A. Bais: Solar dimming and brightening over Thessaloniki, Greece, and Beijing,
18 China. Tellus, 61B, 657-665, doi:10.1111/j.1600-0889.2009.00425.x, 2009.
19
20 Zerefos, C. S., Tourpali, K., Eleftheratos, K., Kazadzis, S., Meleti, C., Feister, U., Koskela, T., and
21 Heikkilä, A.: Evidence of a possible turning point in solar UV-B over Canada, Europe and Japan,
22 Atmos. Chem. Phys., 12, 2469-2477, <https://doi.org/10.5194/acp-12-2469-2012>, 2012.
23
24 Zhao X., Chan, P., and NOAA CDR Program: NOAA Climate Data Record (CDR) of AVHRR
25 Daily and Monthly Aerosol Optical Thickness over Global Oceans, Version 2.0.AOT1, NOAA
26 National Centers for Environmental Information, doi:10.7289/V5SB43PD, 2014.