

# H<sub>2</sub>O<sub>2</sub> modulates the energetic metabolism of the cloud microbiome

N. Wirgot<sup>1</sup>, V. Vinatier<sup>1</sup>, L. Deguillaume<sup>2</sup>, M. Sancelme<sup>1</sup>, and A.-M. Delort<sup>1</sup>

<sup>1</sup>Université Clermont Auvergne, CNRS, Sigma-Clermont, Institut de Chimie de Clermont-Ferrand, 63000 Clermont-Ferrand, France

<sup>2</sup>Université Clermont Auvergne, CNRS, Laboratoire de Météorologie Physique, 63000 Clermont-Ferrand, France

Correspondence to: A.-M. Delort (A-marie.delort@uca.fr)

**Abstract.** Chemical reactions in clouds lead to oxidation processes driven by radicals (mainly HO<sup>•</sup>, NO<sub>3</sub><sup>•</sup> or HO<sub>2</sub><sup>•</sup>) or strong oxidants such as H<sub>2</sub>O<sub>2</sub>, O<sub>3</sub>, nitrate and nitrite. Among those species, hydrogen peroxide plays a central role in the cloud chemistry by driving its oxidant capacity. In cloud droplets, H<sub>2</sub>O<sub>2</sub> is transformed by microorganisms which are metabolically active. Biological activity can therefore impact the cloud oxidant capacity. The present article aims at highlighting the interactions between H<sub>2</sub>O<sub>2</sub> and microorganisms within the cloud system.

First, experiments were performed with selected strains studied as reference isolated from clouds in microcosms designed to mimic the cloud chemical composition, including the presence of light and iron. Biotic and abiotic degradation rates of H<sub>2</sub>O<sub>2</sub> were measured and results showed that biodegradation was the most efficient process together with photo-Fenton process. H<sub>2</sub>O<sub>2</sub> strongly impacted the microbial energetic state as shown by adenosine triphosphate (ATP) measurements in the presence and absence of H<sub>2</sub>O<sub>2</sub>. This ATP depletion was not due to the loss of cell viability. Secondly, correlation studies were performed based on real cloud measurements from 37 clouds samples collected at the PUY station (1465 m a.s.l., France). The results support a strong correlation between ATP and H<sub>2</sub>O<sub>2</sub> concentrations and confirm that H<sub>2</sub>O<sub>2</sub> modulates the energetic metabolism of the cloud microbiome. The modulation of microbial metabolism by H<sub>2</sub>O<sub>2</sub> concentration could thus impact cloud chemistry, in particular the biotransformation rates of carbon compounds and consequently can perturb the way the cloud system is modifying the global atmospheric chemistry.

Keywords: Cloud water, Microorganisms, Hydrogen peroxide, Energetic metabolism, Atmospheric chemistry

## 1 Introduction

The atmosphere is an oxidizing medium where trace gases are transformed/removed by oxidation including methane and other organic compounds, carbon monoxide, nitrogen oxides, and sulfur gases. Evaluating the oxidizing power of the atmosphere is crucial since it controls pollutant formation and fate, aerosol production and greenhouse radiative forcing (Thompson, 1992).

In this context, hydroperoxides (ROOH) contribute to the oxidizing power of the troposphere (Lee et al., 2000; Herrmann et al., 2015) by controlling the cycling of HO<sub>x</sub> radicals (HO<sup>•</sup>, HO<sub>2</sub><sup>•</sup>). They can serve as temporary reservoirs of HO<sub>x</sub> radical since, for example, their photolysis and reactivity will regenerate HO<sup>•</sup> radicals. Among

37 hydroperoxide, hydrogen peroxide is a key gas phase atmospheric chemical species (Vione et al., 2003) with  
38 concentration in the gas phase in the ppb<sub>v</sub> level or less. The atmospheric concentration of H<sub>2</sub>O<sub>2</sub> is impacted by a  
39 variety of meteorological parameters (*e.g.* actinic flux, temperature and relative humidity) and is affected by the  
40 levels of chemical species such as VOCs, CO, O<sub>3</sub>, and NO<sub>x</sub> (Lee et al., 2000). One of the significant parameters  
41 controlling the evolution of H<sub>2</sub>O<sub>2</sub> concentration is the actinic flux intensity. Diurnal and seasonal variations of  
42 hydrogen peroxide are shown by field measurements with higher concentrations during the day and in summer  
43 than during the night and in winter. This is linked to the atmospheric formation of H<sub>2</sub>O<sub>2</sub> that results from a series  
44 of photochemical reactions creating free radicals followed by corresponding radical reactions with appropriate  
45 precursor substances.

46 In the presence of atmospheric liquid water (cloud, fog, rain), H<sub>2</sub>O<sub>2</sub> is rapidly dissolved because of its high  
47 Henry's law constant ( $7.7 \cdot 10^4$  M/atm at 298K; Sander, 2014). In this liquid phase, it is also produced by aqueous  
48 phase reactivity (Möller, 2009). Several field campaigns have reported H<sub>2</sub>O<sub>2</sub> concentrations in atmospheric water  
49 in the  $\mu$ M range (Gunz and Hoffmann, 1990; Marinoni et al., 2011; Deguillaume et al., 2014; Li et al., 2017).  
50 Hydrogen peroxide plays a central role in various important chemical processes in clouds. First, H<sub>2</sub>O<sub>2</sub> is  
51 considered as the most important oxidant for the conversion of sulfite to sulfate for pH lower than 5.5, therefore  
52 contributing significantly to the acidification of clouds and precipitations (Deguillaume et al., 2004; Shen et al.,  
53 2012). Second, the photolysis of H<sub>2</sub>O<sub>2</sub> will lead to an efficient production of the hydroxyl radical HO<sup>•</sup> (Arakaki  
54 et al., 2013) and recent study have shown that this can be a dominant aqueous source (Bianco et al., 2015). They  
55 can also directly oxidize organic compounds in the aqueous phase (Schöne and Herrmann, 2015). Finally, H<sub>2</sub>O<sub>2</sub>  
56 is involved in redox processes leading to the conversion of reactive free radicals and trace metals such as iron  
57 (Kieber et al., 2001; Deguillaume et al., 2005; Hems et al., 2017). Consequently, H<sub>2</sub>O<sub>2</sub> is a key chemical  
58 compound controlling the aqueous phase oxidant capacity and leading to the transformation of inorganic and  
59 organic compounds present in the atmospheric aqueous phase. The resulting inorganic and organic products can  
60 contribute to the aerosol phase when the cloud evaporates leading to climatic effect.

61 A few decades ago, living microorganisms **were observed** in cloud water (Sattler et al., 2001; Amato et al.,  
62 2005, 2007a,b; Wei et al., 2017). Particularly through measurements of adenosine triphosphate (ATP) and  
63 anabolic precursors or nutrient incorporation rates, it has been shown that cloud microorganisms are  
64 metabolically active and play an important role in cloud chemical reactivity (Sattler et al., 2001; Amato et al.,  
65 2007a; Hill et al., 2007; Väitilingom et al., 2012, 2013). Several studies performed on simplified or real  
66 microcosms have demonstrated that cloud microorganisms are able to degrade carbon compounds (Ariya et al.,  
67 2002; Amato et al., 2005, 2007c; Husarova et al., 2011; Väitilingom et al., 2010, 2011, 2013; Matulová et al.,  
68 2014); recent studies have also shown that this could be the case in the air (Krumins et al., 2014).

69 Microorganisms are also in direct interaction with oxidant species in clouds (iron, hydroxyl radical, hydrogen  
70 peroxide, *etc.*). Väitilingom et al. (2013) have demonstrated that microorganisms present in real cloud water are  
71 able to efficiently degrade hydrogen peroxide. This suggests that cloud microorganisms found strategies to  
72 survive and resist stresses encountered in this medium and in particular oxidative stress. In this context, Joly et  
73 al. (2015) have conducted laboratory experiments to investigate the survival of selected strains (bacteria and  
74 yeasts) isolated from cloud waters, in the presence of various concentrations of hydrogen peroxide. The results  
75 showed that the survival rates of the studied strains were not affected by H<sub>2</sub>O<sub>2</sub> exposure. In addition, the strains

76 were exposed to artificial UV-visible light mimicking the natural solar irradiation inside clouds. No significant  
77 impact on the survival of the bacterial strains was observed.

78 These results have been confirmed in real cloud water, including the microbial community and chemical  
79 complexity (iron, H<sub>2</sub>O<sub>2</sub>, *etc.*), incubated in a photo-bioreactor designed to mimic cloud conditions (Vaithilingom  
80 et al., 2013). Thanks to ADP/ATP ratio measurements, reflecting the energetic metabolism of microorganisms,  
81 exposed or not to solar radiation, it has been shown that microorganisms were not impacted by artificial light and  
82 consequently by the generation of radicals from H<sub>2</sub>O<sub>2</sub> photo-reactivity. In addition, H<sub>2</sub>O<sub>2</sub> is efficiently degraded  
83 by catalases and peroxidases involved in oxidative metabolism. Solar light did not modify the degradation rates  
84 of H<sub>2</sub>O<sub>2</sub>, demonstrating that the biological process was not inhibited by UV radiations and radicals.

85 Indeed solar light can indirectly impact the viability of cells by the production of reactive oxygenated species  
86 (ROS) including HO<sup>•</sup> and O<sub>2</sub><sup>•-</sup> radicals. The main sources of these radicals are H<sub>2</sub>O<sub>2</sub> photolysis or Fenton and  
87 photo-Fenton reactions involving iron (Fe) and H<sub>2</sub>O<sub>2</sub>. Most of these compounds can cross the cytoplasmic  
88 membrane by diffusion. Aerobic microorganisms can also produce similar ROS during respiration. These  
89 radicals can potentially damage major cellular components such as proteins, DNA and lipids and lead to cellular  
90 death. Because microorganisms usually are protected against these ROS, they can specifically modify their  
91 metabolism to face oxidative stress taking place in clouds. Therefore, microorganisms utilize various  
92 mechanisms involved in oxidative stress metabolism such as i) the production of pigments and antioxidant  
93 molecules (vitamins, glutathione, *etc.*) which can scavenge radicals or ii) the production of specific enzymes  
94 such as superoxide dismutase which can transform O<sub>2</sub><sup>•-</sup> into H<sub>2</sub>O<sub>2</sub>. H<sub>2</sub>O<sub>2</sub> can be dismutated or reduced  
95 respectively by catalases and other peroxidases (Delort et al., 2017).

96 The studies from Vaithilingom et al. (2013) and Joly et al. (2015) highlighted the interactions between biological  
97 activity and oxidants in clouds. **It is crucial to consider all sinks and sources of H<sub>2</sub>O<sub>2</sub>, especially in atmospheric  
98 chemistry models, since H<sub>2</sub>O<sub>2</sub> impacts many relevant processes in the atmosphere.** In the present work, we  
99 artificially reproduced cloud conditions in microcosms to study the biotic and abiotic transformation of H<sub>2</sub>O<sub>2</sub>  
100 and, conversely, the impact of hydrogen peroxide on the metabolism of cloud microorganisms.

101 For this purpose, we decided to study individually the effect of parameters interacting with H<sub>2</sub>O<sub>2</sub>: UV radiation,  
102 iron and bacteria. Under various experimental conditions, the degradation rates of H<sub>2</sub>O<sub>2</sub> were followed to  
103 highlight how individual parameters control its transformation. Moreover, the impact of H<sub>2</sub>O<sub>2</sub> on the energetic  
104 state of the bacterial cells was evaluated by measuring the ATP concentration over time when the cells were  
105 exposed or not to H<sub>2</sub>O<sub>2</sub>. In order to confirm our laboratory results on the interaction between microorganisms  
106 and H<sub>2</sub>O<sub>2</sub>, we performed a correlation analysis considering bio-physico-chemical parameters measured in real  
107 cloud samples collected at the PUY station. **This work will lead to a better description of the mechanisms linking  
108 biological activity and cloud reactivity.**

## 109 **2 Material and methods**

### 110 **2.1 Bacterial strains and growth conditions**

111 *Pseudomonas graminis*, 13b-3, DQ512786; *Pseudomonas syringae*, 13b-2, DQ512785, *Sphingomonas sp.*, 14b-  
112 5, DQ512789 were grown in 10 mL of R2A medium (Reasoner and Geldreich, 1985) under stirring (200 r.p.m)  
113 at 17°C for approximately 17 h, 24 h or 48 h, depending on the strain. **The three selected bacterial strains**

114 belonging to the Gamma-Proteobacteria (*Pseudomonas*) and Alpha- Proteobacteria classes (*Sphingomonas*)  
115 were isolated from cloud water and are representative of the genera most frequently found in cloud water  
116 samples (Väitilingom et al., 2012) collected at the PUY site.

117 Cells in the exponential growth phase were collected by centrifugation for 3 min at around 10000 g. The  
118 supernatant was removed and the bacterial pellet was suspended and washed twice with an artificial cloud  
119 solution and incubated in microcosms to perform biodegradation experiments (see section 2.2). The bacterial cell  
120 concentration was estimated by optical density at 575 nm to obtain a concentration close to  $10^6$  cell mL<sup>-1</sup>.  
121 Finally, the concentration of cells was precisely determined by flow cytometry analysis (BD FacsCalibur Becton-  
122 Dickinson;  $\lambda_{exc}$  = 488 nm;  $\lambda_{em}$  = 530 nm) using a method based on the addition of a fluorochrome (SYBR-green)  
123 for their counting (Marie et al., 1999).

124

## 125 **2.2. Incubations in microcosms**

126 Microcosms were designed to simulate as much as possible the water phase of cloud waters. They provide the  
127 opportunity to work under artificial solar light condition and also in the presence of microorganisms. The  
128 experiments were performed under bulk conditions as cloud droplets cannot be reproduced in these bioreactors  
129 (Infors HT Multitron II).

130 For irradiation condition the bioreactor was equipped with lamps that emit UV-radiation (Sylvania Reptistar; 15  
131 W; 6500 K) to mimic solar light measured directly in clouds at the PUY station (Fig. SM1). The incubation  
132 flasks were Pyrex crystallizers covered with a Pyrex filter and equipped with Teflon tubes of 8 mm Ø plugged  
133 with sterile cotton, letting air and light pass (see Väitilingom et al. 2013) while for dark conditions they were  
134 amber Erlenmeyer flasks.

135 All incubation flasks contained 100 mL of artificial cloud solution under agitation (130 rpm), its composition  
136 was first described in Väitilingom et al. (2011). This solution was mimicking cloud chemical composition from  
137 cloud samples classified as “marine” following the work from Deguillaume et al. (2014) at the PUY station The  
138 major part of the collected cloud samples were classified as marine (52%) supporting our choice for the artificial  
139 cloud composition. Stock solutions of this artificial cloud medium were prepared with the following  
140 concentrations: 200  $\mu$ M for acetic acid (CH<sub>3</sub>COOH; Acros organics), 145  $\mu$ M for formic acid (HCOOH; Fluka),  
141 30  $\mu$ M for oxalic acid (H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>;Fluka), 15  $\mu$ M for succinic acid (H<sub>6</sub>C<sub>4</sub>O<sub>4</sub>; Fluka), 800  $\mu$ M for ammonium nitrate  
142 (H<sub>4</sub>N<sub>2</sub>O<sub>3</sub>; Fluka), 100  $\mu$ M for magnesium chloride hexahydrate (MgCl<sub>2</sub>, 6H<sub>2</sub>O; Sigma-Aldrich), 50  $\mu$ M for  
143 potassium sulfate (K<sub>2</sub>SO<sub>4</sub>; Fluka), 400  $\mu$ M for calcium chloride dihydrate (CaCl<sub>2</sub>, 2H<sub>2</sub>O; Sigma-Aldrich), 2000  
144  $\mu$ M for sodium chloride (NaCl; Sigma-Aldrich), 1100  $\mu$ M for sodium hydroxide (NaOH; Merck), 315  $\mu$ M for  
145 sulfuric acid (H<sub>2</sub>SO<sub>4</sub>; Sigma-Aldrich). Finally, the obtained solution was adjusted to pH 6 as necessary with a  
146 few drops of the solutions of NaOH or H<sub>2</sub>SO<sub>4</sub> used for the preparation of the marine artificial cloud water  
147 solution and sterilized by filtration (Polyethersulfone membrane, 0.20  $\mu$ m; Fisher Scientific) before use.

148 For biotic conditions, the flasks were inoculated at  $10^6$  bacterial cells per mL. The artificial cloud water solution  
149 was ten times more concentrated than a real cloud water solution in order to stabilize the pH. This was also the  
150 case for bacteria concentration because the bacteria/substrate ratio should be kept identical to that of real cloud.  
151 Indeed, it has been demonstrated that if this ratio is maintained, the degradation rate remains constant

152 (Vaïtilingom et al., 2010). The equipment was sterilized by autoclaving at 121°C for 20 minutes and all  
153 manipulations were performed under sterile conditions.

154 Depending on the conditions, hydrogen peroxide and iron complex (Fe-[EDDS]) were added or not to the  
155 solution in the incubators. These two compounds are present in marine cloud water collected at the PUY station  
156 at average concentrations of 7.5 µM (with a dispersion of mean values ranging from 0.1 – 20.8 µM) for H<sub>2</sub>O<sub>2</sub>  
157 and 0.5 µM (with a dispersion of mean values ranging from BDL. – 4.9) for Fe(III) (Deguillaume et al., 2014).  
158 In the cloud aqueous phase, Fe(III) may be complexed by organic compounds. Recently, it has been  
159 hypothesized that iron can be chelated by other organic ligands of biological origin (Herckes et al., 2013;  
160 Herrmann et al., 2015), and in particular by siderophores (Vinatier et al., 2016) that are ligands characterized by  
161 high complexing constants ( $K > 10^{20}$ ). Fe-[EDDS] was chosen as an iron(III) complex model because this ligand  
162 has a complexing constant for iron very close to the values for siderophores. Moreover, it is known to be stable  
163 at the working pH of 6.0 and because its chemistry has been studied in details by Li et al. (2010).

164 Hydrogen peroxide solution was prepared from a commercial solution (H<sub>2</sub>O<sub>2</sub>, 30%; not stabilized Fluka  
165 Analytical). 1:1 stoichiometry iron complex solution was prepared from iron (III) chloride hexahydrate (FeCl<sub>3</sub>,  
166 6H<sub>2</sub>O; Sigma-Aldrich) and from (S,S)- ethylenediamine-N,N'-disuccinic acid trisodium salt (EDDS, 35% in  
167 water). The hydrogen peroxide solution and the iron complex solution were freshly prepared before each  
168 experiment and the final working concentrations were fixed at 20 µM and 4 µM respectively, in agreement with  
169 the real concentrations detected in samples collected at the PUY station multiplied by a factor ten when median  
170 values measured in marine cloud waters are considered (Deguillaume et al., 2014).

171 In addition, the working temperature was fixed at 17°C which is the average temperature of cloud samples in  
172 summer. For all the incubation conditions, samples were taken at regular intervals, and stored at -20 °C before  
173 analysis.

### 174 **2.3 Analyses**

175 Hydrogen peroxide was quantified with a miniaturised Lazrus fluorimetric assay (Lazrus et al., 1985;  
176 Vaïtilingom et al., 2013). This method is based on a reaction between hydrogen peroxide, Horse Radish  
177 Peroxidase (HRP) and 4-hydroxyphenylacetic acid that produces a fluorescent dimeric compound. Fluorescence  
178 readings (Safire II TECAN<sup>®</sup>;  $\lambda_{exc} = 320$  nm,  $\lambda_{em} = 390$  nm) were made in a 96 well plate format.

179 Bioluminescence was used to analyse adenosine triphosphate (ATP) concentrations (Glomax<sup>®</sup> 20/20 single tube  
180 luminometer from Promega). This technique is based on an enzymatic reaction involving luciferin and  
181 luciferase. The protocol used was adapted from Biothema<sup>®</sup> commercial kit instructions (Koutny et al., 2006).

182 In order to determine the survival rate of microorganisms in the presence of hydrogen peroxide (20 µM), plate-  
183 counts were performed on R2A agar medium at the beginning of each experiment and after 3, 7 and 24 hours of  
184 incubation. Plates were incubated 3 days at 17°C before CFU counts.

### 185 **2.4 Determination of the initial degradation rates of hydrogen peroxide**

186 The processing of data was done with the Origin 6.1 software.

187 The graphs representing the hydrogen peroxide concentration decrease as a function of time were plotted. The  
188 degradation rates have been calculated from the initial slopes (the first five time points i.e. between 0 and 2  
189 hours) normalized with the concentrations of cells. During these two hours no cell growth was observed.

190

## 191 **2.5 Cloud sampling and statistical analysis**

192 Cloud water sampling was performed on the summit of the PUY station (summit of the puy de Dôme, 1465 m  
193 a.s.l., France) which is part of the atmospheric survey networks EMEP, GAW, and ACTRIS. The detachable part  
194 of the impactor was sterilized beforehand by autoclave at 121°C for 20 min and the fixed part was rinsed with  
195 alcohol at 70° and then with sterile water just before sampling.

196 Between 2004 and 2013, 89 cloud events were collected at the PUY station. The origin of these clouds can be  
197 analyzed according to their back trajectories in four sectors (North/West, South/West, West and North/East).  
198 They can be also considered in four different categories considering their chemical composition (marine,  
199 continental, highly marine and polluted) as described in Deguillaume et al. (2014).

200 Various parameters were measured including ATP, bacteria and fungi concentration, inorganic and organic  
201 species concentration ( $\text{H}_2\text{O}_2$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Cl}^-$ , acetate, formate, oxalate,  $\text{Na}^+$ ,  $\text{NH}_4^+$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ), temperature  
202 and pH (see Table SM1 for details). More information about the cloud sample collection is given in Deguillaume  
203 et al. (2014).

204 These data were used in this study to achieve statistical analyses. R software 3.1.2 was used to carry out  
205 principal component analysis (PCA). The data of 37 cloud events (of 89 total) were selected after the constraints  
206 related to this statistical analysis (e.g. the cloud events with more than 10 percent of missing values (parameters)  
207 were not considered) were applied.

208 In addition, statistical significance test was evaluated using PAST software (Hammer et al., 2001). Mean  
209 difference was considered to be statistically significant for a p-value less than 0.05.

## 210 **3 Results**

211 The interactions between  $\text{H}_2\text{O}_2$ , which is one of the major oxidant present in clouds, and microorganisms were  
212 investigated by performing experiments in artificial cloud microcosms but also by considering chemical and  
213 biological parameters measured in real cloud samples over long period at the PUY station.

### 214 **3.1 Experiments in artificial cloud water microcosms**

215 Experiments were conducted in microcosms mimicking cloud conditions in which each important parameters  
216 including  $\text{H}_2\text{O}_2$ , iron, light and the presence of bacteria could be studied individually or in complementarity.

#### 217 **Hydrogen peroxide degradation in artificial cloud water**

218  $\text{H}_2\text{O}_2$  degradation was monitored periodically over a 8 h period. The kinetic profiles were similar for the three  
219 strains. Results obtained with *Pseudomonas graminis* (13b-3) are illustrated in Figure 1 whereas the results  
220 obtained with the other strains are presented for information in Figure SM2.

221 Under abiotic condition (Fig. 1, orange traces), the degradation of hydrogen peroxide is clearly effective in the  
222 presence of artificial solar light and Fe-[EDDS] complex, due to the photo-Fenton reaction, with an initial  
223 degradation rate of  $1.07 \cdot 10^{-9} \text{ mol L}^{-1} \text{ s}^{-1}$  (Table 1(a)). After 150 min this degradation rate decreases in parallel  
224 with EDDS by oxidation occurs (Li et al., 2010). In the presence of the Fe-[EDDS] complex alone and in the  
225 absence of light, hydrogen peroxide is almost not degraded. Indeed, the degradation rate of  $\text{H}_2\text{O}_2$  due to the

226 Fenton reaction is much lower ( $2.23 \cdot 10^{-10} \text{ mol L}^{-1} \text{ s}^{-1}$ ) than the value obtained with the photo-Fenton reaction.  
227 Exposing the microcosm only to our light conditions, the photolysis reaction of  $\text{H}_2\text{O}_2$  is extremely slow ( $1.38 \cdot 10^{-10}$   
228  $\text{ mol L}^{-1} \text{ s}^{-1}$ ) due to the low absorption of  $\text{H}_2\text{O}_2$  in the solar spectrum measured inside a cloud and that was  
229 reproduced by the lamps used for these experiments (Fig. SM1).

230 For the biotic conditions (Fig.1, green traces), the initial biodegradation rates are summarized in Table 1(b).  
231 These results show that, under our experimental conditions, hydrogen peroxide was degraded more efficiently in  
232 the presence of bacteria even if the values obtained stay within the same order of magnitude compared to the  
233 abiotic conditions with artificial light and Fe-[EDDS] complex. *Pseudomonas graminis* (13b-3) and  
234 *Pseudomonas syringae* (13b-2) are the most active strains followed by *Sphingomonas sp* (14b-5). For each  
235 strain, biodegradation rates are within the same order of magnitude without wide variations depending on the  
236 tested conditions, *i.e.* in the presence or absence of artificial light and of Fe-[EDDS] complex.

237 The selected strains all degrade  $\text{H}_2\text{O}_2$  within the same order of magnitude (average value for the three strains and  
238 for the condition with iron and light  $1.76 \cdot 10^{-9} \text{ mol L}^{-1} \text{ s}^{-1}$  and with iron without light  $1.40 \cdot 10^{-9} \text{ mol L}^{-1} \text{ s}^{-1}$ ). In  
239 Vaitilingom et al. (2013), the biodegradation rates of  $\text{H}_2\text{O}_2$  were found within the same order of magnitude  
240 (average value for two distinct clouds with light  $0.98 \cdot 10^{-9} \text{ mol.L}^{-1} \text{ s}^{-1}$  and without light  $0.29 \cdot 10^{-9} \text{ mol L}^{-1} \text{ s}^{-1}$ ). The  
241 results obtained are within the same order of magnitude of values in real cloud environment thereby validating  
242 our microcosm conditions. This demonstrates that under our experimental conditions, the selected strains  
243 degrade  $\text{H}_2\text{O}_2$  like the microbiome of real clouds. In addition it validates our approach to separately analyse the  
244 influence of each parameter (Fe,  $\text{H}_2\text{O}_2$ , light,...) on the microbial energetic state metabolism in artificial marine  
245 cloud solution detailed in the next section.

#### 246 **Impact of the $\text{H}_2\text{O}_2$ on the microbial energetic state in artificial marine cloud solution**

247 The impact of the presence of  $\text{H}_2\text{O}_2$  on the energetic state of the bacterial cells was evaluated by measuring the  
248 time evolution of ATP concentration for the three strains (Fig. 2). The ATP concentration was measured in the  
249 presence (Fig. 2a, b, c - black square) or absence (Fig. 2a, b, c - white square) of  $\text{H}_2\text{O}_2$ . In the absence of  $\text{H}_2\text{O}_2$ , a  
250 strong increase of ATP concentration was observed reflecting an active metabolism of the bacteria. On the  
251 contrary, in the presence of  $\text{H}_2\text{O}_2$ , the results were clearly different and can be described in two phases. In the  
252 first phase, ATP concentration was decreasing while in a second phase it was progressively increasing  
253 (*Pseudomonas graminis*, 13b-3) or stabilizing (*Pseudomonas syringae*, 13b-2, *Sphingomonas sp.*, 14b-5). The  
254 kinetics of ATP concentration evolution and  $\text{H}_2\text{O}_2$  degradation are closely related. As discussed earlier (Fig. 1),  
255 the  $\text{H}_2\text{O}_2$  initially present (20  $\mu\text{M}$ ) was entirely degraded in approximately 3 h (depending on the strain); this  
256 corresponds exactly to the end of the ATP decrease. Complementary experiments were performed with  
257 incubations of the cells in the presence or absence of light and/or iron complex (Fe-[EDDS]) under conditions  
258 similar to that described previously in the presence of  $\text{H}_2\text{O}_2$  alone. The results obtained for the three strains are  
259 reported in Figure SM3 (*Pseudomonas graminis*), Figure SM4 (*Pseudomonas syringae*) and Figure SM5  
260 (*Sphingomonas sp.*).

261 The results show that light and iron complex have no impact on the ATP concentration decrease. The measured  
262 ATP concentration in the presence or absence of artificial light and/or iron(III) complex is similar to that  
263 observed in the presence of  $\text{H}_2\text{O}_2$  alone. The ATP concentration is thus only linked to the presence of  $\text{H}_2\text{O}_2$ .

#### 264 **Impact of $\text{H}_2\text{O}_2$ on the survival of the microbial strains**

265 We also controlled that the decrease of ATP in the presence of H<sub>2</sub>O<sub>2</sub> was not due to cell mortality. Results of the  
266 number of culturable bacteria in the presence or absence H<sub>2</sub>O<sub>2</sub> are shown in Figure 3. The evolution of the cell  
267 concentration was not significantly different when cells were incubated in the presence or absence of hydrogen  
268 peroxide. The decrease of ATP is therefore not linked to a lower concentration of cells but to a modification of  
269 metabolic pathways due to H<sub>2</sub>O<sub>2</sub> presence. The total number of cells increased by a factor 5 to 10 after 24h  
270 showing that bacteria were also able to divide and grow.

### 271 **3.2 Impact of H<sub>2</sub>O<sub>2</sub> on the microbial energetic metabolism in real cloud environment**

272 In the previous section, we showed that H<sub>2</sub>O<sub>2</sub> had a strong impact on the energetic metabolism of cells under our  
273 microcosm conditions. To go further, we looked at the potential impact of H<sub>2</sub>O<sub>2</sub> on microbial energetic states in  
274 real cloud samples by carrying out statistical analyses based on data measured on real cloud water collected at  
275 the PUY station.

276 For this, principal component analysis (PCA) was used. In order to perform this multivariate statistical analysis,  
277 Table SM1 was built.

278 The result of the PCA analysis is presented in Figure 4. The first two dimensions contain practically 50% of the  
279 total inertia (total variance of the data table) reflecting the validity and reliability of the result. The PCA shows  
280 that if we consider all important parameters in the collected cloud samples a strong correlation appears between  
281 ATP and H<sub>2</sub>O<sub>2</sub> concentrations (longer vectors and very close on the PCA). There is no correlation between ATP  
282 concentration and the number of bacteria (vectors practically orthogonal); this shows that H<sub>2</sub>O<sub>2</sub> is linked to the  
283 energetic state of the cells and not to their concentration. Also, there is no correlation between ATP and markers  
284 of pollution such as the pH values, the NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup> and NH<sub>4</sub><sup>+</sup> concentrations or even the temperature that could  
285 impact microbial metabolism.

286 In addition, Spearman rank correlation test (non-parametric test) was performed based on the 37 cloud samples  
287 to confirm the correlation between H<sub>2</sub>O<sub>2</sub> and ATP. The values used for this test are presented in Table SM1. A p-  
288 value of 0.0047 was obtained with a Spearman's coefficient of 0.45 (Zar, 1972). This shows an extremely strong  
289 correlation between H<sub>2</sub>O<sub>2</sub> and ATP, as theoretically the Spearman's coefficient must be greater than 0.27 for 37  
290 observations and the p-value less than 0.05 (significance threshold). To confirm that, ATP depletion due to H<sub>2</sub>O<sub>2</sub>  
291 impact was not linked with the mortality of cells, a Spearman rank correlation test was also performed to  
292 evaluate the correlation between ATP and total microorganisms concentrations (sum of bacteria and fungi  
293 concentrations in Table SM1) (p-value superior to 0.37).

294 Figure 4 suggested that ATP or H<sub>2</sub>O<sub>2</sub> could be also correlated to formate and oxalate since the vectors were  
295 relatively close. A Spearman rank correlation test (non-parametric test) was thus performed based on data  
296 extracted from the 37 cloud samples (Table SM1). A strong correlation was obtained between ATP and formate  
297 (p-value=0.0043, Spearman's coefficient = 0.46), and between H<sub>2</sub>O<sub>2</sub> and formate (p-value = 0.00015,  
298 Spearman's coefficient= 0.58). ATP-oxalate correlation is rather weak (p-value = 0.030, Spearman's  
299 coefficient= 0.36) and much lower than the ATP-H<sub>2</sub>O<sub>2</sub> correlation, similar values were obtained for oxalate and  
300 H<sub>2</sub>O<sub>2</sub> (p-value = 0.035, Spearman's coefficient = 0.35).

#### 301 4 Discussion

302 Our objective was to study in detail the interactions between cloud microorganisms and H<sub>2</sub>O<sub>2</sub>.

303 First we looked at the mechanisms involved in H<sub>2</sub>O<sub>2</sub> transformations under laboratory conditions by isolating  
304 each parameter to determine its impact on H<sub>2</sub>O<sub>2</sub> (artificial light, Fe-[EDDS] complex and bacteria). Degradation  
305 rates of hydrogen peroxide were precisely determined for different microbial strains frequently found in cloud  
306 water samples collected at the PUY site. The results show that all bacterial strains studied under these conditions  
307 degrade H<sub>2</sub>O<sub>2</sub> within the same order of magnitude as abiotic conditions. The degradation rates of H<sub>2</sub>O<sub>2</sub> by  
308 bacteria are not impacted by the presence of light and Fe-[EDDS] and consequently by the generation of HO<sup>•</sup>  
309 radicals. On the opposite, in these laboratory experiments mimicking real cloud conditions, we have shown that  
310 H<sub>2</sub>O<sub>2</sub> has a strong impact on the microbial energetic state of the cells. This strong decrease of ATP concentration  
311 is not linked to the number of cells as bacteria are able to divide and grow in the presence of H<sub>2</sub>O<sub>2</sub>. This reveals  
312 that microorganisms are able to manage the stress induced by H<sub>2</sub>O<sub>2</sub> through their metabolism. It is likely that  
313 they could respond using enzymes involved in H<sub>2</sub>O<sub>2</sub> degradation (*e.g.* catalases, peroxidases, *etc.*) and other  
314 typical antioxidant molecules (glutathione, *etc.*).

315 A few studies reported the decrease of ATP concentration in microorganisms (Perricone et al., 2003), plants  
316 (Tiwari et al., 2002) or mammalian cells (Spragg et al., 1985; Josephson et al., 1991; Sporn and Peters-  
317 Goldenwhen, 1988, Hyslop et al., 1988; Oka et al., 2012) exposed to H<sub>2</sub>O<sub>2</sub>. Fig. 5 illustrates how H<sub>2</sub>O<sub>2</sub> could  
318 affect the concentration of ATP in the cells. First H<sub>2</sub>O<sub>2</sub> could directly inhibit the ATP synthase, a membrane  
319 protein synthesizing ATP from ADP (Tamarit et al. 1998). Second H<sub>2</sub>O<sub>2</sub> could impact different metabolic  
320 pathways which are interconnected including glutathione metabolism, glycolysis, TCA cycle and DNA repair  
321 system. The functioning of the enzymes in these pathways and also the activity of the ATP synthase are  
322 dependent on the redox potential of the cells (NAD<sup>+</sup>/NADH, NADP<sup>+</sup>/NADPH ratios), and as a consequence the  
323 ATP concentration is regulated by this redox potential (Haddock and Jones, 1977, Singh et al., 2007, Oka et al.,  
324 2012). If for instance NAD<sup>+</sup> is depleted when the repair system is activated to avoid potential DNA damages  
325 induced by H<sub>2</sub>O<sub>2</sub>, then ATP is depleted, and finally all the metabolic pathways involving these compounds are  
326 impacted and a complete change in the metabolome can be expected.

327 The measurements performed in microcosms do not reproduce what is really occurring in cloud droplets. First  
328 incubations were performed with artificial cloud water and model strains, nevertheless the obtained results were  
329 consistent with those obtained with real cloud water samples. Second the potential growth of microorganisms  
330 during a cloud event could also modify transformation rates, this is only realistic for long cloud lifetimes (> 24  
331 hours). Finally experiments were performed under bulk conditions and not with individual cloud droplets, only  
332 models can take into account the complexity of cloud conditions, in particular the multiphase aspect of cloud  
333 chemistry. To go further and integrate biodegradation rates in atmospheric chemistry models, complementary  
334 experiments should be performed and biodegradation rates should be expressed as mol<sup>-1</sup>.cell<sup>-1</sup>.h<sup>-1</sup>.

335 However the most important result of this work was to show the correlation between H<sub>2</sub>O<sub>2</sub> concentrations and  
336 ATP concentrations. This result obtained under our microcosm conditions was confirmed using data measured in  
337 real cloud samples that experienced multiphase and real cloud conditions. Indeed, we have shown, thanks to  
338 statistical analyses, that there was also a high correlation between H<sub>2</sub>O<sub>2</sub> and ATP concentrations in real cloud  
339 samples collected under various environmental conditions. We suggest thus that hydrogen peroxide modulates  
340 the global metabolism of cloud microorganisms.

341 Another interesting correlation was obtained between H<sub>2</sub>O<sub>2</sub> and formate as well as ATP with formate. This could  
342 result from different concomitant processes. First formate is the most oxidized carbon molecule before CO<sub>2</sub>  
343 generated from successive oxidations of the organic matter by radicals issued from H<sub>2</sub>O<sub>2</sub>. Second it could reveal  
344 the impact of H<sub>2</sub>O<sub>2</sub> on the C1 metabolism; it is known that C1 compounds can be transformed by cloud  
345 microorganisms (Husàrovà et al., 2011, Vaitilingom et al., 2010, 2011, 2013). In addition Thomas et al. (2016)  
346 report the overproduction of formate in a strain of *Pseudomonas fluorescens* exposed to H<sub>2</sub>O<sub>2</sub>. Indeed formate  
347 contributes to the anti-oxidant strategy of this bacterium to supply NADH which is known to be decreased under  
348 oxidative conditions, formate helps thus to control the cellular redox potential (see Fig. 5)  
349 Finally, this work brings new insights into the interactions between H<sub>2</sub>O<sub>2</sub> and the cloud microbiome and its  
350 potential consequences on cloud chemistry (see Fig. 6).  
351 First it confirms that cloud microorganisms are able to efficiently degrade hydrogen peroxide and potentially  
352 impact the global carbon budget and the oxidant capacity of clouds as already shown in Vaitilingom et al.  
353 (2013). By decreasing H<sub>2</sub>O<sub>2</sub> concentration, radical chemistry is less efficient to degrade the organic matter.  
354 Second we show here for the first time that H<sub>2</sub>O<sub>2</sub> impacts the energetic metabolism of the cloud microbiome and  
355 thus potentially modulates its carbon metabolism. As a consequence it can modify the final transformation of the  
356 organic matter in clouds. This reciprocal interaction between H<sub>2</sub>O<sub>2</sub> and microorganisms and its subsequent  
357 impact on cloud chemistry is clearly dependent on H<sub>2</sub>O<sub>2</sub> concentration.  
358 To go further in the understanding of the modulation of the metabolic pathways (including carbon, nitrogen,  
359 amino-acids or sugars) induced by H<sub>2</sub>O<sub>2</sub>, a combined metabolomic and transcriptomic approach could be used.  
360 The next step could be to integrate biological data in numerical atmospheric models to better quantify  
361 consequence of this modulation on atmospheric chemistry.

362 *Acknowledgments.* N. Wirgot is a recipient of a PhD fellowship from the MESR (French government).  
363 Part of this work was supported by the French ANR program BIOCAP (ANR-13-BS06-0004). The authors are  
364 very grateful to the OPGC/LaMP staff for collecting the cloud samples at the PUY station, see the database of  
365 cloud biological and chemical composition [wwobs.univ-bpclermont.fr/SO/beam/index.php](http://wwobs.univ-bpclermont.fr/SO/beam/index.php).

## 366 **References**

- 367 Amato, P., Ménager, M., Sancelme, M., Laj, P., Mailhot, G., Delort, A.-M.: Microbial population in cloud water  
368 at the Puy de Dôme: implications for the chemistry of clouds, *Atmos. Environ.*, 39, 4143-4153,  
369 doi :10.1016/j.atmosenv.2005.04.002, 2005.
- 370 Amato, P., Parazols, M., Sancelme, M., Mailhot, G., Laj, P., Delort, A. M.: An important oceanic source of  
371 micro-organisms for cloud water at the Puy de Dôme (France), *Atmos. Environ.*, 41, 8253-8263,  
372 doi:10.1016/j.atmosenv.2007.06.022, 2007a.
- 373 Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G., Delort, A. M.: Microorganisms isolated from the  
374 water phase of tropospheric clouds at the Puy de Dôme: major groups and growth abilities at low  
375 temperatures, *FEMS Microbiol. Ecol.*, 59, 242-254, doi:10.1111/j.1574-6941.2006.00199, 2007b.
- 376 Amato, P., Demeer, F., Melaouhi, A., Fontanella, S., Martin-Biesse, A.-S., Sancelme, M., Delort, A.-M.: A fate  
377 for organic acids, formaldehyde and methanol in cloud water: their biotransformation by micro-organisms,

378 Atmos. Chem. Phys., 7, 5253-5276, doi:10.5194/acp-7-4159-2007, 2007c.

379 Arakaki, T.; Anastasio, C.; Kuroki, Y.; Nakajima, H.; Okada, K.; Kotani, Y.; Handa, D.; Azechi, S.; Kimura, T.;  
380 Tshako, A. A general scavenging rate constant for reaction of hydroxyl radical with organic carbon in  
381 atmospheric waters. Environ. Sci. Technol. 2013 , 47 (15), 8196-8203.

382 Ariya, P. A., Nepotchaykh, O., Ignatova, O., Amyot, M.: Microbiological degradation of atmospheric organic  
383 compounds, Geophys. Res. Lett., 29, 34.1-34.4, doi:10.1029/2002GL015637, 2002.

384 Bianco, A., Passananti, M., Perroux, H., Voyard, G., Mouchel-Vallon, C., Chaumerliac, N., Mailhot, G.,  
385 Deguillaume, L., Brigante, M.: A better understanding of hydroxyl radical photochemical sources in cloud  
386 waters collected at the puy de Dôme station - experimental versus modelled formation rates, Atmos. Chem.  
387 Phys., 15, 9191-9202, doi:10.5194/acp-15-9191-2015, 2015.

388 Deguillaume, L., Leriche, M., Desboeufs, K., Mailhot, G., George, C., Chaumerliac, N.: Transition metals in  
389 atmospheric liquid phases: sources, reactivity, and sensitive parameters, Chem. Rev., 105, 9, 3388-3431,  
390 doi:10.1021/cr040649c, 2005.

391 Deguillaume, L., Charbouillot, T., Joly, M., Vaïtilingom, M., Parazols, M., Marinoni, A., Amato, P., Delort, A-  
392 M., Vinatier, V., Flossmann, A., Chaumerliac, N., Pichon, J-M., Houdier, S., Laj, P., Sellegri, K., Colomb,  
393 A., Brigante, M., Mailhot, G.: Classification of clouds sampled at the puy de Dôme (France) based on 10 yr  
394 of monitoring of their physicochemical properties, Atmos. Chem. Phys., 14, 1485-1506, doi:10.5194/acp-14-  
395 1485-2014, 2014.

396 Delort, A.-M., Vaïtilingom, M., Joly, M., Amato, P., Wirgot, N., Lallement, A., Sancelme, M., Matulova, M.,  
397 Deguillaume, L.: Clouds: a transient and stressing habitat for microorganisms, in Microbial Ecology of  
398 Extreme Environments, Ed. Chénard, C. and Lauro, F.M., Springer, Chapter 10, doi: 10.1007/978-3-319-  
399 51686-8\_10, 2017.

400 Gunz, D. W., and Hoffmann, M. R.: Atmospheric chemistry of peroxides: a review, Atmos. Environ. Part A.  
401 General Topics, 24, 1601-1633, doi:10.1016/0960-1686(90)90496-A, 1990.

402 Haddock, B.A., and Jones, C.W.: Bacterial respiration, Bacteriol. Rev., 41, 47-99, 1977.

403 Hammer, Ø., Harper, D.A.T, Ryan, P.D.: PAST: palaeontological statistics software package for education and  
404 data analysis, Palaeontol. Electron., 4, 178, 2001.

405 Herrmann H, Schaefer T, Tilgner A, Styler SA, Weller C, Teich M, et al.: Tropospheric aqueous-phase  
406 chemistry: kinetics, mechanisms, and its coupling to a changing gas phase. Chem. Rev. 115, 4259–334, doi:  
407 10.1021/cr500447k, 2015.

408 Hems, R.F.; Hsieh, J.S.; Slodki, M.A.; Shouming, Z. ; Abbatt, J.P.D. Suppression of OH Generation from the  
409 Photo-Fenton Reaction in the Presence of  $\alpha$ -Pinene Secondary Organic Aerosol Material. Environ. Sci.  
410 Technol. Lett. DOI:10.1021/acs.estlett.7b00381

411 Hill, K. A., Shepson, P. B., Galbavy, E. S., Anastasio, C., Kourtev, P. S., Konopka, A., Stirm, B. H.: Processing  
412 of atmospheric nitrogen by clouds above a forest environment, J. Geophys. Res., 112, doi:  
413 10.1029/2006JD008002, 2007.

414 Husárová, S., Vařtilingom, M., Deguillaume, L., Traikia, M., Vinatier, V., Sancelme, M., Amato, P., Matulová,  
415 M., Delort, A.-M.: Biotransformation of methanol and formaldehyde by bacteria isolated from clouds.  
416 Comparison with radical chemistry, *Atmos. Environ.*, 45, 6093-6102, doi:10.1016/j.atmosenv.2011.06.035,  
417 2011.

418 Hyslop, P.A., Hinshawz, D.B., Halsey, W.A., Schraufstatter, I.U., Sauerhebery, R.D., Roger G. Spraggj, R.G.,  
419 Jackson, J.H., Cochrane, C.G.: Mechanisms of oxidant-mediated cell injury the glycolytic and mitochondrial  
420 pathways of ADP phosphorylation are major intracellular targets inactivated by hydrogen peroxide, *J. Biol.*  
421 *Chem.*, 263, 1665-1675, 1988.

422 Joly, M., Amato, P., Sancelme, M., Vinatier, V., Abrantes, M., Deguillaume, L., Delort, A.-M.: Survival of  
423 microbial isolates from clouds toward simulated atmospheric stress factors, *Atmos. Environ.*, doi:  
424 10.1016/j.atmosenv.2015.07.009, 2015.

425 Josephson, R.A., Silverman, H.S., Lakatta, E.G., Stern, M.D., Zweier, J.L.: Study of the mechanisms of  
426 hydrogen peroxide and hydroxyl free radical-induced cellular injury and calcium overload in cardiac  
427 myocytes, *J. Biol. Chem.*, 266, 2354-2361, 1991.

428 Kieber, R.J., Peake, B., Willey, J.D., Jacobs, B.: Iron speciation and hydrogen peroxide concentrations in New  
429 Zealand rainwater, *Atmos. Environ.*, 35, 6041-6048, doi:10.1016/S1352-2310(01)00199-6, 2001.

430 Kok, G.L.: Measurements of hydrogen peroxide in rainwater, *Atmos. Environ.*, 14, 653-656, doi:10.1016/0004-  
431 6981(80)90048-7, 1980.

432 Koutny, M., Sancelme, M., Dabin, C., Pichon, N., Delort, A.-M., Lemaire, J.: Acquired biodegradability of  
433 polyethylenes containing pro-oxidant additives, *Polym. Degrad. Stabil.*, 91, 1495-1503,  
434 doi:10.1016/j.polymdegradstab.2005.10.007, 2006.

435 Kroll, J. H., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: formation and evolution of low-  
436 volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593-3624, doi:10.1016/j.atmosenv.2008.01.003,  
437 2008.

438 Krumins, V., Mainelis G., Kerkhof, L.J., and Fennell, D.E.: Substrate-dependent rRNA production in an airborne  
439 bacterium. *Environ. Sci. Technol. Lett.*, 9, 376-381, 2014.

440 Lazrus, A. L., Kok, G. L., Gitlin, S. N., Lind, J. A., McLaren, S. E.: Automated fluorimetric method for  
441 hydrogen peroxide in atmospheric precipitation, *Anal. Chem.*, 57, 917-922, doi:10.1021/ac00281a031, 1985.

442 Lee, M., Heikes, B.G., O'Sullivan, D.W.: Hydrogen peroxide and organic hydroperoxide in the troposphere: a  
443 review, *Atmos. Environ.*, 34, 3475-3494, doi:10.1016/S1352-2310(99)00432-X, 2000.

444 Li, J., Mailhot, G., Wu, F., Deng, N.: Photochemical efficiency of Fe(III)-EDDS complex: OH radical  
445 production and 17 $\beta$ -estradiol degradation, *J. Photoch. Photobio. A.*, 212, 1-7,  
446 doi:10.1016/j.jphotochem.2010.03.001, 2010.

447 Li, J., Wang, X., Chen, J., Zhu, C., Li, W., Li, C., Liu, L., Xu, C., Wen, L., Xue, L., Wang, W., Ding, A. and  
448 Herrmann, H.: Chemical composition and droplet size distribution of cloud at the summit of Mount Tai,  
449 China, *Atmospheric Chem. Phys. Discuss.*, 1-21, doi:10.5194/acp-2016-1175, 2017.

450 Marie, D., Brussaard, C.P.D., Partensky, F., Vaultot, D. : Flow cytometric analysis of phytoplankton, bacteria and  
451 viruses, Robinson, J.P., Ed. Curr. Protoc. Cytom., John Wiley & Sons, 11.11, 1-15, 1999.

452 Marinoni, A., Parazols, M., Brigante, M., Deguillaume, L., Amato, P., Delort, A.-M., Laj, P., Mailhot, G.:  
453 Hydrogen peroxide in natural cloud water: sources and photoreactivity, Atmos. Res., 101, 1-2, 256-263,  
454 doi :10.1016/j.atmosres.2011.02.013, 2011.

455 Matulová, M., Husárová, S., Capek, P., Sancelme, M., Delort, A. M.: Biotransformation of various saccharides  
456 and production of exopolymeric substances by cloud-borne *Bacillus sp.* 3B6, Environ. Sci. Technol., 48,  
457 14238-14247, doi:10.1021/es501350s, 2014.

458 Möller, D.: Atmospheric hydrogen peroxide: evidence for aqueous-phase formation from a historic perspective  
459 and one-year measurement campaign, Atmos. Chem. Phys., 43, 5923-5936,  
460 doi:10.1016/j.atmosenv.2009.08.013, 2009.

461 Oka, S.I., Hsu, C.P., Sadoshima, J.: Regulation of cell survival and death by pyridine nucleotides., Circ. Res.,  
462 111, 611-627, doi:10.1161/CIRCRESAHA.111.247932, 2012.

463 Perricone, C.D., Park, S., Imlay, J.A., Weiser, J.N.: Factors contributing to hydrogen peroxide resistance in  
464 *Streptococcus pneumoniae* include pyruvate oxidase (SpxB) and avoidance of the toxic effects of the Fenton  
465 reaction, J. Bacteriol., 185, 6815-6825, doi:10.1128/JB.185.23.6815-6825.2003, 2003.

466 Pruppacher, H. R., and Jaenicke, R.: The processing of water vapor and aerosols by atmospheric clouds, a global  
467 estimate, Atmos. Res, 38, 283-295, doi:10.1016/0169-8095(94)00098-X, 1995.

468 Ravishankara, A.R.: Heterogeneous and multiphase chemistry in the troposphere, Science, 276, 1058-1065,  
469 doi:10.1126/science.276.5315.1058, 1997.

470 Reasoner, D.J., Geldreich, E.E.: A new medium for the enumeration and subculture of bacteria from potable  
471 water, Appl. Environ. Microb., 49, 1-7, 1985.

472 Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, Atmos. Chem. Phys., 15,  
473 4399-4981, doi:10.5194/acp-15-4399-2015, 2015.

474 Sattler, B., Puxbaum, H., Psenner, R.: Bacterial growth in supercooled cloud droplets, Geophys. Res. Lett., 28,  
475 239-242, doi :10.1029/2000GL011684, 2001.

476 Schöne, L. and Herrmann, H.: Kinetic measurements of the reactivity of hydrogen peroxide and ozone towards  
477 small atmospherically relevant aldehydes, ketones and organic acids in aqueous solutions, Atmos. Chem.  
478 Phys., 14, 4503-4514, doi:10.5194/acp-14-4503-2014, 2014.

479 Shen, X., Lee, T., Guo, J., Wang, X., Li, P., Xu, P., Wang, Y., Ren, Y., Wang, W., Wang, T., Li, Y., Carn, S. A., and  
480 Collett, J. L.: Aqueous phase sulfate production in clouds in eastern China, Atmospheric Environment, 62,  
481 502-511, <https://doi.org/10.1016/j.atmosenv.2012.07.079>, 2012.

482 Singh, R., Mailloux, R.J., Puiseux-Dao, S., Appanna, V.D.: Oxidative stress evokes a metabolic adaptation that  
483 favors increased NADPH synthesis and decreased NADH production in *Pseudomonas fluorescens*, J.  
484 Bacteriol., 189, 6665-6675, doi:10.1128/JB.00555-07, 2007.

485 Sporn, P.H. and Peters-Golden, M.: Hydrogen peroxide inhibits alveolar macrophages 5-lipoxygenase  
486 metabolism in association with depletion of ATP, *J. Biol. Chem.*, 263, 14776-14783, 1988.

487 Spragg, R.G., Hinshaw, D.B., Hyslop, P.A., Schraufstatter, I.U., Cochrane, C.G.: Alterations in adenosine  
488 triphosphate and energy charge in cultured endothelial and P388D1 cells after oxidant injury, *J. Clin. Invest.*,  
489 76, 1471-1476, doi:10.1172/JCI112126, 1985.

490 Tamarit, J., Cabiscol, E., Ros, J.: Identification of the major oxidatively damaged proteins in *Escherichia coli*  
491 cells exposed to oxidative stress, *J. Biol. Chem.*, 273, 3027-3032, doi:10.1074/jbc.273.5.3027, 1998.

492 Thomas, S.C., Alhasawi, A., Auger, C., Omri, A., Appanna, V.D.: The role of formate in combatting oxidative  
493 stress, *J. Microbiol.*, 109, 263-271, doi:10.1007/s10482-015-0629-6, 10.1007/s10482-015-0629-6, 2016.

494 Thompson, A. M.: The oxidizing capacity of the Earth's atmosphere: probable past and future changes, *Science*,  
495 256(5060), 1157-1165, doi:10.1126/science.256.5060.1157, 1992.

496 Tiwari, B.S., Belenghi, B., Levine, A.: Oxidative stress increased respiration and generation of reactive oxygen  
497 species, resulting in ATP depletion, opening of mitochondrial permeability transition, and programmed cell  
498 death, *Plant Physiol.*, 128, 1271-1281, doi:10.1104/pp.010999, 2002.

499 Vaïtilingom, M., Amato, P., Sancelme, M., Laj, P., Leriche, M., Delort, A. M.: Contribution of microbial activity  
500 to carbon chemistry in clouds. *Appl. Environ. Microb.*, 76, 23-29, doi:10.1128/AEM.01127-09, 2010.

501 Vaïtilingom, M., Charbouillot, T., Deguillaume, L., Maisonobe, R., Parazols, M., Amato, P.; Sancelme, M.,  
502 Delort, A.-M.: Atmospheric chemistry of carboxylic acids: microbial implication versus photochemistry,  
503 *Atmos. Chem. Phys.*, 11, 8721-8733, doi:10.5194/acp-11-8721-2011, 2011.

504 Vaïtilingom, M., Attard, E., Gaiani, N., Sancelme, M., Deguillaume, L., Flossmann, A. I., Amato, P., Delort, A.  
505 M.: Long-term features of cloud microbiology at the puy de Dôme (France), *Atmos. Environ.*, 56, 88-100,  
506 doi :10.1016/j.atmosenv.2012.03.072, 2012.

507 Vaïtilingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N., Delort, A.-M.:  
508 Potential impact of microbial activity on the oxidant capacity and organic carbon budget in clouds, *P. Natl.*  
509 *Acad. Sci USA*, 110, 559-564, doi:10.1073/pnas.1205743110, 2013.

510 Vione, D, Maurino, V.; Minero, C., Pelizzetti, E.: The atmospheric chemistry of hydrogen peroxide: a review,  
511 *Ann. Chim-Rome*, 93, 477-488, 2003.

512 Wei, M., Xu, C., Chen, J., Zhu, C., Li, J., and Lv, G.: Characteristics of bacterial community in cloud water at  
513 Mt Tai: similarity and disparity under polluted and non-polluted cloud episodes, *Atmos. Chem. Phys.*, 17,  
514 5253-5270, <https://doi.org/10.5194/acp-17-5253-2017>, 2017.

515 Zar, J.H.: Significance testing of the Spearman rank correlation coefficient, *J. Am. Stat. Assoc.*, 67, 578-580,  
516 doi:10.2307/2284441, 1972.

517

518

519 **Table 1: Initial rates of abiotic degradation (a) and of biotic degradation (b) of H<sub>2</sub>O<sub>2</sub> measured in artificial cloud**  
 520 **water. Values are expressed in 10<sup>-9</sup> mol L<sup>-1</sup> s<sup>-1</sup>. Standard errors were calculated.**


(a)	Light + Fe-EDDS]	Fe-[EDDS]	Light
	1.07 ± 0.18	0.22 ± 0.05	0.14 ± 0.08

521

(b)	Light + Fe-[EDDS] + Bacteria	Fe-[EDDS] + Bacteria	Light + Bacteria	Bacteria
<i>Pseudomonas graminis</i> 13b-3	1.55 ± 0.25	1.93 ± 0.18	2.15 ± 0.02	2.07 ± 0.01
<i>Pseudomonas syringae</i> 13b-2	1.75 ± 0.15	1.27 ± 0.04	1.72 ± 0.14	1.18 ± 0.08
<i>Sphingomonas sp.</i> 14b-5	1.97 ± 0.06	1.01 ± 0.21	0.87 ± 0.04	0.76 ± 0.11

522


523




524

525 **Figure 1: Evolution of H<sub>2</sub>O<sub>2</sub> concentration as a function of time (min) under abiotic conditions: Light + Fe-[EDDS]**  
 526 **(orange diamond), Light (orange circle), Fe-[EDDS] (orange square with black cross) and biotic conditions: Light +**  
 527 **Fe-[EDDS] + *Pseudomonas graminis* (13b-3) (green diamond), Fe-[EDDS] + *Pseudomonas graminis* (13b-3) (green**  
 528 **square with black cross), Light + *Pseudomonas graminis* (13b-3) (green circle), *Pseudomonas graminis* (13b-3) (green**  
 529 **triangle). Values shown are averages of triplicates plus/minus one standard deviation. Where error bars are not**  
 530 **visible they are smaller than the symbol.**


531




532




533




534 Figure 2: ATP concentration ( $\mu\text{M}$ ) as a function of time (min) in the presence (red square) or the absence (navy circle)  
535 of  $\text{H}_2\text{O}_2$  for the three strains: (a) *Pseudomonas graminis* (13b-3), (b) *Pseudomonas syringae* (13b-2), (c) *Sphingomonas*  
536 *sp.* (14b-5).  
537 The vertical bar illustrates the time corresponding to the total degradation of  $\text{H}_2\text{O}_2$ .



538




539



540

541 **Figure 3: Bacterial cell numbers measured by plate-counting in the presence (red) and the absence (navy) of H<sub>2</sub>O<sub>2</sub> at**  
 542 **20 μM for the three strains: (a) *Pseudomonas graminis* (13b-3), (b) *Pseudomonas syringae* (13b-2) and (c)**  
 543 ***Sphingomonas sp.* (14b5). Error bars represent standard deviation from the means (n=3).**




544

545 **Figure 4: Variables factor map (PCA) of the 37 cloud events on the plane PC1-PC2 based on 17 variables.**

546

547



548


549 **Figure 5: Hypothetical mechanism that could explain the impact of H<sub>2</sub>O<sub>2</sub> on cell metabolism and ATP concentration.**

550 **Interconnection between ATP synthesis and cellular redox potential (NAD<sup>+</sup>/NADH, NADP<sup>+</sup>/NADPH ratios). NAD<sup>+</sup>**

551 **depletion related to DNA repair system. Adapted from Oka et al. (2012).**

552 -----> **Inhibition of ATP synthase**

553 -.-.-.-.-> **NAD<sup>+</sup> depletion related to DNA repair system**



554

555 **Figure 6: Interaction between  $H_2O_2$  and cloud microorganisms and its potential consequences on atmospheric**  
 556 **chemistry. (1) Cloud microorganisms degrade  $H_2O_2$  thanks to their catalases and peroxidases (oxidative stress**  
 557 **metabolism) as a result it impacts the oxidant capacity of clouds. The concentration of radicals issued from  $H_2O_2$  is**  
 558 **decreased and radical chemistry is less efficient to transform the organic matter. (2)  $H_2O_2$  impacts the energetic**  
 559 **metabolism of microorganisms that react to this stress. The depletion of ATP modulates the global carbon metabolism**  
 560 **of the microorganisms, and consequently the transformation of the organic matter. These processes are modulated by**  
 561 **the  $H_2O_2$  concentration that varies depending on atmospheric scenari.**

562

