

Ice Nucleating Particles in a Coastal Tropical Site

Luis A. Ladino^{1,*}, Graciela B. Raga¹, Harry Alvarez-Ospína², Manuel A. Andino-Enríquez³, Irma Rosas¹, Leticia Martínez¹, Eva Salinas¹, Javier Miranda⁴, Zyanya Ramírez-Díaz¹, Bernardo Figueroa⁵, Cedric Chou⁶, Allan K. Bertram⁶, Erika T. Quintana⁷, Luis A. Maldonado⁸, Agustín García-Reynoso¹, Meng Si⁶, and Victoria E. Irish⁶

¹Centro de Ciencias de la Atmosfera, Universidad Nacional Autónoma de México, México City, México

²Facultad de Ciencias, Universidad Nacional Autónoma de México, México City, México

³School of Chemical Sciences and Engineering, Universidad Yachay Tech, Ecuador

⁴Instituto de Física, Universidad Nacional Autónoma de México, México City, México

⁵Laboratorio de Ingeniería y Procesos Costeros, Instituto de Ingeniería, Universidad Nacional Autónoma de México, Sisal, Yucatán, México

⁶Chemistry Department, University of British Columbia, Vancouver, Canada.

⁷Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, México City, México

⁸Facultad de Química, Universidad Nacional Autónoma de México, México City, México

Correspondence to: Luis A. Ladino (luis.ladino@atmosfera.unam.mx)

Abstract. Atmospheric aerosol particles that can nucleate ice are referred to as ice nucleating particles (INPs). Recent studies have confirmed that aerosol particles emitted by mid- and high-latitude oceans can act as INPs. This very relevant information can be included in climate and weather models to predict the formation of ice in clouds, given that most of them do not consider the oceans as a source of INPs. Very few studies to sample INPs have been carried out in tropical latitudes, and there is a need to evaluate their availability to understand the potential role that marine aerosol may play in the hydrological cycle of tropical regions.

This study presents results from the first measurements obtained during a field campaign conducted in the tropical village of Sisal, located on the coast of the Gulf of Mexico of the Yucatán peninsula in México in January-February 2017, and one of the few datasets currently available at such latitudes (i.e., 21°N). Aerosol particles sampled in Sisal are shown to be very efficient INPs in the immersion freezing mode, with onset freezing temperatures in some cases as high as -3 °C, similar to the onset temperature from *Pseudomonas syringae*. The results show that the INP concentration in Sisal was higher than at other locations sampled with the same type of INP counter. Air masses arriving in Sisal after the passage of cold fronts have, surprisingly, higher INP concentrations than the campaign-average, despite their lower total aerosol concentration.

The high concentrations of INPs at warmer ice nucleation temperatures ($T > -15$ °C) and the supermicron size of the INPs suggest that biological particles may have been a significant contributor

20 to the INP population in Sisal during this study. However, our observations also suggest that at temperatures ranging between $-20\text{ }^{\circ}\text{C}$ and $-30\text{ }^{\circ}\text{C}$ mineral dust particles are the likely source of the measured INPs.

1 Introduction

Clouds are essential to the hydrological cycle of the planet and also play a significant role in the radiative balance of the climate system (Ramanathan et al., 1989; Lohmann and Feichter, 2005; 25 Andreae and Rosenfeld, 2008; Stevens and Feingold, 2009). Cloud formation depends on the presence of cloud condensation nuclei (CCN) and most precipitation from mixed-phase clouds involves also the presence of ice nucleating particles (INPs). Aerosol-cloud interactions are one of the main sources of uncertainty in climate projections as assessed by the Intergovernmental Panel on Climate Change (Stocker et al., 2013), prompting large research efforts from the scientific community in 30 recent years. Nevertheless, the formation and evolution of ice crystals in mixed-phase and cirrus clouds still remain highly uncertain (Seinfeld et al., 2016; Kanji et al., 2017; Field et al., 2017). Several pathways have been proposed as potentially responsible for ice formation: condensation freezing, contact freezing, immersion freezing, and deposition nucleation (Vali et al., 2015). Murray et al. (2012) and Ladino et al. (2013) have suggested that contact freezing and immersion freezing 35 are the most efficient mechanisms leading to ice nucleation in clouds; however, the atmospheric relevance of contact freezing is still under debate (Hobbs and Atkinson, 1976; Ansmann et al., 2005; Cui et al., 2006; Phillips et al., 2007; Seifert et al., 2011; Kanji et al., 2017).

Most of the precipitation from deep convection in the tropics, e.g. in the Inter-tropical Convergence Zone, forms via the ice phase (Mülmenstädt et al., 2015). Given the ice-nucleating potential 40 of a variety of aerosol particles such as mineral dust, biological particles, crystalline salts, carbonaceous particles, and secondary organic aerosol, the main source of INPs at tropical latitudes is highly uncertain (Kanji et al., 2017; Yakobi-Hancock et al., 2014; DeMott et al., 2010). Although it is yet not fully understood what exactly makes an aerosol particle an efficient INP (e.g. its composition, 45 active sites, crystal structure, size, or hygroscopicity), there is evidence that their composition is one of the key factors (Kanji et al., 2017). On a global scale, the large tropospheric concentrations and the good ice nucleating abilities of mineral dust have been widely reported as an important INP source (Hoose and Möhler, 2012; Nenes et al., 2014; Atkinson et al., 2013; Kanji et al., 2017). Bioaerosol has also been identified as very efficient INP (Kanji et al., 2017; Hoose and Möhler, 50 2012; Fröhlich-Nowoisky et al., 2016; Hill et al., 2017), with onset freezing temperatures reported as high as $-2\text{ }^{\circ}\text{C}$ (Yankofsky et al., 1981; Després et al., 2012; Fröhlich-Nowoisky et al., 2015; Wex et al., 2015; Stopelli et al., 2017). Global climate models parameterize cloud droplet and ice crystal formation from observational studies and results from such modelling suggest that on a global scale bioaerosol is not a major source of INPs and therefore, have a lower impact in ice cloud formation

55 in comparison to mineral dust particles (Hoose et al., 2010; Sesartic et al., 2012). However, this
may not be the case on a regional scale (Burrows et al., 2013; Mason et al., 2015a). Marine organic
matter, likely of biological origin, has been suggested to be an important oceanic sources of INPs in
the southern oceans, north Atlantic, and north Pacific (Burrows et al., 2013; Yun and Penner, 2013;
Wilson et al., 2015; Vergara-Temprado et al., 2017). However, the maritime source suggestion was
60 made with little or no data from tropical latitudes.

Important efforts were made during the 1950-70s to understand the role of the oceans in ice cloud
formation (Bigg, 1973; Schnell and Vali, 1975; Schnell, 1975, 1977, 1982; Rosinski et al., 1987,
1988). There is recent new and robust evidence that biological material from the marine environment
could act as efficient INPs (Knopf et al., 2011; Wilson et al., 2015; Mason et al., 2015b; DeMott
65 et al., 2016; Ladino et al., 2016; McCluskey et al., 2017; Irish et al., 2017; Welti et al., 2018). Most
of the past available INP data were obtained from mid- and high-latitudes studies, with tropical
latitudes heavily under represented (Schnell, 1982; Rosinski et al., 1987, 1988; Boose et al., 2016;
Welti et al., 2018; Price et al., 2018). Marine and coastal INP concentration ([INP]) typically ranges
from 10^{-4} L^{-1} to 10^{-1} L^{-1} for temperatures between $-10 \text{ }^{\circ}\text{C}$ and $-25 \text{ }^{\circ}\text{C}$ (Kanji et al., 2017), but
70 have shown to be higher at tropical coastal sites (Rosinski et al., 1988; Boose et al., 2016; Welti
et al., 2018; Price et al., 2018). This large [INP] range may strongly depend on the microbiota
concentration, the marine biological activity, and the organic matter enrichment in the sea surface
microlayer as shown in Wilson et al. (2015).

At marine and coastal sites, a large variety of bacteria have been identified with *Proteobacteria*,
75 *Firmicutes*, and *Bacteroidetes* as the main reported phyla (Després et al., 2012). Also, airborne
fungi are common in both continental and marine environments with *Cladosporium*, *Alternaria*,
Penicillium, *Aspergillus*, and *Epicoccum* the main identified genus (Després et al., 2012). Besides
bacteria and fungal spores, viruses, algae, and pollen have also been identified in the bioaerosol of
marine environments (Després et al., 2012; Fröhlich-Nowoisky et al., 2015; Michaud et al., 2018).
80 Therefore, the concentration, ice nucleating abilities, and variability of tropical bioaerosol need to
be better characterized to quantify their role in cloud formation and precipitation development at
regional level and within the tropical zonal band.

The Yucatan peninsula, surrounded by the Gulf of Mexico to the West and by the Caribbean sea
to the East, with a large variety of tropical vegetation, is a great source of both terrestrial and marine
85 microorganisms (Guzmán, 1982; Videla et al., 2000; Morales et al., 2006). Tropical cyclones (TC)
and cold fronts are some of the meteorological phenomena that seasonally affect the Yucatan penin-
sula every year (Whigham et al., 1991; Landsea, 2007; Knutson et al., 2010). DeLeon-Rodriguez
et al. (2013) show that TC can significantly enhance the concentration of biological particles through-
out the troposphere and can also efficiently transport biological particles far away from their sources.
90 Moreover, Mayol et al. (2017) has shown that ocean and terrestrial microorganisms can be efficiently
transported long distances from their sources over the tropical and subtropical oceans.

This study presents results of the INP concentration as a function of temperature and particle size, and the concentration and composition of biological particles at a tropical coastal site (Sisal, Yucatan), to infer the potential relevance of biological particles in mixed-phase cloud formation and precipitation development.

2 Methods

2.1 Sampling site

Ambient aerosol particles were collected between January 21 and February 02, 2017 in the coastal village of Sisal, located in the northwest corner of the Yucatan peninsula ($21^{\circ}09'55''N$ $90^{\circ}01'50''W$), as shown in Figure 1. Sisal had 1837 inhabitants in 2015 (SEDESOL, 2015), with fishing and tourism recognized as the main economical activities. The closest industry is located approximately 25 km away from the village and the nearest city is Merida, 75 km away.

The instruments used in this study were located on the roof of the Engineering Institute building of the Universidad Nacional Autonoma de Mexico (UNAM, Sisal Campus) which is 50 m from the shoreline and about 1.7 km from the center of Sisal (Figure 1). The roof is 25 m above ground level and directly faces the ocean.

January and February are part of the cold dry season in Mexico, with isolated events of rain associated with cold fronts reaching the deep tropics. The average temperature and relative humidity (RH) during the sampling period were 22.3 ± 3.6 °C and 68.9 ± 6.2 %, respectively.

2.2 Instrumentation

A suite of instrumentation was deployed in Sisal to characterize the aerosol chemical composition, concentration, size distribution, biological content, INPs concentration, and meteorological variables (Table 1). Most instruments were run simultaneously and next to each other (less than 10 m apart) and only wet aerosol particles were sampled (mean RH=69 %). Additionally, none of the instrumentation used an impactor or cyclone ahead of their inlets. The inlets were located around 1.5 m - 2.0 m above the roof surface. The meteorological data was obtained with a meteorological station (Davis, VANTAGE PRO2) placed in a different building approximately 20 m away from the other instruments.

2.2.1 Aerosol concentration and size distribution

The aerosol particle concentration and size distribution was monitored with a condensation particle counter (CPC 3010, TSI) and with an optical particle counter (LasAir II 310A, PMS), respectively. In the CPC, the size of the aerosol particles is increased in a heated saturator/cooled condenser system prior to their detection. The particles grown are directed towards a laser beam and the dispersed light is collected by a photodetector that convert it to particle concentration. Similar to the CPC,

125 aerosol particles in the LasAir are counted by passing them through a laser beam (without any prior
treatment). Based on the pulses (or voltage) and their amplitude the dispersed light by the particles
is then converted to particle concentration and size. The total particle concentration reported by the
CPC was collected every second at a flow rate of 1 L min^{-1} , whereas the aerosol concentration as
function of size (cut-sizes at $0.3 \mu\text{m}$, $0.5 \mu\text{m}$, $1.0 \mu\text{m}$, $5.0 \mu\text{m}$, $10.0 \mu\text{m}$, and $25 \mu\text{m}$) was recorded
130 every 11 s with the LasAir at a flow rate 28.3 L min^{-1} .

2.2.2 Ice nucleating particles

Aerosol particles were collected on hydrophobic glass cover slips (HR3-215; Hampton Research)
with the help of a Micro-Orifice Uniform Deposit Impactor (MOUDI 110R, MSP) to determine INP
concentrations in ambient air. Identical substrate holders as those described in Mason et al. (2015a)
135 were used to keep the glass cover slips at a location on the impaction plate where particle concen-
trations varied by a relatively small amount. The MOUDI has eight stages for particle separation
and collection as a function of their aerodynamic diameter (cut-sizes are $10.0 \mu\text{m}$, $5.6 \mu\text{m}$, $3.2 \mu\text{m}$,
 $1.8 \mu\text{m}$, $1.0 \mu\text{m}$, $0.56 \mu\text{m}$, $0.32 \mu\text{m}$, and $0.18 \mu\text{m}$). The flow through the MOUDI is 30 L min^{-1}
and the typical sampling time was 6 h. It has been recognized that when sampling with a MOUDI
140 under dry conditions (i.e., RH below approximately 60 %), aerosol particles can bounce from the
impaction plates moving to lower stages (Winkler, 1974; Chen et al., 2011; Bateman et al., 2014).
Although this is a known artifact when using this technique, this may not have been an issue in the
current study given that the ambient RH was typically above 67 %. The glass substrates containing
the ambient aerosol particles were stored in petri dishes at $4 \text{ }^\circ\text{C}$ prior to their analysis.

145 The INP concentrations were measured with a cold cell coupled to an optical microscope with an
EC Plan-Neofluar 5 X objective (Axiolab, Zeiss) following the MOUDI-DFT method described by
Mason et al. (2015a). The cold cell-microscope system used here is the same one used in previous
studies (Mason et al., 2015a,b, 2016; DeMott et al., 2016; Si et al., 2018). The following steps
encompass the analysis: i) The samples collected on glass cover slips were placed in the cold cell
150 at room temperature, ii) The cold cell was isolated and kept at $0 \text{ }^\circ\text{C}$, while humid air (RH>100 %) was
injected into the cell to induce liquid droplets formation by water vapor condensation; iii) Dry
air (N_2) was then injected into the cold cell to prevent the newly formed droplets from touching.
This is a key step to minimize the probability of liquid droplets freezing by contact; and iv) Once
droplets sizes and thermodynamic conditions were stable, the cold cell was closed. The activation
155 scans were conducted between $0 \text{ }^\circ\text{C}$ and $-40 \text{ }^\circ\text{C}$ at a cooling rate of $-10 \text{ }^\circ\text{C per minute}$ for particles
collected on stages two to seven. Stage one (cut-size of $10.0 \mu\text{m}$) was not taken into account given
that the aerosol concentration on the glass substrates was typically very low, whereas in stage eight
(cut-size of $0.18 \mu\text{m}$) the number concentration of particles deposited on the glass substrates was
so high that inhibited the proper formation of water drops. The temperature at which each droplet
160 froze was determined by analyzing the video from the CCD camera (XC-ST50, Sony) connected

to the microscope and the data reported by the resistance temperature detector (RTD) located at the center of the cold cell with a ± 0.2 °C uncertainty (Mason et al., 2015b). Homogeneous freezing experiments were performed on laboratory blanks exposed during the preparation of the MOUDI, while heterogeneous freezing experiments were run on ambient particles deposited on the glass cover slips (Figure S1). The [INP] was calculated using the following expression:

$$[INPs(T)] = -\ln\left(\frac{N_u(T)}{N_o}\right) \cdot \left(\frac{A_{deposit}}{A_{DFTV}}\right) \cdot N_o \cdot f_{ne} \cdot f_{nu,0.25-0.10\text{ mm}} \cdot f_{nu,1\text{ mm}}, \quad (1)$$

where $N_u(T)$ is the number of unfrozen droplets at temperature T , N_o the total number of droplets, $A_{deposit}$ the total area of the aerosol deposit on the hydrophobic glass cover slip, A_{DFT} the area of the hydrophobic glass cover slip analyzed in the DFT experiments, V the total volume of air sampled, f_{ne} a correction factor to account for uncertainty associated with the number of nucleation events in each experiment, $f_{nu,0.25-0.10\text{ mm}}$ and $f_{nu,1\text{ mm}}$ a non-uniformity factor which corrects for aerosol deposit inhomogeneity at a scale of 0.25 mm - 0.10 mm, and 1 mm, respectively (Mason et al., 2015a). We refer the readers to Mason et al. (2015a) and Mason et al. (2015b) for more details of the MOUDI-DFT operational principle.

175 2.2.3 Chemical composition

A second eight-stage MOUDI (100NR, MSP) was operated simultaneously to collect aerosol particles for chemical composition analysis with particle sizes ranging from 0.18 μm to 10.0 μm . Particles were collected on 47 mm Teflon filters (Pall Science) for 48 h at a flow rate of 30 L min^{-1} . Filters were weighed prior and after the sampling and stored in petri dishes at 4 °C until they were analyzed. Two different analyses were performed on each filter: elemental composition, followed by ion-cation concentration analysis.

Elemental composition of the aerosol samples was determined by X-ray fluorescence (XRF), using the x-ray spectrometer at Laboratorio de Aerosoles, Instituto de Fisica, UNAM (Espinosa et al., 2012). The samples were mounted on plastic frames with no previous treatment. The analysis was carried out using an Oxford Instrument (Scotts Valley, CA, USA) x-ray tube with an Rh anode and an Amptek (Bedford, MA, USA) Silicon Drift Detector (resolution 140 eV at 5.9 keV). The tube operated at 50 kV and a current of 500 μA , irradiating during 900 s per spectrum. The efficiency of the detection system was measured using a set of thin film standards (MicroMatter Co., Vancouver, Canada). The spectra obtained for the samples were deconvolved with the WinQXAS computer code (IAEA, 1997), and the experimental uncertainties in elemental concentrations were computed according to the method described by Espinosa et al. (2010).

After the XRF analysis, the Teflon filters were analyzed for NO_3^- , SO_4^{2-} , Cl^- , K^+ , Na^+ , Ca^{2+} , Mg^{2+} , and NH_4^+ using a Dionex model ICS-1500, equipped with an electrical conductivity detector, following Chow and Watson (1999). NO_3^- , Cl^- , and SO_4^{2-} were separated using a Thermo Scientific Dionex IonPac AS23-4 μm Analytical Column (4 mm x 250 mm) with Thermo Scientific

Dionex CES 300 Capillary Electrolytic Suppressor module. The injection volume was 1000 μL , mobile phase was 4.5 mM Na_2CO_3 - 0.8 mM NaHCO_3 at 1 mL min^{-1} flow rate. For NH_4^+ , Na^+ , Ca^{2+} , Mg^{2+} , and K^+ , volumes of 1000 μL were injected in a Thermo Scientific Dionex IonPac CS12A Cation-Exchange Column (4 mm x 250 mm) with the Thermo Scientific Dionex CES 300 Capillary Electrolytic Suppressor. The mobile phase was a solution CH_4SO_3 20 mM and 1 mL min^{-1} flow rate.

2.2.4 Biological particles

Air samples were collected using two Quick Take 30 Sample Pump BioStage viable cascade impactor (SKC Inc. USA), which is a one-stage portable battery-powered instrument operated at a constant airflow rate (28.3 L min^{-1}) for a sampling time of 5 min. Petri dishes containing Trypticase soy agar (TSA; BD Bioxon) media, supplemented with 100 mg L^{-1} cycloheximide (Sigma-Aldrich) to prevent fungal growth, were used for capture cultivable total bacteria, and Malt extract agar (EMA; BD Bioxon) for cultivable airborne propagule fungi. The two impactors, one with the TSA and the other one with EMA growing media, were run in parallel. After exposure, the plates were incubated at 37 $^\circ\text{C}$ during 24 h - 48 h for cultivable total bacteria and at 25 $^\circ\text{C}$ during 48 h - 72 h for propagule fungi. After incubation, colonies growing on each plate were counted and concentrations were calculated by taking the sampling rates into account and reported as colony-forming units per cubic meter (cfu m^{-3}) of air. The petri dishes with the grown colonies were stored at 4 $^\circ\text{C}$ prior to their analysis. Fungi were identified to genus level by macroscopic characteristics of the colonies and microscopic examination of the spore structure. Representative bacterial colonies were selected and purified using several transfer steps of single colonies on TSA and checked by Gram-staining and microscopy. Fresh biomass of the bacterial isolates were suspended in 30 % glycerol LB-broth (Alpha Biosciences, Inc.) and stored at -72 $^\circ\text{C}$, for further analyses.

Bacteria isolated from the pure cultures were identified by 16S rRNA sequencing. DNA was extracted using the QIAamp DNA Mini kit (QIAGEN), according to the manufacturers protocol. Partial 16S rRNA gene sequences were amplified by polymerase chain reaction (PCR) using universal bacterial primers 27F (5-AGA GTT TGA TCM TGG CTC AG-3) and 1492R (5-TAC GGY TAC CTT GTT ACG ACT T-3) (Lane, 1991). PCRs were performed in a total volume of 50 μL including 2 μL of bacterial DNA, 35.4 μL of ddH_2O , 5 μL of 10 X buffer, 1.5 μL of MgCl_2 (1.5 mM), 1 μL of dNTPs (10 mM), 0.1 μL of Taq DNA polymerase (5 U μL^{-1}), and 2.5 μL each primer (10 μM). Cycle conditions were as follows: initial denaturation at 94 $^\circ\text{C}$ for 1 min; followed by 35 cycles at 94 $^\circ\text{C}$ for 1 min, 56 $^\circ\text{C}$ for 30 s, 72 $^\circ\text{C}$ for 1.5 min; and a final extension at 72 $^\circ\text{C}$ for 5 min. The PCR products were examined for size and yield using 1.0 % (w/v) agarose gels in TAE buffer. After successful amplification, the obtained products were sequenced using a PRISM 3730 automated sequencer (Applied Biosystem Inc.). DNA sequences were edited and assembled using the Seq-Man and Edit Seq software (Chromas Lite, Technely Slom Pty Ltd. USA). Sequence similarity analysis

was performed using the BLAST software (<http://www.ncbi.nlm.nih.gov/BLAST>).

235 Although specific growing media for actinobacteria were not used in this study, some actinobacteria colonies were able to grow on the TSA petri dishes; therefore, in some cases they were isolated and identified as follows. Genomic DNA was extracted using standard protocols reported previously for actinobacteria (Maldonado et al., 2009). The DNA preparations were then used as template for 16S rRNA gene amplification using the universal set of bacterial primers 27f and 1525r (Lane, 1991). The following components for the PCR mix were employed: 0.5 μL DNA template (for a final concentration of 100 ng μL^{-1} , 5 μL 10X DNA polymerase buffer, 1.5 μL MgCl_2 (50 mM stock solution), 1.25 μL dNTP (10 mM stock mixture), 0.5 μL of each primer (20 μM stock solution), 2.0 240 units of Taq polymerase made up to 50 μL with deionized sterile distilled water.

The PCR amplification was achieved using a Techne 512 gradient machine using the protocol described in Maldonado et al. (2008). The expected product (size approx 1,500 bp) was checked by horizontal electrophoresis (70 V, 40 min) and then purified using the QIAquick PCR purification kit (QIAGEN, Germany) following the manufacturers instructions. Purified 16S rRNA gene PCR products 245 were sent for sequencing to Macrogen (Korea) for the DyeDeoxy Terminator Cycle Sequencing kit (Applied Biosystems). Assembly of each 16S rRNA gene sequence was performed using Chromas (www.technelysium.com.au) and checked manually with the SeaView software (Galtier et al., 1996). Each assembled sequence was compared against two databases, namely, (a) the GenBank database (250 www.ncbi.nlm.nih.gov) by using the BLAST option and (b) the EZCloud (www.ezbiocloud.net) under its EZTaxon option. Both databases generated a list of the closest phylogenetic neighbors to each sequence and the EZTaxon, specifically provided the list of the closest described (type) species. At least 650 bp were employed for the analyzes.

3 Results and Discussion

255 3.1 Aerosol Concentration and Meteorology

Two cold fronts affected Sisal during the sampling period between, January 21 and February 02, 2017, providing different air mass characteristics. The periods affected by each of the fronts are indicated in Figure 2 by vertical grey bars and labeled Cold Front A and Cold Front B, associated with increased wind speed and shifts in wind direction. Figure 2B-D shows the time series of the aerosol particle concentration between January 21 and February 02, 2017. There is a large diurnal variability for the aerosol particle concentration measured by the CPC (particles > 30 nm, Fig. 2B) 260 and the LasAir (particles >300 nm, Fig. 2C), with geometric mean concentrations (assuming log-normal distributions, Limpert et al. (2001)) for the entire sampling period of $758.51 \text{ } \mu\text{g}/\text{m}^3$ and $1.00 \text{ } \mu\text{g}/\text{m}^3$, respectively. From the CPC data shown in Figure 2B, there seems to be a daily cycle with most of the highest concentration taking place between 7 h and 12 h (local time), most notably on days without the influence of cold fronts. The data reported by the CPC and the LasAir 265

indicate that most of the aerosol particles were smaller than 300 nm. A similar result was found by Rosinski et al. (1988) in the Gulf of Mexico (GoM) who found that the aerosol concentration for particles ranging between 0.5 μm and 1.0 μm was three to four orders of magnitude smaller than particles ranging between 0.003 μm and 0.1 μm . A decrease in aerosol particle concentration was observed at the arrival and during the passage of two cold fronts during the sampling period, associated with an increase in horizontal wind speed of at least a factor of three (Fig. 2A). Also note than during the influence of cold front A, the wind direction was almost constant from approximately 270 $^\circ$, while during cold front B the wind direction varied between 270 $^\circ$ and 360 $^\circ$, a more northerly component and a larger influence from the GoM compared to winds associated with cold front A.

Back-trajectories were estimated using the Hysplit model (Stein et al., 2015) from the measurement site. They were run for 72 h for each of the days of the campaign. In the absence of cold fronts A and B, air masses arriving in Sisal had a predominantly continental influence, associated with southerly winds (Figure S2). However, when the cold fronts A and B reached the Yucatan Peninsula, northerly and northwesterly winds prevailed and contributed a more maritime influence. The arrival of the cold fronts was also confirmed by the surface weather maps for January 22 and 29 (Figure S3) provided by the National Oceanic and Atmospheric Administration (NOAA).

Air masses behind both cold fronts, flowing over the GoM, were characterized by lower aerosol particle concentrations than air masses coming from the south to the site. This result agrees well with a large body of evidence indicating that marine air masses have lower aerosol particle concentration than continentally-influenced air masses (Patterson et al., 1980; Fitzgerald, 1991). As for the total aerosol concentration (Figure 2), the number size distributions of the aerosol particles larger than 300 nm were also impacted by the cold fronts. For example, the concentration of particles smaller than 5.0 μm was lower during the passage of the cold fronts (Figure S4). As shown in Figure 3 (and Figure S5), the XRF analysis indicates that although there are small differences in the bulk chemical composition of the aerosol particles is generally comparable in the presence or absence of cold fronts. Note, however, that this is not a completely fair comparison given that sampling time for the chemical analysis was 48 h, while sampling time for determining the influence of the cold air masses was on the order of 36 h. Therefore, the periods denoted as cold fronts contain aerosol particles that may not technically belong to this category.

3.2 Ice Nucleating Particle Concentration ([INP])

A total of 41 samples (8 stages each) were collected during the Sisal field campaign to calculate the [INP] as a function of temperature and particle size. Some of these samples showed a high ice nucleating activity with onset freezing temperatures found to occur at temperatures as high as -3 $^\circ\text{C}$ (Fig. S6). Figure 4 summarizes the [INP] as a function of temperature and particle size for 29 analyzed samples. Due to technical issues it was not possible to analyze the samples collected after January 30th. Figure 4 also shows recent literature data obtained at coastal/marine regions from

DeMott et al. (2016), Welti et al. (2018), and Irish et al. (2019).

At -15 °C the [INP] measured in Sisal are in relatively good agreement with those found at Cabo Verde (Welti et al., 2018) but are one to two orders of magnitude higher than the values reported by Irish et al. (2019) from the Arctic boundary layer, and by DeMott et al. (2016) from sea spray laboratory generated particles and ambient marine boundary layer particles. As temperature decreases from -20 °C to -30 °C there is a better agreement between the Sisal [INP] and data from DeMott et al. (2016). It is important to note that the large variability of the [INP] from Welti et al. (2018) is related to the large amount of data summarized on each dotted line (i.e., from 2009 to 2013).

The “high” [INP] found at -15 °C can be explained in part by the very efficient INPs shown in Figure S6 with sizes ranging from 1.0 μm to 1.8 μm . However, it is important to note that particles with diameters between 1.8 μm and 10 μm also contribute to the total [INP] at warm temperatures. Aerosol particles acting as INPs at -15 °C are usually biological given that other aerosol particles such as metals, crystalline salts, combustion particles (e.g., soot), and organics are not efficient INPs under these conditions. Moreover, for typical atmospheric concentrations of mineral dust, ice nucleation at these temperatures seems to be of secondary importance (Hoose and Möhler, 2012; Murray et al., 2012; Kanji et al., 2017). The potential sources of the measured INPs in Sisal are discussed below.

Figure 5 is based on Mason et al. (2016) and shows the average [INP] for three different temperatures (-15 °C, -20 °C, and -25 °C) at different locations around the globe using the same sampling and analysis methods. The Sisal data corresponds to particle diameters ranging between 0.32 μm and 10 μm ; full information in all size stages was obtained in 16 out of the 29 samples analyzed. At -15 °C the average [INP] in Sisal was lower than Kolby (USA) and Labrador Sea; however, the obtained values are comparable to those found at UBC (Canada), Saclay (France), and Ucluelet (Canada). At -20 °C and -25 °C the average [INP] in Sisal were comparable or higher than at the other locations. As shown by the stars on top of the Sisal bars, the [INP] during the passage of the cold fronts was found to be higher than the average [INP] although the obtained values are within the uncertainty bars. For example, at -15 °C the [INP] increases from 0.33 L^{-1} to 0.59 L^{-1} in the cold air mass after the passage of cold front B. Recalling that the air masses behind cold front B contained a lower aerosol particle concentration, this suggests that the marine particles in that air mass are more efficient INPs than in the air masses with more continental influence. Given that the bulk chemical composition as shown in Figure S5 (and Fig. 3) is comparable before, during, and after the passage of the cold front B, it is possible that the observed differences in the ice nucleating abilities are linked with the biological content in the cold air masses. This is further discussed below.

The majority of the field studies performed to measure the [INP] have been conducted in mid-latitudes; nevertheless, we compare here our observations with the results presented by Rosinski et al. (1988) who measured the [INP] in the condensation freezing mode for particles in the GoM during a cruise between July 20 and August 30, 1986, during mid-summer. The study reports very

340 efficient INPs with onset freezing temperatures as high as $-4\text{ }^{\circ}\text{C}$ for particles with diameters between
0.1 μm and 0.4 μm . On August 6, 1986 (the closest sampling site to Sisal in the GoM) the study
shows that the [INP] at $-15\text{ }^{\circ}\text{C}$ was on the order of 10^{-2} L^{-1} for particles with sizes between 0.1 μm
and 0.4 μm . In contrast, our results indicate that the [INP] at $-15\text{ }^{\circ}\text{C}$ varied between 10^{-1} L^{-1} and
345 10^0 L^{-1} for particles ranging between 0.32 μm and 10 μm . This discrepancy could be attributed
to the differences in the size of the particles sampled and could also be influenced by seasonal
variability. As shown by DeMott et al. (2010) particles larger than 500 nm are the more likely
potential INPs and as stated by Mason et al. (2016) and as shown in Figure 4, super-micron particles
are a large contributor to the INP population. Additionally, the chemical composition of the aerosol
particles collected by Rosinski et al. (1988) indicate that the air masses in the GoM in July-August
350 were significantly influenced by mineral dust particles. African dust episodes reach Florida between
May and October (Lenes et al., 2012), and there have been no reported episodes during the sampling
period of this study (Jan-Feb).

3.2.1 [INP] vs. particle size

Figure 6 shows the mean [INP] concentration as a function of particle size between 0.32 μm and
355 10 μm at four different temperatures ($-15\text{ }^{\circ}\text{C}$, $-20\text{ }^{\circ}\text{C}$, $-25\text{ }^{\circ}\text{C}$, and $-30\text{ }^{\circ}\text{C}$). Note that the INP size
distributions are different for each of the temperatures considered, in contrast with the results from
Mason et al. (2015b) on the Pacific coast of Canada. At $-15\text{ }^{\circ}\text{C}$ the peak [INP] corresponds to parti-
cles ranging between 1.0 μm and 1.8 μm ; this range has been reported as the typical size for airborne
bacteria (Burrows et al., 2009). Similar size distributions were obtained at $-20\text{ }^{\circ}\text{C}$ and $-25\text{ }^{\circ}\text{C}$ with
360 peak concentration for particles ranging in size between 3.2 μm and 5.6 μm . Finally, at $-30\text{ }^{\circ}\text{C}$ the
peak was observed at smaller sizes (i.e., between 1.8 μm and 3.2 μm). The discrepancies between
the present results and those from Mason et al. (2015b) at $-15\text{ }^{\circ}\text{C}$ and $-30\text{ }^{\circ}\text{C}$ could be explained
by differences in air mass history. Although both studies were conducted at coastal locations, the
back-trajectories from the present study indicate that during “normal” days (i.e., 70 % of the time)
365 the sampled air masses had a significant continental contribution (Fig. S2). In contrast, air masses
were mostly maritime in the Mason et al. (2015b) study. Also, it is important to note that although
the cold air masses that reached Sisal behind the cold fronts had crossed the GoM, the aerosol parti-
cles found in them are likely a mixture of particles originated in the US Central Plains and the GoM
(Figs. S2B-C and S5).

370 Figure 6 also shows that most of the INPs are in the supermicron size range, where submicron
particles represent less than 10 % of the total [INP] independent of temperature, in agreement with
Mason et al. (2015b) and Mason et al. (2016). To confirm the size dependence and the importance
of supermicron particles to the [INP] in Sisal, the fraction of particles acting as INPs was calculated
combining the DFT and LasAir data (Fig.7). The [INP] was normalized for four size bins (i.e.,
375 0.3 μm - 0.5 μm , 0.5 μm - 1.0 μm , 1.0 μm - 5.0 μm , and 5.0 μm - 10.0 μm). As expected (from

Figs. 4 and 6), the fraction of particles acting as INPs increases with increasing particle size and with decreasing temperature. This trend is in agreement with the results shown by Si et al. (2018), with the present results being higher. Figure 7 also shows that the fraction of aerosol particles acting as INP is higher when influenced by the cold fronts (black symbols), especially for particles ranging
380 between 1.0 μm and 5.0 μm .

3.3 Identification of the Potential INP Sources

The chemical analyzes of the sampled aerosol particles (for the whole sampling period) indicate that a large fraction of the particle mass (for sizes between 0.18 μm and 10.0 μm) are likely of marine origin (Figures 3 and 8(A-B)). Both techniques, i.e., XRF and HPLC found that the main
385 elements and ions are sodium and chlorine. The low concentrations of Ti, Cu, K, and Zn shows the very low probability of anthropogenic influence at the sampling site. However, although sulfate and ammonium can be emitted by natural sources, their presence, in addition to nitrates, indicate that the influence of anthropogenic activities to the aerosol population is not completely negligible. Finally, the low concentration of Al, Fe, Ca, and Si suggest that mineral dust is not a major contributor
390 of aerosol particle mass during the sampling period. However, although the long-range transport of mineral dust particles from Africa to the Yucatan Peninsula and the GoM is very rare between January and February, mineral dust particles are frequently found in the Caribbean including the GoM (Rosinski et al., 1988; Prospero and Lamb, 2003; Doherty et al., 2008; Kishcha et al., 2014).

Figure 8(C-D) shows the mean mass size distribution for the whole sampling period of the main
395 five elements/ions determined by the XRF and HPLC techniques. For the XRF analyzes Na, Cl, and Ca, have a single peak at 3.2 μm , whereas the S and Mg reported two peaks at 0.32 μm and 3.2 μm . Similar to the XRF results, the HPLC analyzes for Na^+ and Cl^- also showed a single peak at 3.2 μm . SO_4^{2-} , NO_3^- showed two peaks at 0.32 μm and 3.2 μm , whereas for NH_4^+ the peaks were located at 0.32 μm and 5.6 μm . The obtained size distributions are in agreement with those of
400 sea salt type particles as reported elsewhere (O'dowd et al., 2004; Prather et al., 2013).

Although Al, Si, Ca, and Fe were found in low concentrations (Fig. 8A), Tables 2 and S1 suggest that mineral dust particles are an important source of INPs in Sisal at temperatures ranging from -20 $^\circ\text{C}$ to -30 $^\circ\text{C}$. This is in close agreement with the results obtained by Si et al. (2019) in the Canadian high Arctic. Given that mineral dust particles are unlikely the source of the measured INPs
405 above -15 $^\circ\text{C}$ (as suggested by Table 2), and as organics and soot are not typically efficient INPs at temperatures above -15 $^\circ\text{C}$ (Kanji et al., 2017), in addition to the supermicron size of ca. 90 % of the INPs (Fig. 6), bioaerosol is a potential source of the INPs measured at warm temperatures. Note that bioparticles have been shown to efficiently nucleate ice at those high temperatures (Hoose and Möhler, 2012; Murray et al., 2012; Ladino et al., 2013). To confirm the presence of bioparticles
410 around Sisal and to determine their potential role in the ice nucleating abilities of the collected aerosol particles, bacteria and fungi identification was performed.

Samples for viable bacteria and fungi were collected every day at 6:00, 8:00, 10:00, and 12:00 local time. However, a single daily profile was performed between January 22 and 23. Bacteria and fungi colony forming units (cfu) m^{-3} were usually above zero with the highest concentrations found early in the morning (Figure S7). The bacteria and fungi concentrations showed a relatively good correlation ($r=0.55$, $p<0.0005$ not shown) with average values for the whole sampling period of 295 ± 312 cfu m^{-3} and 438 ± 346 cfu m^{-3} , respectively. The bacteria concentration are comparable to the values found by Hurtado et al. (2014) in Tijuana, on the Pacific coast of Mexico (i.e., 230 cfu m^{-3} - 280 cfu m^{-3}). Bacteria and fungi concentrations were found to be lower when the wind was coming from the north in comparison with southern-continental air masses (Figure S8), a behavior similar to the aerosol concentration shown in section 3.2.

Figure 9 shows the time series of the [INP] together with the bacteria and fungal concentrations. Panels B and C show a poor correlation between the bacteria and fungal concentrations with the [INP] with correlation coefficients at -15 °C of 0.12 ($p=0.06$) and 0.36 ($p=0.03$), respectively. This poor correlation can be in part due to the different sampling time of the MOUDI and the biosamplers. Another additional factor is the fact that the reported bacteria and fungi concentrations are only a small fraction of the total population given that the used method is selective to viable microorganisms only. Note that the fraction of detected microorganisms by culture methods are typically ca. 1 % (but can be lower) of the total population (Lighthart, 2000; Burrows et al., 2009). From Figure 9 it is notable that although the bacteria and fungi concentrations were very low on January 29 (i.e., under the influence of the cold front B), the [INP] at -15 °C was comparable to the average value for the entire campaign. It is therefore intriguing if the marine microorganisms brought to Sisal by the cold front B could be efficient INPs.

Table 3 summarizes the identified bacteria before the arrival of cold front A and after the passage of cold fronts A and B. Additionally, Table 4 shows the fungi identification for the whole campaign. To our knowledge this is the first time that airborne viable bacteria, and fungi are identified at this coastal location. Although biological microorganism characterization has been previously conducted in Mexico, those studies focused on health effects mainly (Santos-Burgoa et al., 1994; Guzman, 1998; Maldonado et al., 2009; Frías-De León et al., 2016; Ríos et al., 2016). Note that 76 % of the detected bacteria were Gram positive with *Micrococcus*, *Staphylococcus*, and *Bacillus* as the main identified genus (Figure S9). As shown in Table 3, before the arrival of cold front A (January 21-22), a large variety of bacteria species were found with different typical sources, mostly terrestrial. This is in contrast with the identified species found after the passage of cold fronts A and B. Especially, after cold front B different *Vibrio* species were identified, most of which are typically of marine origin. Recently, Hurtado et al. (2014) found that the most common genera of the bacteria in Tijuana were *Staphylococcus*, *Streptococcus*, *Pseudomonas*, and *Bacillus* in close agreement with the present results.

Regarding fungi, different genus were also identified as shown in Table 4 with *Cladosporium* and

Penicillium as the most frequent ones (51 % and 11 %, respectively) as shown in Figure S9. This is
450 in good agreement with the data reported by Després et al. (2012).

Several studies have shown the good correlation between the concentration of fluorescent biological particles and the [INP]; however, from those studies it is highly uncertain if the good ice nucleating abilities can be attributed to a single microorganism species (Mason et al., 2015b; Twohy et al., 2016). Off-line methods as the one used here have been able to identify from rain water and
455 cloud water specific microorganisms such as *Pseudomonas syringae*, *Micrococcus*, *Staphylococcus*, *Cladosporium*, *Penicillium*, *Aspergillus* among others, with some showing good ice nucleating ability (Amato et al., 2007, 2017; Delort et al., 2010; Failor et al., 2017; Stopelli et al., 2017; Akila et al., 2018).

4 Conclusions

460 Aerosol particles around Sisal (on the northwest coast of the Yucatan peninsula) from January 21 to February 02, 2017 were found to be efficient INPs with onset freezing temperatures as high as $-3\text{ }^{\circ}\text{C}$, similar to the onset freezing temperature of the well known efficient INP *Pseudomonas syringae* (Wex et al., 2015) and Arctic sea surface microlayer organic-enriched waters (Wilson et al., 2015). The results show that the INPs concentrations in Sisal are comparable ($0.33\pm 0.53\text{ L}^{-1}$,
465 $2.20\pm 2.26\text{ L}^{-1}$, and $8.96\pm 7.38\text{ L}^{-1}$ at $-15\text{ }^{\circ}\text{C}$, $-20\text{ }^{\circ}\text{C}$, $-25\text{ }^{\circ}\text{C}$, respectively) and in specific cases even higher than at other locations studied using the same INP counter type, especially under the influence of cold fronts. This is an intriguing result given that the air masses behind the cold front contained lower aerosol particle concentrations. This deserves further analysis given that the Yucatan peninsula and the Caribbean region are impacted regularly by this meteorological phenomenon
470 during the winter and early spring months.

The chemical analyzes performed on the sampled aerosol particles did not indicate the presence of mineral dust particles in high concentrations (the combined mass concentrations of Al, Si, and Fe correspond to 5.1 % of the total particle mass measured by the XRF); however, Al, Si, Ca, and Fe showed high correlation coefficients (r^2 above 0.64 with $p < 0.05$) with the [INP] at temperatures
475 between $-20\text{ }^{\circ}\text{C}$ and $-30\text{ }^{\circ}\text{C}$. From the correlation of the [INP] and the aerosol chemical composition at $-15\text{ }^{\circ}\text{C}$ magnesium was the only element showing a correlation that is statistically significant at the 95 % confidence interval ($p < 0.05$). Although Mg can be found in mineral dust particles in low percentages, it can also be found in marine environments linked to sea spray aerosol (e.g., Savoie and Prospero (1980); Andreae (1982); Casillas-Ituarte et al. (2010)). At $-15\text{ }^{\circ}\text{C}$ the [INP] in Sisal was
480 one to two orders of magnitude higher than the concentrations reported from other coastal/marine regions around the globe. The size of this very efficient INPs was found to be above $1.0\text{ }\mu\text{m}$ with a large contribution to the [INP] of particle ranging from $1.0\text{ }\mu\text{m}$ to $1.8\text{ }\mu\text{m}$. Based on the large [INPs] above $-15\text{ }^{\circ}\text{C}$, the supermicron size of 90 % of the INPs, the presence of marine biological particles

in the cold air masses those of which showed the highest [INP], in addition to the poor correlation
485 shown by the mineral dust tracers with the [INP], we hypothesize that the likely source of the INPs
measured in Sisal at high temperatures are biological particles. Therefore, our results suggest that
continental and maritime biological particles could play an important role in ice cloud formation
and precipitation development in the Yucatan peninsula. Although several bacteria and fungi were
490 identified, it is unknown if any of them were responsible for the observed ice nucleating abilities of
the aerosol around Sisal. As stated by Islebe et al. (2015) both bacteria and fungi need to be properly
documented in the peninsula and the GoM to fully understand their regional importance.

The present results are important for the development of new parametrizations to be incorporated
in climate models given that the currently available parametrizations contain little or no data from
tropical latitudes. However, further similar studies are needed given that the [INP] may vary sea-
495 sonally. Especially, the arrival of mineral dust particles to the GoM and the Caribbean region from
Africa in July-August are expected to impact the [INP] and therefore, ice cloud formation, as shown
by Rosinski et al. (1988) and DeMott et al. (2003).

The quantitative understanding of the importance of biological particles in ice particle formation is
a challenging task for the cloud physics community. As shown here, even when combining biology
500 with chemistry, physics, and meteorology, the results obtained are not as quantitative as would be
desired. Efficient INPs such as those measured in Sisal could be very important for cloud glaciation.
Additionally, they can trigger ice multiplication or secondary ice formation at such high tempera-
tures via the Hallett-Mossop mechanism (Hallett and Mossop, 1974; Field et al., 2017) and impact
precipitation formation. Therefore, further studies are needed in order to improve our current limited
505 understanding of the role that tropical microorganisms could play in ice cloud formation.

Author contributions. LAL and GBR designed the experiments. LAL, HAO, MAE, and BF
carried out the INP and aerosol measurements. IR, LM, ES, EQ, LAM, and AGR analyzed the
biological particles. HAO and JM performed the chemical analyzes. LAL, ZRD, CC, AKB, MS and
VI performed the INP analyzes. LAL wrote the paper, with contributions from all co-authors.

510 *Acknowledgements.* The authors thank E. Garcia, G. Chavez, I. Gavilan, R. Gutierrez, J. Munoz, L. Landeros,
F. Cordoba, W. Gutierrez, M. Garcia, M. Robles, A. Rodriguez, J.C. Pineda, L. Gonzalez, A. Prieto, T. Castro,
M.I. Saavedra, and A. Cruz for their invaluable help. We also thank David S. Valdes from CINVESTAV Merida
for sharing the meteorological data. Finally, we thank the National Oceanic and Atmospheric Administration
(NOAA) for facilitating the use of the surface maps and the Hysplit. This study was financially supported by
515 the Direccion General de Asuntos del Personal Academico (DGAPA) and by the Consejo Nacional de Ciencia
y Tecnologia (Conacyt) through grants PAPIIT IA108417 and IN102818 and I000/781/2106.

References

- Akila, M., Priyamvada, H., Ravikrishna, R., and Gunthe, S. S.: Characterization of bacterial diversity and ice-nucleating ability during different monsoon seasons over a southern tropical Indian region, *Atmos. Environ.*, 520 191, 387–394, doi:10.1016/j.atmosenv.2018.08.026, 2018.
- Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G., and Delort, A.-M.: Microorganisms isolated from the water phase of tropospheric clouds at the Puy de Dôme: major groups and growth abilities at low temperatures, *FEMS Microbiol. Ecol.*, 59, 242–254, doi:10.1111/j.1574-6941.2006.00199.x, 2007.
- Amato, P., Joly, M., Besaury, L., Oudart, A., Taib, N., Moné, A. I., Deguillaume, L., Delort, A.-M., and 525 Debroas, D.: Active microorganisms thrive among extremely diverse communities in cloud water, *PloS one*, 12, e0182869, doi:10.1371/journal.pone.0182869, 2017.
- Andreae, M. and Rosenfeld, D.: Aerosol–cloud–precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth Sci. Rev.*, 89, 13–41, doi:10.1016/j.earscirev.2008.03.001, 2008.
- Andreae, M. O.: Marine aerosol chemistry at Cape Grim, Tasmania, and Townsville, Queensland, *Journal of* 530 *Geophysical Research: Oceans*, 87, 8875–8885, doi:10.1029/JC087iC11p08875, 1982.
- Ansmann, A., Mattis, I., Müller, D., Wandinger, U., Radlach, M., Althausen, D., and Damoah, R.: Ice formation in Saharan dust over central Europe observed with temperature/humidity/aerosol Raman lidar, *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD005000, 2005.
- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., 535 OSullivan, D., and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, *Nature*, 498, 355, doi:10.1038/nature12278, 2013.
- Bateman, A. P., Belassein, H., and Martin, S. T.: Impactor apparatus for the study of particle rebound: Relative humidity and capillary forces, *Aerosol Sci. Technol.*, 48, 42–52, doi:10.1080/02786826.2013.853866, 2014.
- 540 Bigg, E.: Ice nucleus concentrations in remote areas, *J. Atmos. Sci.*, 30, 1153–1157, doi:/10.1175/1520-0469(1973)030<1153:INCIRA>2.0.CO;2, 1973.
- Boose, Y., Sierau, B., García, M. I., Rodríguez, S., Alastuey, A., Linke, C., Schnaiter, M., Kupiszewski, P., Kanji, Z. A., and Lohmann, U.: Ice nucleating particles in the Saharan Air Layer, *Atmos. Chem. Phys.*, 16, 9067–9087, doi:10.5194/acp-16-9067-2016, 2016.
- 545 Burrows, S. M., Elbert, W., Lawrence, M., and Pöschl, U.: Bacteria in the global atmosphere—Part 1: Review and synthesis of literature data for different ecosystems, *Atmos. Chem. Phys.*, 9, 9263–9280, doi:10.5194/acp-9-9263-2009, 2009.
- Burrows, S. M., Hoose, C., Pöschl, U., and Lawrence, M.: Ice nuclei in marine air: biogenic particles or dust?, *Atmos. Chem. Phys.*, 13, 245–267, doi:10.5194/acp-13-245-2013, 2013.
- 550 Casillas-Ituarte, N. N., Callahan, K. M., Tang, C. Y., Chen, X., Roeselová, M., Tobias, D. J., and Allen, H. C.: Surface organization of aqueous MgCl₂ and application to atmospheric marine aerosol chemistry, *Proceedings of the National Academy of Sciences*, 107, 6616–6621, doi:10.1073/pnas.0912322107, 2010.
- Chen, S.-C., Tsai, C.-J., Chen, H.-D., Huang, C.-Y., and Roam, G.-D.: The influence of relative humidity on nanoparticle concentration and particle mass distribution measurements by the MOUDI, *Aerosol Sci. Technol.*, 45, 596–603, doi:10.1080/02786826.2010.551557, 2011.
- 555 Chow, J. C. and Watson, J. G.: Ion chromatography in elemental analysis of airborne particles, *Elemental*

- analysis of airborne particles, 1, 97–137, 1999.
- Cui, Z., Carslaw, K. S., Yin, Y., and Davies, S.: A numerical study of aerosol effects on the dynamics and microphysics of a deep convective cloud in a continental environment, *Journal of Geophysical Research: Atmospheres*, 111, doi:10.1029/2005JD005981, 2006.
- 560 DeLeon-Rodriguez, N., Latham, T. L., Rodriguez-R, L. M., Barazesh, J. M., Anderson, B. E., Beyersdorf, A. J., Ziemba, L. D., Bergin, M., Nenes, A., and Konstantinidis, K. T.: Microbiome of the upper troposphere: species composition and prevalence, effects of tropical storms, and atmospheric implications, *P. Natl. Acad. Sci.*, 110, 2575–2580, doi:10.1073/pnas.1212089110, 2013.
- 565 Delort, A.-M., Vaitilingom, M., Amato, P., Sancelme, M., Parazols, M., Mailhot, G., Laj, P., and Deguillaume, L.: A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes, *Atmos. Res.*, 98, 249–260, doi:10.1016/j.atmosres.2010.07.004, 2010.
- DeMott, P. J., Sassen, K., Poellot, M. R., Baumgardner, D., Rogers, D. C., Brooks, S. D., Prenni, A. J., and Kreidenweis, S. M.: African dust aerosols as atmospheric ice nuclei, *Geophys. Res. Lett.*, 30, doi:10.1029/2003GL017410, 2003.
- 570 DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M., Eidhammer, T., and Rogers, D.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *P. Natl. Acad. Sci.*, 107, 11 217–11 222, doi:10.1073/pnas.0910818107, 2010.
- DeMott, P. J., Hill, T. C., McCluskey, C. S., Prather, K. A., Collins, D. B., Sullivan, R. C., Ruppel, M. J., Mason, R. H., Irish, V. E., Lee, T., et al.: Sea spray aerosol as a unique source of ice nucleating particles, *P. Natl. Acad. Sci.*, 113, 5797–5803, doi:10.1073/pnas.1514034112, 2016.
- 575 Després, V., Huffman, J. A., Burrows, S. M., Hoose, C., Safatov, A., Buryak, G., Fröhlich-Nowoisky, J., Elbert, W., Andreae, M., Pöschl, U., et al.: Primary biological aerosol particles in the atmosphere: a review, *Tellus B*, 64, 15 598, doi:10.3402/tellusb.v64i0.15598, 2012.
- 580 Doherty, O., Riemer, N., and Hameed, S.: Saharan mineral dust transport into the Caribbean: Observed atmospheric controls and trends, *Journal of Geophysical Research: Atmospheres*, 113, doi:10.1029/2007JD009171, 2008.
- Espinosa, A., Miranda, J., and Pineda, J.: Uncertainty evaluation in correlated quantities: application to elemental analysis of atmospheric aerosols, *Rev. Mex. Fis.*, E56, 134–140, 2010.
- 585 Espinosa, A., Reyes-Herrera, J., Miranda, J., Mercado, F., Veytia, M., Cuautle, M., and Cruz, J.: Development of an X-ray fluorescence spectrometer for environmental science applications, *Instr. Sci. Technol.*, 40, 603–617, doi:10.1080/10739149.2012.693560, 2012.
- Failor, K., Schmale Iii, D., Vinatzer, B., and Monteil, C. L.: Ice nucleation active bacteria in precipitation are genetically diverse and nucleate ice by employing different mechanisms, *ISME J.*, 11, 2740, doi:10.1038/ismej.2017.124, 2017.
- 590 Field, P. R., Lawson, R. P., Brown, P. R., Lloyd, G., Westbrook, C., Moisseev, D., Miltenberger, A., Nenes, A., Blyth, A., Choulaton, T., et al.: Secondary Ice Production: Current State of the Science and Recommendations for the Future, *Meteorol. Monogr.*, 58, 7.1–7.20, doi:10.1175/AMSMONOGRAPHS-D-16-0014.1, 2017.
- 595 Fitzgerald, J. W.: Marine aerosols: A review, *Atmos. Environ.*, 25, 533–545, doi:10.1016/0960-1686(91)90050-H, 1991.

- Frías-De León, M. G., Duarte-Escalante, E., del Carmen Calderón-Ezquerro, M., del Carmen Jiménez-Martínez, M., Acosta-Altamirano, G., Moreno-Eutimio, M. A., Zúñiga, G., García-González, R., Ramírez-Pérez, M., and del Rocío Reyes-Montes, M.: Diversity and characterization of airborne bacteria at two health
600 institutions, *Aerobiologia*, 32, 187–198, doi:10.1007/s10453-015-9389-z, 2016.
- Fröhlich-Nowoisky, J., Hill, T. C., Pummer, B. G., Yordanova, P., Franc, G. D., and Pöschl, U.: Ice nucleation activity in the widespread soil fungus *Mortierella alpina*, *Biogeosciences*, 12, 1057, doi:10.5194/bg-12-1057-2015, 2015.
- Fröhlich-Nowoisky, J., Kampf, C. J., Weber, B., Huffman, J. A., Pöhlker, C., Andreae, M. O., Lang-Yona,
605 N., Burrows, S. M., Gunthe, S. S., Elbert, W., et al.: Bioaerosols in the Earth system: Climate, health, and ecosystem interactions, *Atmos. Res.*, 182, 346–376, doi:10.1016/j.atmosres.2016.07.018, 2016.
- Galtier, N., Gouy, M., and Gautier, C.: SEAVIEW and PHYLO_WIN: two graphic tools for sequence alignment and molecular phylogeny, *Bioinformatics*, 12, 543–548, doi:10.1093/bioinformatics/12.6.543, 1996.
- Guzmán, G.: New species of fungi from the Yucatan Peninsula, *Mycotaxon*, 1982.
- 610 Guzman, G. N.: Inventorying the fungi of Mexico, *Biodivers. Conserv.*, 7, 369–384, 1998.
- Hallett, J. and Mossop, S.: Production of secondary ice particles during the riming process, *Nature*, 249, 26, 1974.
- Hill, T. C., DeMott, P., Conen, F., and Möhler, O.: Impacts of Bioaerosols on Atmospheric Ice Nucleation Processes, *Microbiol. Aerosol.*, p. 197, 2017.
- 615 Hobbs, P. V. and Atkinson, D. G.: The concentrations of ice particles in orographic clouds and cyclonic storms over the Cascade Mountains, *Journal of the Atmospheric Sciences*, 33, 1362–1374, doi:10.1175/1520-0469(1976)033<1362:TCOPI>2.0.CO;2, 1976.
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 98179854, doi:10.5194/acp-12-9817-2012, 2012.
- 620 Hoose, C., Kristjánsson, J., and Burrows, S.: How important is biological ice nucleation in clouds on a global scale?, *Environ. Res. Lett.*, 5, 024009, doi:10.1088/1748-9326/5/2/024009, 2010.
- Hurtado, L., Rodríguez, G., López, J., Castillo, J., Molina, L., Zavala, M., and Quintana, P. J.: Characterization of atmospheric bioaerosols at 9 sites in Tijuana, Mexico, *Atmos. Environ.*, 96, 430–436, doi:10.1016/j.atmosenv.2014.07.018, 2014.
- 625 IAEA: Manual for QXAS, International Atomic Energy Agency, p. Vienna, 1997.
- Irish, V. E., Elizondo, P., Chen, J., Chou, C., Charette, J., Lizotte, M., Ladino, L. A., Wilson, T. W., Gosselin, M., Murray, B. J., et al.: Ice-nucleating particles in Canadian Arctic sea-surface microlayer and bulk seawater, *Atmos. Chem. Phys.*, 17, 10583–10595, doi:10.5194/acp-17-10583-2017, 2017.
- Irish, V. E., Hanna, S. J., Willis, M. D., China, S., Thomas, J. L., Wentzell, J. J., Cirisan, A., Si, M., Leaitch,
630 W. R., Murphy, J. G., et al.: Ice nucleating particles in the marine boundary layer in the Canadian Arctic during summer 2014, *Atmospheric Chemistry and Physics*, 19, 1027–1039, 2019.
- Islebe, G. A., Calmé, S., León-Cortés, J. L., and Schmook, B.: Biodiversity and conservation of the Yucatán Peninsula, Springer, 2015.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of
635 ice nucleating particles, *Meteorol. Monogr.*, 58, 1.1–1.33, doi:10.1175/AMSMONOGRAPHS-D-16-0006.1, 2017.

- Kishcha, P., da Silva, A., Starobinets, B., Long, C., Kalashnikova, O., and Alpert, P.: Meridional distribution of aerosol optical thickness over the tropical Atlantic Ocean, *Atmos. Chem. Phys. Discuss.*, 14, 23 309–23 339, doi:10.5194/acpd-14-23309-2014, 2014.
- 640 Knopf, D., Alpert, P., Wang, B., and Aller, J.: Stimulation of ice nucleation by marine diatoms, *Nat. Geosci.*, 4, 88, doi:10.1038/NGEO1037, 2011.
- Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P., Srivastava, A., and Sugi, M.: Tropical cyclones and climate change, *Nat. Geosci.*, 3, 157, doi:10.1038/ngeo779, 2010.
- 645 Ladino, L., Stetzer, O., and Lohmann, U.: Contact freezing: a review of experimental studies, *Atmos. Chem. Phys.*, 13, 9745–9769, doi:10.5194/acp-13-9745-2013, 2013.
- Ladino, L., Yakobi-Hancock, J., Kilthau, W., Mason, R., Si, M., Li, J., Miller, L., Schiller, C., Huffman, J., Aller, J., et al.: Addressing the ice nucleating abilities of marine aerosol: A combination of deposition mode laboratory and field measurements, *Atmos. Environ.*, 132, 1–10, doi:10.1016/j.atmosenv.2016.02.028, 2016.
- 650 Landsea, C.: Counting Atlantic tropical cyclones back to 1900, *Eos*, 88, 197–202, doi:10.1029/2007EO180001, 2007.
- Lane, D.: 16S/23S rRNA sequencing. In *Nucleic acid techniques in bacterial systematics*. (Eds E Stackebrandt, M Goodfellow) pp. 115–175, 1991.
- Lenes, J., Prospero, J., Landing, W., Virmani, J., and Walsh, J.: A model of Saharan dust deposition to the eastern Gulf of Mexico, *Marine Chemistry*, 134, 1–9, doi:10.1016/j.marchem.2012.02.007, 2012.
- 655 Lighthart, B.: Mini-review of the concentration variations found in the alfresco atmospheric bacterial populations, *Aerobiologia*, 16, 7–16, 2000.
- Limpert, E., Stahel, W. A., and Abbt, M.: Log-normal distributions across the sciences: keys and clues: on the charms of statistics, and how mechanical models resembling gambling machines offer a link to a handy way to characterize log-normal distributions, which can provide deeper insight into variability and probability normal or log-normal: that is the question, *BioScience*, 51, 341–352, doi:10.1641/0006-3568(2001)051[0341:LNDATS]2.0.CO;2, 2001.
- 660 Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–737, doi:10.5194/acp-5-715-2005, 2005.
- 665 Maldonado, L. A., Stach, J. E., Ward, A. C., Bull, A. T., and Goodfellow, M.: Characterisation of micromonosporae from aquatic environments using molecular taxonomic methods, *Anton. Leeuw.*, 94, 289–298, doi:10.1007/s10482-008-9244-0, 2008.
- Maldonado, L. A., Fragoso-Yáñez, D., Pérez-García, A., Rosellón-Druker, J., and Quintana, E. T.: Actinobacterial diversity from marine sediments collected in Mexico, *Anton. Leeuw.*, 95, 111–120, doi:10.1007/s10482-008-9294-3, 2009.
- 670 Mason, R., Chou, C., McCluskey, C., Levin, E., Schiller, C., Hill, T., Huffman, J., DeMott, P., and Bertram, A.: The micro-orifice uniform deposit impactor–droplet freezing technique (MOUDI-DFT) for measuring concentrations of ice nucleating particles as a function of size: improvements and initial validation, *Atmos. Meas. Tech.*, 8, 2449–2462, doi:10.5194/amt-8-2449-2015, 2015a.
- 675 Mason, R., Si, M., Li, J., Chou, C., Dickie, R., Toom-Sauntry, D., Pöhlker, C., Yakobi-Hancock, J., Ladino, L., Jones, K., et al.: Ice nucleating particles at a coastal marine boundary layer site: correlations with aerosol type

- and meteorological conditions, *Atmos. Chem. Phys.*, 15, 12 547–12 566, doi:10.5194/acp-15-12547-2015, 2015b.
- 680 Mason, R., Si, M., Chou, C., Irish, V., Dickie, R., Elizondo, P., Wong, R., Brintnell, M., Elsasser, M., Lassar, W., et al.: Size-resolved measurements of ice-nucleating particles at six locations in North America and one in Europe, *Atmos. Chem. Phys.*, 16, 1637–1651, doi:10.5194/acp-16-1637-2016, 2016.
- Mayol, E., Arrieta, J. M., Jiménez, M. A., Martínez-Asensio, A., Garcias-Bonet, N., Dachs, J., González-Gaya, B., Royer, S.-J., Benítez-Barrios, V. M., Fraile-Nuez, E., et al.: Long-range transport of airborne microbes over the global tropical and subtropical ocean, *Nat. Commun.*, 8, 201, doi:10.1038/s41467-017-00110-9, 685 2017.
- McCluskey, C. S., Hill, T. C., Malfatti, F., Sultana, C. M., Lee, C., Santander, M. V., Beall, C. M., Moore, K. A., Cornwell, G. C., Collins, D. B., et al.: A Dynamic Link between Ice Nucleating Particles Released in Nascent Sea Spray Aerosol and Oceanic Biological Activity during Two Mesocosm Experiments, *J. Atmos. Sci.*, 74, 151–166, doi:10.1175/JAS-D-16-0087.1, 2017.
- 690 Michaud, J., Luke, T., Drishti, K., Josh, E., Alexander, R., Zhenjiang, Z., Christopher, L., Kevin, P., Charlotte, B., Francesca, M., Farooq, A., Rob, K., Michael, B., Christopher, D., and Kimberly, P.: Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-atmosphere mesocosm, *Nat. Commun.*, 18, 1–10, doi:10.1038/s41467-018-04409-z, 2018.
- Morales, J. L., Cantillo-Ciau, Z. O., Sánchez-Molina, I., and Mena-Rejón, G. J.: Screening of antibacterial and antifungal activities of six marine macroalgae from coasts of Yucatan peninsula, *Pharm. Biol.*, 44, 632–635, 695 doi:10.1080/13880200600897569, 2006.
- Mülmenstädt, J., Sourdeval, O., Delanoë, J., and Quaas, J.: Frequency of occurrence of rain from liquid-, mixed-, and ice-phase clouds derived from A-Train satellite retrievals, *Geophys. Res. Lett.*, 42, 6502–6509, doi:10.1002/2015GL064604, 2015.
- 700 Murray, B., O'sullivan, D., Atkinson, J., and Webb, M.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519–6554, doi:10.1039/C2CS35200A, 2012.
- Nenes, A., Murray, B., and Bougiatioti, A.: Mineral dust and its microphysical interactions with clouds, in: *Mineral Dust*, pp. 287–325, Springer, 2014.
- O'dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and 705 Putaud, J.-P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676, doi:10.1038/Nature02959, 2004.
- Patterson, E., Kiang, C., Delany, A., Wartburg, A., Leslie, A., and Huebert, B.: Global measurements of aerosols in remote continental and marine regions: Concentrations, size distributions, and optical properties, *J. Geophys. Res.*, 85, 7361–7376, doi:10.1029/JC085iC12p07361, 1980.
- 710 Phillips, V. T., Donner, L. J., and Garner, S. T.: Nucleation processes in deep convection simulated by a cloud-system-resolving model with double-moment bulk microphysics, *Journal of the atmospheric sciences*, 64, 738–761, doi:10.1175/JAS3869.1, 2007.
- Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., et al.: Bringing the ocean into the laboratory to probe the chemical 715 complexity of sea spray aerosol, *P. Natl. Acad. Sci.*, 110, 7550–7555, doi:10.1073/pnas.1300262110, 2013.
- Price, H., Baustian, K., McQuaid, J., Blyth, A., Bower, K., Choularton, T., Cotton, R., Cui, Z., Field, P.,

- Gallagher, M., et al.: Atmospheric Ice-Nucleating Particles in the Dusty Tropical Atlantic, *J. Geophys. Res.*, 123, 2175–2193, doi:10.1002/2017JD027560, 2018.
- Prospero, J. M. and Lamb, P. J.: African droughts and dust transport to the Caribbean: Climate change implications, *Science*, 302, 1024–1027, doi:10.1126/science.1089915, 2003.
- Ramanathan, V., Cess, R., Harrison, E., Minnis, P., Barkstrom, B., Ahmad, E., and Hartmann, D.: Cloud-radiative forcing and climate: Results from the Earth Radiation Budget Experiment, *Science*, 243, 57–63, doi:10.1126/science.243.4887.57, 1989.
- Ríos, B., Torres-Jardón, R., Ramírez-Arriaga, E., Martínez-Bernal, A., and Rosas, I.: Diurnal variations of airborne pollen concentration and the effect of ambient temperature in three sites of Mexico City, *Int. J. Biometeorol.*, 60, 771–787, doi:10.1007/s00484-015-1061-3, 2016.
- Rosinski, J., Haagenson, P., Nagamoto, C., and Parungo, F.: Nature of ice-forming nuclei in marine air masses, *J. Aerosol Sci.*, 18, 291–309, doi:10.1016/0021-8502(87)90024-3, 1987.
- Rosinski, J., Haagenson, P., Nagamoto, C., Quintana, B., Parungo, F., and Hoyt, S.: Ice-forming nuclei in air masses over the Gulf of Mexico, *J. Aerosol Sci.*, 19, 539–551, doi:10.1016/0021-8502(88)90206-6, 1988.
- Santos-Burgoa, C., Rosas, I., and Yela, A.: Occurrence of airborne enteric bacteria in Mexico City, *Aerobiología*, 10, 39–45, 1994.
- Savoie, D. and Prospero, J.: Water-soluble potassium, calcium, and magnesium in the aerosols over the tropical North Atlantic, *Journal of Geophysical Research: Oceans*, 85, 385–392, doi:10.1029/JC085iC01p00385, 1980.
- Schnell, R.: Ice nuclei produced by laboratory cultured marine phytoplankton, *Geophys. Res. Lett.*, 2, 500–502, doi:10.1029/GL002i011p00500, 1975.
- Schnell, R.: Ice nuclei in seawater, fog water and marine air off the coast of Nova Scotia: Summer 1975, *J. Atmos. Sci.*, 34, 1299–1305, doi:10.1175/1520-0469(1977)034<1299:INISFW>2.0.CO;2, 1977.
- Schnell, R.: Airborne ice nucleus measurements around the Hawaiian Islands, *J. Geophys. Res.*, 87, 8886–8890, doi:10.1029/JC087iC11p08886, 1982.
- Schnell, R. and Vali, G.: Freezing nuclei in marine waters, *Tellus*, 27, 321–323, doi:10.1111/j.2153-3490.1975.tb01682.x, 1975.
- SEDESOL: Secretaria de desarrollo social: Catalogo de localidades, <http://www.microrregiones.gob.mx/catloc/contenido.aspx?refnac=310380004>, 2015.
- Seifert, P., Ansmann, A., Gross, S., Freudenthaler, V., Heinold, B., Hiebsch, A., Mattis, I., Schmidt, J., Schnell, F., Tesche, M., et al.: Ice formation in ash-influenced clouds after the eruption of the Eyjafjallajökull volcano in April 2010, *Journal of Geophysical Research: Atmospheres*, 116, doi:10.1029/2011JD015702, 2011.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., et al.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *P. Natl. Acad. Sci.*, 113, 5781–5790, doi:10.1073/pnas.1514043113, 2016.
- Sesartic, A., Lohmann, U., and Storelvmo, T.: Bacteria in the ECHAM5-HAM global climate model, *Atmos. Chem. Phys.*, 12, 8645–8661, doi:10.5194/acp-12-8645-2012, 2012.
- Si, M., Irish, V. E., Mason, R. H., Vergara-Temprado, J., Hanna, S. J., Ladino, L. A., Yakobi-Hancock, J. D., Schiller, C. L., Wentzell, J. J., Abbatt, J. P., et al.: Ice-nucleating ability of aerosol particles and possible sources at three coastal marine sites, *Atmospheric Chemistry and Physics*, 18, 15 669–15 685, doi:10.5194/

acp-18-15669-2018, 2018.

- Si, M., Evoy, E., Yun, J., Xi, Y., Hanna, S. J., Chivulescu, A., Rawlings, K., Veber, D., Platt, A., Kunkel, D., et al.: Concentrations, composition, and sources of ice-nucleating particles in the Canadian High Arctic during spring 2016, *Atmospheric Chemistry and Physics*, 19, 3007–3024, doi:10.5194/acp-19-3007-2019, 2019.
- Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.
- 765 Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, *Nature*, 461, 607, doi:10.1038/nature08281, 2009.
- Stocker, T. F., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.: *Climate change 2013: the physical science basis. Intergovernmental panel on climate change, working group I contribution to the IPCC fifth assessment report (AR5)*, New York, 2013.
- 770 Stopelli, E., Conen, F., Guilbaud, C., Zopfi, J., Alewell, C., and Morris, C. E.: Ice nucleators, bacterial cells and *Pseudomonas syringae* in precipitation at Jungfraujoch, *Biogeosciences*, 14, 1189–1196, doi:10.5194/bg-14-1189-2017, 2017.
- Twohy, C. H., McMeeking, G. R., DeMott, P. J., McCluskey, C. S., Hill, T. C., Burrows, S. M., Kulkarni, G. R., Tanarhte, M., Kafle, D. N., and Toohey, D. W.: Abundance of fluorescent biological aerosol particles at temperatures conducive to the formation of mixed-phase and cirrus clouds, *Atmos. Chem. Phys.*, 16, 8205–8225, doi:10.5194/acp-16-8205-2016, 2016.
- 775 Vali, G., DeMott, P., Möhler, O., and Whale, T.: A proposal for ice nucleation terminology, *Atmos. Chem. Phys.*, 15, 10263–10270, doi:10.5194/acp-15-10263-2015, 2015.
- Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., DeMott, P. J., et al.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmos. Chem. Phys.*, 17, 3637, doi:10.5194/acp-17-3637-2017, 2017.
- 780 Videla, H. A., Guimet, P. S., and de Saravia, S. G.: Biodeterioration of Mayan archaeological sites in the Yucatan Peninsula, Mexico, *Int. Biodeter. Biodegr.*, 46, 335–341, doi:10.1016/S0964-8305(00)00106-2, 2000.
- 785 Welti, A., Müller, K., Fleming, Z. L., and Stratmann, F.: Concentration and variability of ice nuclei in the subtropic, maritime boundary layer, *Atmos. Chem. Phys.*, 18, 5307–5320, doi:10.5194/acp-18-5307-2018, 2018.
- Wex, H., Augustin-Bauditz, S., Boose, Y., Budke, C., Curtius, J., Diehl, K., Dreyer, A., Frank, F., Hartmann, S., Hiranuma, N., et al.: Intercomparing different devices for the investigation of ice nucleating particles using Snomax® as test substance, *Atmos. Chem. Phys.*, 15, 1463–1485, doi:10.5194/acp-15-1463-2015, 2015.
- 790 Whigham, D. F., Olmsted, I., Cano, E. C., and Harmon, M. E.: The impact of Hurricane Gilbert on trees, litterfall, and woody debris in a dry tropical forest in the northeastern Yucatan Peninsula, *Biotropica*, pp. 434–441, doi:10.2307/2388263, 1991.
- Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Burrows, S. M., Carslaw, K. S., Huffman, J. A., Judd, C., Kilhau, W. P., et al.: A marine biogenic source of atmospheric ice-nucleating particles, *Nature*, 525, 234, doi:10.1038/nature14986, 2015.
- 795

Winkler, P.: Relative humidity and the adhesion of atmospheric particles to the plates of impactors, *J. Aerosol Sci.*, 5, 235–240, doi:10.1016/0021-8502(74)90058-5, 1974.

800 Yakobi-Hancock, J. D., Ladino, L. A., and Abbatt, J. P.: Review of Recent Developments and Shortcomings in the Characterization of Potential Atmospheric Ice Nuclei: Focus on the Tropics, *Revista de Ciencias*, 17, 15–34, doi:10.25100/rc.v17i3.476, 2014.

Yankofsky, S., Levin, Z., Bertold, T., and Sandlerman, N.: Some basic characteristics of bacterial freezing nuclei, *J. Appl. Meteorol.*, 20, 1013–1019, doi:10.1175/1520-0450(1981)020<1013:SBCOBF>2.0.CO;2, 1981.

805 Yun, Y. and Penner, J. E.: An evaluation of the potential radiative forcing and climatic impact of marine organic aerosols as heterogeneous ice nuclei, *Geophys. Res. Lett.*, 40, 4121–4126, doi:10.1002/grl.50794, 2013.




Fig. 1. Map showing the sampling location. The red star shows the location of the Engineering Institute building where the sampling took place (GoogleMaps).




Fig. 2. Time evolution of wind and aerosol particle concentration time series for the entire campaign (21 January - 02 February, 2017) . A) Time series of the wind speed (yellow) and wind direction (green), B) Particle concentration measured by the CPC, C) Particle concentration measured by the LasAir full size range (0.3 μm to 25 μm), and D) Particle concentration measured by the LasAir for particles >500 nm (0.5 μm to 25 μm). Grey areas denote the periods affected by cold fronts (A and B). Each tick mark on the x-axis corresponds to midnight local time.




Fig. 3. Time series of the ambient aerosol mass concentration and bulk chemical composition as measured by the XRF. Each sample was collected for 48 h starting at 12:00 h local time.




Fig. 4. Summary of INP concentrations as a function of temperature and particle size (solid symbols). Total [INP] are represented by the grey triangles, whereas the brown asterisks, light blue dotted lines, and purple stars are data from DeMott et al. (2016), Welti et al. (2018), and Irish et al. (2019), respectively. The upper and lower detection limits of the MOUDI-DFT are 30 L^{-1} and 0.01 L^{-1} , respectively.

Table 1. List of the measured variables and the corresponding instrumentation.

Measured Variable	Instrument
INP concentration	MOUDI-DFT (Mason et al., 2015a)
Aerosol concentration	Condensation particle counter (CPC, TSI 3010)
Coarse aerosol size distribution	LasAir Optical particle counter (MSP)
Chemical composition	X-Ray fluorescence and High-performance liquid chromatography
Bacterial and fungal concentration	Biostage impactor (SKC)
Meteorology	Weather station (Davis)

Table 2. Correlation coefficients (r^2) of the average chemical composition and the average [INP] per sample at $-15 \text{ }^\circ\text{C}$, $-20 \text{ }^\circ\text{C}$, $-25 \text{ }^\circ\text{C}$, and $-30 \text{ }^\circ\text{C}$. Bold text highlights the r^2 with $p < 0.05$ (Table S1) for each temperature.

Temperature	Na	Mg	Al	Si	P	S	Cl	K	Ca	Ti	Mn	Fe	Cu	Zn
$-15 \text{ }^\circ\text{C}$	0.02	0.70	0.08	0.15	0.01	0.02	0.07	0.02	0.08	0.21	0.61	0.31	0.03	0.25
$-20 \text{ }^\circ\text{C}$	0.18	0.02	0.44	0.86	0.24	0.33	0.27	0.77	0.64	0.19	0.47	0.74	0.35	0.26
$-25 \text{ }^\circ\text{C}$	0.54	0.00	0.33	0.45	0.27	0.45	0.02	0.40	0.89	0.06	0.40	0.49	0.17	0.07
$-30 \text{ }^\circ\text{C}$	0.53	0.00	0.65	0.74	0.46	0.56	0.00	0.65	0.79	0.02	0.34	0.81	0.06	0.03




Fig. 5. Mean INP number concentrations at droplet freezing temperatures of -15 °C (light gray), -20 °C (dark gray), and -25 °C (black). The blue and red stars represent the mean INP concentration during the cold fronts A+B and cold front B, respectively. Uncertainties are given as the standard uncertainty of the mean (adapted from Mason et al. (2016)).




Fig. 6. Mean INP concentration as a function of aerosol particle size at A) -15 °C, B) -20 °C, C) -25 °C, and D) -30 °C. Uncertainties are given as the standard uncertainty of the mean.

Table 3. Bacterial isolation for top) Jan 21-22, middle) Cold front A, bottom) Cold front B. ^a, Isolated on TSA media; ^b, Isolated on GYM media.

Phylum	Genus/species	Source
Actinobacteria	^a <i>Kocuria palustris</i>	Soil, rhizoplane
	^a <i>Micrococcus</i> spp	Water, soil, dust, and skin
Firmicutes	^a <i>Rhodococcus corynebacteroides</i>	Soil, water and eukaryotic cells
	^a <i>Staphylococcus kloosii</i>	Human and animal skin
	^a <i>Staphylococcus lugdunensis</i>	Human and animal skin
	^a <i>Staphylococcus nepalensis</i>	Mucocutaneous zones of humans and animals
	^a <i>Staphylococcus arlettae</i>	Animal skin, mucosal zones, polluted water
	^a <i>Staphylococcus epidermidis</i>	Human skin, mucosal microbiota
	^a <i>Bacillus aryabhattai</i>	Upper atmosphere, rhizosphere
	^a <i>Bacillus gibsonii</i>	Alkaline soil
	^a <i>Bacillus aeris</i>	Soil
	^a <i>Staphylococcus lentus</i>	Soil
Alphaproteobacteria	^a <i>Sphingomonas mucosissima</i>	Water and soil
Actinobacteria	^a <i>Micrococcus</i> spp	Water, soil, dust, and skin
Firmicutes	^a <i>Bacillus oceanisediminis</i>	Marine sediments
Gammaproteobacteria	^a <i>Proteus mirabilis</i>	Water and soil
	^a <i>Pseudomonas stutzeri</i>	Soil
Actinobacteria	^{a,b} <i>Micrococcus</i> spp	Water, soil, dust, and skin
	^a <i>Micrococcus lentus</i>	Soil, dust, water and air
Firmicutes	^b <i>Micrococcus yunnanensis</i>	Roots of <i>Polyspora axillaris</i>
	^b <i>Streptomyces</i> spp	Cosmopolitan
	^a <i>Bacillus</i> spp	Cosmopolitan
	^a <i>Bacillus niacini</i>	Soil
	^a <i>Bacillus subtilis</i>	Soil, gut commensal in ruminants and humans
	^a <i>Planomicrobium koreense</i>	Fermented seafood
	^a <i>Staphylococcus</i> spp	Human and animal skin, mucous zones, soils
	^b <i>Solibacillus isronensis</i>	Air
	^a <i>Staphylococcus equorum</i>	Human and animal skin
	Gammaproteobacteria	^a <i>Pseudomonas reactants</i>
^a <i>Vibrio alginolyticus</i>		Marine
^a <i>Vibrio natriegens</i>		Marine
^a <i>Vibrio neocaledonicus</i>		Marine
^a <i>Vibrio parahaemolyticus</i>		Marine
^a <i>Zobellella</i> sp.		Marine and estuarine environments




Fig. 7. Fraction of aerosol particles acting as an INP ($[INP] / N_{Tot}$) as a function of particle size at $-15^{\circ}C$, $-20^{\circ}C$, and $-25^{\circ}C$, respectively. N_{Tot} refers to the number of aerosol particles in a given size range measured by the LasAir. The solid colored symbols represent the entire campaign, while the black symbols represent the samples collected under the influence of the cold fronts.

Table 4. Fungal identification on EMA media for the whole sampling period

Phylum	Genus	Source
Dothideomycetes	<i>Alternaria</i>	Dead plants, soil, foods, air, indoor environments, decaying organic matter, indoor bioaerosols, on animal systems and in freshwater and marine habitats.
	<i>Cladoporium</i>	
	<i>Drechslera</i>	
Euascomycetes	<i>Curvularia</i>	Dead plants, soil, foods, air, indoor environments, decaying organic matter, indoor bioaerosols, on animal systems and in freshwater and marine habitats.
Eurotiomycetes	<i>Aspergillus</i>	
	<i>Penicillium</i>	
Leotiomycetes	<i>Monilia</i>	Dead plants, soil, foods, air, indoor environments, decaying organic matter, indoor bioaerosols, on animal systems and in freshwater and marine habitats.
Sordariomycetes	<i>Fusarium</i>	
Zygomycetes	<i>Rhizopus</i>	




Fig. 8. A) Mean mass concentration of 14 detected elements for the collected aerosol particles using XRF, B) Mean mass concentration of 8 detected ions for the collected aerosol particles using HPLC, C) and D) Mean mass size distribution of the main five detected elements/ions with the XRF and HPLC, respectively. This results are the average for the whole sampling period.




Fig. 9. A) Time series of the [INP] at -15 °C (blue), -20 °C (brown), and -25 °C (yellow), B) Time series of the [INP] at -15 °C (blue) together with bacteria concentration (red), and C) Time series of the [INP] at -15 °C (blue) together with fungal concentration (black). Each X-axis tick corresponds to 06:00 local time. The horizontal uncertainty bars indicate the time span the MOUDI-DFT measurements i.e., 6 h. Grey areas denote the periods affected by the entrance of a cold front (A and B).