

# **Responses to Referees' Comments**

## **Effects of Arctic stratospheric ozone changes on spring precipitation in the northwestern United States (acp-2018-414)**

Xuan Ma, Fei Xie, Jianping Li, Xinlong Zheng, Wenshou Tian,  
Ruiqiang Ding, Cheng Sun, and Jiankai Zhang

December 2018

## Response to the Editor

*1. Based on the detailed reviewer comments, the authors have expanded and very much improved their revised manuscript. I am very grateful to the reviewers for their advice. However, while referee #2 concludes that all previous comments have been sufficiently addressed and recommends publication of the manuscript with only a few minor modifications, referee #1 is not convinced and suggests rejection of the manuscript. The main concern of referee #1 is that the model responses over the northwestern United States due to stratospheric ozone forcing are not statistically significant and the response due to ozone changes alone is of a different sign as the observed correlation. I agree with referee #1 on this point. However, I do feel that this study contains valuable results that justify publication in Atmos. Chem. Phys.*

*In order to be acceptable for publication, I ask the authors to take the additional comments of the referees into account. In particular also to arrive at a more balanced and appropriate discussion of the role of ASO vs. SST variations in forcing of northwestern US precipitation anomalies. The revised discussion should in particular reflect the concerns raised by referee #1. As suggested by referee #2, a more explicit link to the coupled model study of Xie et al. (2017) would help to improve the discussion of ASO vs. SST forcing.*

**Response: We are very grateful to the editor for the positive comments on our work and for giving us another chance to address the problems that the reviewers felt were still present.**

**The figure showing the key model result in this study has been redrawn. Actually, zonal wind anomalies at 200 hPa and 500 hPa are significant at the 95% confidence level over the western United States. In terms of the physical connection, the significant U anomalies at 200 hPa and 500 hPa over the western United States are likely to cause significant changes in precipitation in the western United States.**

**To further confirm the possible influence of ASO on precipitation in the northwestern United States, a transient experiment (1955–2005) based on the atmosphere–ocean coupled WACCM4 model (this links to the study of Xie et al. 2017) has been added in the revised manuscript to answer the question of whether the ASO by itself can cause the Victoria SST mode in the North Pacific**

and rainfall anomalies in the northwestern United States. The answer is yes.

The leading relationship between ASO and precipitation in the northwestern United States is evident in observations, time-slice experiments, and the transient experiment with specified ozone. All these conclusions show that our results are reasonable and credible.

More details are given in the Responses to reviewers 1 and 2.

*2. The statement that dynamical coupling affects primarily the Atlantic sector while ozone changes affect primarily the Pacific sector does not seem to be well supported by other studies. I suggest to de-emphasize this point unless there is specific and robust evidence.*

**Response:** Thanks for the comment.

In the Response to reviewer 1, we show a recent result from Ivy et al. (2017), who investigated the effect of ASO on Northern Hemisphere surface climate using only observations. A very strong and significant ASO-related signal can be found in the North Pacific sector in their Figure 3a and b. It agrees well with our result that ASO can influence the North Pacific sector. The papers (Smith et al. (2014) and Calvo et al. (2015)) pointed out by reviewer 1 only use model simulations. Our result is based on both observations and simulations.

In addition, we show that the ASO-related signal in the North Atlantic sector occurs in March; the ASO-related signal in the North Pacific sector occurs in April. In this study, we focus on the influence of ASO on circulation anomalies in April. This is why our result is different from those of Smith et al. (2014) and Calvo et al. (2015), who focus on the March–April average. If we use the March–April circulation, we also find significant ASO-related anomalies in the North Atlantic.

The statement in the original manuscript is indeed not precise. In the revised manuscript, this statement has been amended. However, our results are well supported by other studies. We have given a very detailed answer to this question. Please refer to the Response to Reviewer 1.

*3. Please take into account also the additional minor comments of both referees. In addition, a reference to the HadISST data set should be included. Why is in one case (Fig. 8) HadISST used and in another case (Fig. 9) CESM SST forcing data?*

**Response:** Thanks for the comment. The reference (Rayner et al. 2003) has been cited in the revised manuscript. The CESM SST forcing data are based on HadISST data but they have different resolution. The SST anomalies in Figure 9 are used to force the simulation. Thus, CESM SST forcing data are used.

**Reference:**

Rayner, N. A., D. E. Parker, E. B. Horton, C. K. Folland, L. V. Alexander, and D. P. Rowell, 2003: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *J. Geophys. Res.*, 108, 4407.

## Response to Referee 1

*I thank the authors for their consideration of the reviewers' comments. They have largely improved many of the cosmetic issues with the original manuscript, but many of the major scientific flaws of the original manuscript remain. Furthermore, the statistical significance contours added to the model results show that the key model results of this study that are foundation for the authors' conclusions in the abstract are not statistically significant. So, unfortunately, I cannot recommend publication of this manuscript for the following reasons:*

**Response:** Once again, we thank the reviewer for taking the time to assess the manuscript, highlight important issues, and provide helpful comments and suggestions to improve the manuscript. Although the reviewer suggested rejecting the manuscript, we have given new and detailed responses to the reviewer's new comments to address the problems that the reviewer felt were still present in the study. We hope to get another chance and that the reviewer can judge our work again.

Figure 10, the key model result in this study, has been redrawn (shown below as Figure R1) to show the differences between the R5 experiment (forced by ASO+SST anomalies) and R4 experiment in terms of April precipitation (mm/day) and zonal wind at 200, 500, and 850 hPa. To make the significant region clearer, this map focuses on North America. The black (white) dots denote significance at the 90% (95%) confidence level in this figure. Actually, the zonal wind anomalies at 200 hPa and 500 hPa are significant at the 95% confidence level over the western United States (Figure R1b and c) and all the anomalies (including U and precipitation anomalies) are significant at the 90% confidence level in the western United States. In terms of physical connection, the significant U anomalies at 200 hPa and 500 hPa (Figure R1b and c, 95% confidence level) in the western United States are likely to cause significant changes in precipitation in the northwestern United States.

In order to further confirm the possible influence of ASO on precipitation in the northwestern United States, a transient experiment R8 (1955–2005) based on the atmosphere–ocean coupled WACCM4 model has been added in the revised manuscript to confirm that the ASO by itself can cause the Victoria SST mode in

the North Pacific and rainfall anomalies in the northwestern United States. Note that the ozone forcing in the experiment is specified; it is derived from the CMIP5 ensemble mean ozone output. Please refer to R8 in Table 1 in the revised manuscript for a detailed description of the experiment. Figure R2 shows the correlation coefficients between the specified March ASO variations and simulated April 500 hPa U, SST, and precipitation anomalies for the period 1955–2005. The significant and leading effects of the specified ASO anomalies on 500 hPa U, the Victoria mode in the North Pacific, and rainfall anomalies in the northwestern United States are well captured (Fig. R2). As the ozone forcing in the experiment is specified, the relationships between ASO and U and SST and precipitation could only be caused by ASO influencing U, and then U influencing SST and precipitation; the ASO changes are completely independent of the polar vortex.

The leading relationship between ASO and precipitation in the northwestern United States can be found in observations, time-slice experiments (R1–7), and a transient experiment with specified ozone (R8). Table 1 in the revised manuscript gives details of experiments R1–8. All these results strengthen the main conclusion of the article.

Figure R2 and relevant content have been added in the revised manuscript. Please see Lines 336–355 and Figure 12.





Figure R1. Differences between experiments R5 and R4 in terms of April (a) precipitation

(mm/day) and (b–d) zonal wind at 200, 500, and 850 hPa, respectively. The black (white) dots denote significance at the 90% (95%) confidence level. A detailed description of the experiments may be found in Table 1 in the revised manuscript.



**Figure R2.** Correlation coefficients between the specified March ASO variations and simulated anomalies of April U (a), SST (b), and precipitation (c) for the period 1955–2005 based on the transient experiment R8. Regions above the 95% confidence level are dotted. The seasonal cycle and linear trend were removed from all quantities before correlation.

1. *Figure 10, which presents the authors' central conclusion (both Arctic ozone and SST forcings are necessary to reproduce the observed signals), does not actually show statistically significant precipitation and circulation signatures over the Pacific Northwest. In my view, this casts some serious doubt on the robustness of the observed relationship documented in this study, if models cannot faithfully reproduce the observed relationship. The most statistically significant features in the model results are in Fig. 7, which are driven by ozone anomalies only and are of the*

*opposite sign to those in the observations.*

**Response: Thanks for the comment. Please refer to above Response.**

2. *The authors are incorrect in stating that midwinter stratospheric anomalies couple down to the troposphere and impact the North Atlantic sector, but that springtime stratospheric anomalies instead impact the North Pacific sector (lines 67–71). Smith et al. (2014) and Calvo et al. (2015) have shown that Arctic stratospheric ozone anomalies primarily affect the Atlantic sector during spring (see Fig. 4 of Smith et al. 2014; see Figs. 3–4 of Calvo et al. 2015). As I stated in my original review, the authors need to provide a convincing explanation for why their patterns are focused in the North Pacific sector, instead of the North Atlantic sector (as would be anticipated from many previous studies).*

**Response: Thanks for the question. The statement in Lines 67–71 in the original manuscript is not precise. It has been modified in the revised manuscript.**

**Figure 4 of Smith et al. (2014) indeed showed that the ASO significantly influences only the North Atlantic sector. However, they obtained their results from a model that does not contain complete stratospheric processes—the CAM3 model. This issue is also pointed out by Calvo et al. (2015), who repeated the study using the WACCM4 model. In figure 3 of Calvo et al. (2015), the influence of ASO is significant in both the North Atlantic sector and the North Pacific sector (Figure R3 below). Note that the results from Smith et al. (2014) and Calvo et al. (2015) are based entirely on simulations. Recently, Ivy et al. (2017) investigated the effect of ASO on Northern Hemisphere surface climate using observations. A very strong and significant ASO-related signal can be found in the North Pacific sector in Figure 3a and b of Ivy et al. (2017) (shown as Figure R4 below). It agrees well with our result that ASO can influence the North Pacific sector.**

**Figure R5a, c, and e shows the relationship between March ASO and March 500 hPa U anomalies based on composite, correlation and regression analyses, while Figure R5b, d and f shows the corresponding relationship between March ASO and April 500 hPa U anomalies. Strong and significant ASO-related signals can be found in the North Pacific and North Atlantic sectors in March. However, the strong and significant ASO-related signal can only be found in the North Pacific sector in April. Ivy et al. (2017) investigated the effect of ASO on March–**

April average circulation. This is why they obtained significant ASO-related signals in both the North Pacific and North Atlantic. In Figure R6, we show the relationship between March ASO and March–April 500 hPa U anomalies. Significant ASO-related signals are also found in both the North Pacific and North Atlantic.

The above analysis at least suggests that ASO may affect the circulation in both the North Pacific and North Atlantic sectors. However, this study focuses on the impact of March ASO on April precipitation in the northwestern United States; ASO is more likely to influence precipitation in the northwestern United States by influencing circulation over the North Pacific Ocean. Therefore, in our study, we mainly analyze the influence of March ASO on the April circulation over the North Pacific Ocean rather than its effect on the circulation over the North Atlantic Ocean.

In the revised manuscript, this statement has been revised as follows (please see Lines 67–71):

*“Similar to the effects of winter stratospheric dynamical processes on the tropospheric North Atlantic Oscillation and the incidence of extreme weather events (Baldwin and Dunkerton, 2001; Black et al., 2005, 2006, 2009), the depletion of spring ASO can cause circulation anomalies that influence the tropospheric North Atlantic and North Pacific sectors.”*






Figure R3. Figure 3 of Calvo et al. (2015). The composite differences of zonal mean zonal wind ( $\text{m s}^{-1}$ ) at 500 hPa between the April and May averages of the LOW and HIGH ozone years during 1985–2005. Stippling indicates significant differences at the 95% confidence level.



**Figure R4.** Figure 3 of Ivy et al. (2017). Differences (colors) in March–April zonal wind at 500 hPa ( $\text{m s}^{-1}$ ) and sea level pressure (hPa) from MERRA between years with low March ozone and years with high March ozone. The contour lines show the climatological mean winds in the top panel; contours are every  $5 \text{ m s}^{-1}$  ( . . . ,  $-10$ ,  $-5$ ,  $5$ ,  $10$ , . . . ). Hatching denotes differences that are statistically significant at the 95% level.



**Figure R5.** (a) Differences in composite March 500 hPa U wind anomalies between positive and negative ASO anomaly events. The ASO anomaly events are selected based on Table 2 in

the manuscript. (c) Correlation coefficients between March ASO index and March 500 hPa U wind variations. (e) Regression coefficients of March U wind variations regressed onto the March ASO. ASO is taken from SWOOSH and U winds from NCEP2 for 1984–2016. Before performing the analysis, the seasonal cycle and linear trend were removed from the original dataset. Dots denote significance at the 95% confidence level, according to Student's *t*-test. (b, d and f), Same as (a, c and e), but for March ASO and April 500 hPa U anomalies.




Figure R6. Same as Figure R5a, c and e, but between March ASO and March–April 500 hPa U anomalies.

**Reference:**

- Calvo, N., Polvani, L. M., and Solomon, S.: On the surface impact of Arctic stratospheric ozone extremes, *Environ. Res. Lett.*, **10**, 094003, doi:10.1088/1748-9326/10/9/094003, 2015.
- Ivy, D. J., Solomon, S., Calvo, N., and Thompson, D. W.: Observed connections of Arctic stratospheric ozone extremes to Northern Hemisphere surface climate, *Environ. Res. Lett.*, **12**, 024004, doi:10.1088/1748-9326/aa57a4, 2017.
- Smith, K. L. and Polvani, L. M.: The surface impacts of Arctic stratospheric ozone anomalies, *Environ. Res. Lett.*, **9**, 074015, doi:10.1088/1748-9326/9/7/074015, 2014.

*More minor concerns:*

3. *The authors' argument that Arctic stratospheric ozone anomalies are more important than stratospheric polar vortex variability for Pacific Northwest precipitation is very misleading in my view. They have added Fig. 12 to address this issue, but they compare February vortex variability with April precipitation, but March ozone with April precipitation. For this to be a true comparison, they need to compare March vortex variability with April precipitation (or February ozone with April precipitation).*


**Response:** Thanks for the comment. Figure 12 (it is Fig. 13 in the revised manuscript) in the original manuscript was included to answer a question from reviewer 2, who wanted to check whether the stratospheric polar vortex variability in late winter might be a more important factor in leading spring precipitation variations in the northwestern United States than the March ASO. Figure 12 shows that the variations of the spring precipitation in the northwestern United States are more closely related to the ASO than to the late winter polar vortex.

It is indeed misleading to argue that Arctic stratospheric ozone anomalies are more important than stratospheric polar vortex variability for influencing precipitation over the northwestern United States. We have revised the description of Figure 12 to express the problem more clearly. Please see Lines 396–408 in the revised manuscript.

The variations of the March polar vortex are the intermediate step linking March ASO anomalies and April precipitation changes in the northwestern United States. Therefore, the correlation between the March polar vortex and April precipitation variations is strong.

Here, we divide the variations of the March polar vortex strength (SPV) (Zhang et al. 2018) into two parts: one related to ASO ( $SPV_{ASO}$ ) and one unrelated to ASO ( $SPV_{NO\_ASO}$ );  $SPV = SPV_{ASO} + SPV_{NO\_ASO}$ . Figure R7 shows that the correlation coefficients between March  $SPV_{ASO}$  and April precipitation variations are significant in the northwestern United States, but are not significant between March  $SPV_{NO\_ASO}$  and April precipitation variations. Figure R7 illustrates that, as far as the relationship between March polar vortex and April precipitation variations is concerned, only the March polar vortex variations related to ASO are closely linked to precipitation variations in the

northwest United States.



**Figure R7.** (a) Correlation coefficients between the March  $-SPV_{ASO}$  (the strength index of SPV is defined by Zhang et al. 2018) and April precipitation variations. (b) As for (a), but between March  $-SPV_{NO\_ASO}$  and April precipitation variations.  $SPV_{ASO}$  and  $SPV_{NO\_ASO}$  are calculated by regressing the March SPV index onto the March ASO index and removing the regressed values from the original SPV. Dots denote significance at the 95% confidence level, according to Student's  $t$ -test. The long-term linear trend and seasonal cycle in all variables were removed before the correlation analysis. The ASO data are from SWOOSH, and precipitation from GPCP.

4. The SST anomalies associated with Pacific Northwest precipitation variations and ASO variations closely resemble those associated with ENSO and the PDO (see Fig. R5 in authors' responses). While the authors have presented some correlation analyses in their responses, I'm not entirely convinced that the observed signals in this manuscript aren't being aliased by concurrent ENSO and PDO variability. It would be nice to show that the observed signals still exist if only neutral ENSO years are sampled, as very large El Niño and La Niña events occur within their composites (Table 2).

**Response:** Thanks for the comment. It is a good idea to check whether the

precipitation anomalies in the northwest United States are related to ENSO. Figure R8 shows the differences in composite April precipitation anomalies between positive and negative ASO anomaly events during neutral ENSO years. The precipitation anomalies in the northwestern United States agree well with the previous result. Removing the effect of ENSO events completely does not affect the results. Please note that these anomalies are not significant. However, this should be due to the small number of cases.




Figure R8. Differences in composite April precipitation anomalies between positive and negative ASO anomaly events during ENSO neutral years in 1984–2016. The seasonal cycle and linear trend were removed from the original dataset. The ASO anomaly events are selected based on Table R1.

Table R1. Selected ASO anomaly events during neutral ENSO years for 1984–2016.

Positive March ASO anomaly events	Negative March ASO anomaly events
2001, 2004, 2010	1995, 1996

## Response to Referee 2

*Having reviewed the initial submitted version of the manuscript, I will not provide a summary of the manuscript here. My only significant concern with the revised version is that the authors have not completely shown that the circulation and precipitation anomalies are caused by ASO anomalies alone. The new Figure 12 comparing correlations with SPV and ASO, if anything, strengthens the idea that anomalies in ozone and anomalies in the stratospheric circulation reinforce each other. I am not sure this point really comes through in the tone of the discussion. But at some point it becomes an argument about semantics and I think the reader can decide for themselves.*

*I think the authors have sufficiently addressed the concerns raised by the reviewers on the first version here. I might add that the work presented here is much more easily accepted and put into context when combined with the coupled model simulations presented in Xie et al. (2017). The authors may wish to develop a more direct link with the Xie et al. work and stress that the coupled model simulations presented in Xie et al. (2017) were able to produce SST anomalies that were somewhat like the Victoria mode anomalies from the observed SSTs used here. This point of what causes the SST anomalies, which are imposed in the experiments here, is an integral part of the argument about the role of ASO anomalies and should be strengthened here.*


**Response:** Once again, we thank the reviewer for taking the time to assess the manuscript and we sincerely appreciate the reviewer's new comments. We have revised the manuscript carefully according to the reviewer's comments and suggestions.

Following the comment, a transient experiment (1955–2005) based on the atmosphere–ocean coupled WACCM4 model, which is shown in Xie et al. (2017), has been added in the revised manuscript to confirm that the ASO by itself can cause the Victoria SST mode in the North Pacific and rainfall anomalies in the northwestern United States. Note that the ozone forcing in the experiment is specified; it is derived from the CMIP5 ensemble mean ozone output. Please refer to R8 in Table 1 in the revised manuscript for a detailed description of the experiment. Figure RR1 shows the correlation coefficients between the specified

**March ASO variations and simulated April 500 hPa U, SST, and precipitation anomalies for the period 1955–2005. The significant and leading effects of the specified ASO anomalies on 500 hPa U, the Victoria mode in the North Pacific, and rainfall anomalies in the northwestern United States are well captured (Fig. RR1). As the ozone forcing in the experiment is specified, the relationships between ASO and U and SST and precipitation could only be caused by ASO influencing U, and then U influencing SST and precipitation; the ASO changes are completely independent of the polar vortex.**

**The leading relationship between ASO and precipitation in the northwestern United States can be found in observations, time-slice experiments (R1–7), and a transient experiment with specified ozone (R8). Table 1 in the revised manuscript gives details of experiments R1–8. All these results strengthen the main conclusion of the article.**

**Figure RR1 and relevant content have been added in the revised manuscript. Please see Lines 336–355 and Figure 12.**



**Figure RR1.** Correlation coefficients between the specified March ASO variations and simulated anomalies of April U (a), SST (b), and precipitation (c) for the period 1955–2005 based on the transient experiment R8. Regions above the 95% confidence level are dotted. The seasonal cycle and linear trend were removed from all quantities before correlation.

*Minor comments:*

*Line 74-75: I think ‘Comparing with the effect of the winter stratospheric dynamical processes...’ should perhaps be ‘Similar to the effects of winter stratospheric dynamical processes...’*

**Response:** Thanks for the suggestion. The sentence has been revised according to this comment. Please see Line 67 in the revised paper.

*Lines 154-156 are difficult to understand, particularly because the WACCM4-GHG scheme has not been defined.*

**Response: Thanks for the comment. The following sentence has been added in the revised manuscript. See Lines 147–149 in the revised manuscript.**

***“WACCM4-GHG: The chemistry is specified in this scheme; i.e., the volume mixing ratios of forcings of O<sub>3</sub>, CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, CFC11, CFC12 and so on are prescribed.”***

*Lines 353 – 355 ‘Second, the ASO changes generate SST anomalies over the North Pacific...’ should really have a reference to Xie et al. (2017).*

**Response: Thanks. The reference has been added.**

*Lines 402 – 405 ‘Such circulation anomalies force an anomalous cyclone in the western United States in the middle and upper troposphere, which likely enhances cold and dry airflow from the North American continent to the North Pacific...’ At least in the climatological monthly average the winds over the northwestern United States continue to be from the west, bringing moisture in from the Pacific. The anomalous cyclone is not a reversal of the winds, but a reduction in the winds. I do not believe you have shown any westerly transport of dry air from the North American continent to the North Pacific – this would require winds from the east. Not just a reduction in westerly winds. I believe a much more justifiable interpretation of the changes is a reduction in the transport of moisture inland from the Pacific.*


**Response: Thanks very much for the suggestion. This sentence and the relevant content have been modified in the revised suggestion as follows (See Lines 229–234 and Lines 368–373):**

***“The climatological wind over the northwestern United States blows from west to east, bringing moisture from the Pacific to the western United States. Such circulation anomalies force an anomalous cyclone in the western United States in the middle and upper troposphere, which reduces the climatological wind. It would decrease the water vapor concentration in the air over the northwestern United States.”***

*Figure 8 – Can the authors check that the revised Figure 8 actually shows*

significance at the 95% level. To my eye Figure 8 in the revised version looks identical to Figure 10 in the original submission where significance was plotted at the 90% level. The significance is denoted with dots and I am not able to find a single dot that has changed between the original Figure 10 and the revised Figure 8 – even in places where a single, isolated dot appears by itself.

**Response: Thanks for checking this. In Figure 10 of the original submission, the significance was plotted at the 95% confidence level. Only this figure used a 95% confidence level, but the other figures used a 90% confidence level in the original submission. We are very sorry for the misunderstanding caused by this inconsistency. Please see Figure RR2 below; this was Figure 10 of the original submission.**



690

691 **Figure 10.** (a) Correlation coefficients between regional precipitation (43°–50°N,  
692 115°–130°W) and SST variations in April for 1984–2016. (b) Correlation coefficients  
693 between March ASO ( $\times -1$ ) and April SST variations for 1984–2016. Dots denote  
694 significance **at the 95% confidence level**, according to Student's  $t$ -test. Before  
695 performing the analysis, the seasonal cycle and linear trend were removed from the  
696 original data. ASO data are from SWOOSH, precipitation from NCEP2, and SST  
697 from HadSST.

**Figure RR2. Figure 10 in the original submission.**

# Effects of Arctic stratospheric ozone changes on spring precipitation in the northwestern United States

Xuan Ma<sup>1</sup>, Fei Xie<sup>1\*</sup>, Jianping Li<sup>1,2</sup>, Xinlong Zheng<sup>1</sup>, Wenshou Tian<sup>3</sup>,  
Ruiqiang Ding<sup>4</sup>, Cheng Sun<sup>1</sup>, and Jiankai Zhang<sup>3</sup>

<sup>1</sup>*College of Global Change and Earth System Science, Beijing Normal University, Beijing, China*

<sup>2</sup>*Laboratory for Regional Oceanography and Numerical Modeling, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China*

<sup>3</sup>*College of Atmospheric Sciences, Lanzhou University, Lanzhou, China*

<sup>4</sup>*State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China*

Submitted as an Article to: *Atmospheric Chemistry and Physics*

4 December 2018

\* Corresponding author:

Dr. Fei Xie, Email: xiefei@bnu.edu.cn.

删除的内容: <sup>1</sup>State Key Laboratory of Earth Surface Processes and Resource Ecology and College

删除的内容: 13 September

带格式的: 字体颜色: 自动设置

## Abstract

Using observations and reanalysis, we find that changes in April precipitation variations in the northwestern US are strongly linked to March Arctic stratospheric ozone (ASO). An increase in ASO can result in enhanced westerlies in the high and low latitudes of the North Pacific but weakened westerlies in the mid-latitudes. The anomalous circulation over the North Pacific can extend eastward to western North America, ~~decreasing the water vapor concentration in the air over the northwestern United States~~ and enhancing downwelling in the northwestern US, which results in decreased precipitation there, and vice versa for the decrease in ASO. Model simulations using WACCM4 support the statistical analysis of observations and reanalysis data, and further reveal that the ASO influences circulation anomalies over the northwestern US in two ways. Stratospheric circulation anomalies caused by the ASO changes can propagate downward to the troposphere in the North Pacific and then eastward to influence the strength of the circulation anomalies over the northwestern US. In addition, sea surface temperature anomalies over the North Pacific, which may be related to the ASO changes, would cooperate with the ASO changes to modify the circulation anomalies over the northwestern US. Our results suggest that ASO variations could be a useful predictor of spring precipitation changes in the northwestern US.

删除的内容: facilitating

删除的内容: flow of a dry and cold airstream from the middle of North America to the North Pacific

带格式的: 字体颜色: 自动设置

## **1. Introduction**

Stratospheric circulation anomalies can affect tropospheric climate via chemical–radiative–dynamical feedback processes (Baldwin and Dunkerton, 2001; Graf and Walter, 2005; Cagnazzo and Manzini, 2009; Ineson and Scaife, 2009; Thompson et al., 2011; Reichler et al., 2012; Karpechko et al., 2014; Kidston et al., 2015; Li et al., 2016; Zhang et al., 2016; Wang et al., 2017). Since stratospheric ozone can influence stratospheric temperature and circulation via the atmospheric radiation balance (Tung, 1986; Haigh, 1994; Ramaswamy et al., 1996; Forster and Shine, 1997; Pawson and Naujokat, 1999; Solomon, 1999; Randel and Wu, 1999, 2007; Labitzke and Naujokat, 2000; Gabriel et al. 2007; Gillett et al. 2009; McCormack et al. 2011), the impact of ozone on tropospheric climate change has recently received widespread attention (e.g., Nowack et al. 2015, 2017, 2018).

In recent decades, Antarctic stratospheric ozone has decreased dramatically due to the increase in anthropogenic emissions of ozone depleting substances (Solomon, 1990, 1999; Ravishankara et al., 1994, 2009). Numerous studies have found that the decreased Antarctic ozone has contributed substantially to climate change in the Southern Hemisphere. The Southern Hemisphere circulation underwent a marked change during the late 20th century, with a slight poleward shift of the westerly jet (Thompson and Solomon, 2002; Archer and Caldeira, 2008). The poleward circulation shift would cause surface temperature anomalies by affecting localized wind patterns and associated thermal advection (Son et al., 2010; Thompson et al. 2011; Feldstein, 2011). Subsequent studies concluded that Antarctic ozone depletion is responsible for at least 50% of the circulation shift (Lu et al., 2009; Son et al., 2010; McLandress et al., 2011; Polvani et al., 2011; Hu et al., 2013; Gerber and Son, 2014; Waugh et al., 2015). In addition, the poleward displacement of the westerly jet has

been linked to an extension of the Hadley cell (Son et al., 2009, 2010; Min and Son, 2013) and variations in mid- to high-latitude precipitation during austral summer; i.e., increased rainfall in the subtropics and high latitudes and reduced rainfall in the mid-latitudes of the Southern Hemisphere (Son et al., 2009; Feldstein, 2011; Kang et al., 2011; Polvani et al., 2011). The changes in Antarctic ozone are not only related to the displacement of the westerly jet in the Southern Hemisphere, but also affect its intensity. Thompson and Solomon (2002) argued that Antarctic ozone depletion can also enhance westerly winds via the strong radiative cooling effect and thermal wind relationship. The westerly winds are enhanced from the stratosphere to the mid-latitude troposphere in the case of wave–mean flow interaction (Son et al., 2010; Thompson et al., 2011), thereby accelerating circumpolar currents in the mid-latitudes. Moreover, changes in subtropical drought, storm tracks and ocean circulation in the Southern Hemisphere are also closely related to Antarctic ozone variations (Yin, 2005; Russell et al., 2006; Son et al., 2009; Polvani et al., 2011; Bitz and Polvani, 2012).

The variations in Arctic stratospheric ozone (ASO) in the past five decades are quite different from those of Antarctic stratospheric ozone, as the multi-decadal loss of ASO is much smaller than that of Antarctic stratospheric ozone (WMO, 2011). However, sudden stratospheric warming in the Arctic (Randel, 1988; Charlton and Polvani, 2007; Manney et al., 2011; Manney and Lawrence, 2016) means that the year-to-year variability in ASO has an amplitude equal to or even larger than that of Antarctic stratospheric ozone. Thus, the effect of ASO on Northern Hemisphere climate change has also become a matter of concern.

~~Similar to the effects of winter stratospheric dynamical processes on the tropospheric North Atlantic Oscillation and the incidence of extreme weather events~~

带格式的: 字体颜色: 自动设置

删除的内容: Comparing with

删除的内容: effect

删除的内容: the

删除的内容: (NAO)

(Baldwin and Dunkerton, 2001; Black et al., 2005, 2006, 2009), the depletion of spring ASO can cause circulation anomalies that influence the tropospheric North Atlantic and North Pacific sectors. Cheung et al. (2014) used the UK Met Office operational weather forecasting system and Karpechko et al. (2014) used ECHAM5 simulations to investigate the relationship between extreme Arctic ozone anomalies in 2011 and tropospheric climate. Smith and Polvani (2014) used an atmospheric global climate model to reveal a significant influence of ASO changes on tropospheric circulation, surface temperature, and precipitation when the amplitudes of the forcing ASO anomaly in the model are larger than those historically observed. Subsequently, using a fully coupled chemistry–climate model, Calvo et al. (2015) again confirmed that changes in ASO can produce robust anomalies in Northern Hemisphere temperature, wind, and precipitation. Furthermore, the effects of ASO on the Northern Hemisphere climate can be seen in observations. Ivy et al. (2017) presented observational evidence for the relationship between ASO and tropospheric climate, revealing that the maximum daily surface temperature anomalies in spring (March–April) in some regions of the Northern Hemisphere occurred during years with low ASO in March. Xie et al. (2016, 2017a, 2017b) demonstrated that the tropical climate can also be affected by ASO. They pointed out that stratospheric circulation anomalies caused by March ASO changes can rapidly extend to the lower troposphere and then propagate horizontally to the North Pacific in about 1 month, influencing the North Pacific sea surface temperature (SST) in April. The induced SST anomalies (Victoria Mode) associated with the circulation anomalies can influence El Niño–Southern Oscillation (ENSO) and tropical rainfall over a timescale of ~20 months.

As shown above, a large number of observations and simulations have shown that ASO variations have a significant impact on Northern Hemisphere tropospheric

刪除的內容: Oscillation

climate, but few studies have focused on regional characteristics. Xie et al. (2018) found that the ASO variations could significantly influence rainfall in the central China, since the circulation anomalies over the North Pacific caused by ASO variations can extend westward to China. This motivates us to investigate whether the circulation anomalies extend eastward to affect the precipitation in North America. In this study, we find a strong link between ASO and precipitation in the northwestern US in spring. We focus on analyzing the characteristics of the impact of ASO on precipitation in the northwestern US in spring and the associated mechanisms. The remainder of this manuscript is organized as follows. Section 2 describes the data and numerical simulations, and section 3 discusses the relationship between the ASO anomalies and precipitation variations in the northwestern US, as well as the underlying mechanisms. The results of simulations are presented in section 4, and conclusions are given in section 5.

## **2. Data and simulations**

The ASO variations is defined as the Arctic stratospheric ozone averaged over the latitude of 60°–90°N at an altitude of 100–50 hPa after removing the seasonal cycle and trend. Ozone values used in the present analysis are derived from the Stratospheric Water and OzOne Satellite Homogenized (SWOOSH) dataset (Davis et al., 2016), which is a collection of stratospheric ozone and water vapor measurements obtained by multiple limb sounding and solar occultation satellites over the previous 30 years. Monthly mean ozone data from SWOOSH (1984–2016) is zonal-mean gridded dataset at a horizontal resolution of 2.5° (latitude: 89°S to 89°N) and vertical pressure range of 31 levels from 316 hPa to 1 hPa. Another set of ozone data is taken from Global Ozone Chemistry and Related trace gas Data Records for the Stratosphere (GOZCARDS, 1984–2013) project (Froidevaux et al., 2015) based on

high quality data from past missions (e.g., SAGE, HALOE data) and ongoing missions (ACE-FTS and Aura MLS). It is also a zonal-mean dataset with a meridional resolution of  $10^\circ$ , extending from the surface to 0.1 hPa (25 levels).

In addition, two sets of global precipitation reanalysis datasets are employed in this study: monthly mean precipitation data constructed by the Global Precipitation Climatology Project (GPCP), which is established by the World Climate Research program (WCRP) in 1986 aiming to observe and estimate the spatial and temporal global precipitation (Huffman et al., 1997), with a resolution of  $2.5^\circ$  latitude/longitude grid for the analysis period 1984–2016; global terrestrial rainfall dataset derived from the Global Precipitation Climatology Centre (GPCC) based on quality-controlled data from 67200 stations world-wide, with a resolution of  $1.0^\circ$  latitude/longitude grid. In addition, SST is taken from the UK Met Office Hadley Centre for Climate Prediction and Research SST (HadSST, [Rayner et al., 2003](#)). Other atmospheric datasets including monthly-mean wind and [vertical velocity](#) fields for the period 1984–2016 are obtained from the NCEP/Department of Energy (DOE) Reanalysis 2 (NCEP-2), regarded as an updated NCEP/NCAR Reanalysis Project (NCEP-1).

We use the Whole Atmosphere Community Climate Model version 4 (WACCM4), a part of the National Center for Atmospheric Research's Community Earth System Model (CESM), version 1.0.6, to investigate precipitation response in the northwestern United States to the ASO anomalies. WACCM4 encompasses the Community Atmospheric Model version 4 (CAM4) and as such includes all of its physical parameterizations (Neale et al., 2013). It uses a system made up of four components, namely atmosphere, ocean (specified SST), land, and sea ice (Holland et al., 2012) and has detailed middle-atmosphere chemistry. This improved version of

删除的内容: .

删除的内容: geopotential height

WACCM uses a finite-volume dynamical core, and it extends from the surface to approximately 145 km geometric altitude (66 levels), with a vertical resolution of about 1 km in the tropical tropopause layer and the lower stratosphere. Note that the simulations in the present paper are disable interactive chemistry as WACCM4-GHG scheme (Garcia et al., 2007) with a  $1.9^\circ \times 2.5^\circ$  horizontal resolution. WACCM4-GHG: The chemistry is specified in this scheme, i.e., the volume mixing ratios of forcings of O<sub>3</sub>, CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, CFC11, CFC12 and so on are prescribed. More information can be seen in Marsh et al. (2013). The model's radiation scheme uses these conditions: fixed greenhouse gas (GHG) values (averages of emissions scenario A2 of the Intergovernmental Panel on Climate Change (WMO, 2003) over the period 1995–2005). The prescribed ozone forcing used in the experiments is a 12-month seasonal cycle averaged over the period 1995–2005 from CMIP5 ensemble mean ozone output. The Quasi Biennial Oscillation (QBO) phase signals with a 28-month fixed cycle are included in WACCM4 as an external forcing for zonal wind.

Seven time-slice experiments (R1–R7) and a transient experiment with specified ozone (R8) are designed to investigate the precipitation changes in the northwestern US due to the ASO anomalies. Details of the eight experiments are given in Table 1. Seven time-slice experiments (R1–R7) are run for 33 years, with the first 3 years excluded for the model spin-up and only the last 30 years are used. The transient experiment (R8) is run for 51 years.

删除的内容: seven

删除的内容: All the

删除的内容: for analysis

### **3. Response of precipitation in the northwestern US to ASO anomalies in spring**

Since the variations in ASO are most obvious in March due to the Arctic polar vortex break down (Manney et al., 2011), previous studies have reported that the ASO

changes in March have the strongest influence on the Northern Hemisphere (Ivy et al., 2017; Xie et al., 2017a). In addition, these studies pointed out that the changes in ASO affect the tropospheric climate with a lead of about 1–2 months, which is similar to the troposphere response to the Northern Hemisphere sudden stratospheric warmings (Baldwin and Dunkerton 2001; Black et al., 2005, 2006, 2009) and Southern Hemisphere stratospheric ozone depletion (Thompson and Solomon 2002); the relevant mechanisms have been investigated in detail by Xie et al. (2017a). We therefore show in Fig. 1 the correlation coefficients between ASO variations in March from SWOOSH and GOZCARDS data, and precipitation anomalies in April from GPCP and GPCP data over western North America. In all cases in Fig. 1 the March ASO changes are significantly anti-correlated with April precipitation anomalies in the northwestern US (mainly in Washington and Oregon), implying that positive spring ASO anomalies are associated with less spring precipitation in the northwestern US, and vice versa for the negative spring ASO anomalies. Note that since this kind of feature appears in the northwestern US, Fig. 1 shows only the west side of North America.

The correlation coefficients between March ASO variations and precipitation anomalies (January to December are in the same year) in the northwestern US are shown in Fig. 2. The correlation coefficients between March ASO variations and April precipitation anomalies in the northwestern US are the largest and are significant at the 95% confidence level. Note that the correlation coefficients between March ASO variations and July precipitation anomalies are also significant. The impact of March ASO on precipitation in the northwestern US in summer and the associated mechanisms are different from those considered in this study (not shown) and will be presented in another paper, but will not be investigated further here. March ASO

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

changes are not significantly correlated with simultaneous (March) precipitation variations (Fig. 2), illustrating that the ASO changes lead precipitation anomalies by about 1 month. Since the results from four sets of observations show a common feature, and SWOOSH and GPCP data span a longer period, only SWOOSH ozone and GPCP precipitation are used in the following analysis.

The above statistical analysis shows a strong negative correlation between March ASO variations and April precipitation anomalies in the northwestern US, meaning that the ASO can be used to predict changes in spring precipitation in the northwestern US. The process and underlying mechanism that are responsible for the impact of ASO anomalies on precipitation changes need further analysis.

Figure 3 shows the correlation coefficients between March ASO anomalies and April zonal wind variations at 200, 500, and 850 hPa, respectively. The spatial distribution of significant correlation coefficients over the North Pacific exhibits a tripolar mode with a zonal distribution at 200 and 500 hPa; i.e. a positive correlation in the high and low latitudes in the North Pacific and a negative correlation in mid-latitudes. This implies that the increase in ASO can result in enhanced westerlies in the high and low latitudes of the North Pacific but weakened westerlies in the mid-latitudes, corresponding to the weakened Aleutian Low in April, and vice versa for the decrease in ASO. The Aleutian Low acts as a bridge connecting variations in ASO and circulation anomalies over the North Pacific (Xie et al., 2017a). At 850 hPa, the anomalous circulation signal in the low latitudes of the North Pacific has weakened and disappeared. It is evident that the anomalous changes in the zonal wind over the North Pacific can extend westward to East Asia. Xie et al. (2018) identified the effect of spring ASO changes on spring precipitation in China. Note that the

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

weakened westerlies in the mid-latitudes and the enhanced westerlies at low latitudes can also extend eastward to the western United States. This kind of circulation anomaly corresponds to two barotropic structures; i.e., an anomalous anticyclone in the Northeast Pacific and a cyclone in the southwestern United States at 500 hPa and 200 hPa. Coincidentally, the northwestern United States is located to the north of the intersection of the anticyclone and cyclone, corresponding to convergence of the airflow at high levels, which may lead to downwelling in the northwestern United States, and vice versa for negative March ASO anomalies.

To further validate our inference regarding the response of the circulation in the western United States to ASO changes, we analyze the differences between April horizontal wind anomalies during positive and negative March ASO anomaly events at 200, 500, and 850 hPa (Fig. 4). As in the increased ASO case, the difference shows an anomalous anticyclone in the Northeast Pacific and an anomalous cyclone in the southwestern United States. The climatological wind over the northwestern United States blows from west to east, bringing moisture from the Pacific to the western United States. Such circulation anomalies force an anomalous cyclone in the western United States in the middle and upper troposphere, which reduces the climatological wind. It would decrease the water vapor concentration in the air over the northwestern United States. In addition, the northwestern United States is located to the north of the intersection of the anticyclone and cyclone, suggesting downwelling flow in the region.

Figure 5a shows a longitude–latitude cross-section of differences in April vertical velocity anomalies averaged over 1000–500 hPa between positive and negative March ASO anomaly events. When the March ASO increases, anomalous downwelling is

删除的内容: This kind of circulation anomaly

删除的内容: southwestern

删除的内容: enhances cold and dry airflow

删除的内容: North American continent

删除的内容: North Pacific, reducing

删除的内容: western United States and possibly reducing April precipitation in the

found in the northwestern United States (115°–130° W). This situation may inhibit precipitation in the northwestern United States in April. Figure 5b depicts the longitude–height cross-section of differences in April vertical velocity averaged over 43°–50°N between positive and negative March ASO anomaly events, which further shows an anomalous downwelling over the northwestern United States when the ASO increases. Based on the above analysis, the circulation anomalies in the northwestern United States associated with positive March ASO anomalies may inhibit the formation of local precipitation in April, and vice versa for that with negative March ASO anomalies.

#### **4. Simulations of the effect of ASO variations on precipitation in the northwestern US during spring**

Using observations and reanalysis data, we investigated the relationship between March ASO and April precipitation in the northwestern US and revealed the underlying mechanisms in section 3. In this section, we use WACCM4 simulations (see section 2) to confirm the above conclusions. First, we check the model performance in simulating precipitation over western North America. Figure 6 shows the April precipitation climatology over the region 95°–140°W, 30°–63°N from the control experiment R1 (Table 1) and from GPCP for the period 1995–2005. The model simulates a center of high precipitation over the west coast of North America (Fig. 6a). It is clear that the spatial distribution of the simulated precipitation climatology is similar to that calculated by GPCP (Fig. 6b).

Figure 7a displays the differences in April precipitation between experiments R3 and R2. The pattern of simulated April precipitation anomalies forced by ASO changes in western North America (Fig. 7a) is different from that observed (Fig. 1);

i.e., the increased March ASO forces an increase in precipitation in the northwestern United States. The differences in April zonal wind at 200, 500, and 850 hPa between experiments R3 and R2 are shown in Fig. 7b, c, and d, respectively. The simulated pattern of April zonal wind anomalies in western North America (Fig. 7b, c and d) shifted a little further to the north than in the observations (Fig. 3). Comparing the global pattern of simulated April zonal wind anomalies with the observations, it is surprising to find that the positions of simulated zonal wind anomalies over the Northeast Pacific and western North America are shifted northward. This results in the simulated precipitation anomalies over western North America also shifting northward, so that a decrease in precipitation on the west coast of Canada in April is found in Fig. 7a. This explains why we find the pattern of simulated April precipitation anomalies in the North America (Fig. 7a) is nearly opposite to that observed (Fig. 1). Figure 7 shows that the results of the model simulation in which we only change the ASO forcing do not reflect the real situation of April precipitation anomalies in the northwestern United States, with a shift in position compared with observations. This leads us to consider whether other factors interact with March ozone to influence April precipitation in the northwestern United States.

Previous studies have found that the North Pacific SST has a significant effect on precipitation in the United States (e.g., Namias, 1983; Ting and Wang, 1997; Wang and Ting, 2000; Barlow et al., 2001; Lau et al., 2002; Wang et al., 2014). Figure 8a shows the correlation coefficients between regional averaged ( $43^{\circ}$ – $50^{\circ}$ N,  $115^{\circ}$ – $130^{\circ}$ W) precipitation anomalies and SST variations in April. Interestingly, the results show that the distribution of correlation coefficients over the North Pacific has a meridional tripole structure, which is referred to as the Victoria Mode SST anomaly pattern. Xie et al. (2017a) demonstrated that the ASO has a lagged impact on the sea

删除的内容: northwestern United States

surface temperature in the North Pacific mid–high latitudes based on observation and simulation. They showed that stratospheric circulation anomalies caused by ASO changes can rapidly extend to the lower troposphere in the high latitudes of the Northern Hemisphere. The circulation anomalies in the high latitudes of the lower troposphere take about a month to propagate to the North Pacific mid-latitudes and then influence the North Pacific SST. Figure 8b shows the correlation coefficients between March ASO (multiplied by  $-1$ ) and April SST variations. The pattern in Fig. 8b is in good agreement with that in Fig. 8a. It is further found that removing the Victoria Mode signal from the time series of precipitation in the northwestern United States reduces the correlation coefficient between March ASO anomalies and filtered April precipitation variations in the northwestern United States to  $-0.40$  (the correlation coefficient is  $-0.55$  for the original time series, see Fig. 2), but it remains significant. Figure 8 indicates that the ASO possibly influences precipitation anomalies in the northwestern United States in two ways. First, the stratospheric circulation anomalies caused by the ASO changes can propagate downward to the North Pacific troposphere and eastward to influence precipitation over northwestern United States. Second, the ASO changes generate SST anomalies over the North Pacific that act as a bridge for ASO to affect precipitation in the northwestern United States (Xie et al., 2017a). The SST anomalies caused by ASO change likely interact with the direct changes in atmospheric circulation driven by the ASO change to jointly influence precipitation in the northwestern United States. Experiments R2 and R3 do not include the effects of SST, which may explain why the results of the model simulation in which we only change the ASO forcing do not reflect the observed precipitation anomalies in the northwestern United States (Fig. 7).

Two sets of experiments (R4 and R5) that include the joint effects of ASO and

删除的内容: .

带格式的: 字体颜色: 自动设置

SST change (Fig. 9) are added. Details of the experiments are given in Table 1. Figure 10 shows the differences in April precipitation and zonal wind between experiments R5 and R4. It is clear that the simulated changes in precipitation in the northwestern United States (Fig. 10a) are in good agreement with the observed anomalies shown in Fig. 1; i.e., the increase in March ASO forces a decrease in April precipitation in the northwestern United States. In addition, the spatial distributions of simulated zonal wind anomalies (Fig. 10b–d) are consistent with the observations (Fig. 3). Overall, the simulated precipitation and circulation in R4 and R5 are no longer shifted northward and are closer to the observations.

To further emphasize the importance of the joint effects of ASO and ASO-related SST anomalies on precipitation in the northwestern United States, we investigate whether the spring Victoria Mode-like SST anomalies alone could force the observed precipitation anomalies in the northwestern United States. Two sets of experiments are performed here (R6 and R7), in which only April SST anomalies over the North Pacific have been changed (Fig. 9). Details of the experiments are given in Table 1. Figure 11 shows the differences in April precipitation and zonal wind between experiments R7 and R6. The simulated precipitation anomalies over the west coast of the United States (Fig. 11a) are much weaker, and the simulated circulation anomalies (Fig. 11b–d) are quite different from those in Fig. 3. This suggests that the ASO-related North Pacific SST anomalies alone cannot force the observed precipitation anomalies in the northwestern United States, but that the combined effect of ASO and ASO-related North Pacific SST anomalies is required (Fig. 10).

In order to further confirm the possibly influence of ASO on precipitation in the northwestern United States, a transient experiment (1955–2005) based on the

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

atmosphere-ocean coupled WACCM4 model is added to confirm whether the ASO by itself can cause the Victoria SST mode in the North Pacific and rainfall anomalies in the northwestern United States. Note that the ozone forcing in the experiment is specified which is derived from the CMIP5 ensemble mean ozone output. Please refer to R8 in Table 1 for a detailed description of the experiment. Figure 12 shows the correlation coefficients between the specified March ASO variations and simulated April 500 hPa U, SST, and precipitation anomalies for the period 1955–2005. The significant and leading effects of the specified ASO anomalies on 500 hPa U, the Victoria mode in the North Pacific, and rainfall anomalies in the northwestern United States are well captured (Fig. 12). As the ozone forcing in the experiment is specified, the relationships between ASO and U and SST and precipitation could only be caused by ASO influencing U, and then U influencing SST and precipitation; the ASO changes are completely independent of polar vortex. The leading relationship between ASO and precipitation in the northwestern United States can be found in observations, time-slice experiments (R1–7), and a transient experiment with specified ozone (R8). Thus, we have shown that the relationship between March ASO and April precipitation in the northwestern US in the observations and the underlying mechanisms can be verified by WACCM4.

带格式的: 字体颜色: 自动设置

## 5. Discussion and summary

Many observations and simulations have shown that ASO variations have a significant impact on Northern Hemisphere tropospheric climate, but few studies have focused on regional characteristics. Using observations, reanalysis datasets, and WACCM4, we have shown that the March ASO changes have a significant effect on April precipitation in the northwestern United States (mainly in Washington and Oregon) with a lead of 1–2 months. When the March ASO is anomalously high, April

删除的内容: spring

precipitation decreases in the northwestern United States, and vice versa for low ASO.

During positive ASO events, the zonal wind changes over the North Pacific exhibit a tripolar mode with a zonal distribution; i.e., enhanced westerlies in the high and low latitudes of the North Pacific, and weakened westerlies in the mid-latitudes. The anomalous wind can extend eastward to North America, causing anomalous circulation in western North America. The climatological wind over the northwestern United States blows from west to east, bringing moisture from the Pacific to western United States. Such circulation anomalies force an anomalous cyclone in the western United States in the middle and upper troposphere, which reduces the climatological wind. It would decrease the water vapor concentration in the air over the northwestern United States. At the same time, downwelling in the northwestern US is enhanced. The two processes possibly decrease April precipitation in the northwestern US. When the March ASO decreases, the effect is just the opposite.

The WACCM4 model is used to confirm the statistical results of observations and the reanalysis data. The results of the model simulation in which we only change the ASO forcing do not reflect the observed precipitation anomalies in the northwestern United States in April; i.e., the pattern of simulated April precipitation and circulation anomalies in the western North America shifted a little further to the north than observed. It is found that SST anomalies over North Pacific caused by ASO changes are likely to interact with ASO changes to jointly influence precipitation in the northwestern United States. Thus, the ASO influences precipitation anomalies over the northwestern United States in two ways. First, the stratospheric circulation anomalies caused by the ASO change can propagate downward to the North Pacific troposphere and directly influence precipitation over

删除的内容: likely enhances cold and dry airflow from

删除的内容: North American continent to the North Pacific, reducing

the northwestern United States. Second, the ASO changes generate SST anomalies over the North Pacific that act as a bridge, (Xie et al., 2017a), allowing the ASO changes to affect precipitation in the northwestern United States.

删除的内容: ,

It is well known that the spring ASO variations are related to changes in the winter Arctic stratospheric vortex (SPV). The strength of the winter SPV can affect spring ASO, and then the ASO affects tropospheric teleconnection and precipitation in the northwestern United States (indirect effect of SPV). The strength of the winter SPV may also have a direct leading effect on tropospheric teleconnection (Baldwin and Dunkerton, 2001; Black et al., 2005, 2006, 2009) and precipitation in the northwestern United States, in spring. A question arises here, whether the stratospheric polar vortex variability in late winter can be a better factor for leading spring precipitation variations in the northwestern United States than the spring ASO anomalies? Figure 13 shows the correlation coefficients between the February SPV (multiplied by -1) index and April 200 hPa zonal wind and precipitation variations (Fig. 13a and b), and between March ASO and April 200 hPa zonal wind and precipitation (Fig. 13c and d). The SPV index is defined as the strength of the stratospheric polar vortex, following Zhang et al. (2018). It is found that the relationship between the strength of February SPV and the variations in 200 hPa zonal wind and precipitation is significant (Fig. 13a and b), indicating indirect or direct effects of winter SPV on spring tropospheric climate. However, the relationship isn't stronger than that between March ASO and April 200 hPa zonal wind and precipitation (Fig. 13c and d). In this study, we try to state that the ASO changes

删除的内容: . Figure 12

删除的内容: 12a

删除的内容: 12c

删除的内容: Although they are similar,

删除的内容: ASO variations are much closer than

删除的内容: the stratospheric polar vortex to

删除的内容: . That indicates

删除的内容: and

删除的内容: Since the coupling between dynamical and radiative processes in spring is strong, the connection

删除的内容: winter SPV

删除的内容: spring tropospheric circulation seems weaker than that between the spring ASO

删除的内容: tropospheric circulation.

could influence precipitation in the northwestern United States, emphasizing the influence of stratospheric ozone on tropospheric regional climate. As for the effect of coupling between dynamical and radiative processes in spring on precipitation is an interesting question that deserves further investigation.

**Acknowledgments.** Funding for this project was provided by the National Natural Science Foundation of China (41630421, 41790474 and 41575039). We acknowledge ozone datasets from the SWOOSH and GOZCARDS; precipitation from GPCP and GPCP; Meteorological fields from NCEP2, SST from the UK Met Office Hadley Centre, and WACCM4 from NCAR.

带格式的: 英语(英国)

删除的内容: China Meteorological Administration,

## References:

Archer, C. L. and Caldeira, K.: Historical trends in the jet streams, *Geophys. Res. Lett.*, 35, L08803, doi:10.1029/2008GL033614, 2008.

带格式的: 字体颜色: 自动设置

Baldwin, M. P. and Dunkerton, T. J.: Stratospheric harbingers of anomalous weather regimes, *Science*, 294, 581-584, doi:10.1126/science.1063315, 2001.

带格式的: 字体颜色: 自动设置

Barlow, M., Nigam, S., and Berbery, E. H.: ENSO, Pacific decadal variability, and US summertime precipitation, drought, and stream flow, *J. Climate*, 14, 2105-2128, doi:10.1175/1520-0442(2001)014<2105:EPDVAU>2.0.CO;2, 2001.

带格式的: 字体颜色: 自动设置

Bitz, C. M. and Polvani, L. M.: Antarctic climate response to stratospheric ozone depletion in a fine resolution ocean climate model, *Geophys. Res. Lett.*, 39, L20705, doi:10.1029/2012GL053393, 2012.

Black, R. X., Mcdaniel, B. A., and Robinson, W. A.: Stratosphere Troposphere Coupling during Spring Onset, *J. Climate*, 19, 4891-4901, doi:10.1175/Jcli3907.1, 2005.

带格式的: 字体颜色: 自动设置

Black, R. X., and Mcdaniel, B. A.: The Dynamics of Northern Hemisphere Stratospheric Final Warming Events, *J. Atmos. Sci.*, 64, 2932-2946, doi:10.1175/Jas3981.1, 2006.

带格式的: 字体颜色: 自动设置

Black, R. X. and Mcdaniel, B. A.: SubMonthly polar vortex variability and stratosphere-troposphere coupling in the Arctic, *J. Climate*, 22, 5886-5901, doi:10.1175/2009JCLI2730.1, 2009.

带格式的: 字体颜色: 自动设置

Cagnazzo, C. and Manzini, E.: Impact of the Stratosphere on the Winter Tropospheric Teleconnections between ENSO and the North Atlantic and European Region, *J.*

Climate, 22, 1223–1238, doi:10.1175/2008JCLI2549.1, 2009.

带格式的: 字体颜色: 自动设置

Calvo, N., Polvani, L. M., and Solomon, S.: On the surface impact of Arctic stratospheric ozone extremes, *Environ. Res. Lett.*, 10, 094003, doi:10.1088/1748-9326/10/9/094003, 2015.

Charlton, A. J. and Polvani, L. M.: A new look at stratospheric sudden warmings. Part I: Climatology and modeling benchmarks, *J. Climate*, 20, 449–469, doi:10.1175/JCLI3996.1, 2007.

带格式的: 字体颜色: 自动设置

Cheung, J. C. H., Haigh, J. D., and Jackson, D. R.: Impact of EOS MLS ozone data on medium-extended range ensemble weather forecasts, *J. Geophys. Res.*, 119, 9253–9266, doi:10.1002/2014JD021823, 2014.

带格式的: 字体颜色: 自动设置

Davis, S. M., Rosenlof, K. H., Hassler, B., Hurst, D. F., Read, W. G., Vomel, H., Selkirk, H., Fujiwara, M., and Damadeo, R.: The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) database: a long-term database for climate studies, *Earth Syst Sci Data*, 8, 461–490, doi:10.5194/essd-8-461-2016, 2016.

Feldstein, S. B.: Subtropical Rainfall and the Antarctic Ozone Hole, *Science*, 332, 925–926, doi:10.1126/science.1206834, 2011.

Forster, P. M. D. and Shine, K. P.: Radiative forcing and temperature trends from stratospheric ozone changes, *J. Geophys. Res.*, 102, 10841–10855, doi:10.1029/96JD03510, 1997.

Froidevaux, L., Anderson, J., Wang, H.-J., Fuller, R. A., Schwartz, M. J., Santee, M. L., Livesey, N. J., Pumphrey, H. C., Bernath, P. F., Russell III, J. M., and

- McCormick, M. P.: Global Ozone Chemistry And Related trace gas Data records for the Stratosphere (GOZCARDS): methodology and sample results with a focus on HCl, H<sub>2</sub>O, and O<sub>3</sub>, *Atmos. Chem. Phys.*, 15, 10471–10507, doi:10.5194/acp-15-10471-2015, 2015.
- Gabriel, A., Peters, D., Kirchner, I., and Graf, H. F.: Effect of zonally asymmetric ozone on stratospheric temperature and planetary wave propagation, *Geophys. Res. Lett.*, 34, L06807, doi:10.1029/2006GL028998, 2007.
- Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulation of secular trends in the middle atmosphere, 1950–2003, *J. Geophys. Res.*, 112, D09301, doi:10.1029/2006JD007485, 2007.
- Gerber, E. P. and Son, S. W.: Quantifying the Summertime Response of the Austral Jet Stream and Hadley Cell to Stratospheric Ozone and Greenhouse Gases, *J. Climate*, 27, 5538–5559, doi:10.1175/JCLI-D-13-00539.1, 2014.
- Gillett, N. P., Scinocca, J. F., Plummer, D. A., and Reader, M. C.: Sensitivity of climate to dynamically-consistent zonal asymmetries in ozone, *Geophys. Res. Lett.*, 36, L10809, doi:10.1029/2009GL037246, 2009.
- Graf, H. F. and Walter, K.: Polar vortex controls coupling of North Atlantic Ocean and atmosphere, *Geophys. Res. Lett.*, 32, L01704, doi:10.1029/2004GL020664, 2005.
- Haigh, J. D.: The Role of Stratospheric Ozone in Modulating the Solar Radiative Forcing of Climate, *Nature*, 370, 544–546, doi:10.1038/370544a0, 1994.
- Holland, M. M., Bailey, D. A., Briegleb, B. P., Light, B., and Hunke, E.: Improved

Sea Ice Shortwave Radiation Physics in CCSM4: The Impact of Melt Ponds and Aerosols on Arctic Sea Ice, *J. Climate*, 25, 1413–1430, doi:10.1175/JCLI-D-11-00078.1, 2012.

Hu, Y., Tao, L., and Liu, J.: Poleward expansion of the Hadley circulation in CMIP5 simulations, *Adv. Atmos. Sci.*, 30, 790–795, doi:10.1007/s00376-012-2187-4, 2013.

Huffman, G. J., Adler, R. F., Arkin, P., Chang, A., Ferraro, R., Gruber, A., Janowiak, J., McNab, A., Rudolf, B., and Schneider, U.: The Global Precipitation Climatology Project (GPCP) Combined Precipitation Dataset, *B. Am. Meteorol. Soc.*, 78, 5–20, doi:10.1175/1520-0477(1997)078<0005:TGPCPG>2.0.Co;2, 1997.

Ineson, S. and Scaife, A. A.: The role of the stratosphere in the European climate response to El Nino, *Nat. Geosci.*, 2, 32–36, doi:10.1038/NGEO381, 2009.

Ivy, D. J., Solomon, S., Calvo, N., and Thompson, D. W.: Observed connections of Arctic stratospheric ozone extremes to Northern Hemisphere surface climate, *Environ. Res. Lett.*, 12, 024004, doi:10.1088/1748-9326/aa57a4, 2017.

Kang, S. M., Polvani, L. M., Fyfe, J. C., and Sigmond, M.: Impact of Polar Ozone Depletion on Subtropical Precipitation, *Science*, 332, 951–954, doi:10.1126/science.1202131, 2011.

Karpechko, A. Y., Perlwitz, J., and Manzini, E.: A model study of tropospheric impacts of the Arctic ozone depletion 2011, *J. Geophys. Res.*, 119, 7999–8014, doi:10.1002/2013JD021350, 2014.

Kidston, J., Scaife, A. A., Hardiman, S. C., Mitchell, D. M., Butchart, N., Baldwin, M.

P., and Gray, L. J.: Stratospheric influence on tropospheric jet streams, storm tracks and surface weather, *Nat. Geosci.*, 8, 433–440, doi:10.1038/NGEO2424, 2015.

Labitzke, K. and Naujokat, B.: The lower Arctic stratosphere in winter since 1952, *Sparc Newsletter*, 15, 11–14, 2000.

Lau, K. M., Kim, K. M., and Shen, S. S.: Potential predictability of seasonal precipitation over the U.S. from canonical ensemble correlation predictions, *Geophys. Res. Lett.*, 29, 1–4, doi:10.1029/2001GL014263, 2002.

Li, F., Vikhliayev, Y. V., Newman, P. A., Pawson, S., Perlwitz, J., Waugh, D. W., and Douglass, A. R.: Impacts of Interactive Stratospheric Chemistry on Antarctic and Southern Ocean Climate Change in the Goddard Earth Observing System, Version 5 (GEOS-5), *J. Climate*, 29, 3199–3218, doi:10.1175/JCLI-D-15-0572.1, 2016.

Lu, J., Deser, C., and Reichler, T.: Cause of the widening of the tropical belt since 1958, *Geophys. Res. Lett.*, 36, L03803, doi:10.1029/2008GL036076, 2009.

Manney, G. L., Santee, M. L., Rex, M., Livesey, N. J., Pitts, M. C., Veefkind, P., Nash, E. R., Wohltmann, I., Lehmann, R., Froidevaux, L., Poole, L. R., Schoeberl, M. R., Haffner, D. P., Davies, J., Dorokhov, V., Gernandt, H., Johnson, B., Kivi, R., Kyrö, E., Larsen, N., Levelt, P. F., Makshtas, A., McElroy, C. T., Nakajima, H., Parrondo, M. C., Tarasick, D. W., von der Gathen, P., Walker, K. A., and Zinoviev, N. S.: Unprecedented Arctic ozone loss in 2011, *Nature*, 478, 469–475, <https://doi.org/10.1038/nature10556>, 2011.

带格式的: 字体颜色: 自动设置

Manney, G. L. and Lawrence, Z. D.: The major stratospheric final warming in 2016: dispersal of vortex air and termination of Arctic chemical ozone loss, *Atmos. Chem. Phys.*, 16, 15371–15396, doi:10.5194/acp-16-15371-2016, 2016.

Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J. F., Calvo, N., and Polvani, L. M.: Climate Change from 1850 to 2005 Simulated in CESM1(WACCM), *J. Climate*, 26, 7372–7391, doi:10.1175/JCLI-D-12-00558.1, 2013.

McCormack, J. P., Nathan, T. R., and Cordero, E. C.: The effect of zonally asymmetric ozone heating on the Northern Hemisphere winter polar stratosphere, *Geophys. Res. Lett.*, 38, 1–5, doi:10.1029/2010GL045937, 2011.

McLandress, C., Shepherd, T. G., Scinocca, J. F., Plummer, D. A., Sigmond, M., Jonsson, A. I., and Reader, M. C.: Separating the dynamical effects of climate change and ozone depletion. Part II: Southern Hemisphere troposphere, *J. Climate*, 24, 1850–1868, doi:10.1175/2010JCLI3958.1, 2011.

Min, S. K. and Son, S. W.: Multimodel attribution of the Southern Hemisphere Hadley cell widening: Major role of ozone depletion, *J. Geophys. Res.*, 118, 3007–3015, doi:10.1002/jgrd.50232, 2013.

Namias, J.: Some causes of U.S. drought, *J. Clim. Appl. Meteorol.*, 22, 30–39, doi:10.1175/1520-0450(1983)022<0030:Scousd>2.0.Co;2, 1983.

Neale, R. B., Richter, J., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., and Zhang, M. H.: The Mean Climate of the Community Atmosphere Model (CAM4) in Forced SST and Fully Coupled Experiments, *J. Climate*, 26, 5150–5168, doi:10.1175/JCLI-D-12-00236.1, 2013.

带格式的: 字体颜色: 自动设置

Nowack, P. J., Abraham, N. L., Maycock, A. C., Braesicke, P., Gregory, J. M., Joshi, M. M., Osprey, A., and Pyle, J. A.: A large ozone-circulation feedback and its implications for global warming assessments, *Nat. Clim. Change*, 5, 41–45, doi:10.1038/NCLIMATE2451, 2015.

Nowack, P. J., Braesicke, P., Abraham, N. L., and Pyle, J. A.: On the role of ozone feedback in the ENSO amplitude response under global warming, *Geophys. Res. Lett.*, 44, 3858–3866, doi:10.1002/2016GL072418, 2017.

Nowack, P. J., Abraham, N. L., Braesicke, P., and Pyle, J. A.: The impact of stratospheric ozone feedbacks on climate sensitivity estimates, *J. Geophys. Res.*, 123, 4630–4641, doi:10.1002/2017JD027943, 2018.

Pawson, S. and Naujokat, B.: The cold winters of the middle 1990s in the northern lower stratosphere, *J. Geophys. Res.*, 104, 14209–14222, doi:10.1029/1999JD900211, 1999.

Polvani, L. M., Waugh, D. W., Correa, G. J., and Son, S.-W.: Stratospheric ozone depletion: The main driver of twentieth-century atmospheric circulation changes in the Southern Hemisphere, *J. Climate*, 24, 795–812, doi:10.1175/2010JCLI3772.1, 2011.

Ramaswamy, V., Schwarzkopf, M. D., and Randel, W. J.: Fingerprint of ozone depletion in the spatial and temporal pattern of recent lower-stratospheric cooling, *Nature*, 382, 616–618, doi:10.1038/382616a0, 1996.

Randel, W. J.: The Seasonal Evolution of Planetary-Waves in the Southern-Hemisphere Stratosphere and Troposphere, *Quarterly Journal of the*

带格式的: 字体颜色: 自动设置

Royal Meteorological Society, 114, 1385–1409, doi:10.1002/qj.49711448403, 1988.

Randel, W. J. and Wu, F.: Cooling of the arctic and antarctic polar stratospheres due to ozone depletion, *J. Climate*, 12, 1467–1479, doi:10.1175/1520-0442(1999)012<1467:COTAAA>2.0.Co;2, 1999.

Randel, W. J. and Wu, F.: A stratospheric ozone profile data set for 1979-2005: Variability, trends, and comparisons with column ozone data, *J. Geophys. Res.*, 112, D06313, doi:10.1029/2006JD007339, 2007.

Ravishankara, A. R., Turnipseed, A. A., Jensen, N. R., Barone, S., Mills, M., Howard, C. J., and Solomon, S.: Do hydrofluorocarbons destroy stratospheric ozone?, *Science*, 263, 71–75, doi:10.1126/science.263.5143.71, 1994.

Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N<sub>2</sub>O): the dominant ozone-depleting substance emitted in the 21st century, *Science*, 326, 123–125, doi:10.1126/science.1176985, 2009.

[Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., and Rowell, D. P.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, \*J. Geophys. Res.\*, 108, 4407, 2003.](#)

[Reichler, T., Kim, J., Manzini, E., and Kroger, J.: A stratospheric connection to Atlantic climate variability, \*Nat. Geosci.\*, 5, 783–787, doi:10.1038/NGEO1586, 2012.](#)

Russell, J. L., Dixon, K. W., Gnanadesikan, A., Stouffer, R. J., and Toggweiler, J. R.:

带格式的: 字体颜色: 自动设置

The Southern Hemisphere westerlies in a warming world: Propping open the door to the deep ocean, *J. Climate*, 19, 6382–6390, doi:10.1175/JCLI3984.1, 2006.

Smith, K. L. and Polvani, L. M.: The surface impacts of Arctic stratospheric ozone anomalies, *Environ. Res. Lett.*, 9, 074015, doi:10.1088/1748-9326/9/7/074015, 2014.

Solomon, S.: Antarctic ozone: Progress towards a quantitative understanding, *Nature*, 347, 354, doi:10.1038/347347a0, 1990.

Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, doi:10.1029/1999RG900008, 1999.

Son, S.-W., Tandon, N. F., Polvani, L. M., and Waugh, D. W.: Ozone hole and Southern Hemisphere climate change, *Geophys. Res. Lett.*, 36, L15705, doi:10.1029/2009GL038671, 2009.

Son, S.-W., Gerber, E. P., Perlwitz, J., Polvani, L. M., Gillett, N. P., Seo, K.-H., Eyring, V., Shepherd, T. G., Waugh, D., Akiyoshi, H., Austin, J., Baumgaertner, A., Bekki, S., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M. P., Cugnet, D., Dameris, M., Dhomse, S., Frith, S., Garny, H., Garcia, R., Hardiman, S. C., Jöckel, P., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Nakamura, T., Morgenstern, O., Pitari, G., Plummer, D. A., Pyle, J., Rozanov, E., Scinocca, J. F., Shibata, K., Smale, D., Teyssède, H., Tian, W., and Yamashita, Y.: Impact of stratospheric ozone on Southern Hemisphere circulation change: A multimodel assessment, *J. Geophys. Res.*, 115, D00M07, doi.org/10.1029/2010JD014271,

2010.

Thompson, D. W. J. and Solomon, S.: Interpretation of recent Southern Hemisphere climate change, *Science*, 296, 895–899, doi:10.1126/science.1069270, 2002.

Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nature Geosci.*, 4, 741–749, doi:10.1038/NGEO1296, 2011.

Ting, M. and Wang, H.: Summertime US Precipitation Variability and Its Relation to Pacific Sea Surface Temperature, *J. Climate*, 10, 1853–1873, doi:10.1175/1520-0442(1997)010<1853:SUSPVA>2.0.CO;2, 1997.

Tung, K. K.: On the Relationship between the Thermal Structure of the Stratosphere and the Seasonal Distribution of Ozone, *Geophys. Res. Lett.*, 13, 1308–1311, doi:10.1029/GL013i012p01308, 1986.

Wang, F., Yang, S., Higgins, W., Li, Q. P., and Zuo, Z. Y.: Long-term changes in total and extreme precipitation over China and the U.S. and their links to oceanic-atmospheric features, *Int. J. Climatol.*, 34, 286–302, doi:10.1002/joc.3685, 2014.

Wang, H. and Ting, M. F.: Covariabilities of winter US precipitation and Pacific Sea surface temperatures, *J. Climate*, 13, 3711–3719, doi:10.1175/1520-0442(2000)013<3711:Cowusp>2.0.Co;2, 2000.

Wang, L., Ting, M., Kushner, P. J.: A robust empirical seasonal prediction of winter NAO and surface climate, *Sci. Rep.*, 7, 279, 2017.

Waugh, D. W., Garfinkel, C. I., and Polvani, L. M.: Drivers of the Recent Tropical Expansion in the Southern Hemisphere: Changing SSTs or Ozone Depletion?, *J. Climate*, 28, 6581–6586, doi:10.1175/JCLI-D-15-0138.1, 2015.

WMO: Scientific Assessment of Ozone depletion: 2002. In: Global Ozone Research and Monitoring Project, Report No. 47, Geneva, 498 pp., 2003.

WMO: Scientific Assessment of Ozone Depletion: 2010. WMO Tech. Note 52, World Meteorological Organization, Geneva, Switzerland, 516 pp., 2011.

Xie, F., Li, J., Tian, W., Fu, Q., Jin, F.-F., Hu, Y., Zhang, J., Wang, W., Sun, C., Feng, J., Yang, Y., and Ding, R.: A connection from Arctic stratospheric ozone to El Niño-Southern Oscillation, *Environ. Res. Lett.*, 11, 124026, doi:10.1088/1748-9326/11/12/124026, 2016.

Xie, F., Li, J., Zhang, J., Tian, W., Hu, Y., Zhao, S., Sun, C., Ding, R., Feng, J., Yang, Y.: Variations in North Pacific sea surface temperature caused by Arctic stratospheric ozone anomalies, *Environ. Res. Lett.*, 12, 114023, doi:10.1088/1748-9326/aa9005, 2017a.

Xie, F., Zhang, J., Sang, W., Li, Y., Qi, Y., Sun, C., and Shu, J.: Delayed effect of Arctic stratospheric ozone on tropical rainfall, *Atmos. Sci. Lett.*, 18, 409–416, 2017b.

Xie, F., Ma, X., Li, J., Huang, J., Tian, W., Zhang, J., Hu, Y., Sun, C., Zhou, X., Feng, J., Yang, Y.: An advanced impact of Arctic stratospheric ozone changes on spring precipitation in China, *Clim. Dyn.*, 51, 4029–4041, doi:10.1007/s00382-018-4402-1, 2018.

删除的内容: <https://doi.org/>

带格式的: 字体颜色: 自动设置

Yin, J. H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate, *Geophys. Res. Lett.*, 32, L18701, doi:10.1029/2005GL023684, 2005.

Zhang, J. K., Tian, W. S., Chipperfield, M. P., Xie, F., and Huang, J. L.: Persistent shift of the Arctic polar vortex towards the Eurasian continent in recent decades, *Nat. Clim. Change.*, 6, 1094–1099, doi:10.1038/nclimate3136, 2016.

Zhang J. K., Tian, W. S., Xie, F., Chipperfield, M. P., Feng, W. H., Son, S-W., Abraham, N. L., Archibald, A. T., Bekki, S., Butchart, N., Deushi, M., Dhomse, S., Han, Y. Y., Jöckel, P., Kinnison, D., Kirner, O., Michou, M., Morgenstern, O., O'Connor, F. M., Pitari, G., Plummer, D. A., Revell, L. E., Rozanov, E., Visioni, D., Wang, W. K., Zeng, G.: Stratospheric ozone loss over the Eurasian continent induced by the polar vortex shift, *Nat. Commun.*, 9, 206, doi.org/10.1038/s41467-017-02565-2, 2018.

**Table 1.** CESM-WACCM4 experiments with various specified ozone and SST forcing.

Exp*1	Specified ozone and SST forcing	Other forcing
		Fixed solar constant, fixed greenhouse gas (GHG) values (averages of emissions scenario A2 of the Intergovernmental Panel on Climate Change (WMO, 2003) over the period 1995–2005), volcanic aerosols (from the Stratospheric Processes and their Role in Climate (SPARC) Chemistry–Climate Model Validation (CCMVal) REF-B2 scenario recommendations), and QBO phase signals with a 28-month zonal wind fixed cycle.
R1	Time-slice run as the control experiment used case F_2000_WACCM_SC. The specified ozone forcing is a 12-month cycle of monthly ozone averaged from 1995 to 2005. The specified SST forcing is a 12-month cycle of monthly SST averaged from 1995 to 2005.	
R2	Same as R1, except that the March ozone in the region 30°–90°N at 300–30 hPa*2 is decreased by 15% compared with R1.	Same as R1
R3	Same as R1, except that March ozone in the region 30°–90°N at 300–30 hPa is increased by 15% compared with R1.	Same as R1
R4	Same as R2, except that SST anomalies in the region 0°–70°N and 120°E–90°W related to negative ASO anomalies*3 is added in the SST forcing in April.	Same as R1
R5	Same as R3, except that SST anomalies in the region 0°–70°N and 120°E–90°W related to positive ASO anomalies*4 is added in the SST forcing in April.	Same as R1
R6	Same as R1, except that SST anomalies in the region 0°–70°N and 120°E–90°W related to negative ASO anomalies*3 is added in the SST forcing in April.	Same as R1

R7	Same as R1, except that SST anomalies in the region 0°–70°N and 120°E–90°W related to positive ASO anomalies*4 is added in the SST forcing in April.	Same as R1
R8	<u>Transient run using case B_1955–2005 WACCM SC CN in CESM. E<sub>1</sub> is a historical simulation covering the period 1955–2005. Note that the specified ozone forcing for 1955–2005 was derived from the CMIP5 ensemble mean ozone output. The specified ozone forcing was named ghg forcing 1955-2005 CMIP5 En sMean.c140414.nc, and can be downloaded at <a href="https://svn-ccsm-inputdata.cgd.ucar.edu/trunk/inputdata/atm/waccm/ub/ghg_forcing_1955-2005_CMIP5_Ens_Mean.c140414.nc">https://svn-ccsm-inputdata.cgd.ucar.edu/trunk/inputdata/atm/waccm/ub/ghg_forcing_1955-2005_CMIP5_Ens_Mean.c140414.nc</a></u>	<u>All natural and anthropogenic external forcings for R8 based on observation and from original CESM input data.</u>

\*1Integration time for time-slice runs is 33 years and for transient run is 51 years.


\*2To avoid the effect of the boundary of ozone change on the Arctic stratospheric circulation simulation, the replaced region (30°–90°N, 300–30 hPa) was larger than the region used to define the ASO index (60°–90°N, 100–50 hPa).

\*3For SST anomalies, see Fig. 9a.


\*4For SST anomalies, see Fig. 9b.

**Table 2.** Selected positive and negative years for March ASO anomaly events based on SWOOSH data for the period 1984–2016. Positive and negative March ASO anomaly events are defined using a normalized time series of March ASO variations from 1984 to 2016. Values larger than 1 standard deviation are defined as positive March ASO anomaly events, and those below  $-1$  standard deviation are defined as negative March ASO anomaly events.


Positive March ASO anomaly events	Negative March ASO anomaly events
1998, 1999, 2001, 2004, 2010	1993, 1995, 1996, 2000, 2011



**Figure 1.** Correlation coefficients between March ASO and April precipitation variations calculated from SWOOSH (a, b) and GOZCARDS (c, d) ozone, and GPCC (a, c) and GPCP (b, d) rainfall for the period 1984–2016. Dots denote significance at the 95% confidence level, according to Student’s t-test. The long-term linear trend and seasonal cycle in all variables were removed before the correlation analysis.




**Figure 2.** (a) Correlation coefficients between March ASO index and precipitation anomalies in the northwestern US ( $43^{\circ}$ – $50^{\circ}$ N,  $115^{\circ}$ – $130^{\circ}$ W) for each month calculated from SWOOSH (a, b) and GOZCARDS (c, d) ozone, and GPCC (a, c) and GPCP (b, d) rainfall for the period 1984–2016. The dashed blacked lines refer to the correlation coefficient that is significance at 95% confidence level. The long-term linear trend and seasonal cycle were removed from the original datasets before calculating the correlation coefficients.




**Figure 3.** Correlation coefficients between March ASO index and April zonal wind variations (m/s, from NCEP2) from 1984 to 2016 at 200 hPa (a), 500 hPa (b), and 850 hPa (c). Dots denote significance at the 95% confidence level, according to Student's  $t$ -test. Blue square is the area shown in Fig. 1. Before performing the analysis, the seasonal cycle and linear trend were removed from the original datasets.

带格式的: 字体颜色: 自动设置

删除的内容: ASO data is from SWOOSH.



**Figure 4.** Differences in composite April winds (vectors, m/s, from NCEP2) between positive and negative ASO anomaly events at 200 hPa (a), 500 hPa (b), and 850 hPa (c) for 1984–2016. Colored regions are statistically significant at the 90% (light yellow) and 95% (dark yellow) confidence levels. The seasonal cycle and linear trend were removed from the original dataset. The ASO anomaly events are selected based on Table 2.




**Figure 5.** (a) Longitude–latitude cross-section of differences in composite April vertical velocity anomalies (averaged over 1000–500 hPa) between positive and negative ASO anomaly events for 1984–2016. (b) Longitude–height cross-section of differences in composite April vertical velocity anomalies (averaged over 43°–50°N) between positive and negative ASO anomaly events from 1984 to 2016. Blue is upward motion and red is downward motion. Dots denote significance at the 95% confidence level. Before performing the analysis, the seasonal cycle and linear trend were removed from the original dataset. The ASO anomaly events are selected based on Table 2. The vertical velocity (Pa/s) dataset is from NCEP2.

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置


带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置




**Figure 6.** (a) Spatial distribution of April precipitation (mm/day) climatology in the control experiment (R1). (b) Same as (a), but precipitation from the GPCP for the period 1995–2005. For details of specific experiments, see Table 1.

带格式的: 字体颜色: 自动设置



**Figure 7.** Differences between experiments R3 and R2 in terms of April (a) precipitation (mm/day) and (b–d) zonal wind at 200, 500, and 850 hPa, respectively. Dots denote significance at the 95% confidence level.

删除的内容: <对象>




**Figure 8.** (a) Correlation coefficients between regional precipitation ( $43^{\circ}$ – $50^{\circ}$ N,  $115^{\circ}$ – $130^{\circ}$ W) and SST variations in April for 1984–2016. (b) Correlation coefficients between March ASO ( $\times -1$ ) and April SST variations for 1984–2016. Dots denote significance at the 95% confidence level, according to Student's  $t$ -test. Before performing the analysis, the seasonal cycle and linear trend were removed from the original data. ASO data are from SWOOSH, precipitation from GPCP, and SST from HadSST.

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

删除的内容: is




**Figure 9.** (a) Composite SST anomalies during negative ASO anomaly events. (b) Composite SST anomalies during positive ASO anomaly events. The ASO anomaly events are selected based on Table 2. SST data are from CESM SST forcing data.

带格式的: 字体颜色: 自动设置

删除的内容: is

带格式的: 字体颜色: 自动设置




**Figure 10.** Same as Fig. 7, but for the difference between experiments R5 and R4.

删除的内容: <对象>

带格式的: 字体颜色: 自动设置


带格式的: 字体颜色: 自动设置




**Figure 11.** Same as Fig. 7, but for the difference between experiments R7 and R6.

删除的内容: <对象>

带格式的: 字体颜色: 自动设置



**Figure 12.** Correlation coefficients between the specified March ASO variations and simulated anomalies of April U (a), SST (b), and precipitation (c) for the period 1955–2005 based on the transient experiment R8. Regions above the 95% confidence level are dotted. The seasonal cycle and linear trend were removed from all quantities before correlation.



**Figure 13.** (a) Correlation coefficients between the February  $-SPV$  ( $10^5 \text{ K m}^2 \text{ kg}^{-1} \text{ s}^{-1}$ ) index defined by Zhang et al. (2018) and April zonal wind variations at 200 hPa for 1984–2016. (b) Correlation coefficients between February  $-SPV$  index and April precipitation variations. (c) and (d) As for (a) and (b), but between March ASO and April 200 hPa zonal wind and April precipitation variations. Dots denote significance at the 95% confidence level, according to Student's  $t$ -test. The long-term linear trend and seasonal cycle in all variables were removed before the correlation analysis. The ASO data are from SWOOSH, zonal wind from NCEP2, and precipitation from GPCP.

带格式的: 字体颜色: 自动设置

删除的内容: 12

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

带格式的: 字体颜色: 自动设置

删除的内容: is

带格式的: 字体颜色: 自动设置

带格式的: 字体: 非加粗

