



# Unraveling the role of silicon in atmospheric aerosol secondary formation: A new conservative tracer for aerosol chemistry

Dawei Lu<sup>1</sup>, Jihua Tan<sup>2</sup>, Xuezhi Yang<sup>1,2</sup>, Xu Sun<sup>1</sup>, Qian Liu<sup>1,2,3</sup>, Guibin Jiang<sup>1,2</sup>

<sup>1</sup>State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

<sup>2</sup>College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>Institute of Environment and Health, Jiangnan University, Wuhan 430056, China

Correspondence to: Qian Liu ([qianliu@rcees.ac.cn](mailto:qianliu@rcees.ac.cn))

**Abstract.** Aerosol particles are ubiquitous in the atmosphere and affect the quality of human life through their climatic and health effects. The formation and growth of aerosol particles involve extremely complex reactions and processes. Due to limited research tools, the sources and chemistry of aerosols are still not fully understood, and until now they are normally investigated by using chemical species of secondary aerosols (e.g.,  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ , SOC) as tracers. Here we investigated the role of silicon (Si), a ubiquitous but relatively inert element, during the secondary aerosol formation process. We analyzed the correlation of Si in airborne fine particles ( $\text{PM}_{2.5}$ ) collected in Beijing—a typical pollution region—with the secondary chemical species and secondary particle precursors (e.g.,  $\text{SO}_2$  and  $\text{NO}_x$ ). The total mass of Si in  $\text{PM}_{2.5}$  was found to be uncorrelated with the secondary aerosol formation process, which suggested that Si is a new conservative tracer for aerosol chemistry. This finding enables the accurate estimation of secondary aerosol contribution to  $\text{PM}_{2.5}$  by using Si as a single tracer rather than normally used multiple chemical tracers. In addition, we show that the correlation analysis of secondary aerosols with the Si isotopic composition of  $\text{PM}_{2.5}$  can further reveal the sources of the precursors of secondary aerosols. Therefore, Si may open a new method for aerosol chemistry studies and pollution control policy development.

## 1 Introduction

Atmospheric particulate pollution is a global environmental issue that seriously threatens human health and sustainable development. The fine particulate matter with size less than  $2.5\ \mu\text{m}$  ( $\text{PM}_{2.5}$ ) is of greatest concern because it poses significant risks to human health and is a major cause of haze phenomenon (Pope et al., 2009; Pope et al., 2002). Understanding the sources and chemistry of aerosols is critical for air pollution control. The aerosol particles can be directly emitted from primary sources (primary particles) or secondarily formed from gaseous precursors (e.g.,  $\text{SO}_2$ ,  $\text{NO}_x$ ,  $\text{NH}_3$ , and volatile organic compounds (VOCs)) through complex chemical reactions and processes in the atmosphere (secondary particles) (Zhang et al., 2012b; Zhang et al., 2015). Particularly, secondary aerosol (SA) is of great importance as it contributes to a large portion of particulate pollution in most pollution regions (e.g., China and India) (Huang et al., 2014). However, until now, the formation mechanism of secondary aerosols is not fully understood and it is difficult to accurately estimate



secondary aerosols due to the extreme complexity of aerosol chemistry and limited research tools. The traditional method is based on the combined estimation of secondary inorganic aerosol (SIA) and secondary organic aerosol (SOA) using multiple chemical components indicative of secondary chemistry, e.g.  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$  for SIA (Bi et al., 2007), and organic carbon (OC) and elemental carbon (EC) for SOA (Docherty et al., 2008). However, the accuracy of the method is still controversial and measurements of these multiple tracers are laborious and may cause large uncertainties to the results (Schmid et al., 2001; Watson et al., 2005).

As the second highest abundant element in the Earth's crust, silicon (Si) is ubiquitous in terrestrial systems including in atmospheric aerosols. The roles of Si in continental and marine environments have been well clarified in biogeochemical studies (Basile-Doelsch, 2006). However, as a mismatch of the ubiquitousness of Si, the research on atmospheric Si is rather limited (Bzdek et al., 2014; Lu et al., 2018). We note that Si is a special element compared with other high-abundance ones (e.g., O, C, N, and S). It is relatively inert that normally forms bonds only with oxygen (in  $\text{Si}^{\text{IV}}$ ) and does not form volatile compounds in the natural environment (Savage et al., 2014). In fact, over 90% of Si in the Earth's crust is composed of nonvolatile silicate minerals that should not be able to serve as gaseous precursors for secondary aerosols (an exception is synthetic organosilicons that will be discussed later). Therefore, Si may provide a different route to aerosol chemistry from that using traditional tracers.

In this study, we aimed to clarify the role of Si during the secondary aerosol formation process in the atmosphere. Based on the nature of Si, we hypothesize that Si in  $\text{PM}_{2.5}$  is unaffected by the secondary particle formation process and predominantly emitted from primary sources (as depicted in Fig. 1). Therefore, the particular properties of Si (i.e., high abundance and chemical inactiveness) may make it a new conservative tracer for aerosol chemistry studies. To specify this point, we herein show the estimation of secondary aerosols based on the dilution effect of Si during the secondary aerosol formation process by using Si as a single tracer rather than normally used multiple chemical tracers.

## 2 Methodology

### 2.1 Chemicals and reagents

The standard reference material of urban atmospheric particulate matter (NIST-1648a) and the Si isotope standard (NIST SRM-8546) were purchased from the National Institute of Standards and Technology (Gaithersburg, MD). The Si isotope standard IRMM-017 was purchased from the Institute for Reference Materials and Measurements (GEEL, Belgium). The element calibration standard solution was from Agilent (Santa Clara, CA). Sodium hydroxide was from Beijing Ruikai Co. (Beijing, China). Nitric acid was from Merck (Darmstadt, Germany). Hydrochloric acid was from Beijing Chemicals Works (Beijing, China). Hydrogen peroxide was from Sinopharm Chemical Reagent Co. (Shanghai, China). Ultrapure water (18.3  $\text{M}\Omega\cdot\text{cm}$ ) obtained from a Milli-Q Gradient system (Millipore, Bedford) was used.



## 2.2 Sampling of PM<sub>2.5</sub> samples

The PM<sub>2.5</sub> and the primary source samples were collected around the Beijing region, China, as described in a previous report (Lu et al., 2018). Briefly, the PM<sub>2.5</sub> samples were collected around Beijing on random haze days ( $n = 100$ ) in 2013 by using a low-volume air sampler (Partisol 2025i, Thermo Fisher, USA) at a flow rate of 16.7 L/min. The PM<sub>2.5</sub> samples were collected onto Whatman 3-5 Teflon membrane filters ( $\varnothing = 47$  mm, Maidstone, UK) or Munktell MK360 quartz filters ( $\varnothing = 90$  mm, Maidstone, UK) and weighed by the giant gravimetric balance method (Yang et al., 2015). Different filters were chosen depending on target analytes, e.g., the analysis of Si and water soluble inorganic ions must be performed with Teflon filters, while quartz filters should be used for OC and EC analysis.

## 2.3 Sample preparation procedures

- 10 For the analysis of Si concentration and isotopic composition, the samples were digested and purified as reported previously (Georg et al., 2006; Zambardi and Poitrasson, 2011). Briefly, the sample was first dried in a silver crucible in a muffle furnace at 1000 K for 10 min. Then, high-purity solid NaOH was added to the sample at a ratio of 1:20 and the mixture was heated at 1000 K for another 10 min followed by cooling down to room temperature. The obtained fusion cake was dissolved with 2 mL of water followed by 24 h incubation, and then the solution was acidized by an HCl solution to pH ~2.
- 15 To eliminate the interference from matrix, the samples were purified by cation exchange column chromatography. The cation resin (Dowex 50WX8, 200-400 mesh) in H<sup>+</sup> form was packed to a 1.8 mL resin bed in a BioRad column and then rinsed with HCl, HNO<sub>3</sub> solution, and water until the eluate reached pH ~7. Afterwards, 1 mL of the sample solution was loaded to the resin and eluted with 2 mL of water. Because Si in the solution is present mostly in the form of non-ionic Si(OH)<sub>4</sub> and anionic H<sub>3</sub>SiO<sub>4</sub><sup>-</sup> (Georg et al., 2006), cationic species could be removed from Si species by the cation exchange resin. The recovery during the column purification procedures with the IRMM-017 was >95%. The recovery of Si during the whole sample preparation procedures with the NIST-1648 was 89.9-96.2%.
- 20

## 2.4 Measurement of Si concentration

The concentration of Si was measured on an Agilent 8800 inductively coupled plasma mass spectrometer (Santa Clara, CA, USA). The clean blank membrane filters have also been analyzed using the same procedures to subtract the background signals from the samples.

25

## 2.5 Estimation of secondary aerosols in PM<sub>2.5</sub> by using the Si-dilution method

Providing that the total Si in PM<sub>2.5</sub> ( $\text{Si}_{\text{PM}_{2.5}}$ ) keeps unchanged during the secondary growth of PM<sub>2.5</sub>, we can obtain

$$m_{\text{PM}_{2.5}} = m_{\text{pri}} + m_{\text{sec}} \quad (1)$$

$$\text{Si}_{\text{PM}_{2.5}} = m_{\text{PM}_{2.5}} \times C_{\text{PM}_{2.5}} = m_{\text{pri}} \times C_{\text{pri}} \quad (2)$$



where  $m_{\text{PM}_{2.5}}$ ,  $m_{\text{pri}}$ , and  $m_{\text{sec}}$  represent the mass of  $\text{PM}_{2.5}$ , primary particles, and secondary particles, respectively;  $\text{Si}_{\text{PM}_{2.5}}$  represent the total Si mass in  $\text{PM}_{2.5}$ ;  $C_{\text{PM}_{2.5}}$  and  $C_{\text{pri}}$  represent Si abundance in  $\text{PM}_{2.5}$  and primary particles, respectively. Thus, the contribution of secondary aerosols to  $\text{PM}_{2.5}$  ( $f_{\text{sec}}$ ) can be estimated by the Eq. (3):

$$f_{\text{sec}} = \frac{m_{\text{sec}}}{m_{\text{PM}_{2.5}}} = 1 - \frac{C_{\text{PM}_{2.5}}}{C_{\text{pri}}} \quad (3)$$

- 5 The  $C_{\text{PM}_{2.5}}$  can be obtained by direct measurement of the collected  $\text{PM}_{2.5}$  samples, and the  $C_{\text{pri}}$  can be estimated from the mixed Si abundances of primary sources:

$$C_{\text{pri}} = \sum(C_i \times f_i) \quad (4)$$

where  $C_i$  and  $f_i$  represent the Si abundance and mass fraction of different primary sources ( $f_i = m_i / \sum m_i$ ). In Eq. (4), the  $C_i$  can be obtained by measuring the collected primary source samples. In order to obtain  $f_i$ , we need to know the emission amount ( $m_i$ ) of each source. The  $m_i$  of anthropogenic sources in the Beijing region was adopted from the Multi-resolution Emission Inventory for China (MEIC) database (MEIC; Li et al., 2017). For the  $m_i$  of natural sources (i.e., dusts), it can be calculated based on the isotopic mass balance of Si in  $\text{PM}_{2.5}$ :

$$\delta^{30}\text{Si}_{\text{PM}_{2.5}} = \delta^{30}\text{Si}_{\text{pri}} = \sum(\delta^{30}\text{Si}_i \times C_i \times f_i) / C_{\text{pri}} \quad (5)$$

where  $\delta^{30}\text{Si}_{\text{PM}_{2.5}}$  represents Si isotopic composition of  $\text{PM}_{2.5}$  and  $\delta^{30}\text{Si}_i$  represents Si isotopic signatures of different primary sources. The  $m_i$  values for different primary sources used in the calculation are given in Table 1.

## 2.6 Uncertainty analysis in the estimation of secondary aerosols

The uncertainty of secondary aerosol contribution estimated by the Si-dilution method was obtained by the error propagation calculation. The general formula for the error propagation calculation is as follows (Ku, 1966; Larsen et al., 2012):

$$S_f = \sqrt{\sum \left( \frac{\partial f}{\partial x_i} \times S_i \right)^2} \quad (6)$$

20 where  $S_f$  represents the standard deviation of the function  $f$ ,  $x_i$  represents the variables in the function  $f$ ,  $S_i$  represents the standard deviation of  $x_i$ . The Eq. (6) was applied to each step of the calculation of secondary aerosol contribution ( $f_{\text{sec}}$ ) to obtain the uncertainty of the final result. Briefly, the uncertainty of  $f_{\text{sec}}$  is dependent on several variables in the calculation, including the measured Si abundance in  $\text{PM}_{2.5}$  ( $C_{\text{PM}_{2.5}}$ ), Si abundance ( $C_i$ ) and mass fraction of primary sources ( $f_i$ ), and Si isotopic composition of  $\text{PM}_{2.5}$  ( $\delta^{30}\text{Si}_{\text{PM}_{2.5}}$ ) and primary sources ( $\delta^{30}\text{Si}_i$ ). The errors of  $C_{\text{PM}_{2.5}}$ ,  $C_i$ ,  $\delta^{30}\text{Si}_{\text{PM}_{2.5}}$ , and  $\delta^{30}\text{Si}_i$  can be directly obtained in the experimental measurements. The  $f_i$  is given by the MEIC database (MEIC; Li et al., 2017). The MEIC is a public emission inventory database that has been widely applied and well validated in air pollution research in China (Jiang et al., 2015; Zheng et al., 2015; Guan et al., 2014; Lin et al., 2014; Zhang et al., 2012a), but it does not include error information for the data. So, the error of  $f_i$  was not considered (i.e., assuming that the relative emission ratios of primary sources are steady). Based on the method described above, the uncertainty of the annual mean  $f_{\text{sec}}$  in 2013 was calculated to be 26.1%.



### 3 Results and discussion

#### 3.1 Correlation of Si with the secondary species in PM<sub>2.5</sub>

To verify the role of Si as depicted in Fig. 1, we analyzed Si abundance and secondary aerosol tracers (e.g., NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, OC, and EC) in PM<sub>2.5</sub> samples collected around Beijing, China, a typical particulate pollution region, on random haze days ( $n = 63$ ) in 2013, in which year the particulate pollution in this region reached an unprecedentedly high level (Huang et al., 2014). We found that the PM<sub>2.5</sub> concentration showed a clear seasonal trend (higher in spring/winter than in summer/autumn) within the range of 28.4-337.1 μg/m<sup>3</sup>, and all secondary species in PM<sub>2.5</sub> showed a similar seasonal trend with the PM<sub>2.5</sub> concentration (Fig. S1, Supplement). However, the total Si in PM<sub>2.5</sub> (expressed as Si<sub>PM<sub>2.5</sub></sub> relative to air volume in μg/m<sup>3</sup>) did not show any seasonal trends (Fig. S2, Supplement).

Furthermore, Fig. 2 shows the correlation of PM<sub>2.5</sub> and its Si content with the secondary species in PM<sub>2.5</sub>. It can be seen that the PM<sub>2.5</sub> concentration was highly linearly correlated with the secondary species ( $P < 0.01$ ). These chemical species are directly indicative of secondary chemistry: sulfate is mainly converted from atmospheric SO<sub>2</sub> primarily emitted from coal combustion (Seinfeld and Pandis, 2006), nitrate originates from NO<sub>x</sub> emitted mainly from vehicle exhaust and power plants (Seinfeld and Pandis, 2006), and secondary organic carbon (SOC), as an indicator of SOA, derives from complex gaseous precursors (Hallquist et al., 2009). Specifically, PM<sub>2.5</sub> showed a higher correlation with SIA ( $P < 0.001$ ) than with SOA ( $P < 0.01$ ) probably due to the higher contribution of SIA to SA than SOA (Fig. S3, Supplement). However, as a stark contrast to the significant correlation in Fig. 2a-d, the Si<sub>PM<sub>2.5</sub></sub> showed no significant correlation with any of the secondary species ( $P > 0.14$ ); either, no obvious trend in Si<sub>PM<sub>2.5</sub></sub> was observed with the increase of secondary species. These results strongly evidenced that Si is inactive during the secondary chemical process of aerosols.

#### 3.2 Correlation of Si with the secondary precursors and relative humidity

To further demonstrate the role of Si in secondary aerosols, we analyzed the correlation of Si<sub>PM<sub>2.5</sub></sub> with the secondary precursors (SO<sub>2</sub> and NO<sub>x</sub>) and relative humidity (RH). RH is an important meteorological parameter that affects the secondary aerosol formation (Tang et al., 2016). As shown in Fig. 3a-c, PM<sub>2.5</sub> was positively correlated with the atmospheric concentration of SO<sub>2</sub> and NO<sub>x</sub> ( $P < 0.001$ ) and RH ( $P < 0.05$ ). Noteworthy, all PM<sub>2.5</sub> samples yielded an OC/EC ratio  $> 2$  (Fig. 3d), which indicated the generation of secondary aerosols (Chow et al., 1994). While for Si<sub>PM<sub>2.5</sub></sub>, it was not significantly correlated with any of the secondary precursors or RH ( $P > 0.3$ ; Fig. 3e-g), again demonstrating that the Si<sub>PM<sub>2.5</sub></sub> was not affected by the secondary chemistry of aerosols. Furthermore, we also found that the Si<sub>PM<sub>2.5</sub></sub> was not correlated with the PM<sub>2.5</sub> concentration (Fig. S4, Supplement).



### 3.3 Effects of synthetic organosilicons and dry deposition

Based on the aforementioned results, we infer that Si should mainly be present in primary particles or act as nuclei during the formation of secondary particles. This was consistent with the non-volatile nature of most Si-containing compounds in the nature environment. An exception is synthetic low-mass organic Si compounds (e.g., methylsiloxanes (Schweigkofler and Niessner, 1999; Ahrens et al., 2014; Xu et al., 2012)) that may be able to transform into aerosol nanoparticles via hydroxyl radicals-mediated oxidative cleavage of Si-C bonds in the upper atmosphere (Atkinson, 1991; Sommerlade et al., 1993; Atkinson et al., 1995; Wu and Johnston, 2017). However, previous studies on the impact of low-mass organosilicons on the air quality have demonstrated that organosilicons do not contribute to the lower atmosphere aerosol formation (Graiver et al., 2003; Sommerlade et al., 1993). As a result, the U.S. Environmental Protection Agency (EPA) has excluded organosilicons from the regulation concerning restriction of VOCs in the atmosphere. Recently, Janecek et al. also found that the oxidative product concentration of cyclic volatile methylsiloxanes was ~10-40 times lower than their parent compounds in the U.S. (Janecek et al., 2017). While, another study by Wu et al. reported that secondary aerosol yields of siloxane oxidation might reach 15% (Wu and Johnston, 2017), suggesting that the contribution of organosilicons to secondary aerosols may vary in different regions. Thus, it may be necessary to consider the secondary Si production to determine whether it should be included in the secondary aerosol calculation. Note that the secondary Si production depends on the atmospheric concentration of both organosilicons and hydroxyl radicals. Up to now, no data about atmospheric organosilicons in the studied region was reported. Despite that, some recent studies reported the measurement and modeling results of atmospheric hydroxyl radicals in the North China Plain (Tan et al., 2017; Tham et al., 2016). The daily maximum concentration of atmospheric hydroxyl radicals was in the range of  $(5 - 15) \times 10^6 \text{ cm}^{-3}$  ( $\sim 10^{-10} \text{ mol m}^{-3}$ ). Since hydroxyl radical is a crucial condition for the siloxane oxidation, the extremely low concentration of hydroxyl radicals in the atmosphere could greatly limit the secondary Si production from siloxanes in the studied region, suggesting that the secondary Si production could be negligible compared with the high-abundance Si in primary particles. This deduction has also been experimentally verified by the irrelevance of  $\text{Si}_{\text{PM}_{2.5}}$  with the secondary chemistry of aerosols in Figs. 2 and 3.

We also considered the influence of dry deposition on the Si abundance in  $\text{PM}_{2.5}$ . The dry deposition is a size-dependent process that can cause coarse particles to fall more quickly than fine particles. However, it was found that the average Si abundance did not change significantly when the particle size was smaller than  $2.5 \mu\text{m}$  (Tan et al., 2016). That is to say, for fine particles ( $< 2.5 \mu\text{m}$ ), the dry deposition rate would not significantly affect the Si abundance. Thus, the Si-dilution effect with  $\text{PM}_{2.5}$  should be predominantly controlled by the secondary aerosol formation.

### 3.4 Secondary aerosol estimate using the Si-dilution method

The special role of Si in  $\text{PM}_{2.5}$  enables us to gain insights into the secondary particles in  $\text{PM}_{2.5}$  by using Si as a new conservative tracer. Here, as an example, we show the estimation of secondary aerosol contribution to  $\text{PM}_{2.5}$ , which was thought as a complicated and difficult task. Providing that the  $\text{Si}_{\text{PM}_{2.5}}$  keeps unchanged during the secondary growth of  $\text{PM}_{2.5}$ ,



the secondary formation can cause a dilution of the Si abundance in  $PM_{2.5}$ . Thus, it is possible to estimate the secondary aerosol contribution to  $PM_{2.5}$  ( $f_{sec}$ ) simply by the dilution factor (see Eq. (3) in the Sect. 2.5). To this end, the theoretical Si abundance in primary particles ( $C_{pri}$ ) and the final Si abundance in  $PM_{2.5}$  ( $C_{PM_{2.5}}$ ) need to be known. The  $C_{PM_{2.5}}$  can be directly measured with the collected  $PM_{2.5}$  samples. In this study, an annual mean value of 1.56% ( $n = 63$ ) was obtained. For the  $C_{pri}$ , it can derive from the mixed Si abundance of primary sources based on individual Si abundance and emission amounts of primary sources: the former can be measured with the collected primary source samples, and the latter can be given by a public emission inventory (e.g., MEIC database). It should be noted that the emission inventory does not include the emission amount of natural primary sources (i.e., dusts) (MEIC), which can be estimated by the isotopic mass balance of Si in  $PM_{2.5}$  based on the emission data of other sources and Si isotopic signatures of primary sources (see Table 1) (Lu et al., 2018). The whole estimation procedures are shown in Fig. 4a. In this way, the theoretical mean Si abundance in primary particles was calculated to be 7.51%. That is, the theoretical Si abundance in primary particles (7.51%) was diluted to be 1.56% by the secondary aerosol formation. Based on these values, the mean contribution of secondary particles to  $PM_{2.5}$  on haze days in 2013 in the studied region was easily calculated to be 79.2% (Fig. 4b). In the same way, we have also estimated the secondary aerosol contribution to  $PM_{2.5}$  in different seasons (Fig. 4c). Notably, the method also keeps the possibility of being modified to be applicable for the cases when the secondary Si production is not negligible by including the secondary Si production in the mass balance calculations.

### 3.5 Comparison with the traditional method

The uncertainty of the estimate result could be obtained by the error propagation calculation (Ku, 1966; Larsen et al., 2012) from the errors of variables used in the calculation (see the Sect. 2.6). In this way, the uncertainty of the annual mean SA contribution in 2013 was calculated to be 26.1%. As a comparison, the traditional method uses much more variables (e.g.,  $NH_4^+$ ,  $NO_3^-$ ,  $SO_4^{2-}$ , EC, and OC) in the calculation, which may bring in huge uncertainties to the result. For example, Schmid et al. reported that the RSD in the EC measurement could reach 36.6–45.5% (Schmid et al., 2001), which would cause large uncertainties in calculating the ratio of OC/EC and consequently cause the SOC to be biased by up to 64% (Guo et al., 2014). Meanwhile, the approach to estimate SOA by multiplying the SOC by an empirical coefficient is still controversial (Docherty et al., 2008). Therefore, considering that the Si concentration and isotopic composition can be accurately and precisely measured by ICP-MS, the present method should be more straightforward and precise than the traditional method that uses multiple secondary tracers.

We also compared the estimation results of the Si-dilution method with the traditional multi-tracer method (Table S1, Supplement). As shown in Fig. 4, the SA contribution estimated by the Si-dilution method was slightly higher than that obtained by the traditional method. Such a difference can be explained by the different strategies of the methods: the traditional method is based on limited rather than all active secondary species in  $PM_{2.5}$ , and thus the SA may be prone to be underestimated; while, the Si-dilution method only uses an inactive species (i.e., Si), and possible loss of Si-containing



components during the sample pretreatment procedures may cause the result to be overestimated. Despite that, the results between the two methods were actually very close (taking into account the uncertainties), thus proving the accuracy of the Si-dilution method.

### 3.6 Sources of secondary particle precursors revealed by Si isotopic signatures

5 In addition to the SA formation estimation, the correlation analysis of Si isotopic composition of  $PM_{2.5}$  with the SA can further reveal the sources of the precursors of SA. Note that the Si isotopic composition is independent of Si abundance of  $PM_{2.5}$ , and it can actually reflect the primary sources of  $PM_{2.5}$ , because different primary sources have different Si isotopic signatures (e.g., coal burning and industrial emission are  $^{30}\text{Si}$ -depleted sources, while vehicle emission and dusts are  $^{30}\text{Si}$ -enriched sources) (Lu et al., 2018). From Fig. S5, Supplement, the secondary aerosols were negatively correlated with the  
10  $\delta^{30}\text{Si}$  of  $PM_{2.5}$ , suggesting that  $^{30}\text{Si}$ -depleted primary sources (e.g., coal burning and industrial emission) contributed more precursors to secondary aerosols than  $^{30}\text{Si}$ -enriched sources. Specifically, from Fig. 5a, the  $\delta^{30}\text{Si}$  showed a decline trend with the increase in  $\text{SO}_4^{2-}$  concentration ( $P = 0.006$ ), suggesting that the  $^{30}\text{Si}$ -depleted primary sources contributed more precursor (i.e.,  $\text{SO}_2$ ) to the  $\text{SO}_4^{2-}$  species than  $^{30}\text{Si}$ -enriched sources. This was consistent with the fact that  $\text{SO}_2$  mainly derives from coal burning that is a  $^{30}\text{Si}$ -depleted source (Lu et al., 2018). The  $\text{NH}_3^+$  and SOC concentration was also negatively correlated  
15 with the  $\delta^{30}\text{Si}$  (Fig. 5c and 5d), suggesting that the precursors of these secondary species were also largely contributed by  $^{30}\text{Si}$ -depleted primary sources. However, the  $\text{NO}_3^-$  concentration did not show any correlation with the  $\delta^{30}\text{Si}$  ( $P = 0.63$ ; Fig. 5b). This could be explained by the fact that the sources of  $\text{NO}_3^-$  were more uncertain and involved both  $^{30}\text{Si}$ -enriched (e.g., vehicle exhaust) and  $^{30}\text{Si}$ -depleted primary sources (e.g., power plants) (Seinfeld and Pandis, 2006). Accordingly, we show that the Si isotopic composition not only indicates the primary sources, but also can reveal the sources of precursors of  
20 secondary species of  $PM_{2.5}$ .

### 3.7 Implications for air pollution control policies

Regardless of the different sampling date, the SA contribution obtained here (79.2% for the entire 2013 and 88.7% for Jan 2013; Fig. 4) was overall higher than that in a previous report (30-77% for four cities in China in Jan 2013) (Huang et al., 2014). Noteworthy, these results have been verified by two independent methods (Fig. 4). Our results suggest that the  
25 secondary aerosol contribution to Beijing's severe haze pollution, although has been known to be very high (Huang et al., 2014), might still be underestimated in the past. Thus, more strict policies should be enforced to reduce the emission of secondary particle precursors including  $\text{SO}_2$ ,  $\text{NO}_x$ , and VOCs for haze pollution control, especially for  $\text{NO}_x$  and VOCs which have not aroused full attention in current pollution control strategies (Wang et al., 2013). Furthermore, the  $^{30}\text{Si}$ -depleted primary sources (e.g., coal burning and industrial emission) should also be priorly regulated because they make a great  
30 contribution to both primary particles and secondary particle precursors.



#### 4 Conclusions

In summary, we have investigated the role of Si during aerosol secondary formation and shown that the high abundance and chemical inactiveness of Si make it a new conservative tracer to investigate the aerosol formation process. Based on the Si-dilution effect, we have proposed a new method to estimate the secondary aerosol contribution to PM<sub>2.5</sub> by using Si as a single tracer. In addition, the sources of the precursors of secondary aerosols can be revealed by the correlation analysis of secondary aerosols with the Si isotopic composition of PM<sub>2.5</sub>. Overall, the unique role of Si may open up a new strategy for aerosol research. Notably, some measures may further improve the accuracy of the method, such as increasing the size and representativeness of primary source sample sets. Considering the inter-regional transport of PM<sub>2.5</sub>, future efforts will also be made to include primary sources and emission inventory of adjacent cities/regions in the calculation.

#### 10 Supplement

The Supplement related to this article is available online at <https://doi.org/10.5194/acp-2018-914>.

#### Author contributions

QL designed the research; GJ supervised the project; DL performed most of experiments; JT measured the concentration of secondary species and calculated the secondary aerosol contribution using the traditional method; XY helped with the measurements; XS measured the atmospheric concentration of SO<sub>2</sub> and NO<sub>x</sub> and RH; QL and DL analyzed the data and wrote the paper.

#### Competing interests

The authors declare that they have no conflict of interest.

#### Acknowledgements

20 This work was financially supported by the National Natural Science Foundation of China (No. 91543104 and 21825403), the Chinese Academy of Sciences (XDB14010400, QYZDB-SSW-DQC018), and the National Basic Research Program of China (2015CB931903, 2015CB932003).



## References

- Ahrens, L., Harner, T., and Shoeib, M.: Temporal variations of cyclic and linear volatile methylsiloxanes in the atmosphere using passive samplers and high-volume air samplers, *Environ. Sci. Technol.*, 48, 9374-9381, 10.1021/es502081j, 2014.
- Atkinson, R.: Kinetics of the gas-phase reactions of a series of organosilicon compounds with OH and NO<sub>3</sub> Radicals and O<sub>3</sub> at 297 ± 2 K, *Environ. Sci. Technol.*, 25, 863-866, Doi 10.1021/Es00017a005, 1991.
- Atkinson, R., Tuazon, E. C., Kwok, E. S. C., Arey, J., Aschmann, S. M., and Bridier, I.: Kinetics and products of the gas-phase reactions of (CH<sub>3</sub>)<sub>4</sub>Si, (CH<sub>3</sub>)<sub>3</sub>SiCH<sub>2</sub>OH, (CH<sub>3</sub>)<sub>3</sub>SiOSi(CH<sub>3</sub>)<sub>3</sub> and (CD<sub>3</sub>)<sub>3</sub>SiOSi(CD<sub>3</sub>)<sub>3</sub> with Cl atoms and OH radicals, *J. Chem. Soc. Faraday Trans.*, 91, 3033-3039, Doi 10.1039/Ft9959103033, 1995.
- Basile-Doelsch, I.: Si stable isotopes in the Earth's surface: A review, *J. Geochem. Explor.*, 88, 252-256, 10.1016/j.gexplo.2005.08.050, 2006.
- Bi, X. H., Feng, Y. C., Wu, J. H., Wang, Y. Q., and Zhu, T.: Source apportionment of PM<sub>10</sub> in six cities of northern China, *Atmos. Environ.*, 41, 903-912, 10.1016/j.atmosenv.2006.09.033, 2007.
- Bzdek, B. R., Horan, A. J., Pennington, M. R., Janecek, N. J., Baek, J., Stanier, C. O., and Johnston, M. V.: Silicon is a frequent component of atmospheric nanoparticles, *Environ. Sci. Technol.*, 48, 11137-11145, 10.1021/es5026933, 2014.
- Chow, J. C., Watson, J. G., Fujita, E. M., Lu, Z. Q., Lawson, D. R., and Ashbaugh, L. L.: Temporal and spatial variations of PM<sub>2.5</sub> and PM<sub>10</sub> aerosol in the Southern California air quality study, *Atmos. Environ.*, 28, 2061-2080, Doi 10.1016/1352-2310(94)90474-X, 1994.
- Docherty, K. S., Stone, E. A., Ulbrich, I. M., DeCarlo, P. F., Snyder, D. C., Schauer, J. J., Peltier, R. E., Weber, R. J., Murphy, S. M., Seinfeld, J. H., Grover, B. D., Eatough, D. J., and Jimenez, J. L.: Apportionment of Primary and Secondary Organic Aerosols in Southern California during the 2005 Study of Organic Aerosols in Riverside (SOAR-1), *Environ. Sci. Technol.*, 42, 7655-7662, 10.1021/es8008166, 2008.
- Georg, R. B., Reynolds, B. C., Frank, M., and Halliday, A. N.: New sample preparation techniques for the determination of Si isotopic compositions using MC-ICPMS, *Chem. Geol.*, 235, 95-104, 10.1016/j.chemgeo.2006.06.006, 2006.
- Graiver, D., Farminer, K. W., and Narayan, R.: A review of the fate and effects of silicones in the environment, *J. Polym. Environ.*, 11, 129-136, Doi 10.1023/A:1026056129717, 2003.
- Guan, D. B., Klasen, S., Hubacek, K., Feng, K. S., Liu, Z., He, K. B., Geng, Y., and Zhang, Q.: Determinants of stagnating carbon intensity in China, *Nat. Clim. Change*, 4, 1017-1023, 10.1038/Nclimate2388, 2014.
- Guo, S., Hu, M., Guo, Q. F., and Shang, D. J.: Comparison of Secondary Organic Aerosol Estimation Methods, *Acta Chim Sinica*, 72, 658-666, 10.6023/A14040254, 2014.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prevot, A. S. H., Seinfeld, J. H., Surratt, J. D.,



- Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155-5236, DOI 10.5194/acp-9-5155-2009, 2009.
- Huang, R. J., Zhang, Y., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., El Haddad, I., and Prevot, A. S.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218-222, 10.1038/nature13774, 2014.
- Janecek, N. J., Hansen, K. M., and Stanier, C. O.: Comprehensive atmospheric modeling of reactive cyclic siloxanes and their oxidation products, *Atmos. Chem. Phys.*, 17, 8357-8370, 2017.
- 10 Jiang, X. J., Zhang, Q., Zhao, H. Y., Geng, G. N., Peng, L. Q., Guan, D. B., Kan, H. D., Huo, H., Lin, J. T., Brauer, M., Martin, R. V., and He, K. B.: Revealing the Hidden Health Costs Embodied in Chinese Exports, *Environ. Sci. Technol.*, 49, 4381-4388, 10.1021/es506121s, 2015.
- Ku, H. H.: Notes on the use of propagation of error formulas, *J. Res. Natl. Bureau Stand.*, 70C, 263-273, 1966.
- Larsen, B. R., Gilardoni, S., Stenstrom, K., Niedzialek, J., Jimenez, J., and Belis, C. A.: Sources for PM air pollution in the Po Plain, Italy: II. Probabilistic uncertainty characterization and sensitivity analysis of secondary and primary sources, *Atmos. Environ.*, 50, 203-213, 10.1016/j.atmosenv.2011.12.038, 2012.
- 15 Li, M., Zhang, Q., Kurokawa, J., Woo, J. H., He, K. B., Lu, Z. F., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y. F., Hong, C. P., Huo, H., Jiang, X. J., Kang, S. C., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, 10.5194/acp-17-935-2017, 2017.
- 20 Lin, J. T., Pan, D., Davis, S. J., Zhang, Q., He, K. B., Wang, C., Streets, D. G., Wuebbles, D. J., and Guan, D. B.: China's international trade and air pollution in the United States, *Proc. Natl. Acad. Sci. U. S. A.*, 111, 1736-1741, 10.1073/pnas.1312860111, 2014.
- Lu, D., Liu, Q., Yu, M., Yang, X., Fu, Q., Zhang, X., Mu, Y., and Jiang, G.: Natural Silicon Isotopic Signatures Reveal the Sources of Airborne Fine Particulate Matter, *Environ. Sci. Technol.*, 52, 1088-1095, 2018.
- 25 Multi-resolution Emission Inventory for China. <http://meicmodel.org/>.
- Pope, C. A., 3rd, Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *JAMA*, 287, 1132-1141, 2002.
- Pope, C. A., 3rd, Ezzati, M., and Dockery, D. W.: Fine-particulate air pollution and life expectancy in the United States, *N. Engl. J. Med.*, 360, 376-386, 10.1056/NEJMsa0805646, 2009.
- 30 Savage, P. S., Armytage, R. M. G., Georg, R. B., and Halliday, A. N.: High temperature silicon isotope geochemistry, *Lithos*, 190, 500-519, 10.1016/j.lithos.2014.01.003, 2014.



- Schmid, H., Laskus, L., Abraham, H. J., Baltensperger, U., Lavanchy, V., Bizjak, M., Burba, P., Cachier, H., Crow, D., Chow, J., Gnauk, T., Even, A., ten Brink, H. M., Giesen, K. P., Hitzengerger, R., Hueglin, C., Maenhaut, W., Pio, C., Carvalho, A., Putaud, J. P., Toom-Sauntry, D., and Puxbaum, H.: Results of the "carbon conference" international aerosol carbon round robin test stage I, *Atmos. Environ.*, 35, 2111-2121, Doi 10.1016/S1352-2310(00)00493-3, 2001.
- 5 Schweigkofler, M., and Niessner, R.: Determination of siloxanes and VOC in landfill gas and sewage gas by canister sampling and GC-MS/AES analysis, *Environ. Sci. Technol.*, 33, 3680-3685, Doi 10.1021/Es9902569, 1999.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2<sup>nd</sup> Ed, Wiley, 2006.
- Sommerlade, R., Parlar, H., Wrobel, D., and Kochs, P.: Product analysis and kinetics of the gas-phase reactions of selected  
10 organosilicon compounds with OH radicals using a smog chamber-mass spectrometer system, *Environ. Sci. Technol.*, 27, 2435-2440, Doi 10.1021/Es00048a019, 1993.
- Tan, J., Duan, J., Zhen, N., He, K., and Hao, J.: Chemical characteristics and source of size-fractionated atmospheric particle in haze episode in Beijing, *Atmos. Res.*, 167, 24-33, 2016.
- Tan, Z., Fuchs, H., Lu, K., Hofzumahaus, A., Bohn, B., Broch, S., Dong, H., Gomm, S., Häsel, R., He, L., Holland, F., Li,  
15 X., Liu, Y., Lu, S., Rohrer, F., Shao, M., Wang, B., Wang, M., Wu, Y., Zeng, L., Zhang, Y., Wahner, A., and Zhang, Y.: Radical chemistry at a rural site (Wangdu) in the North China Plain: observation and model calculations of OH, HO<sub>2</sub> and RO<sub>2</sub> radicals *Atmos. Chem. Phys.*, 17, 663-690, 2017.
- Tang, M., Cziczo, D. J., and Grassian, V. H.: Interactions of water with mineral dust aerosol: Water adsorption, hygroscopicity, cloud condensation, and ice nucleation, *Chem. Rev.*, 116, 4205-4259, 10.1021/acs.chemrev.5b00529, 2016.
- 20 Tham, Y. J., Wang, Z., Li, Q., Yun, H., Wang, W., Wang, X., Xue, L., Lu, K., Ma, N., Bohn, B., Li, X., Kecorius, S., Größ J., Shao, M., Wiedensohler, A., Zhang, Y., and Wang, T.: Significant concentrations of nitryl chloride sustained in the morning: investigations of the causes and impacts on ozone production in a polluted region of northern China, *Atmos. Chem. Phys.*, 16, 14959-14977, 2016.
- Tian, S. L., Pan, Y. P., and Wang, Y. S.: Size-resolved source apportionment of particulate matter in urban Beijing during  
25 haze and non-haze episodes, *Atmos. Chem. Phys.*, 16, 1-19, 2016.
- Wang, Y., Zhang, Q. Q., He, K., Zhang, Q., and Chai, L.: Sulfate-nitrate-ammonium aerosols over China: response to 2000-2015 emission changes of sulfur dioxide, nitrogen oxides, and ammonia, *Atmos. Chem. Phys.*, 13, 2635-2652, 10.5194/acp-13-2635-2013, 2013.
- Watson, J. G., Chow, J. C., and Chen, L.-W. A.: Summary of organic and elemental carbon/black carbon analysis methods  
30 and intercomparisons, *Aerosol Air Qual. Res.*, 5, 65-102, 2005.
- Wu, Y., and Johnston, M. V.: Aerosol formation from OH oxidation of the volatile cyclic methyl siloxane (cVMS) decamethylcyclopentasiloxane, *Environ. Sci. Technol.*, 51, 4445-4451, 2017.



- Xu, L., Shi, Y. L., Wang, T., Dong, Z. R., Su, W. P., and Cai, Y. Q.: Methyl siloxanes in environmental matrices around a siloxane production facility, and their distribution and elimination in plasma of exposed population, *Environ. Sci. Technol.*, 46, 11718-11726, 10.1021/es3023368, 2012.
- Yang, J., Fu, Q., Guo, X., Chu, B., Yao, Y., Teng, Y., and Wang, Y.: Concentrations and seasonal variation of ambient PM<sub>2.5</sub> and associated metals at a typical residential area in Beijing, China, *Bull. Environ. Contaminat. Toxicol.*, 94, 232-239, 2015.
- 5 Zambardi, T., and Poitrasson, F.: Precise determination of silicon isotopes in silicate rock reference materials by MC-ICP-MS, *Geostand. Geoanal. Res.*, 35, 89-99, 10.1111/j.1751-908X.2010.00067.x, 2011.
- Zhang, Q., He, K. B., and Huo, H.: Cleaning China's air, *Nature*, 484, 161-162, 2012a.
- Zhang, R., Khalizov, A., Wang, L., Hu, M., and Xu, W.: Nucleation and growth of nanoparticles in the atmosphere, *Chem. Rev.*, 112, 1957-2011, 10.1021/cr2001756, 2012b.
- 10 Zhang, R., Wang, G., Guo, S., Zamora, M. L., Ying, Q., Lin, Y., Wang, W., Hu, M., and Wang, Y.: Formation of urban fine particulate matter, *Chem. Rev.*, 115, 3803-3855, 10.1021/acs.chemrev.5b00067, 2015.
- Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, *Atmos. Chem. Phys.*, 15, 2031-2049, 10.519
- 15



## Tables

**Table 1. Parameters for primary sources used in the estimation of secondary aerosol contribution to PM<sub>2.5</sub>.**

	Coal combustion	Industrial emission	Biomass burning	Vehicle exhaust	Dusts
Emission mass ( $m_i$ ; Mg/year)	43813 <sup>a</sup>	34111 <sup>a</sup>	27476 <sup>b</sup>	5379 <sup>a</sup>	65199 <sup>c</sup>
Si abundance ( $C_i$ ) <sup>d</sup>	8.06%	0.79%	1.01%	0.2%	14%
Si isotopic signature ( $\delta^{30}\text{Si}_i$ ) <sup>d</sup>	-2.4‰	-1.4‰	-0.50‰	1.0‰	-0.28‰

<sup>a</sup> Data from the MEIC database (MEIC).


<sup>b</sup> Since the MEIC database does not include data for biomass burning, the  $m_i$  of biomass burning was estimated from the data of other sources according to source apportionment results of 2013 reported previously (Tian et al., 2016).

<sup>c</sup> In the calculation, soil, construction, and urban fugitive dusts are treated as a single source considering their similarity in Si abundance and Si isotopic signatures (Lu et al., 2018). The  $m_i$  of dusts was calculated by using isotopic mass balance of Si in PM<sub>2.5</sub> (see Methodology).

<sup>d</sup> The Si abundance and natural Si isotopic signatures of primary sources around Beijing are adopted from a previous study (Lu et al., 2018).




## Figures




**Figure 1.** Scheme showing the role of Si during the secondary formation process of aerosol particles. *Note:* this scheme only reflects the role of Si but does not show all reactions involved in the secondary aerosol formation.

5




**Figure 2.** Correlation analysis of PM<sub>2.5</sub> concentration and its Si content (Si<sub>PM<sub>2.5</sub></sub>) with the secondary species in PM<sub>2.5</sub>. (a)-(d), Correlation of PM<sub>2.5</sub> concentration with SO<sub>4</sub><sup>2-</sup> (a), NO<sub>3</sub><sup>-</sup> (b), NH<sub>4</sub><sup>+</sup> (c), and SOC (d). (e)-(h), Correlation of Si<sub>PM<sub>2.5</sub></sub> with SO<sub>4</sub><sup>2-</sup> (e), NO<sub>3</sub><sup>-</sup> (f), NH<sub>4</sub><sup>+</sup> (g), and SOC (h).


5



**Figure 3.** Correlation analysis of PM<sub>2.5</sub> concentration and Si<sub>PM2.5</sub> with typical secondary aerosol precursors and relative humidity (RH). (a)-(d), Correlation of PM<sub>2.5</sub> concentration with SO<sub>2</sub> (a), NO<sub>x</sub> (b), RH (c) and OC/EC ratio (d). (d)-(f), Correlation of Si<sub>PM2.5</sub> with SO<sub>2</sub> (e), NO<sub>x</sub> (f), and RH (g) and OC/EC ratio (h). The red dotted lines in (d) and (h) represent OC/EC = 2, which can be used to indicate the formation of secondary aerosols at OC/EC > 2 (Chow et al., 1994).



**Figure 4.** Comparison of the SA estimate results between the Si-dilution method and the traditional method. **(a)**, A flow chart showing how the  $f_{sec}$  was obtained by the Si-dilution method. **(b)**, Annual mean SA contribution to  $PM_{2.5}$  in Beijing. **(c)**, Seasonal mean SA contribution in Beijing. *Note:* the error bars for the Si-dilution method include both method uncertainty and the variations of samples on different dates (see Sect. 2.6); while for the traditional method the error bars only represent the variations of samples on different dates (without method uncertainty). Please see text for a more detailed discussion on the uncertainties.



**Figure 5.** Correlation analysis of Si isotopic composition ( $\delta^{30}\text{Si}$ ) of  $\text{PM}_{2.5}$  with the secondary species ( $\text{SO}_4^{2-}$  (a),  $\text{NO}_3^-$  (b),  $\text{NH}_4^+$  (c), and SOC (d)).