

## Response to Comment by Editor

**Co-Editor Decision: Reconsider after major revisions** (27 Sep 2019) by Pedro Jimenez-Guerrero

Comments to the Author:

Dear authors:

Please address the comments by the reviewers before the manuscript can be considered for final publication.

Non-public comments to the Author:

Dear authors:

One of the reviewers raise very important concerns regarding the scientific knowledge gain and the skill of the simulations. I share some of these concerns. The limitations pointed out by the reviewer are well established and supported, so I would beg you to address them before the manuscript can be considered for final publication.

### **Reply:**

**Thank you for your consideration of our manuscript and for your comments and suggestions. We have carefully considered yours and the additional comments from the two reviewers, responded to their comments and suggestions in the attached point-by-point responses, and incorporated revisions in the manuscript to address the shortcomings they noted. In particular to address the Reviewer #2's concern on the possible impacts of the model grid resolution on resolving STT dynamics and events, we conducted two additional sets of WRF model simulations over the contiguous U.S. using 36-km and 12-km grid spacing, respectively. We analyzed and compared the vertical and temporal variations in the estimated potential vorticity (PV) fields at the locations of the ozonesonde sites which are the focus of the analyses; the results are summarized in three additional figures: S7-S9 in the supplemental information and associated brief discussion in the manuscript main text. As detailed in our response to reviewer #2, we find that for simulations at the 3 grid resolutions examined (108-km, 36-km, and 12-km), time-height variations in PV at these locations are largely similar and so is the estimated altitude of the tropopause (i.e., altitude of 2PVU) which is used in our analysis. As shown by the similarity of the altitude of the 2PVU estimated across the 3 different resolutions, the interpretation of STT events in our analysis is not strongly influenced by the lack of resolution in our original 108-km calculations, as speculated by Reviewer #2. This is not completely surprising since all model calculations employed assimilation of analyzed meteorological fields (constrained by prior observations) in the model's UTLS. Consequently, the STT dynamics and capturing of specific events using the PV criteria are similar. Finally, a comparison of O<sub>3</sub> at the model top layer also show good correspondence in the magnitude of O<sub>3</sub> at the model top using**

PV estimates from the 3 different resolutions. As shown by the additional information in the new Figures S7-S9, the robustness of our results and interpretations are not strongly influenced by perceived lack in horizontal grid resolution. We thank the reviewer for raising this important consideration, addressing which has helped strengthen the analysis presented and the associated robustness of the approach and results. We believe that these additions and revisions to the manuscript now address all the points raised by the reviewers as well as your concerns.

We trust that our response to the reviewer comments and the revisions incorporated in the manuscript meet with your criteria for acceptance. Thank you for your consideration of our manuscript for publication in *Atmospheric Chemistry and Physics*.

## Response to Referee Comment 1 by Anonymous Referee #1

The manuscript has been improved significantly after revision especially for the discussions on tropopause, O3/PV tracer, and O3/PV ratios. The authors have addressed all my comments and I'm satisfied with the authors' responses. However, there are still a couple of technical issues needed to be corrected.

### **Reply:**

**We thank the reviewer for providing helpful and constructive comments. We have revised our manuscript according to the reviewer's technical comments and suggestions. We believe that these revisions address all points raised by the reviewer. Our point-by-point responses are provided below, and revisions are indicated in blue in the revised manuscript.**

Technical errors:

Page 5, line 22, "...have been reported (Hoering et al., 1991)", no need "

### **Reply:**

**We thank the reviewer for catching the typo. The extra quote mark has been removed.**

Page 13, line 25, "These stratospheric intrusion are..."

### **Reply:**

**We have corrected as "These stratospheric intrusions are".**

Page 14, line 28-32, "tropospheric o3 column mass" is not a very accurate term here. same for Figure 10.

### **Reply:**

**We have revised this term as "tropospheric O<sub>3</sub> column".**

## Response to Referee Comment 2 by Anonymous Referee #2

The manuscript improved a lot, thanks for that!

Nevertheless, the issue with the horizontal resolution is still a major weakness of this article. As detailed below (item 2+3), both articles cited in my first review do not, at least from my point of view, support the hypothesis of this article (top page 5), that the CMAQ model with 108 km grid spacing is able to capture SST events sufficiently. Additionally, I am not convinced that the provided evaluation shows that the SST events are captured correctly: the averaged SST maybe, but not the events. As the result can change considerably simply due to employing by decreasing the horizontal resolution, the scientific gain from this study is very, very limited. I understand, that it is an issue of computer resources, nevertheless, some studies can simply not be performed, if the resources are not available.

### **Reply:**

**We thank the reviewer for providing helpful and constructive comments. We have revised our manuscript according to the reviewer's technical comments and suggestions. In particular, to address the reviewer's concern on lack of adequate resolution in our model calculations, we have now also included results from new WRF simulations over the continental U.S. employing a finer resolution to examine the possible impacts of resolution on representing dynamics for the SST events analyzed at the locations of the ozonesonde launches – the results and changes incorporated in the manuscript are further detailed in response to the reviewer's specific comments below. We believe that these revisions address all points raised by the reviewer. Our point-by-point responses are provided below, and revisions are indicated in blue in the revised manuscript.**

There are some remaining issues I'd like to point to:

1. Title: the title sounds a little bit weird. Leaving out the first words it reads “stratospheric intrusion of tropospheric ozone [...]”. This seems wrong. I suggested to delete the word “tropospheric” from the title, as “trans-pacific transport” includes, at least for me, that it is a tropospheric process.

### **Reply:**

**Following the reviewer's suggestion we have revised the title as follows:**

**“Modeling Stratospheric Intrusion and Trans-Pacific Transport on Tropospheric Ozone using Hemispheric CMAQ during April 2010: Part 1. Model Evaluation and Air Mass Characterization for Stratosphere-Troposphere Transport”**

2. p. 4 ll. 27-30: In my opinion the results of Cristofanelli et al. are summarized in a wrong way. The important outcome is, that the models with lagrangian transport scheme were able to capture SST, but the Eulerians were not. Unfortunately, the models with lagrangian transport schemes had a resolution of 1x1 degree and the Eulerian models had a much coarser resolution. But the Cristofanelli publication does not provide any evidence, that an Eulerian model in 1x1 degree would be able to capture SST.

**Reply:**

**We thank the reviewer for this comment and have re-examined the results and discussion in Cristofanelli et al. We have revised this discussion to reflect that their results suggested that models with a lagrangian transport scheme were able to capture SST. The revised sentence is as follows:**

**“STT was analyzed by combining analysis of data from a measurement network and predictions from total of seven model simulations over Europe, and reported the advantages of lagrangian models in capturing the STT (Cristofanelli et al., 2003).”**

3. p. 4 ll. 30-32: The study of Gery (2003), as pointed out explicitly within the article, is only valid for the Unified model of the UK Met Office (UM). Gery (2003) explicitly points out that the results very much depend on the convection and diffusion parametrisation used in the respective model. Therefore one can not simply assume that all models are able to capture SST events in a horizontal resolution, just because the UM is able to do so. In principle each model needs to be evaluated from which horizontal resolution on it is able to capture SST reasonable. I do not know such an evaluation for CMAQ driven by WRF.

**Reply:**

**While the results of Gery (2003) are indicative of the resolution impacts, we acknowledge that they should be viewed as results from a single model for a specific region and that the combined impacts of transport formulation, parameterizations, and grid resolution should be examined in each application.**

**To address the reviewer’s concerns, we have conducted two additional WRF simulations over contiguous U.S. (CONUS) domain with 36 and 12 km grid resolutions but with all other physics configuration identical to that of the original 108 km simulation. The analyzed time-height curtain plot of PV is shown below and also included as Figure S7 in the revised supplemental material. As can be seen in these curtain plots, generally modeled PV fields at different resolutions are largely similar . To further quantify the impacts of differences across these resolutions on interpretation of SST, differences in the time-height PV profiles calculated by 36 km–108 km and 12 km–108 km are shown in Figs. S8 in the supplemental material. These results clarified that the higher (lower) PV at upper (lower)**


altitudes is enhanced (weakened) by increasing horizontal grid resolution. The difference from the 108 km simulation was much clearer in the 12 km simulation. As also shown by the similarity of the altitude of the 2PVU estimated across the 3 different resolutions, the interpretation of STT events in our analysis is not strongly influenced by the lack of resolution in our original 108-km calculations. This is not completely surprising since all model calculations employ assimilation of analyzed meteorological fields in the model's UTLS. Consequently, the STT dynamics and capturing of specific events using the PV criteria are similar. Finally, since in our chemical transport calculations we scale O<sub>3</sub> in the model's UTLS using the estimated PV fields, in Figure S9 of the revised manuscript we also present comparisons of the estimated O<sub>3</sub> in the uppermost layer based on the O<sub>3</sub>/PV relation used in this study. These comparisons also show good correspondence in the magnitude of O<sub>3</sub> at the model top using PV estimates from the 3 different resolutions.

Collectively, we believe that these additional analyses should address the reviewer's concerns on the robustness of our results due to the horizontal grid resolution employed. As shown by the additional information in the new Figures S7-S9, the robustness of our results and interpretations are not strongly influenced by perceived lack in resolution as speculated by the reviewer. We nevertheless thank the reviewer for raising this important consideration, addressing which has helped strengthen the analysis presented and the associated robustness of the approach and results.


The added figures are Figs. S7- S9 in the supplemental material, and the relevant sentences in the main manuscript have been modified as follows:

“To investigate the effect of horizontal grid resolution on the representation of STT, additional WRF simulations were conducted over CONUS domain with 36 and 12 km horizontal grid resolutions, and temporal and vertical variations in simulated PV fields across the different resolutions (108, 36, and 12 km) were compared. These results are shown in Fig. S7 in the supplemental material. Generally, the modeled PV fields estimated with different horizontal grid resolutions showed similar features. The differences are displayed in Fig. S8 in the supplemental material. It was revealed that higher (lower) PV at upper (lower) altitude is enhanced (weakened) by increasing horizontal grid resolution. As expected, larger differences are noted between the 108 km and 12 km fields than those between the 108 km and 36 km fields. Although the enhancement of PV at upper altitudes could lead to increase in estimated O<sub>3</sub> through the O<sub>3</sub>/PV relationship used in the model, no systematic differences are noted in the estimated O<sub>3</sub> in the model's UTLS across the three resolutions, at least at the ozonesonde observation sites where our analysis is focused. A comparison of the altitude of 2 PVU which is used to diagnose the tropopause is also plotted in Fig. S8. As noted by the similarity of the altitude of the 2PVU across the 3

different resolutions, the interpretation of STT events is not strongly influenced by the horizontal resolution employed in this study. This is because all model calculations employ assimilation of analyzed meteorological fields in the model's UTLS, resulting in comparable representation of STT events. Finally, a comparison of estimated  $O_3$  at the model top-layer based on the  $O_3/PV$  relation (Xing et al., 2016) by using different PV simulated from different horizontal grid resolutions is illustrated through scatter-plots in Fig. S9 in the supplemental material. These comparisons indicate good correspondence in the magnitude of  $O_3$  at the model top using PV estimates from the 3 different resolutions. At the Boulder site (Fig. S9 (b)), the use of finer grid resolutions could sometimes lead to higher  $O_3$  concentrations. Collectively, the comparisons in Figure S7-S9 suggest that the 108 km horizontal grid resolution in H-CMAQ modeling system in conjunction with the physics and data assimilation options employed in the driving WRF model can capture the variability in the PV fields and associated STT  $O_3$  impacts.”





**Figure S7. Curtain plots of modeled PV by (left) WRF over northern hemisphere with a 108 km horizontal grid resolution, (center) WRF over CONUS domain with a 36 km horizontal grid resolution, and (right) WRF over CONUS domain with a 12 km horizontal grid resolution at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI) during April 2010.**





**Figure S8. Curtain plots of (right) the difference of modeled PV calculated from 36 km–108 km, (center) from 12 km–108 km, and (right) modeled PV lines of 2 PVU by 108 km (red), 36 km (dark orange), and 12 km (light orange) at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI) during April 2010.**



(a) Trinidad Head





(b) Boulder





(c) Huntsville



(d) Wallops Island



(e) Rhode Island



**Figure S9. Correspondence of estimated O<sub>3</sub> concentration based on O<sub>3</sub>/PV relation at the uppermost layer as (left) 108 km vs. 36 km and (right) 108 km vs. 12 km at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI). Plots are hourly data during April 2010 (total number is 720).**

4. p. 9 ll. 19-30: Instead of mentioning the biomass burning as the first and thus most important reason, I would guess, that the underrepresentation of the STT events due to the coarse resolution is the major reason.

**Reply:**

**We have added the following sentence to indicate the horizontal grid resolution as a possible reason:**

**“Another possible reason may stem from the use of a coarse horizontal resolution.”**

**Because we cannot judge the priority of these reasons (based on the analysis presented in response to comment #3), we just mention the possible reasons here.**

5. p. 11 ll. 11-13: As above, I opt that a higher resolution in the horizontal would be much more promising than increasing vertical layering. (By the way, Gery (2003) points out, that increasing vertical resolution in his simulations decreased the vertical transport, which is contrary to the statement in the present article.)

**Reply:**

**As detailed in our response to comment #3, the analysis of WRF simulations at 36-km and 12-km horizontal resolutions do not necessarily suggest significant increase in skill in the diagnosis of STT events relative to our base calculation. Based on the discussion of these results in the revised manuscript we now have removed this earlier somewhat speculative sentence: “This is a hemispheric modeling system but the finer horizontal resolution will be another way to improve this.”.**

6. Section 3.2: SST events are connected with PV streamers and the stratospheric air mass is transported downward in the wake of the front, thus there is a non-negligible horizontal motion involved. I would expect that a high amount of stratospheric influenced air is discarded, just be characterizing the air column-wise from top to bottom.

**Reply:**

**As we have illustrated in our response to comment#3., the simulated dynamics associated with STT over the CONUS appear to be largely similar across the 3 different horizontal resolutions, primarily because of the assimilation of meteorological data in the UTLS. The slight enhancement of PV at higher altitude could lead to the increased O<sub>3</sub> concentrations during some times and might influence mid-tropospheric O<sub>3</sub> in some cases, though systematic differences in air mass motion were not detected based on the PV analysis.**

7. Fig.8: This flow-chart is not really a flow-chart. Somehow the information that the flowchart "cycles" over all columns or for one column only (and that the analysis is columnwise) needs to be

included. More importantly it should be indicated that, if the above grid box is denoting a stratospheric air mass, the vertical index is decrease (one box down in the column) and the analysis is continued for the next lower box. Currently, the same analysis would be repeated over and over again for the same grid box, as no index change is depicted in the flowchart. Last but not least, an outcome for each grid box, indicating stratospheric or not, should be explicitly shown.

**Reply:**

**This flowchart has been revised to explicitly show that repetition is conducted over layers from number 43 to number 1 (surface) by including the loop chart.**

Minor Issues

- p. 4 last sentence: what is the “tracer amount across tropopause”? Are you omitting the word transport here somehow?

**Reply:**

**This sentence has been revised as “the tracer transport across the tropopause”.**

- p. 5, ll. 20-22: This is not a sentence.

**Reply:**

**This sentence has been revised as “The tropopause altitude can be also diagnosed by the traditional approach based on the lapse rate (i.e., thermal tropopause) defined by World Meteorological Organization (WMO) (WMO, 1992), and comparisons with that diagnosed using PV (i.e., dynamical tropopause) have been reported (Hoering et al., 1991).”**

- p. 13, ll. 3/4: “The flowchart the air mass ...” ??? Is not a sentence.

**Reply:**

**We have corrected this typo as “The flowchart of the established air mass characterization method”**

# Modeling Stratospheric Intrusion and Trans-Pacific Transport on Tropospheric Ozone using Hemispheric CMAQ during April 2010: Part 1. Model Evaluation and Air Mass Characterization for Stratosphere-Troposphere Transport

5

Syuichi Itahashi<sup>1</sup>, Rohit Mathur<sup>2</sup>, Christian Hogrefe<sup>2</sup>, and Yang Zhang<sup>3</sup>

<sup>1</sup> Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry (CRIEPI), 1646 Abiko, Abiko, Chiba 270-1194, Japan

10 <sup>2</sup> Environmental Protection Agency (EPA), Computational Exposure Division, National Exposure Research Laboratory, Office of Research and Development, Research Triangle Park, NC 27711, U.S.A.

<sup>3</sup> Department of Marine, Earth, and Atmospheric Sciences (MEAS), North Carolina State University (NCSSU), Campus Box 8208, Raleigh, NC 27695, U.S.A.

*Correspondence to:* Syuichi Itahashi (isyuichi@criepi.denken.or.jp)

## Abstract.

15 **Stratospheric intrusion and** trans-Pacific transport has been recognized as a potential source of tropospheric ozone over the U.S.A. The state-of-the-science Community Multiscale Air Quality (CMAQ) Modeling System has recently been extended for hemispheric-scale modeling applications (referred to as H-CMAQ). In this study, H-CMAQ is applied to study the **stratospheric intrusion and trans-Pacific transport** during April 2010. The results will be presented in two continuous papers. In this part 1 paper, model evaluation for tropospheric ozone (O<sub>3</sub>) is presented. Observations at the surface, by ozonesondes and airplane, and by satellite across the northern hemisphere are used to evaluate the model performance for O<sub>3</sub>. H-CMAQ is able to capture surface and boundary layer (defined as surface to 750 hPa) O<sub>3</sub> with a normalized mean bias (NMB) of -10%; however, a systematic underestimation with an NMB up to -30% is found in the free troposphere (defined as 750-250 hPa). In addition, a new air mass characterization method is developed to distinguish influences of stratosphere-troposphere transport (STT) from the effects of photochemistry on O<sub>3</sub> levels. This method is developed based on the ratio of O<sub>3</sub> and an inert tracer indicating stratospheric O<sub>3</sub> to examine the importance of photochemistry, and sequential intrusion from upper layer. During April 2010 as monthly average, the relationship between surface O<sub>3</sub> mixing ratios and estimated stratospheric air masses in the troposphere show a slight negative slope, indicating that high surface O<sub>3</sub> values are primarily affected by other factors (i.e., emissions), whereas this relationship shows a slight positive slope at elevated sites, indicating that STT has a possible impact at elevated sites. STT shows large day-to-day variations, and STT impacts can either originate from the same air mass over the entire U.S.A. with an eastward movement found during early April, or stem from different air masses at different locations indicated during late April. Based on this newly established air mass characterization technique, this study can contribute to

20

25

30

understand the role of STT, and also the implied importance of emissions leading to high surface O<sub>3</sub>. Further research focused on emissions is discussed in a subsequent part 2 paper.

## 1 Introduction

Tropospheric ozone (O<sub>3</sub>) is a secondary air pollutant produced by a chain of reactions involving photochemical oxidation of volatile organic compounds (VOCs) in the presence of nitrogen oxides (NO<sub>x</sub>) (Haagen-Smit and Fox, 1954). Ozone plays a key role in tropospheric chemistry by controlling the oxidizing capacity through the production of hydroxyl (OH) radicals, and is an important greenhouse gas throughout the troposphere (Logan, 1985). Ground level O<sub>3</sub> poses significant risks to human health and therefore many countries regulate it as a criteria pollutant with an ambient air quality standard. In the U.S.A., the National Ambient Air Quality Standard (NAAQS) for O<sub>3</sub> is based on the annual 4<sup>th</sup> highest maximum daily 8-10 h concentration (MD8O3) averaged over three years and its threshold values have been decreasing from 80 ppbv in 1997 to 75 ppbv in 2008, and 70 ppbv in 2015 (EPA, 2018). Long-term trends of rural O<sub>3</sub> during 1990-2010 revealed significant O<sub>3</sub> decreases in the eastern U.S.A. during spring and summer whereas no significant O<sub>3</sub> decrease was found in the western U.S.A. during spring (Cooper et al., 2012). Analysis of trends in surface O<sub>3</sub> levels between 1998 and 2013 showed that the highest O<sub>3</sub> concentration in the U.S.A. has been reduced in response to substantial decline of precursor (Simon et al., 2015). It was also 15 shown that the O<sub>3</sub> concentration on low O<sub>3</sub> days have increased and led to the narrowing of the O<sub>3</sub> concentration range across the U.S.A.

From the viewpoint of global air quality changes, the dramatic variation of anthropogenic emissions in East Asia (Itahashi et al., 2013, 2014, 2015), may impact atmospheric composition at not only the local and regional scales, but also the global scale. By combining trajectory analysis with detailed chemical and meteorological data, it was suggested that the 20 emissions were lifted into the free troposphere over Asia and then transported to North America in about 5-8 days (Jaffe et al., 1999). Trans-Pacific transport has been studied over the past decade because of its potential impact on rising background O<sub>3</sub> concentrations (Cooper et al., 2010). Asian contributions to surface O<sub>3</sub> levels in the U.S.A. require an additional challenge to meeting more stringent NAAQS for O<sub>3</sub> (Fiore et al., 2002). A typical case of trans-Pacific transport occurred during the so-called “perfect dust storm” during April 2001, transporting Asian dust to North America (Huebert et al., 2003). From an air 25 pollutant perspective, it was reported that the impact of Asian emissions increased background concentration of O<sub>3</sub> by 1 ppbv (2.5%) on a monthly average basis and up to 2.5 ppbv on a daily average basis over the western U.S.A. in April 2001 (Wang et al., 2009). Background O<sub>3</sub> levels entering western North America in spring have increased by approximately 10 ppbv between 1984 and 2002 based on a compilation of observations over the west coast of the U.S.A., and the possible cause for this increase was thought to be Asian emission trends (Jaffe et al., 2003). Asian air pollution can enhance surface O<sub>3</sub> mixing 30 ratios by 5-7 ppbv over western North America in April-May 2006, and the doubled Asian anthropogenic emissions increase during 2000-2006 was estimated to have the impact by 1-2 ppbv (Zhang et al., 2008). The global model simulation assuming the tripling of Asian anthropogenic emissions from 1985 to 2010 indicated an increase O<sub>3</sub> mixing ratios by 2-6 ppbv in the

western U.S.A. and by 1-3 ppb in the eastern U.S.A. on a monthly-mean basis, with the maximum effect occurring in April-June; this increase was suggested to more than offset the benefits of 25% domestic reduction in the western U.S.A. (Jacob et al., 1999). Based on the Emission Database for Global Atmospheric Research (EDGAR) version 4.3.1, anthropogenic emissions of NO<sub>x</sub> and VOCs from China are estimated to have increased by 3.2 and 2.1 times during 1985-2010, respectively (Crippa et al., 2016), which is generally consistent with the assumption by Jacob et al. (1999).

The occurrence of trans-Pacific transport can be inferred from variations in the jet stream related to La Niña and El Niño. The springtime Asian outflow may be enhanced following an El Niño winter due to the eastward extension of the atmospheric circulation over the Pacific-North America sector and the southward shift of the subtropical jet stream (Koumoutsaris et al., 2008; Lin et al., 2015). According to the NOAA Climate Prediction Center (CPC), 2009–2010 wintertime was influenced by strong El Niño conditions (NOAA, 2018). Because of the favorable condition for trans-Pacific transport, it was reported that Asian dust reached North America on at least five occasions during April 2010 (Uno et al., 2011). During May-June 2010, the Asian enhancement to MD8O<sub>3</sub> in the western U.S.A. was estimated to reach 8-15 ppbv in high-elevation regions during strong trans-Pacific transport events (Lin et al., 2012a).

Another process affecting tropospheric O<sub>3</sub> is stratosphere-to-troposphere transport (STT), which is known to be a significant contributor to the tropospheric O<sub>3</sub> budget (Lelieveld and Dentener, 2000). The tightening of the O<sub>3</sub> NAAQS and a continuous decrease of anthropogenic emissions have led to an increased focus on STT. On one hand, stratospheric intrusion of O<sub>3</sub> was found to be below 20 ppbv during March-October 2001 over the entire U.S.A (Fiore et al., 2003). On the other hand, a total of thirteen events were identified during April-June 2010 when stratospheric intrusion impacts reached 20-40 ppbv while accounting for 50-60% of total O<sub>3</sub> at fifteen high-elevation (> 1.4 km above mean sea level; m.s.l.) sites in the western U.S.A. (Lin et al., 2012b). From the perspective of interannual variability, springtime stratospheric intrusions may be enhanced following a La Niña winter due to a meandering of the jet stream, and a large variability in terms of magnitude and frequency have been shown from 1990 to 2012 (Lin et al., 2015). The fraction of O<sub>3</sub> in the troposphere that originates from the stratosphere is still uncertain due to its strong dependence on season and location which affect tropopause heights and is therefore still an area of active research (Mathur et al., 2017).

Table 1 summarizes these studies that provided the motivation for evaluating the impacts of both precursor emissions and STT on tropospheric O<sub>3</sub>. April 2010 is selected as the study period because enhancement of trans-Pacific transport is expected during the 2009-2010 El Niño winter. Along with the gradual reduction of precursor emissions of NO<sub>x</sub> and VOCs in the U.S.A., a gradual decrease of MD8O<sub>3</sub> mixing ratios can be expected and showed a decreasing trend by 0.4%/year; however, mean MD8O<sub>3</sub> mixing ratios in 2010 showed a local maximum and the number of NAAQS threshold exceedances was larger than usual as shown in Fig. S1 in the supplemental material. The variation in monthly mean and percentile distribution of observed MD8O<sub>3</sub> during 2010 are shown in Fig. S2 in the supplemental material. Although high MD8O<sub>3</sub> concentration for 95<sup>th</sup> percentiles and the number of NAAQS exceedances were found during summer time, it is also apparent that mean MD8O<sub>3</sub> during April 2010 higher than any other month. Lower MD8O<sub>3</sub> concentration for 5<sup>th</sup> and 25<sup>th</sup> percentiles were also noted as comparatively high during April 2010, indicating widespread enhancement of low-level O<sub>3</sub> further suggesting the possible

impacts of trans-Pacific transport on O<sub>3</sub> levels across the U.S.A. during this month. This period has already been the subject of other studies (e.g., Uno et al., 2011; Lin et al., 2012a), however, the methods used in this study to investigate the impacts of trans-Pacific transport differ from previous studies. The objective of this study is to better understand the relative contributions of precursor emissions from East Asia and the U.S.A. because the trans-Pacific transport has been recognized as an important factor. Previous studies primarily focused on Asian impacts on the western U.S.A., while this study investigates impacts across the entire U.S.A. In addition, some stratospheric intrusion events have been reported during spring 2010 (Lin et al., 2012b), therefore this period is suitable to examine not only trans-Pacific transport but also stratospheric intrusion, both processes may contribute to the observed high O<sub>3</sub> episodes in the U.S.A. Examination of the impacts of both processes will shed light on the formation mechanisms underlying such high O<sub>3</sub> episodes, thus improving our understanding of their relative importance in leading to these high O<sub>3</sub> episodes. The results of this work will be presented in two parts. Part 1 paper focuses on characterizing the influence of stratosphere-troposphere transport on O<sub>3</sub> distribution in the lower to middle troposphere. A sequential Part 2 manuscript focuses on the contributions of emissions leading to higher O<sub>3</sub> mixing ratio through Trans-Pacific transport. In this part 1 paper, we present the model evaluation and introduce a new method to identify and characterize periods during which surface and lower tropospheric O<sub>3</sub> may be influenced by stratospheric intrusions. This manuscript is organized as follows. In section 2, the modeling system and simulation set up are described, details on the surface, ozonesonde, airplane, and satellite observations used to evaluate the model performance are presented, and evaluation protocols are defined. In section 3, the analysis of model results and comparisons with observations are documented and the newly developed air mass characterization method is introduced and applied to investigate stratospheric intrusions. Finally, the conclusion section includes limitations of this work, future perspectives, and a brief introduction to part 2.

20

## 2 Methodology

### 2.1 Modeling System and Simulation Set Up

The model used in this work is the Community Multiscale Air Quality (CMAQ) version 5.2 extended for hemispheric applications (H-CMAQ) (Mathur et al., 2017). To investigate the impact of emissions from East Asia, H-CMAQ is configured to cover the entire Northern Hemisphere, utilizing a horizontal discretization of 187×187 grid points with a grid spacing of 108 km. The information of longitude and latitude is presented in Fig. S3 in the supplemental material. While the use of finer horizontal grid spacing can better resolve the STT processes, it will substantially increase computational demands. [STT was analyzed by combining analysis of data from a measurement network and predictions from total of seven model simulations over Europe, and reported the advantages of the Lagrangian models in capturing the STT \(Cristofanelli et al., 2003\).](#) In terms of the [Eurelian model](#), another study over Europe investigated the cross-tropopause transport in terms of resolution and diffusion coefficient using horizontal resolutions of 2°×2°, 1°×1°, and 0.5°×0.5° and showed that the simulation with the 2°×2° resolution has difficulty to capture [the tracer transport across the tropopause](#) (Gray, 2003). Based on these findings and the

model evaluation results (see Section 3.1) in this work, using a grid resolution of 108 km provides a good compromise between numerical accuracy and computational constraints. The terrain-following vertical coordinate utilizes 44 layers of variable thickness to resolve the model vertical extent between the surface and 50 hPa based on the extension of the previous 35 layers system (Mathur et al., 2017). The revised layer structure using 44 layers with significantly finer resolution above the boundary layer (BL) better represents long-range transport in the free-troposphere (FT) as well as STT processes, and influences from cloud mixing on both the sub-grid and resolved scales. As indicated in Mathur et al. (2017), the 44 layer configuration employed in the H-CMAQ configuration helps better capture dynamics in the vicinity of the tropopause and reduce excessive diffusion relative to coarser vertical resolution configurations. The emission inputs are based on the Hemispheric Transport of Air Pollution version 2 (HTAP2) modeling experiments, and the detailed description can be found in previous studies (Janssens-Maenhout et al., 2015; Pouliot et al., 2015; Galmarini et al., 2017; Hogrefe et al., 2018). The lightning emissions are prescribed using climatological averages as estimated in the Global Emission Inventory Activity (GEIA) as dataset (Price et al., 1997). For gas-phase chemistry, cb05e51 is used (Appel et al., 2017). This gas-phase mechanism includes the condensed halogen chemistry that leads to O<sub>3</sub> loss in marine environments (Sarwar et al., 2015). For aerosol chemistry, aero6 with nonvolatile primary organic aerosol (POA) (Simon and Bhawe, 2012) is adopted. The boundary conditions of H-CMAQ are taken from the clean tropospheric background values with updates to the physical and chemical sinks for organic nitrate species (Mathur et al., 2017).

Potential vorticity (PV) has been shown to be a robust indicator of air mass exchange between the stratosphere and the troposphere. The value of PV itself generally increases with altitude, and previous studies suggested that a value of 2 PVU (1 PVU = 10<sup>-6</sup> m<sup>2</sup> K kg<sup>-1</sup> s<sup>-1</sup>) is an indicator of stratospheric air (Hoskins et al., 1985; Wernli and Bourqui, 2002; Itoh and Narazaki, 2016). Through this study, the tropopause is diagnosed by 2 PVU. [The tropopause altitudes can be also diagnosed by the traditional approach based on the lapse rate \(i.e., thermal tropopause\) defined by World Meteorological Organization \(WMO\) \(WMO, 1992\), and the comparison with that diagnosed using PV \(i.e., dynamical tropopause\) have been reported \(Hoering et al., 1991\).](#) As shown in Figure S4, estimated tropopause altitudes averaged over April 2010 using PV in this work and the traditional approach of WMO are overall similar with below 10 km over high-latitude region and above 16 km over low-latitude region. PV shows a strong positive correlation with O<sub>3</sub> (Danielsen, 1968), and modeling studies have used this correlation to develop scaling factors that specify O<sub>3</sub> in the modeled upper troposphere/lower stratosphere (UTLS) based on estimated PV. The reported O<sub>3</sub>/PV ratios exhibited a wide range from 20 to 100 ppbv/PVU with differences in locations, altitude and season (e.g., Ebel et al, 1991; Carmichael et al, 1998; McCaffery et al, 2004). To account for the seasonal, latitudinal and altitude dependencies in the O<sub>3</sub>/PV relationship, a dynamic O<sub>3</sub>/PV function was developed to consider latitude, altitude, and time based on 21-year ozonesonde records from the World Ozone and Ultraviolet Radiation Data Centre (WOUDC) and corresponding PV values from WRF-CMAQ simulations across the northern hemisphere from 1990 to 2010 and is used in H-CMAQ (Xing et al., 2016). This parameterization of O<sub>3</sub>/PV is constructed at three topmost vertical levels of 58, 76, and 95 hPa fitted as a 5<sup>th</sup> order polynomial function, and applicable between the range of 50 and 100 hPa. Based on this new parameterization, it was demonstrated that UTLS O<sub>3</sub> agreed much better with observation in terms of its magnitude

and seasonality (Xing et al., 2016). Mathur et al. (2017) further demonstrated improvements in representation of seasonal variations in surface O<sub>3</sub> using the parameterization. To track stratospheric air masses, the O<sub>3</sub> estimated using the O<sub>3</sub>/PV relationship in the three layers listed above is also added as a chemically-inert tracer species in the H-CMAQ simulations as O<sub>3</sub>PV tracer. The O<sub>3</sub>PV tracer undergoes the same transport, scavenging, and deposition processes as O<sub>3</sub>, but its mixing ratios are not affected by chemical production or loss processes.

The meteorological fields are simulated by the Weather Research and Forecasting (WRF) model version 3.6.1 using the same vertical configuration as H-CMAQ. WRF simulation started from 1 March 2009 with more than one year of spin-up time prior to the analysis period of April 2010. The WRF model is configured to use the rapid radiative transfer model for global climate models (RRTMG) radiation scheme for both longwave and shortwave (Iacono et al., 2008), Morrison double-moment scheme (Morrison et al., 2009) and Grell convective parameterization (Grell 1993; Grell and Devenyi, 2002) for microphysics and cumulus parameterization, and Mellor-Yamada-Janjic scheme for planetary boundary layer (Janjic et al., 1994). Wind, temperature, and water vapor fields are nudged towards NCEP/NCAR final analysis data for all layers, these analysis data have 1 degree spatial and 6 h temporal resolution (NCEP, 2018). The WRF meteorological fields are converted to the format required by H-CMAQ using MCIP version 4.3 (Otte and Pleim, 2010), and then used for the H-CMAQ simulation. Relative humidity (RH) can also be used to diagnose stratospheric air masses because the stratosphere is characterized by dry air. CMAQ used the meteorological fields simulated by WRF and calculated RH based on the improved Magnus form approximation for saturation vapor pressure (Alduchov and Eskridge, 1996), and internally set the maximum value on 99% and minimum value on 0.5%. The CMAQ simulation started from 1 March 2010 and initialized with three-dimensional chemical fields from prior model simulations for 2010 by Hogrefe et al. (2018); March is discarded as a spin-up period and April is used as the analysis period. The O<sub>3</sub>PV tracer is also initialized by this prior model simulations of Hogrefe et al. (2018).

## 2.2 Observations and Evaluation Protocols

### 2.2.1 Ground-based Surface O<sub>3</sub> Observations

The northern hemispheric modeling domain and ground-based observations used in this study are shown on the map in Fig. 1. Global ground-based surface O<sub>3</sub> observations were obtained from the World Data Centre for Greenhouse Gases (WDCGG; shown as red circles in Fig. 1). For the study period of April 2010, this dataset contained 52 sites in North America, Europe, and several remote locations with only limited coverage over Asia (WDCGG, 2018). To overcome this limitation, surface O<sub>3</sub> observations are also obtained from the Acid Deposition Monitoring Network in East Asia (EANET) program which provides measurements at 12 sites in Japan, 3 sites in the South Korea, 1 site in Russia, and 4 sites in Thailand. However, the observed data are only available on a daily-mean basis for Russia, and a monthly-mean basis for South Korea and Thailand. Therefore, the only EANET monitors used in this study (EANET, 2018) are those located in Japan; these 9 sites with available data for April 2010 are shown as green triangles in Fig 1. In addition, surface O<sub>3</sub> observations over the U.S.A. were obtained from the Clean Air Status and Trends Network (CASTNET) and are shown as blue squares in Fig. 1. CASTNET monitors

(CASTNET, 2018) are located mostly in rural and remote areas, which makes them appropriate for comparison to O<sub>3</sub> fields from the coarse resolution H-CMAQ simulations. CASTNET data are available at 81 sites during April 2010. MD8O<sub>3</sub> values for April 2010 are calculated from the hourly observations at these WDCGG, EANET, and CASTNET stations.

### 5 2.2.2 Ozonesondes

An evaluation of simulated vertical O<sub>3</sub> profiles is needed to analyze the model's ability to capture the behavior of aloft O<sub>3</sub>. To this end, we obtained ozonesonde data distributed by the WOUDC as well as additional ozonesonde soundings available over the U.S.A. and Greenland that are collected and distributed by the National Oceanic and Atmospheric Administration Earth System Research Laboratory (NOAA-ESRL) (NOAA, 2018a). The total number of available ozonesonde sites during April 2010 was 33 (locations shown as yellow stars in Fig. 1). The data for Hilo and Boulder are available in both the WOUDC and NOAA-ESRL database; the NOAA-ESRL data are used because they include information on uncertainties of the O<sub>3</sub> measurements. Detailed information for each site, including country, site name, latitude (°N), longitude (°E), elevation (m a.s.l.), and the number of launches during April 2010, is provided in Table 2. There are 6 sites located in the U.S.A., 10 sites in Canada, 5 sites over Asia, and 12 sites over the Europe. In addition to measured O<sub>3</sub> mixing ratios, observed RH vertical profiles are used to evaluate the model performance.

### 2.2.3 Airplane

In addition to ozonesonde data to evaluate the vertical O<sub>3</sub> distribution, observations from research aircraft for three sites located in the U.S.A. (Cape May, New Jersey; Homer, Illinois; and Southern Great Plains, Oklahoma) are available from NOAA-ESRL (NOAA, 2018b) for April 2010. Because the observations at Cape May and Homer are only available for a single day during April 2010, we only used the NOAA-ESRL aircraft data at Southern Great Plains which is shown as a gray diamond in Fig. 1. A total of seven flights were conducted at this site during April 2010. In addition to O<sub>3</sub> mixing ratios, RH was used to evaluate the model performance.

### 25 2.2.4 Satellite

Tropospheric column O<sub>3</sub> observed by the Ozone Monitoring Instrument (OMI) onboard the National Aeronautics and Space Administration (NASA) Earth Observing System Aura satellite is used in this study. The methodology to estimate the tropospheric column has been developed (Ziemke et al., 2006) and consists of taking the differences between total column O<sub>3</sub> observed by OMI and stratospheric column O<sub>3</sub> observed by the Microwave Limb Sounder (MLS). The monthly-mean tropospheric O<sub>3</sub> column data are available between 60°S and 60°N (NASA, 2018a). Because this tropospheric column O<sub>3</sub> data are monthly-mean data, in order to take into account the daily missing data by OMI, total column data (OMTO3d) are utilized

to obtain the information on daily missing data in order to compare with the model (NASA, 2018b). This total column data are the products of averaging only good quality flag of level-2 swath data and then gridded into  $1 \times 1$  degree. Such an approach considering daily deficit data has also been applied in previous study (e.g., Chatani et al., 2014). To diagnose the tropopause in the model, PV with a value of 2 PVU is used as threshold. This diagnosis is applied above the boundary layer to avoid the misdiagnosis near the surface due to the high value of PV caused by turbulence.

### 2.2.5 Evaluation Protocol

To evaluate model performance, the Pearson's correlation coefficient (R) with student's *t*-test is used for assessing the statistical significance level. The normalized mean bias (NMB) and the normalized mean error (NME) are calculated using the following equations (e.g., Zhang et al., 2006);

$$R = \frac{\sum_1^N |(O_i - \bar{O})(M_i - \bar{M})|}{\sqrt{\sum_1^N (O_i - \bar{O})^2} \sqrt{\sum_1^N (M_i - \bar{M})^2}} \quad (1)$$

$$\text{NMB} = \frac{\sum_1^N (M_i - O_i)}{\sum_1^N O_i} \quad (2)$$

$$\text{NME} = \frac{\sum_1^N |M_i - O_i|}{\sum_1^N O_i} \quad (3)$$

where, N is the total observation number,  $O_i$  and  $M_i$  represent each individual observation and model value respectively, and  $\bar{O}$  and  $\bar{M}$  represent the arithmetical mean of observations and model values respectively. Based on a compilation of model evaluation reports, Emery et al. (2017) suggested threshold values of  $R > 0.75$ ,  $\text{NMB} < \pm 5\%$ , and  $\text{NME} < 15\%$  as performance goal, and threshold values of  $R > 0.50$ ,  $\pm 5\% < \text{NMB} < \pm 15\%$ , and  $15\% < \text{NME} < 25\%$  as performance criteria for 1-hr  $\text{O}_3$  or MD803 simulated by regional-scale air quality models. Although these recommendations were developed for regional-scale air quality models and suggested to apply over time-space averaging scales of no longer than one month and no more than 1000 km, these three criteria are applied in this work to judge the performance of the April 2010 H-CMAQ simulations due to the lack of other commonly-accepted model performance criteria for hemispheric or global scale  $\text{O}_3$  simulations. Evaluation of surface  $\text{O}_3$  simulated by global models indicated a somewhat loose threshold might be required because of the use of a coarse grid resolution (Zhang et al., 2012; He et al., 2015a, 2015b).

### 3 Simulation Results and Discussion

#### 3.1 Model Evaluation

A scatterplot of modeled vs. observed MD8O3 at WDCGG, EANET, and CASTNET, sites during April 2010 is shown in Fig. 2 using colors and symbols that are consistent with Fig. 1. A summary of the statistical analysis is provided in Table 3. Almost all of the EANET (green triangles) and CASTNET (blue squares) MD8O3 data pairs were within the 1:2 lines across the entire O<sub>3</sub> mixing ratio range. The comparison of H-CMAQ values with EANET observations over Asia shows an R value of 0.49 which is statistically significant at a level of  $p < 0.001$ , an NMB of  $-12.6\%$  and an NME of  $20.6\%$  (Table 3). A comparison to CASTNET observations over the U.S.A. shows that the mean observed and modeled values are close with an NMB of  $-0.9\%$ , and an NME of  $12.6\%$ , and that R had a value of 0.61 with  $p < 0.001$ . This indicates that the H-CMAQ simulations captured the CASTNET observational data within the model criteria performance suggested by Emery et al. (2017). A comparison to WDCGG data across the Northern Hemisphere shows an R of 0.49 with  $p < 0.001$ , an NMB of  $-19.3\%$ , and an NME of  $23.7\%$ . The mean model value is approximately 10 ppbv less than the mean of the observations. This feature is also evident in the scatter-plot shown in Fig. 2. While observed values reach more than 100 ppbv of MD8O3, the corresponding model values are only about half of these high MD8O3 mixing ratios, indicated by a clustering of WDCGG pairs on the 2:1 line. To investigate this model underestimation further, the spatial distributions of monthly mean modeled and observed O<sub>3</sub> mixing ratios are examined in Fig. 3 which shows observed high mixing ratios over eastern Europe. In particular, the four WDCGG monitors at Kosetice, Czech Republic ( $15.08^{\circ}\text{E}$ ,  $49.58^{\circ}\text{N}$ , 534 m a.s.l.), K-puszt, Hungary ( $19.55^{\circ}\text{E}$ ,  $46.97^{\circ}\text{N}$ , 125 m a.s.l.), Rucava, Latvia ( $21.17^{\circ}\text{E}$ ,  $56.16^{\circ}\text{N}$ , 18 m a.s.l.), and Zoseni, Latvia ( $25.54^{\circ}\text{E}$ ,  $57.08^{\circ}\text{N}$ , 182 m a.s.l.) measured MD8O3 mixing ratios larger than 100 ppbv by 14, 10, 5, and 4 days, respectively, during April 2010. An analysis of data collected during an airborne measurement campaign during 15-18, April 2010 over Siberia reported enhanced O<sub>3</sub> mixing ratios influenced by long-range transport, biomass burning plumes, and stratospheric intrusion (Berchet et al., 2013). Since the biomass burning emissions used in the current H-CMAQ simulations are based on climatological averages rather than year-specific events, the model underestimation may at least partially be due to the representation of these emissions. [Another possible reason may stem from the use of a coarse horizontal resolution.](#) From the viewpoint of meteorology, the blocking events over European Russia during spring-summer 2010 were reported, and positive anomaly of O<sub>3</sub> total column over the regions adjacent to the anticyclones (i.e., Europe) were analyzed (Sitnov et al., 2017). Removing data from these four sites from the analysis yields model performance metrics of an R of 0.63 with  $p < 0.001$ , an NMB of  $-14.4\%$ , and an NME of  $19.5\%$ ; which are comparable to performance at the EANET sites. Aside from the underestimation of high observed MD8O3 mixing ratios at these four European sites, H-CMAQ generally captured the WDCGG observations. Summarizing the model evaluation with surface observations, it is confirmed that model reasonably captures MD8O3 almost within model performance criteria of Emery et al. (2017).

To investigate the vertical profiles of O<sub>3</sub>, ozonesonde and airplane data are used in this study. In Fig. 4 and Fig. S5 in the supplemental material, time-height cross-sections (“curtain plots”) of hourly modeled O<sub>3</sub>, O3PV, and RH values during

April 2010 are shown at the location of the ozonesonde sites in the U.S.A., i.e., at Hilo, HI, Trinidad Head, CA, Boulder, CO, Huntsville, AL, Wallops Island, VA, and Rhode Island, RI. The plots also show contour lines of modeled PV for PV values of 1.0, 1.5, 2.0, 2.5, and 3.0 PVU with the red thick lines indicating a value of 2.0 PVU that can be used to diagnose the tropopause. Generally, O<sub>3</sub> and O3PV mixing ratios are very similar in the upper layers, especially above the 2.0 PVU line, indicating that O<sub>3</sub> mixing ratio in these layers are dominated by stratospheric air mass. Below the tropopause as diagnosed by the PV = 2.0 PVU line, O<sub>3</sub> mixing ratios are generally higher than O3PV mixing ratios, suggesting that O<sub>3</sub> was photochemically produced in the troposphere. On the other hand, instances of O<sub>3</sub> mixing ratios lower than O3PV mixing ratios are indicative of photochemical loss. One typical example of such photochemical loss can be seen at Hilo (Fig. S5 (a)). At that location, O<sub>3</sub> mixing ratios are less than 30 ppbv below 2 km whereas the O3PV mixing ratios are larger than 40 ppbv. A likely driver of this strong O<sub>3</sub> loss is the halogen chemistry in marine environments implemented in H-CMAQ (Sarwar et al., 2015) because Hilo site is surrounded by ocean. The impact of photochemical processes is further discussed in Section 3.2. Ozone mixing ratios at the level of the tropopause as diagnosed by the 2 PVU lines generally are around 100 ppbv (light blue colors in Fig. 4). Although high values of PV are typically seen in the upper layer above 10 km, it should be noted that high PV can occasionally also be found in the lower troposphere where could be associated with convection. RH values are below 10% (white colors in Fig. 4) above the tropopause and steeply increased near or below the tropopause as diagnosed by the 2.0 PVU lines. Based on the rough estimation shown in the curtain plots of Fig. 4, RH values at the level of the tropopause are typically on the order of 30-40%. In Fig. 4, the launch times of available ozonesonde measurements are indicated by yellow stars and we discuss below the comparison of model profiles to measurements from these launches.

The vertical profiles of observed and modeled O<sub>3</sub> and RH, as well as modeled O3PV and PV are shown in Fig. 5 and Fig. S6 in the supplemental material. In this figure, vertical red lines corresponding to a PV value of 2.0 PVU are inserted as index of stratospheric air masses, and the diagnosed stratosphere is colored with purple. A quantitative comparison between simulations and observations is conducted by averaging the observations onto the vertical grid spacing used by the model. The vertical layers are then assigned to three vertical ranges based on typical pressure values, i.e., the boundary layer (surface to approximately 750 hPa), the free troposphere (approximately 750-250 hPa), and the upper model layers (approximately 250-50 hPa) following the same approach used in our previous study (Hogrefe et al., 2018). Furthermore, the statistical analysis is performed separately for the three regions of U.S.A. and Canada, Asia, and Europe and the three layers ranges defined above. As shown in Fig. 4, O<sub>3</sub> and O3PV show similar variation in the upper model layer; however, O<sub>3</sub> is greater than O3PV near the tropopause indicated by 2.0 PVU, and this suggests the presence of photochemical production near the tropopause. Results of this statistical analysis for O<sub>3</sub> mixing ratios are shown in Table 3 and reveal that over all three regions the model performed the best for the boundary layer in terms of NMB and NME. The observed mean boundary layer values of around 45 ppbv over the three regions are well captured by the model. Over the U.S.A. and Canada, model performance in the boundary layer satisfies the performance criteria for all three metrics of R, NMB, and NME, and over Asia and Europe, NMB and NME also satisfy the performance criteria whereas R is less than 0.5. Compared to the results for the boundary layer, the model tends to underestimate the observed O<sub>3</sub> mixing ratios in the free troposphere and the upper model layers. In the free troposphere, the

mean observed value is around 80 ppbv while the mean model value is below 60 ppbv. As a result, NMB values are greater than -15% and NME values are greater than 20%. This underestimation is also present in the upper model layers; the mean observed values of 500 – 1000 ppbv are consistently underestimated by about 100 ppbv by the model across the three regions as shown in Table 3. R values tend to increase from the boundary layer to the free troposphere and the upper model layers due to model's ability to capture the increase of O<sub>3</sub> mixing ratios with height. The higher R values in the free troposphere, a region where impacts of photochemistry on O<sub>3</sub> variability is smaller, also suggest greater confidence in the model dynamics, which drive O<sub>3</sub> variations in this part of the atmosphere. Table S1 in the supplemental material shows the statistical results that are obtained when grouping stations into latitude ranges. The results indicate that model performance is similar to that shown in Table 3 and discussed above. These results suggest that although the revision of the dynamic-PV approach described in Xing et al. (2016) led to improved results compared to earlier implementations of the scaling approach, there is a need for further refinement of the approach to better capture high mixing ratios of stratospheric O<sub>3</sub>. Using a finer vertical resolution for the upper layers and extending the model top beyond 50 hPa to cover larger portions of the stratosphere could be potential strategies to address this need. ~~This is a hemispheric modeling system but the finer horizontal resolution will be another way to improve this.~~ In addition, the uncertainty of the lightning emissions prescribed as climatological averages in the current simulations may also contribute to the underestimation of O<sub>3</sub> in the free troposphere.

To investigate the effect of horizontal grid resolution on the representation of STT, additional WRF simulations were conducted over CONUS domain with 36 and 12 km horizontal grid resolutions, and temporal and vertical variations in simulated PV fields across the different resolutions (108, 36, and 12 km) were compared. These results are shown in Fig. S7 in the supplemental material. Generally, the modeled PV fields estimated with different horizontal grid resolutions showed similar features. The differences are displayed in Fig. S8 in the supplemental material. It was revealed that higher (lower) PV at upper (lower) altitude is enhanced (weakened) by increasing horizontal grid resolution. As expected, larger differences are noted between the 108 km and 12 km fields than those between the 108 km and 36 km fields. Although the enhancement of PV at upper altitudes could lead to increase in estimated O<sub>3</sub> through the O<sub>3</sub>/PV relationship used in the model, no systematic differences are noted in the estimated O<sub>3</sub> in the model's UTLS across the three resolutions, at least at the ozonesonde observation sites where our analysis is focused. A comparison of the altitude of 2 PVU which is used to diagnose the tropopause is also plotted in Fig. S8. As noted by the similarity of the altitude of the 2 PVU across the 3 different resolutions, the interpretation of STT events is not strongly influenced by the horizontal resolution employed in this study. This is because all model calculations employ assimilation of analyzed meteorological fields in the model's UTLS, resulting in comparable representation of STT events. Finally, a comparison of estimated O<sub>3</sub> at the model top-layer based on the O<sub>3</sub>/PV relation (Xing et al., 2016) by using different PV simulated from different horizontal grid resolutions is illustrated through scatter-plots in Fig. S9 in the supplemental material. These comparisons indicate good correspondence in the magnitude of O<sub>3</sub> at the model top using PV estimates from the 3 different resolutions. At the Boulder site (Fig. S9 (b)), the use of finer grid resolutions could sometimes lead to higher O<sub>3</sub> concentrations. Collectively, the comparisons in Figure S7-S9 suggest that the 108 km horizontal

grid resolution in H-CMAQ modeling system in conjunction with the physics and data assimilation options employed in the driving WRF model can capture the variability in the PV fields and associated STT O<sub>3</sub> impacts.

Although RH is a diagnostic variable, it may also provide an indication for stratospheric air mass, it is thus included in the model evaluation. As expected, Fig. 5 shows that RH has higher values and large variations in the troposphere and lower values in the stratosphere. For the analysis of modeled vertical profiles, model results of maximum and minimum values within  $\pm 2$  hours from observation time are also shown, and the range of RH showed large variations at lower altitude. Table 4 summarizes the statistical analysis divided into the three regions and three vertical domains for RH. It was found that the model generally overestimates RH over all regions and all three layers ranges. Although the NME value seems high for the upper layers, this is caused by the low absolute values of RH. The mean absolute differences between observed and modeled RH values are 1-2 % over the U.S.A. and Canada and Europe, and 8% over Asia. In Table S2 of the supplemental material, the RH results in Table 4 are presented for different latitude bands in the same fashion as the O<sub>3</sub> results in Table S1. Results show that model performance is similar to that discussed for Table 3. The systematic positive bias of RH occurs despite using analysis nudging for wind, temperature, and water vapor in the WRF simulations. Positive bias in predicted RH is also found in meteorological simulations performed for AQMEII (Vautard et al., 2012).

The tropopause diagnosed by PV = 2.0 PVU is located around 10-12 km at five ozonesonde sites in the U.S.A. except Hilo where it is located around 16 km. Observations in late April show instances of tropopause heights at or below 6 km (e.g., 27 April at Trinidad Head (Fig. 5 (b)), 29 April at Boulder (Fig. 5 (c)), and 27 April at Huntsville (Fig. 5 (d))). These cases illustrate large impacts of episodic STT, with observed O<sub>3</sub> mixing ratios steeply increasing from 100 ppbv at around 6 km to over 500 ppbv at around 8 km. The profiles obtained from the H-CMAQ simulations do not capture this steep increase and only show a gradual increase. This finding further supports a need for further refinement of representing stratospheric high O<sub>3</sub> mixing ratios as discussed above in the context of Table 3. In terms of RH, observed RH values show a sudden decline from around 60% to near 0% at Trinidad Head and Huntsville, whereas modeled RH values show a gradual decrease with large temporal variations. This contributes to the modeled positive RH bias shown in Table 4.

The comparison of model 3D O<sub>3</sub> structure at Southern Great Plains, Oklahoma, with research aircraft measurements is illustrated in Fig. 6. At this site, the curtain plot of modeled O<sub>3</sub> is shown for the entire month of April 2010 in the top row and zoomed inserts for the seven observational times indicated by gray diamonds above those plots are shown in the second row with each box showing airplane observations overlaid on H-CMAQ values. For O<sub>3</sub>, observed and modeled mixing ratios increased from about 30 ppbv at 1 km to about 55 ppbv at 5 km, except for flight #1 which shows persistently high mixing ratios of 50-60 ppbv throughout this altitude range. However, observed high mixing ratios of O<sub>3</sub> over 70 ppbv during flight #5, 6, and 7 are not captured by H-CMAQ. In contrast to O<sub>3</sub>, observed and modeled RH generally decreased from 1 km to 5 km as shown in rows 3 and 4. Overestimation in model RH is noted for flights #3, 4, and 6 above 3 km. Considering the profiles of O<sub>3</sub> and RH, flight # 6 might be a case of STT because observed RH is less than 10% and observed O<sub>3</sub> mixing ratios exceed 75 ppbv; however, the model fails to reproduce this behavior, the tropopause as diagnosed by the PV = 2.0 PVU locates near 10 km. The profile data averaged over all airplane ascents and descents are plotted in the bottom panel of Fig. 6, and statistical

analysis of these profiles is included in Table 3 for O<sub>3</sub> and Table 4 for RH. Similar to the evaluation results for ozonesondes, the model could reasonably capture observed O<sub>3</sub> and RH profiles, but O<sub>3</sub> mixing ratios are generally underestimated and RH is overestimated.

The observed and modeled tropospheric column O<sub>3</sub> are compared in Fig. 7. The observed latitudinal gradients in tropospheric column O<sub>3</sub> with values greater than 40 D.U. over mid-latitudes, column values around 30 D.U. over high- and low-latitudes, and values below 20 D.U. over the Pacific Ocean near the equator are captured well by H-CMAQ. To illustrate the differences between observations and simulations, the normalized bias is also shown in Fig. 7. This normalized bias map shows model tropospheric column O<sub>3</sub> overestimation over Russia and Africa and a slight underestimation over the Pacific Ocean. While the comparison with surface observations from WDCGG shows model underestimation at four sites over eastern Europe, the model slightly overestimates tropospheric column O<sub>3</sub> in this region. In addition, the model underestimation especially in the free-troposphere is noted through comparison with ozonesonde measurements (Table 3); however, this comparison showed model overestimation. The evaluation of satellite data compared to ozonesonde exhibited scattered correspondence and slight overestimation by satellite derived column O<sub>3</sub>. Therefore the model performance could differ from that for column O<sub>3</sub>. The results of the statistical analysis for tropospheric column O<sub>3</sub> are also listed in Table 3. The mean of observed and modeled tropospheric column O<sub>3</sub> across Northern Hemisphere is close on average, with an R of 0.65, an NMB of 4.7%, and an NME of 13.5%. The performance of tropospheric column O<sub>3</sub> judged based on the evaluation protocol developed for mixing ratios, suggests that the model satisfies the performance criteria proposed by Emery et al. (2017).

### 3.2 Air Mass Characterization Method

In order to characterize whether O<sub>3</sub> in a given air mass is dominated by photochemistry or stratospheric intrusion, and further estimate the impacts of STT, a new air mass characterization method is established here. [The flowchart of the established air mass characterization method](#) is illustrated in Fig. 8. The method relies on modeled O<sub>3</sub>PV/O<sub>3</sub> ratio to calculate the intensity of photochemistry. Because the top layer is set to 50 hPa in these H-CMAQ simulations, the uppermost layer (layer number is 44) is always regarded as stratospheric air mass in this method. For all layers below (i.e., layer 43 down to the lowermost layer), the importance of photochemistry is determined based on the ratio of the O<sub>3</sub>PV and O<sub>3</sub> mixing ratios. As noted in the discussion related to Figs. 4 and 5, if the O<sub>3</sub> mixing ratio is higher than the O<sub>3</sub>PV mixing ratio, it implies that photochemical production affected the air mass, and vice versa. Therefore, a O<sub>3</sub>PV/O<sub>3</sub> ratio of less (more) than 1.0, is classified as photochemical production (destruction), and a value near 1.0 can be classified as weakly impacted by photochemistry. The O<sub>3</sub>PV/O<sub>3</sub> ratio is illustrated in Fig. 9 (left) and Fig. S10 (left) in the supplemental material, wherein locations and times colored as orange (blue) represent air masses influenced by photochemical production (destruction), while ratios near 1.0 (range from 0.9 to 1.1) are colored as white. In Fig. 9 and Fig. S10, horizontal lines indicating 750, 500, and 250 hPa are also shown. The next step in the classification scheme is to determine whether an air mass is of stratospheric origin. The concept of a sequential intrusion from upper layers to lower layers is considered. When the grid cell directly above is also diagnosed as stratospheric

air mass, the grid is determined as being dominated by stratospheric air mass. Applying this concept of a sequential stratospheric air mass intrusion is repeatedly used in the air mass characterization scheme to determine whether an air mass is dominated by photochemistry or stratospheric intrusion. It is important to note that characterizing a grid cell as being dominated by a process does not mean that other processes do not impact O<sub>3</sub> mixing ratios as well. For example, O<sub>3</sub> in a grid cell near the tropopause can be dominated by stratospheric air mass, but it can also be affected by photochemical production and destruction. Similarly, although O<sub>3</sub> in a grid cell near the surface layer is often dominated by photochemical processes, it can also be affected by stratospheric air mass.

An illustration of applying this method to illustrate the stratospheric intrusion for the six ozonesonde sites in the U.S.A. is presented in Fig. 9 (right) and Fig. S10 (right) in the supplemental material. Above 750 hPa, the stratospheric air mass is dominated and varied day-to-day with downward movement reach to near 750 hPa. These stratospheric intrusions are clearly found in some cases, and the stratospheric air mass reaches to the surface during early to middle April at Trinidad Head (Fig. 9(a)) and early April at Boulder (Fig. 9(b)). It should be noted that since the classification scheme is based on the most dominant process, a grid cell classified as being dominated by photochemistry can still be influenced by stratospheric air. Therefore, these estimated impacts of stratospheric air masses on the troposphere can be viewed as a lower bound.

15

### 3.3 Investigation of Stratospheric Intrusion

The relationship between the model estimated stratospheric contribution to the total tropospheric O<sub>3</sub> column and observed surface O<sub>3</sub> levels at CASTNET sites is investigated in Fig. 10 on monthly average. The grid classified as stratospheric air mass is used to calculate O<sub>3</sub> column from stratospheric intrusion, and the stratospheric air mass in troposphere is estimated. The troposphere is defined as all layers below 250 hPa for the purpose of this analysis. Using data from all CASTNET locations, the relationship shows a slight negative slope by R value of 0.25 with non-significance ( $p > 0.05$ ), indicating that the influence of stratospheric air decreased with increasing surface MD8O<sub>3</sub> mixing ratios. To further focus on this relationship at elevated sites in the U.S.A., the analysis is repeated using data from sites with an elevation higher than 1000 m as listed in Table S3 in the supplemental material. The result shows a slight positive slope by R value of 0.14 with non-significance ( $p > 0.05$ ), which indicates that at elevated sites, STT has a possible effect on surface mixing ratio values. The finding of a negative slope using the entire dataset over the U.S.A. is consistent with a previous investigation focused on relatively-polluted areas over the western U.S.A. such as the Central Valley, Southern California, and Las Vegas (Lin et al., 2012b). For elevated sites, they also reported a positive slope indicating higher contributions of stratospheric air masses during periods of elevated surface O<sub>3</sub>. The reason for the relatively weak relation found in this study seems to stem from differences in simulated stratospheric O<sub>3</sub> mixing ratios. Lin et al. (2012b) used the fully-coupled stratosphere-troposphere chemistry model GFDL AM3 which tended to overestimate O<sub>3</sub> mixing ratios; therefore, they employed a bias correction approach (assuming that when the estimated stratospheric contribution exceeds the model bias, the bias is caused entirely by excessive stratospheric O<sub>3</sub>) for estimating the stratospheric impacts on surface O<sub>3</sub>. On the other hand, the H-CMAQ simulations analyzed in this study tends to underestimate

tropospheric O<sub>3</sub> levels, especially during STT events, which may suggest that its estimates of stratospheric contributions to high surface O<sub>3</sub> events may also be too low.

The monthly-averaged spatial distribution of the stratospheric air mass is shown in Fig. 11. It indicates higher stratospheric impacts over high-latitude region, and 5-25% on monthly-average over western U.S.A. This is the result of monthly-average and it is well known that the stratospheric intrusion is event occurred on short-term and then lead to the high surface concentration. Time series of daily averaged stratospheric air mass in troposphere at five ozonesonde sites at contiguous U.S.A. is also shown in Fig. 11. These time series reveal large temporal variations of stratospheric air mass in troposphere. On a monthly-mean basis during April 2010, air masses classified as being dominated by stratospheric intrusion contribute about 5-25% to the total [tropospheric O<sub>3</sub> column](#) at five of the ozonesonde sites and 25% at Trinidad Head. However, on specific days, O<sub>3</sub> masses from the stratosphere contribute up to 50-90% of the total [tropospheric O<sub>3</sub> column](#).

The previous section introduces an approach to identify cases when stratospheric air masses impact tropospheric O<sub>3</sub>. Results of the O<sub>3</sub> column mass analysis identify periods in early, middle, and late April 2010 that are affected by stratospheric intrusions over the contiguous U.S.A., and in this section these events are further analyzed. Daily maps of the spatial patterns of the percentage of [tropospheric O<sub>3</sub> column](#) diagnosed as being of stratospheric origin during (i) early and (ii) late April are presented in Fig. 12 while maps for mid-April are presented in Fig. [S11](#) in the supplemental material. On 5 April, a large impact from the stratosphere was seen over the western U.S.A. (indicated as point S<sub>A</sub> on the map) and covered Trinidad Head where the contribution of O<sub>3</sub> from the stratosphere to the tropospheric O<sub>3</sub> mass is greater than 50%. This air mass moved eastward on 6 April when the impact at Boulder is also greater than 50%. During 7-8 April, this air mass was located over the central U.S.A., and then moved further to the east, with contributed at Huntsville on 9 April. Finally, this air mass further moved towards the northeast U.S.A. with impact at Wallops Island and Rhode Island, respectively. The stratospheric impacts in early April are associated with an air mass movement from west to east within 5 days; corresponding to an average speed of about 8-9 m/sec. Compared to the early April case, the case in late April is different. On 25 April, large impacts from the stratosphere were found over western (marked as S<sub>B1</sub>) and eastern (marked as S<sub>B2</sub>) Canada. The S<sub>B1</sub> air mass moved into the western U.S.A. on 27 April, and had large impacts from 27 to 30 April at Trinidad Head, affected Boulder on 29 and 30 April, and finally moved southwestern U.S.A. in a U-shaped pattern on 30 April. Another air mass of S<sub>B2</sub> located over eastern Canada on 26 April moved slowly southward and impacted Wallops Island and Rhode Island, then moved eastward. Thus, for the late April case, stratospheric air was present in different air masses impacting different locations on different days rather than a single air mass simply moving from west to east as in the early April case. Contrasting the early and late April cases illustrates that different synoptic flow scenarios influence how stratospheric air can impact tropospheric O<sub>3</sub> columns over the U.S.A. The impacts of STT during the middle of April are shown in Fig. [S11](#) in the supplemental material. From 12-15 April, tropospheric O<sub>3</sub> columns over the western (i.e., Trinidad Head) and eastern (i.e., Wallops Island and Rhode Island) U.S.A. were dominated by stratospheric intrusion, and these impacts were almost disappeared after 17 April. Previous studies (e.g., Lin et al., 2012b) estimated thirteen STT events during April-June 2010, and April 7, 9, 12-15, 21-23, and 28-29 2010 were reported as STT events. Our investigations based on the air mass characterization method matches with these earlier findings. The impact of

STT at Huntsville has been investigated by combining ozonesonde and ozone lidar data (Kuang et al., 2012). In their report, the period of 27-29 April was associated with STT. However, STT estimation was limited above 750 hPa (Fig. 9). This is due to the model difficulty to capture the STT event itself as revealed from the comparison with ozonesonde dataset (Fig. 5). As discussed in model evaluation, finer horizontal and vertical resolution could be potential way to improve STT representation.

5

#### 4 Conclusions

In this study, the regional chemical transport model CMAQ recently extended for hemispheric applications, the H-CMAQ, is applied to investigate trans-Pacific transport during April 2010. This part 1 manuscript presents results from comprehensive evaluation and a new air mass characterization method based on these results and ozonesonde measurements.

10 The comparison of modeled and observed O<sub>3</sub> at the surface shows a very good performance with NMBs around -10%. Comparisons of vertical O<sub>3</sub> distributions against ozonesonde and aircraft-based observations show that the model can capture well O<sub>3</sub> variations within boundary layer similar to those at the surface, although systematic underestimations of free troposphere O<sub>3</sub> occur with NMBs up to -30%, especially during events that are characterized to have strong STT during late April. Modeled RH exhibits a positive bias with NMBs of +10% or greater at all altitudes. Comparisons of modeled  
15 tropospheric O<sub>3</sub> column with satellite observations suggest that the model can represent the general feature with lower column O<sub>3</sub> over the Pacific Ocean near equator and higher column in the mid-latitudes.

Using ozonesonde measurements, the relationship between PV and RH is examined to characterize stratospheric air masses. The PV-RH relation indicates that PV of 2.0 PVU (1 PVU = 10<sup>-6</sup> m<sup>2</sup> K kg<sup>-1</sup> s<sup>-1</sup>) generally corresponds to RH values of 30-40%. A new air mass characterization method is further developed based on the ratio of modeled O<sub>3</sub> and stratospheric O<sub>3</sub>  
20 tracer mixing ratios to examine the relative importance of photochemistry, and PV and RH to examine the stratospheric air mass. The estimated STT impacts show significant day-to-day variations both in the magnitude of the contribution and the origin of the air mass. The relationship between surface O<sub>3</sub> levels and estimated stratospheric air mass in troposphere exhibits a slight negative slope, indicating that at most locations, high surface O<sub>3</sub> mixing ratios typically result from other factors (e.g., emissions). In contrast, at elevated sites the relationship exhibits a slight positive slope, indicating a steady STT contribution  
25 to O<sub>3</sub> levels at these locations.

Despite the use of a coarse horizontal grid resolution for H-CMAQ simulations in this work, it was found that the model performance at the surface and in the boundary layer are reasonably within the model performance criteria suggested from regional-scale applications. However, this work also shows that the model has difficulty capturing higher O<sub>3</sub> mixing ratios in the free troposphere through the comparison with ozonesonde data. This result suggests a need for model  
30 improvements to accurately represent the STT process. This study focuses on April 2010, and monthly or seasonal behavior of STT, and interannual variations based on the long-term trend analysis should also be considered for future study. The Part 2 paper will focus on other factors that affect surface O<sub>3</sub> mixing ratio, namely emissions, also examine the relative importance of NO<sub>x</sub> and VOCs.

### **Code availability**

Source code for version 5.2 of the CMAQ model can be downloaded from <https://github.com/USEPA/CMAQ/tree/5.2>. For further information, please visit the US Environmental Protection Agency website for the CMAQ system:  
5 <https://www.epa.gov/cmaq>.

### **Data availability**

The observational datasets used in this study are available from their respective websites: <http://ds.data.jma.go.jp/gmd/wdogg/> (WDCGG), <http://www.eanet.asia/index.html> (EANET), and <https://www.epa.gov/castnet> (CASTNET) for surface observation network, <https://woudc.org/home.php> (WOUDC) and <https://www.esrl.noaa.gov/gmd/ozwv/ozsondes/> (NOAA ESRL) for ozonesonde, <https://www.esrl.noaa.gov/gmd/ozwv/aircraft/index.html> (NOAA-ESRL) for airplane, [https://acd-ext.gsfc.nasa.gov/Data\\_services/cloud\\_slice/index.html](https://acd-ext.gsfc.nasa.gov/Data_services/cloud_slice/index.html) (NASA). Last Access: 31 August 2018.  
10

### **Competing interests**

15 The authors declare that they have no conflict of interest.

### **Disclaimer**

The views expressed in this paper are those of the authors and do not necessarily reflect the views or policies of the U.S. Environmental Protection Agency.

20

### **Author contributions**

Syuichi Itahashi performed the analysis of observation and model simulation and prepared the manuscript with contributions from all co-authors. Rohit Mathur and Christian Hogrefe contributed to establish the hemispheric modeling application for this study and prepared the emission dataset, initial condition, and lateral boundary condition from previous long-term simulation

results. Yang Zhang contributed to the literature review of trans-Pacific transport and refined this research through simulation designs, and results interpretation.

## Acknowledgement

- 5 The authors are grateful the observation dataset of surface (WDCGG, EANET, and CASTNET), ozonesonde (WOUDC and NOAA ESRL), airplane (NOAA ESRL), and satellite (NASA). YZ acknowledges support from the 2017-2018 NC State Internationalization Seed Grant and the 2019-2020 NC State Kelly Memorial Fund for US-Japan Scientific Cooperation.

## References

- 10 Alduchov, O. A. and Eskridge, R. E.: Improved Magnus form approximation of saturation vapor pressure, *Journal of Applied Meteorology*, 35, 601–609, 1996
- Appel, K.W., Napelenok, S., Foley, K.M., Pye, H.O.T., Hogrefe, C., Luecken, D.J., Bash, J.O., Roselle, S.J., Pleim, J.E., Foroutan, H., Hutzell, W., Pouliot, G., Sarwar, G., Sarwar, G., Fahey, K., Gantt, B., Gilliam, R.C., Kang, D., Mathur, R., Schwede, D., Spero, T., Wong, D.C., & Young, J.: Overview and evaluation of the Community Multiscale Air Quality  
15 (CMAQ) model version 5.1. *Geosci. Model Dev.*, 10, 1703–1732, doi:10.5194/gmd-10-1703-2017, 2017
- Berchet, A., Paris, J-D., Ancellet, G., Law, K. S., Stohl, A., Nédélec, P., Arshinov, M. Y., Belan, B. D., and Ciais, P.: Tropospheric ozone over Siberia in spring 2010: remote influences and stratospheric intrusion, *Tellus B: Chemical and Physical Meteorology*, 65:1, 19688, doi:10.3402/tellusb.v65i0.19688, 2013.
- Carmichael, G., Uno, I., Phadnis, M. J., Zhang, Y., and Sunwoo, Y.: Tropospheric ozone production and transport in the  
20 springtime in east Asia, *Journal of Geophys. Res.*, 103, D9, 10649–10671, 1998.
- Chatani, S., Amann, M., Goel, A., Hao, J., Klimont, Z., Kumar, A., Mishra, A., Sharma, S., Wang, S. X., Wang, Y. X., and Zhao, B.: Photochemical roles of rapid economic growth and potential abatement strategies on tropospheric ozone over South and East Asia in 2030, *Atmos. Chem. Phys.*, 14, 9259–9277, doi:10.5194/acp-14-9259-2014, 2014.
- Cooper, O. R., Gao, R.-S., Tarasick, D., Leblanc, T., and Sweeney, C.: Long-term ozone trends at rural ozone monitoring sites  
25 across the United States, 1990-2010, *Journal of Geophys. Res.*, 117, D22307, doi:10.1029/2012JD018261, 2012.
- Cooper, O. R., Parrish, D. D., Stohl, A., Trainer, M., Nédélec, P., Thouret, V., Cammas, J. P., Oltmans, S. J., Johnson, B. J., Trasock, D., Leblanc, T., McDermid, I. S., Jaffe, D., Gao, R., Stith, J., Ryerson, T., Aikin, K., Campos, T., Weinheimer, A., and Avery, M. A.: Increasing springtime ozone mixing ratios in the free troposphere over western North America, *Nature*, 463, 21, doi:10.1038/nature08708, 2010.

- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R., and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmos. Chem. Phys.*, 16, 3825–3841, doi:10.5194/acp-16-3825-2016, 2016.
- 5 Cristofanelli, P., Bonasoni, P., Collins, W., Feichter, J., Forster, C., James, P., Kentarchos, A., Kubik, P. W., Land, C., Meloan, J., Roelofs, G. J., Siegmund, P., Sprenger, M., Schnabel, C., Stohl, A., Tobler, L., Tositti, L., Trickl, T., and Zanis, P.: Stratosphere-to-troposphere transport: A model and method evaluation, *Journal of Geophys. Res.*, 108, D12, 8525, doi:10.1029/2002JD002600, 2003.
- Danielsen, E. F.: Stratosphere-Troposphere exchange based on radioactivity, ozone and potential vorticity, *Journal of Atmospheric Sciences*, 25, 502-518, 1968.
- 10 Ebel, A., Hass, H., Jakobs, H., J., Laubem M., Memmesheimer, M., Oberreuter, A., Geiss, H., and Kuo, Y.-H.: Simulation of ozone intrusion caused by a tropopause fold and cut-off low, *Atmos. Environ.*, 25A, 10, 2131–2144, 1991.
- Emery, C., Liu, Z., Russell, A. G., Odman, M. T., Yarwood, G., and Kumar, N.: Recommendations on statistics and benchmarks to assess photochemical model performance, *J. of the Air and Waste Manage. Assoc.*, 67, 582–598, doi:10.1080/10962247.2016.1265027, 2017.
- 15 Fiore, A. M., Jacob, D. J., Bey, I., Yantosca, R. M., Field, B. D., Fusco, A. C., and Wilkinson, J. G.: Background ozone over the United States in summer: Origin, trend, and contribution to pollution episodes, *Journal of Geophys. Res.*, 107, D15, 4275, doi:/10.1029/2001JD000982, 2002.
- Fiore, A., Jacob, D. J., Liu, H., Yantosca, R. M., Fairlie, T. D. and Li, Q.: Variability in surface ozone background over the United States: Implications for air quality policy, *Journal of Geophys. Res.*, 108, D24, 4787, doi:/10.1029/2003JD003855, 2003.
- 20 Galmarini, S., Koffi, B., Solazzo, E., Keating, T., Hogrefe, C., Schulz, M., Benedictow, A., Griesfeller, J. J., Janssens-Maenhout, G., Carmichael, G., Fu, J., and Dentener, F.: Technical note: Coordination and harmonization of the multi-scale multi-model activities HTAP2, AQMEII3, and MICS-Asia3: simulations, emission inventories, boundary conditions, and model output formats, *Atmos. Chem. Phys.*, 17, 1543–1555, doi:10.5194/acp-17-1543-2017, 2017.
- 25 Gray, S. L.: A case study of stratosphere to troposphere transport: The role of convective transport and the sensitivity to model resolution, *Journal of Geophys. Res.*, 108, D18, 4590, doi:10.1029/2002JD003317, 2003.
- Grell, G. A.: Prognostic evaluation of assumptions used by cumulus parameterizations, *Month. Wet. Rev.*, 121, 764–787, 1993.
- Grell, G. A. and Devenyi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophys. Res. Lett.*, 29, 14, 1613, doi:10.1029/2002GL015311, 2002.
- 30 Haagen-Smit, A. J. and Fox, M. M.: Photochemical Ozone Formation with Hydrocarbons and Automobile Exhaust, *Air Repair*, 4, 3, 105-136, doi:10.1080/00966665.1954.10467649, 1954.
- He, J., Zhang, Y., Tilmes, S., Emmons, L., Lamarque, J.-F., Glotfelty, T., Hodzic, A., and Vitt, F.: CESM/CAM5 Improvement and Application: Comparison and Evaluation of Updated CB05\_GE and MOZART-4 Gas-Phase Mechanisms and

- Associated Impacts on Global Air Quality and Climate, *Geosci. Model Dev.*, 8, 3999–4025, doi:10.5194/gmd-8-3999-2015, 2015.
- He, J., Zhang, Y., Glotfelty, T., He, R., Bennartz, R., Rausch, J. and Sartelet, K.: Decadal Simulation and Comprehensive Evaluation of CESM/CAM5.1 with Advanced Chemistry, Aerosol Microphysics, and Aerosol-Cloud Interactions, *Journal of Advances in Modeling Earth Systems*, 7 (1), 110–141, doi:10.1002/2014MS000360, 2015.
- Hoering, M. P., Schaack, T. K., and Lenzen, A. J.: Global objective tropopause analysis, *Mon. Wet. Rev.*, 119, 1816–1831, 1991.
- Hogrefe, C., Liu, P., Pouliot, G., Mathur, R., Roselle, S., Flemming, J., Lin, M., and Park, R. J.: Impacts of different characterizations of large-scale background on simulated regional-scale ozone over the continental United States, *Atmos. Chem. Phys.*, 18, 3839–3864, doi:10.5194/acp-18-3839-2018, 2018.
- Hoskins, B. J., McIntyre, M. E., and Robertson, A. W.: On the use and significance of isentropic potential vorticity maps, *Quarterly Journal of the Royal Meteorological Society*, 111, 877–946, 1985.
- Huebert, B. J., Bates, T., Russell, P. B., Shi, G., Kim, Y. J., Kawamura, K., Carmichael, G., and Nakajima, T.: An overview of ACE-Asia: Strategies for quantifying the relationships between Asian aerosols and their climatic impacts, *Journal of Geophys. Res.*, 108, D23, doi:10.1029/2003JD003550, 2003.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.*, 113, D13103, doi:10.1029/2008JD009944, 2008.
- Itahashi, S., Uno, I., and Kim, S-T.: Seasonal source contributions of tropospheric ozone over East Asia based on CMAQ-HDDM, *Atmos. Environ.*, 70, 204–217, doi:10.1016/j.atmosenv.2013.01.026, 2013.
- Itahashi, S., Uno, I., Irie, H., Kurokawa, J., and Ohara, T.: Regional modeling of tropospheric NO<sub>2</sub> vertical column density over East Asia during the period 2000-2010: comparison with multisatellite observations, *Atmos. Chem. Phys.*, 14, 3623–3635, doi:10.5194/acp-14-3623-2014, 2014.
- Itahashi, S., Hayami, H., and Uno, I.: Comprehensive study of emission source contributions for tropospheric ozone formation over East Asia, *J. Geophys. Res.*, 120, 331–358, doi:10.1002/2014JD022117, 2015.
- Itoh, H., and Narazaki, Y.: Fast descent routes from within or near the stratosphere to the surface at Fukuoka, Japan, studied using <sup>7</sup>Be measurements and trajectory calculations, *Atmos. Chem. Phys.*, 16, 6241–6161, doi:10.5194/acp-16-6241-2016, 2016.
- Jacob, D. J., Logan, J. A., and Murti, P.: Effect of rising Asia emissions on surface ozone in the United States, *Geophys. Res. Lett.*, 26, 14, 2175–2178, 1999.
- Jaffe, D., Anderson, T., Covert, D., Kotchenruther, R., Trost, B., Danielson, J., Simpson, W., Berntsen, T., Karlsdottir, S., Blake, D., Harris, J., Carmichael, G., and Uno, I.: Transport of Asian air pollution to North America, *Geophys. Res. Lett.*, 26, 6, 711–714, 1999.

- Jaffe, D., Price, H., Parrish, D., Glodstein, A., and Harris, J.: Increasing background ozone during spring on the west coast of North America, *Geophys. Res. Lett.*, 30, 12, 1613, doi:10.1029/2003GL017024, 2003.
- Janjic, Z.: The step-mountain Eta coordinate Model: Further developments of the convection, viscous sublayer, and turbulence closure schemes, *Month. Wet. Rev.*, 122, 927–945, 1994.
- 5 Janssens-Maenhout, G., Crippa, M., Guizzardi, F., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmuller, R., Danier van der Gon, H., Kuenen, J.J.P., Kilmont, Z., Frost, G., Darras, S., Koffi, B. and Li, M.: HTAP\_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution. *Atmos. Chem. Phys.* 15: 11411–11432, doi:10.5194/acp-15-11411-2015, 2015.
- Koumoutsaris, S., Bey, I., Generoso, S., and Thouret, V.: Influence of El Niño–Southern Oscillation on the interannual variability of tropospheric ozone in the northern midlatitudes, *Journal of Geophys. Res.*, 113, D19301, doi:10.1029/2007JD009753, 2008.
- 10 Kuang, S., Newchurch, M. J., Burris, J., Wang, L., Knupp, K., and Huang, G.: Stratosphere-to-troposphere transport revealed by ground-based lidar and ozonesonde at a midlatitude site, *J. of Geophys. Res.*, 117, D18305, doi:10.1029/2012JD017695, 2012.
- 15 Lelieveld, J. and Dentener, F.J.: What controls tropospheric ozone? *Journal of Geophysical Research* 105, 3531–3551, 2000.
- Lin, M., Fiore, A. M., Horowitz, L. W., Cooper, O. R., Naik, V., Holloway, J., Johnson, B. J., Middlebrook, A. M., Oltmans, S. J., Pollack, I. B., Ryerson, T. B., Warner, J. X., Wiedinmyer, C., Wilson, J., and Wyman, B.: Transport of Asian ozone pollution into surface air over the western United States in spring, *J. Geophys. Res.*, 117, D00V07, doi:10.1029/2011JD016961, 2012a.
- 20 Lin, M., Fiore, A. M., Cooper, O. R., Horowitz, L. W., Langford, A. O., Levy II, H., Johnson, B. J., Naik, V., Oltmans, S. J., and Senff, C. J.: Springtime high surface ozone events over the western United States: Quantifying the role of stratospheric intrusions, *J. Geophys. Res.*, 117, D00V22, doi:10.1029/2012JD018151, 2012b.
- Lin, M., Fiore, A. M., Horowitz, L. W., Langford, A. O., Oltmans, S. J., Tarasick, D., and Rieder, H. E.: Climate variability modulates western US ozone air quality in spring via deep stratospheric intrusions, *Nature Communications*, 6, 7105, doi:10.1038/ncomms8105, 2015.
- 25 Logan, J. A.: Tropospheric ozone: Seasonal behavior, trends and anthropogenic influences, *J. Geophys. Res.*, 90, 10463–10482, 1985.
- Mathur, R., Xing, J., Gilliam, R., Sarwar, G., Hogrefe, C., Pleim, J., Poliot, G., Roselle, S., Spero, T. L., Wong, D. C., and Young, J.: Extending the Community Multiscale Air Quality (CMAQ) modeling system to hemispheric scales: overview of process considerations and initial applications, *Atmos. Chem. Phys.*, 17, 12449–12474, doi:10.5194/acp-17-12449-2017, 2017.
- 30 McCaffery, S. J., McKeen, S. A., Hsie, E.-Y., Parrish, D. D., Cooper, O. R., Holloway, J. S., Hubler, G., Fehsenfeld, F. C., and Trainer, M.: A case study of stratosphere-troposphere exchange during the 1996 North Atlantic Regional Experiment, *Journal of Geophys. Res.*, 109, D14103, doi:10.1029/2003JD004007, 2004.

- Morrison, H., Thompson, G., and Tatarskii, V.: Impacts of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: comparison of one- and two-moment schemes. *Monthly Weather Rev.* 137, 991–1007, 2009
- Otte, T. L. and Pleim, J. E.: The meteorology-chemistry interface processor (MCIP) for the CMAQ modeling system: updates through MCIPv3.4.1, *Atmos. Chem. Phys.* 3, 243–256, 2010.
- 5 Pouliot, G., Denier van der Gon, H. A. C., Kuenen, J., Zhang, J., Moran, M. D., and Maker, P. A.: Analysis of the emission inventories and model-ready emission datasets of Europe and North America for phase 2 of the AQMEII project, *Atmos. Environ.*, 115, 345–360, doi:10.1016/j.atmosenv.2014.10.061, 2015.
- Price, C., Penner, J., and Prather, M.: NO<sub>x</sub> from lightning 1: Global distribution based on lightning physics, *J. Geophys. Res.*, 10 102, 5929–5941, 1997.
- Sarwar, G., Gantt, B., Schwede, D., Foley, K., Mathur, R., and Saiz-Lopez, A.: Impact of enhanced ozone deposition and halogen chemistry on tropospheric ozone over the Northern Hemisphere, *Environ. Sci. Tech.*, 49, 9203–9211, doi:10.1021/acs.est.5b01657, 2015.
- Simon, H., and Bhave, P. V.: Simulating the degree of oxidation in atmospheric organic particles, *Environ. Sci. Technol.*, 15 46, 331–339. doi:10.1021/es202361w, 2012.
- Simon, H., Reff, A., Wells, B., Xing, J., and Frank, N.: Ozone trends across the United States over a period of decreasing NO<sub>x</sub> and VOC emissions, *Environ. Sci. Technol.*, 49, 186–195. doi:10.1021/es504514z, 2015.
- Sitnov, S. A., Mikhov, I. I., and Lupo, A. R.: Ozone, water vapor, and temperature anomalies associated with atmospheric blocking events over Eastern Europe in spring-summer 2010, *Atmos. Environ.*, 164, 180–194, 20 2017. doi:10.1016/j.atmosenv.2017.06.004, 2017.
- Uno, I., Eguchi, K., Yumimoto, K., Liu, Z., Hara, Y., Sugimoto, N., Shimizu, A., and Takemura, T.: Large Asian dust layers continuously reached North America in April 2010, *Atmos. Chem. Phys.*, 11, 7333–7341, doi:10.5194/acp-11-7333-2011, 2011.
- Vautard, R., Moran, M. D., Solazzo, E., Gilliam, R. C., Matthias, V., Bianconi, R., Chemel, C., Ferreira, J., Geyer, B., Hansen, 25 A. B., Jericevic, A., Prank, M., Segers, A., Silver, J. D., Werhahn, J., Wolke, R., Rao, S. T., and Galmarini, S.: Evaluation of the meteorological forcing used for the Air Quality Model Evaluation International Initiative (AQMEII) air quality simulations, *Atmos. Environ.*, 53, 15–37, doi:10.1016/j.atmosenv.2011.10.065, 2012.
- Wang, K., Zhang, Y., Jang, C., Phillips, S., and Wang, B.: Modeling intercontinental air pollution transport over the trans-Pacific region in 2001 using the Community Multiscale Air Quality modeling system, *Journal of Geophys. Res.*, 114, 30 D04307, doi:10.1029/2008JD010807, 2009.
- Wernli, H., and Bourqui, M.: A Lagrangian “1-year climatology” of (deep) cross-tropopause exchange in the extratropical Northern Hemisphere, *Journal of Geophys. Res.*, 107, D2, 4021, doi:10.1029/2001JD000812, 2002.
- World Meteorological Organization: International Meteorological Vocabulary, Secretariat of the World Meteorological Organization, Geneva, 2nd edn., 1992.

- Xing, J., Mathur, R., Pleim, J., Hogrefe, C., Gan, C.-M., Wong, D. C., Wei, C., Gilliam, R., and Pouliot, G.: Observations and modeling of air quality trends over 1990-2010 across the Northern Hemisphere: China, the United States and Europe, *Atmos. Chem. Phys.*, 15, 2723–2747, doi:10.5194/acp-15-2723-2015, 2015.
- 5 Xing, J., Mathur, R., Pleim, J., Hogrefe, C., Wang, J., Gan, C.-M., Sarwar, G., Wong, D. C., and McKeen, S.: 3-D O<sub>3</sub> distributions in chemistry transport models using a potential vorticity-based parameterization. *Atmos. Chem. Phys.*, 16, 10865–10877, doi:10.5194/acp-16-10865-2016, 2016.
- Zhang, L., Jacon, D. J., Boersma, K. F., Jaffe, D. A., Olson, J. R., Bowman, K. W., Worden, J. R., Thompson, A. M., Avery, M. A., Cohen, R. C., Dibb, J. E., Flock, F. M., Fuelberg, H. E., Huey, L. G., McMillan, W. W., Singh, H. B., and Weinheimer, A. J.: Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: an integrated analysis using satellite, aircraft, ozonesonde, and surface observation, *Atmos. Chem. Phys.*, 8, 6117–6136, doi:10.5194/acp-8-6117-2008, 2008.
- 10 Zhang, Y., Liu, P., Pun, B., and Seigneur, C.: A Comprehensive Performance Evaluation of MM5-CMAQ for the Summer 1999 Southern Oxidants Study Episode, Part-I. Evaluation Protocols, Databases and Meteorological Predictions, *Atmos. Environ.*, 40, 4825–4838, doi:10.1016/j.atmosenv.2005.12.043, 2006.
- 15 Zhang, Y., Karamchandani, P., Glotfelty, T., Streets, D. G., Grell, G., Nenes, A., Yu, F.-Q., and Bennartz, R.: Development and Initial Application of the Global-Through-Urban Weather Research and Forecasting Model with Chemistry (GU-WRF/Chem), *Journal of Geophys. Res.*, 117, D20206, doi:10.1029/2012JD017966, 2012.
- Ziemke, J. R., Chandra, S., Duncan, B. N., Froidevaux, L., Bharita, P. K., Levelt, P. F., and Waters, J. W.: Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's  
20 Chemical Transport Model, *J. of Geophys. Res. Atmosphere*, 111, D19303, doi:10.1029/2006JD007089, 2006.

## Webpages

- Acid Deposition Monitoring Network in East Asia (EANET): <http://www.eanet.asia/index.html>, Last Access: 19 April 2018.
- Clean Air Status and Trends Network (CASTNET), U.S. Environmental Protection Agency Clean Air Markets Division:  
25 <https://www.epa.gov/castnet>, Last Access: 26 April 2018.
- Environmental Protection Agency (EPA), Ozone Pollution: <https://www.epa.gov/ozone-pollution/table-historical-ozone-national-ambient-air-quality-standards-naaqs>, Last Access: 1 April 2018.
- National Centers for Environmental Prediction (NCEP), National Center for Atmospheric Research (NCAR), Final (FNL) Operational Global Analysis data: <https://rda.ucar.edu/datasets/ds083.2/>, Last Access: 15 October 2018.
- 30 National Oceanic and Atmospheric Administration (NOAA), Climate Prediction Center (CPC): <http://www.cpc.ncep.noaa.gov>, Last Access: 10 October 2018.

National Oceanic and Atmospheric Administration (NOAA), Earth System Research Laboratory (ESRL), Global Monitoring Division (GMD), ESRL/GMD Ozonesondes: <https://www.esrl.noaa.gov/gmd/ozwv/ozsondes/>, Last Access: 31 August 2018a.

5 National Oceanic and Atmospheric Administration (NOAA), Earth System Research Laboratory (ESRL), Global Monitoring Division (GMD), ESRL/GMD Tropospheric Aircraft Ozone Measurement Program: <https://www.esrl.noaa.gov/gmd/ozwv/aircraft/index.html>, Last Access: 31 August 2018b.

National Aeronautics and Space Administration (NASA), Goddard Space Flight Center, Chemistry and Dynamics Branch, [https://acd-ext.gsfc.nasa.gov/Data\\_services/cloud\\_slice/index.html](https://acd-ext.gsfc.nasa.gov/Data_services/cloud_slice/index.html), Last Access: 31 August 2018a.

10 National Aeronautics and Space Administration (NASA), Goddard Earth Sciences Data and Information Services Center (GES DISC), [https://disc.gsfc.nasa.gov/datasets/OMTO3d\\_003/summary](https://disc.gsfc.nasa.gov/datasets/OMTO3d_003/summary), Last Access: 31 August 2018b.

World Data Centre for Greenhouse Gases (WDCGG): <http://ds.data.jma.go.jp/gmd/wdcgg/>, Last Access: 26 April 2018.






Figure 1. Geographical mapping of the (left) surface and (right) aloft observational sites used in this study. Detailed information about the ozonesonde observation sites is provided in Table 1.



**Figure 2. Scatter-plot between observations and H-CMAQ simulations for surface MD8O3 during April 2010. Reference lines are provided at ratios of 2:1, 1:1 and 1:2. The symbols and colors used to represent the different surface observational datasets are consistent with Fig. 1.**



**Figure 3. Monthly-mean H-CMAQ O<sub>3</sub> mixing ratios overlaid with WDCGG surface observations, and zoom-in panels over (left) Asia overlaid with EANET surface observations, and (right) U.S. overlaid with CASTNET surface observations.**




Figure 4. Curtain plots of modeled (left)  $O_3$ , (center)  $O_3PV$ , and (right)  $RH$  at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), and (c) Huntsville (AL) during April 2010. Yellow stars indicate the time of available ozonesonde measurements. Contour lines of modeled PV are also inserted for contours of 1.0, 1.5, 2.0, 2.5, and 3.0 PVU with thick red lines denoting the 2.0 PVU contour as an index to diagnose the tropopause.

5






Figure 5. Vertical profiles of observed and modeled O<sub>3</sub> and RH at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), and (c) Huntsville (AL). Also see Figure 4 for ozonesonde measurement times. For modeled O<sub>3</sub> and RH, the hourly result corresponding to the ozonesonde measurement time is shown by circles, and the maximum and minimum model results within ±2 hours of the measurement time are shown by whiskers. For observed O<sub>3</sub> at Hilo and Boulder, the range of uncertainties of the O<sub>3</sub> observations is shown by whiskers. Modeled O<sub>3</sub>PV and PV are also shown. Modeled PV profiles are plotted in red, and vertical lines corresponding to a PV value of 2 PVU are inserted as an index of the tropopause, and the layer range diagnosed as stratospheric air mass is colored in purple.




**Figure 6. Curtain plot and vertical profiles of O<sub>3</sub> and RH at the aircraft observational site: Southern Great Plains, Oklahoma. Gray diamonds indicate the times of the seven aircraft flights, and results for these flights are shown in the expanded boxes for 5-h time windows overlaid with observations. Observed ascent and descent profile data are averaged into 100-m grid resolution, and profiles of the mean and standard deviations are shown. Modeled ascent and descent data corresponding to the observation times are averaged on original modeled layers, and the mean and standard deviations are shown.**




**Figure 7. Tropospheric ozone column of (a) satellite observation, (b) H-CMAQ simulation, and (c) their differences shown as normalized bias. Areas filled in gray colored indicate missing observations.**


5




**Figure 8.** Flowchart of the air mass characterization scheme. See the text for additional details.



**Figure 9.** Curtin plot of model-diagnosed air mass characterization for (left) O<sub>3</sub>PV/O<sub>3</sub> and (right) stratospheric air mass at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), and (c) Huntsville (AL) during April 2010.




5 **Figure 10. Relationship between observed MD8O3 at the surface and the estimated stratospheric air mass contributed to total tropospheric O<sub>3</sub> column as monthly-average during April 2010. The points are color coded based on elevation of each site. (a) All CASTNET sites (except Alaska), and (b) elevated CASTNET sites defined as having an elevation greater than 1000 m (see also Table S3).**





**Figure 11.** Spatial distributions of monthly-averaged stratospheric air mass contributions to total tropospheric O<sub>3</sub> column over the U.S. during April 2010. Yellow stars indicate the ozonesonde observational sites, and the daily variations are plotted.

5

(i) Early April



(ii) Late April



5 **Figure 12.** Spatial distributions of day-to-day variations of stratospheric air mass contributions to total **tropospheric O<sub>3</sub>** column over the U.S. during (i) early, and (ii) late April 2010. Yellow stars indicate the ozonesonde observational sites. Red character strings (**S<sub>A</sub>** and **S<sub>B1</sub>-S<sub>B2</sub>**) denote the different stratospheric air masses discussed in the text.

**Table 1. Literature review of previous studies on estimated impacts of (upper) emissions from Asia, and (lower) stratospheric intrusion.**

Time Period	Estimated impacts	Location	Method	Reference
April 2001	1 ppbv (monthly mean), up to 2.5 ppbv (daily mean)	western U.S.	model (process analysis)	Wang et al. (2009)
April-May 2002	10 ppbv increase from April-May 1984	5 west coast U.S. sites	observation (linear regression)	Jaffe et al. (2003)
April-May 2006	5-7 ppbv (17 April-15 May 2006; INTEX-B)	western North America	model (zero-out <sup>a</sup> )	Zhang et al. (2008)
	increase the above influence by 1-2 ppbv	western North America	model (doubling NO <sub>x</sub> emissions from eastern Asia)	
	2-5 ppbv (17 April-15 May 2006; INTEX-B)	eastern North America	model (zero-out <sup>a</sup> )	
April-June 2010	2-6 ppbv increase from April-June 1985	western U.S.	model (tripling Asian emissions from 1985)	Jacob et al. (1999)
	1-3 ppbv increase from April-June 1985	eastern U.S.	model (tripling Asian emissions from 1985)	
May-June 2010	8-15 ppbv on specific events	high-elevation regions	model (zero-out)	Lin et al. (2012a)
April-June 2010	4.7±2.4 ppbv (three-month mean)	15 high-elevation western U.S. sites	model (zero-out)	Lin et al. (2012b)
March-October 2001	below 20 ppbv	U.S.	tagged O <sub>3</sub> <sup>b</sup>	Fiore et al. (2003)
April-May 1990-2012	10-25 ppbv	22 high-elevation western U.S. sites	tropopause tracer <sup>c</sup>	Lin et al. (2015)
April-June 2010	22.3±11.5 ppbv (three-month mean)	15 high-elevation western U.S. sites	tropopause tracer <sup>c</sup>	Lin et al. (2012b)

Note: <sup>a</sup>: Estimate the impact from the difference between the standard simulation and a simulation with eastern Asian anthropogenic sources shut off. <sup>b</sup>: This tagged method divides simulated O<sub>3</sub> into individual O<sub>3</sub> tracer to track O<sub>3</sub> produced in different region. <sup>c</sup>: This tracer method account for STT contribution to O<sub>3</sub> using e90 tracer, which differentiates tropospheric air mass based on the globally uniform surface source and 90-day folding lifetime; both have been spun-up for three years.

**Table 2. Details of the ozonesonde dataset used in this study.**

Country	Site name	Longitude (°)	Latitude (°)	Elevation (m a.s.l.)	Data source	# of launch
USA (HI)	Hilo	-155.05	19.72	10	NOAA ESRL	3
USA (CA)	Trinidad Head	-124.15	41.06	36	NOAA ESRL	5
USA (CO)	Boulder	-105.20	39.95	1743	NOAA ESRL	4
USA (AL)	Huntsville	-86.65	34.73	203	NOAA ESRL	5
USA (VA)	Wallops Island	-75.47	37.93	13	WOUDC	6
USA (RI)	Rhode Island	-71.42	41.49	21	NOAA ESRL	2
Canada (BC)	Kelonwa	-119.4	49.94	456	WOUDC	4
Canada (AB)	Edmonton	-114.1	53.54	766	WOUDC	4
Canada (SK)	Bratt's Lake	-104.7	50.2	580	WOUDC	4
Canada (NU)	Resolute	-94.97	74.71	46	WOUDC	2
Canada (MB)	Churchill	-94.07	58.74	30	WOUDC	4
Canada (NU)	Eureka	-85.94	79.98	10	WOUDC	3
Canada (ON)	Egbert	-79.78	44.23	252	WOUDC	4
Canada (NS)	Yarmouth	-66.11	43.87	9	WOUDC	3
Canada (NU)	Alert	-62.34	82.49	75	WOUDC	3
Canada (NL)	Goose Bay	-60.36	53.3	36	WOUDC	3
Vietnam	Hanoi	105.8	21.01	7	WOUDC	1
China	Hong Kong	114.17	22.31	66	WOUDC	4
Japan	Naha	127.69	26.21	28	WOUDC	3
Japan	Tateno	140.13	36.06	31	WOUDC	3
Japan	Sapporo	141.33	43.06	26	WOUDC	3
Greenland	Summit	-38.46	72.58	3211	NOAA ESRL	4
Ireland	Valentia	-10.25	51.93	14	WOUDC	3
Spain	Madrid	-3.58	40.47	631	WOUDC	3
UK	Lerwick	-1.19	60.14	80	WOUDC	5
Belgium	Uccle	4.35	50.8	100	WOUDC	12
Netherland	De Bilt	5.18	52.1	4	WOUDC	8
Switzerland	Payerne	6.57	46.49	491	WOUDC	12
German	Hohenpeissenberg	11.0	47.8	976	WOUDC	18
Norway	Ny Alesund	11.95	78.93	11	WOUDC	6
Czech	Praha	14.44	50.0	304	WOUDC	10
Poland	Legionowo	20.97	52.4	96	WOUDC	4
Turkey	Ankara	32.86	39.97	890	WOUDC	2

Note: Parenthesis after the country name indicates the state.

**Table 3. Statistical analysis of modeled O<sub>3</sub> concentration using surface, ozonesonde, aircraft, and satellite observations.**

	N	Mean		R	NMB	NME
		Observation	Model			
<b>Surface</b>						
–WDCGG	1498	53.9	43.5	0.49 <sup>***</sup>	–19.3%	23.7%
–CASTNET	2316	53.4	52.9	0.61 <sup>***</sup>	–0.9%	12.6%
–EANET	240	56.2	49.1	0.49 <sup>***</sup>	–12.6%	20.6%
<b>Ozonesonde</b>						
<b>USA and Canada</b>						
–boundary layer	1016	46.0	42.7	0.70 <sup>***</sup>	–7.1%	16.7%
–free troposphere	893	87.7	59.6	0.79 <sup>***</sup>	–32.1%	33.8%
–upper model layer	512	905.1	770.3	0.91 <sup>***</sup>	–14.9%	30.2%
<b>Asia</b>						
–boundary layer	283	44.2	47.4	0.44 <sup>***</sup>	7.1%	24.5%
–free troposphere	207	70.7	59.0	0.43 <sup>***</sup>	–16.5%	21.5%
–upper model layer	124	529.8	399.2	0.94 <sup>***</sup>	–24.6%	34.4%
<b>Europe</b>						
–boundary layer	1478	47.6	46.8	0.42 <sup>***</sup>	–1.6%	17.9%
–free troposphere	1368	78.2	57.2	0.76 <sup>***</sup>	–26.8%	29.1%
–upper model layer	817	1015.7	894.3	0.94 <sup>***</sup>	–11.9%	24.2%
<b>Aircraft</b>						
–from surface upto 6 km	128	55.9	45.3	0.74 <sup>***</sup>	–19.0%	19.1%
<b>Satellite</b>						
–Tropospheric column	28020	33.2	34.7	0.65 <sup>***</sup>	4.7%	13.5%

Note: The unit of mean for observations and simulations is ppbv except satellite observation expressed as D.U. Maximum daily 8 hour average ozone (MD8O<sub>3</sub>) is used for surface observational data of WDCGG, CASTNET, and EANET. Corresponded hourly modeled O<sub>3</sub> is used for ozonesonde data. 2-4 hours averaged hourly modeled O<sub>3</sub> is used for aircraft data to fully cover each observation time. Significance levels by Students' t-test for correlation coefficients between observations and simulations are remarked as \*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001, and lack of a mark indicates no significance.

**Table 4. Statistical analysis of modeled RH using ozonesonde and aircraft observations.**

	N	Mean		R	NMB	NME
		Observation	Model			
<b>Ozonesonde</b>						
<b>USA and Canada</b>						
-boundary layer	1016	57.70	67.07	0.73***	16.2%	24.5%
-free troposphere	881	39.16	43.31	0.83***	10.8%	29.7%
-upper model layer	398	7.81	8.72	0.79***	11.6%	62.3%
<b>Asia</b>						
-boundary layer	283	65.89	79.63	0.45***	20.8%	28.7%
-free troposphere	184	46.26	51.52	0.38***	11.4%	47.7%
-upper model layer	43	18.96	26.47	0.63***	39.6%	67.2%
<b>Europe</b>						
-boundary layer	1485	63.84	68.92	0.73***	8.0%	17.1%
-free troposphere	1368	36.14	42.82	0.80***	18.5%	32.6%
-upper model layer	679	7.13	9.56	0.91***	34.1%	56.1%
<b>Aircraft</b>						
-troposphere	126	41.66	52.04	0.84***	24.9%	28.7%

Note: Significance levels by Students' t-test for correlation coefficients between observations and simulations are remarked as \*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001, and lack of a mark indicates no significance. 5-hour averaged hourly modeled relative humidity is used for ozonesonde data. 2-4 hours averaged hourly modeled relative humidity is used for aircraft data to fully cover each observation time, and original aircraft data are averaged into 100-m resolution to be compared with model.

*Supplemental of*

**Modeling Trans-Pacific Transport and Stratospheric Intrusion of Tropospheric Ozone using Hemispheric CMAQ during April 2010: Part 1. Model Evaluation and Air Mass Characterization for Stratosphere-Troposphere Transport**

Syuichi Itahashi et al.

*Correspondence to:* Syuichi Itahashi (isyuichi@criepi.denken.or.jp)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.







Figure S1. Long-term trends from 2005 to 2015 as before and after 5 years comparison to 2010. (Top) Mean (blue color, left-axis) and maximum and minimum (black color, right-axis) MD8O3 on April. (Center) Number of total observations (black color, left-axis) and exceedance of NAAQS (dark red color, right-axis; 75 ppbv is used as a criterion as 2010) on April. (Bottom) Annual NO<sub>x</sub> (purple) and VOCs (green) emissions in the U.S.A. except wildfire (<https://www.epa.gov/air-emissions-inventories/air-pollutant-emissions-trends-data>).




**Figure S2. (Top) Monthly mean and percentiles of MD8O3 on 2010. (Bottom) Number of total observations (black color, left-axis) and exceedance of NAAQS (dark red color, right-axis; 75 ppbv is used as a criterion as 2010) on 2010.**




**Figure S3. The information of longitude and latitude in H-CMAQ modelling system.**




**Figure S4. Estimated tropopause altitude averaged over April 2010 by (left) the dynamic approach using PV in this work and (right) the thermal approach using the lapse rate.**




**Figure S5.** Curtain plots of modeled (left) O<sub>3</sub>, (center) O<sub>3</sub>PV, and (right) RH at U.S. ozonesonde sites of (a) Hilo (HI), (b) Wallops Island (VA), and (c) Rhode Island (RI) during April 2010. Yellow stars indicate the time of available ozonesonde measurements. Contour lines of modeled PV are also inserted for contours of 1.0, 1.5, 2.0, 2.5, and 3.0 PVU with thick red lines denoting the 2.0 PVU contour as an index to diagnose the tropopause. See also Figure 4.



**Figure S6.** Vertical profiles of observed and modeled O<sub>3</sub> and RH at U.S. ozonesonde sites of (a) Hilo (HI), (b) Wallops Island (VA), and (c) Rhode Island (RI). Also see Figure S5 for ozonesonde measurement times. For modeled O<sub>3</sub> and RH, the hourly result corresponding to the ozonesonde measurement time is shown by circles, and the maximum and minimum model results within  $\pm 2$  hours of the measurement time are shown by whiskers. For observed O<sub>3</sub> at Hilo and Boulder, the range of uncertainties of the O<sub>3</sub> observations is shown by whiskers. Modeled O3PV and PV are also shown. Modeled PV profiles are plotted in red, and vertical lines corresponding to a PV value of 2 PVU are inserted as an index of the tropopause, and the layer range diagnosed as stratospheric air mass is colored in purple. See also Figure 5.





**Figure S7. Curtain plots of modeled PV by (left) WRF over northern hemisphere with a 108 km horizontal grid resolution, (center) WRF over CONUS domain with a 36 km horizontal grid resolution, and (right) WRF over CONUS domain with a 12 km horizontal grid resolution at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI) during April 2010.**





**Figure S8.** Curtain plots of (right) the difference of modeled PV calculated from 36 km–108 km, (center) from 12 km–108 km, and (right) modeled PV lines of 2 PVU by 108 km (red), 36 km (blue), and 12 km (sky blue) at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI) during April 2010.



(a) Trinidad Head





(b) Boulder



(c) Huntsville



(d) Wallops Island



(e) Rhode Island







Figure S9. Correspondence of estimated O<sub>3</sub> concentration based on O<sub>3</sub>/PV relation at the uppermost layer as (left) 108 km vs. 36 km and (right) 108 km vs. 12 km at U.S. ozonesonde sites of (a) Trinidad Head (CA), (b) Boulder (CO), (c) Huntsville (AL), (d) Wallops Island (VA), and (e) Rhode Island (RI). Plots are hourly data during April 2010 (total number is 720).



**Figure S10.** Curtin plot of model-diagnosed air mass characterization for (left) O3PV/O3 and (right) stratospheric air mass at U.S. ozonesonde sites of (a) Hilo (HI), (b) Wallops Island (VA), and (c) Rhode Island (RI) during April 2010. See also Figure 9.



**Figure S11.** Spatial distributions of day-to-day variations of stratospheric air mass contributions to total tropospheric O<sub>3</sub> column over the U.S. during middle April 2010. Yellow stars indicate the ozonesonde observational sites. See also Figure 11.

**Table S1. Statistical analysis of modeled O<sub>3</sub> mixing ratios to ozonesonde as latitude dependence.**

	N	Mean		R	NMB	NME
		Observation	Model			
Ozonesonde						
< 40°N sites						
–boundary layer	517	47.12	49.13	0.55 <sup>***</sup>	4.3%	21.6%
–free troposphere	483	81.07	59.55	0.75 <sup>***</sup>	–26.5%	29.6%
–upper model layer	283	538.86	461.51	0.90 <sup>***</sup>	–14.4%	33.8%
40°–50°N sites						
–boundary layer	907	49.20	48.95	0.37 <sup>***</sup>	–0.5%	16.6%
–free troposphere	831	77.97	58.12	0.76 <sup>***</sup>	–25.5%	29.0%
–upper model layer	488	932.12	845.30	0.92 <sup>***</sup>	–9.3%	26.9%
50°–60°N sites						
–boundary layer	966	46.64	43.37	0.55 <sup>***</sup>	–7.0%	16.9%
–free troposphere	776	78.39	56.28	0.78 <sup>***</sup>	–28.2%	29.3%
–upper model layer	461	1026.12	861.52	0.94 <sup>***</sup>	–16.0%	25.1%
> 60°N sites						
–boundary layer	387	40.30	37.06	0.56 <sup>***</sup>	–8.0%	20.3%
–free troposphere	378	92.86	60.65	0.78 <sup>***</sup>	–34.7%	35.8%
–upper model layer	221	1260.21	1061.29	0.94 <sup>***</sup>	–15.8%	25.3%

Note: Corresponded hourly modeled O<sub>3</sub> mixing ratio is used for the comparison with ozonesonde data. Significance levels by Students' t-test for correlation coefficients between observations and simulations are remarked as \*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001, and lack of a mark indicates no significance. Ozonesonde observational sites located on < 40°N, 40°–50°N, 50°–60°N, > 60°N, respectively, contain 9, 9, 9, and 6 sites.

**Table S2. Statistical analysis of modeled relative humidity to ozonesonde as latitude dependence.**

	N	Mean		R	NMB	NME
		Observation	Model			
<b>Ozonesonde</b>						
<b>&lt; 40°N sites</b>						
–boundary layer	517	61.70	70.69	0.63***	14.6%	26.0%
–free troposphere	469	36.38	42.22	0.72***	16.1%	42.3%
–upper model layer	148	13.81	16.64	0.56***	20.5%	77.2%
<b>40°–50°N sites</b>						
–boundary layer	907	59.74	70.99	0.51***	18.8%	27.5%
–free troposphere	810	38.73	43.47	0.77***	12.5%	34.1%
–upper model layer	339	9.45	10.77	0.91***	13.9%	44.2%
<b>50°–60°N sites</b>						
–boundary layer	973	60.79	65.28	0.85***	7.4%	14.5%
–free troposphere	776	36.04	41.90	0.85***	16.2%	27.6%
–upper model layer	461	6.64	9.27	0.93***	39.7%	56.8%
<b>&gt; 60°N sites</b>						
–boundary layer	387	69.36	73.82	0.81***	6.4%	15.4%
–free troposphere	378	42.46	49.42	0.80***	16.4%	29.8%
–upper model layer	172	2.65	4.12	0.74***	55.5%	102.5%

Note: Corresponded hourly modeled O<sub>3</sub> mixing ratio is used for the comparison with ozonesonde data. Significance levels by Students' t-test for correlation coefficients between observations and simulations are remarked as \*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001, and lack of a mark indicates no significance. Ozonesonde observational sites located on < 40°N, 40°–50°N, 50°–60°N, > 60°N, respectively, contain 9, 9, 9, and 6 sites.

**Table S3. Elevated CASTNET sites in an alphabetical order.**

ID	Site name	State	Longitude (°)	Latitude (°)	Elevation (m a.s.l.)
BBE401	Big Band NP	TX	-103.178	29.303	1052
CAN407	Canyonlands NP	UT	-109.821	38.458	1809
CHA467	Chiricahua NM	AZ	-109.389	32.009	1570
CNT169	Centennial	WY	-106.240	41.365	3175
CON186	Converse Station	CA	-116.913	34.194	1718
GRB411	Great Basin NP	NV	-114.216	39.005	2060
GRC474	Grand Canyon NP	AZ	-112.184	36.059	2073
GTH161	Gothic	CO	-106.986	38.956	2915
JOT403	Joshua Tree NP	CA	-116.389	34.070	1244
LAV410	Lassen Volcanic NP	CA	-121.576	40.540	1756
MEV405	Mesa Verde NP	CO	-108.490	37.198	2165
PAL190	Palo Duro	TX	-101.665	34.881	1053
PET427	Petrified Forest	AZ	-109.892	34.823	1723
PND165	Pinedake	WY	-109.788	42.929	2386
PNF126	Cranberry	NC	-82.045	36.105	1216
ROM206	Rocky Mountain NP Collocated	CO	-105.546	40.278	2742
ROM406	Rocky Mountain NP	CO	-105.546	40.278	2743
SHN418	Shenandoah NP	VA	-78.435	38.523	1073
WNC429	Wind Cave NP	SD	-103.484	43.558	1292
YEL408	Yellowstone NP	WY	-110.400	44.565	2430
YOS404	Yosemite NP	CA	-119.706	37.713	1605

Note: Elevated sites defined as sites with an elevation higher than 1000 m a.s.l. (above sea level). The available sites during April 2010 are listed.