

Interactive comment on “The Structure of Turbulence and mixed-phase Cloud Microphysics in a Highly Supercooled Altocumulus Cloud” by Paul A. Barrett et al.

Paul A. Barrett et al.

paul.barrett@metoffice.gov.uk

Received and published: 7 January 2020

We thank the reviewer for their helpful comments and suggestions.

The reviewer comments are in *italics*, author response in plain text, and modified manuscript text in **boldface**.

C1

1 Minor Comments

I dont believe the authors make a strong connection between the increasing negative skewness of w with increasing distance beneath cloud top and what they assert is driving this, specifically longwave radiative cooling at cloud top. This assertion is stated both in the abstract and in the conclusions and while I agree their conceptual model fits well with that presented in other papers, such as Schmidt et al (2014) for instance, they really dont provide any evidence or even an explanation why radiative cooling should result in w profiles as those observed.

Long-wave Radiative Cooling and skewness. We have added comment in the discussion relating to the Hogan 2009 study which observed the skewness profile of nocturnal stratocumulus to be similar to the profile we observed. This is the basis for our assertion that the source of the turbulence is at cloud top.

The negatively skewed vertical velocity distribution (Fig. 6) is similar to that found in nocturnal (Nicholls, 1989). Hogan et al. (2009) (Fig. 13 (b)) found a similar stratocumulus (Nicholls, 1989; Hogan et al., 2003a), with a profile of skewness for LWRC driven nocturnal stratocumulus clouds using ground based measurements, in contrast with the profiles of skewness that were obtained when surface heating driven cumulus clouds were overhead. The turbulence kinetic...

Last sentence in the abstract: ...used to evaluate numerical simulations of altocumulus clouds at all scales from eddy resolving to climate. The authors are mixing up spatial scales (eddy-resolving) with temporal (climate). I understand the point they are trying to make, but this should be reworded in my opinion.

Abstract text clarified to account for spatial and temporal scale confusion.

These high resolution in situ measurements support previous remotely-sensed observations from both ground based and space borne instruments, and could be used to evaluate numerical model simulations of altocumulus clouds at spa-

C2

tial scales from eddy resolving to global numerical weather prediction models and climate simulations.

Line 93: I'm pretty sure the thermometer did not respond at 32 Hz, even though measurements may have been reported at this high rate. Authors should provide actual response rate of the instrument. Also, many of the measurements were conducted in cloud. Were corrections made for wetting of the element? Or how were in-cloud (liquid) temperatures determined?

Some of the temperature measurements were in fact collected in cloud. We expect that the low liquid water contents (less than 0.05 g m^{-3}) in the cloud result in a negligible impact on the cooling of the non-de-iced sensor housing. A review by Sinkevich and Lawson (2005) of temperature measurements in convective clouds would seem to confirm this. Figure 4 of that paper shows the difference a radiometric measurement of temperature and an immersion sensor as a function of liquid water content. For the liquid water contents that we observed the temperature difference or $0.05 \text{ }^\circ\text{C}$ is an order of magnitude lower than the quoted uncertainty of our temperature sensors.

There was no evidence of contamination on this sensor housing due to the presence of condensed liquid water when compared against the deiced sensor.

The reviewer notes that the temperature sensor is unlikely to respond as fast as 32 Hz even though it is recorded at the rate. We agree and in-fact here we are reporting data at 1 Hz and have amended the text to reflect this.

Temperature measurements were provided recorded at 32 Hz by a non-de-iced Goodrich Type 102 platinum-resistance thermometer and reported at 1 Hz.

We do however use the turbulent wind data at recorded at 32 Hz. We computed power spectra from one of the geometrically level flight legs and found that all three wind components follow a $-5/3$ power law out to at least 10 Hz, possibly closer to the Nyquist frequency of 16 Hz. We reprocessed the wind components after applying a smoothing

C3

window (to give data at a frequency of less than 10 Hz) and found that the TKE was not impacted by more than 5 percent, with skewness values not impacted.

Page 4 and variety of locations in manuscript regarding discussion of using the CIP15 (and CIP100) to determine NI (ice crystal concentrations). Some number of 30-60 (or maybe even 90) micron particles measured by the CIP15 are likely liquid droplets. First, some liquid in these larger size ranges might exist and the CIP15 does not provide sufficient resolution to distinguish between liquid and ice for such small particles. Second, smaller, out-of-focus droplets will appear larger. This could lead to an over-estimation of the ice crystal concentration nearer cloud top where liquid droplets are largest. How do the results change if the analysis were to

We refer to the CIP15 number concentrations for ice in multiple places throughout the text. Whilst we agree that some artefacts may be present in the CIP15 ice data at sizes between 60 and 90 microns, when we only consider the CIP15 data for particles larger than 90 microns we find only minimal difference in the statistics of the vertical profiles of ice number concentrations, that do not alter the conclusions. As part of the quality control process we looked at the histograms on number concentrations from CIP15 and CIP100 and considered individual channels from CIP15. The figure below (Fig. 1) shows the distribution of particle concentrations from CIP15 (green) and CIP100 (orange). Data from CIP15 are clearly bimodal, with the higher concentration mode having number concentrations greater than 0.5 L^{-1} . Plotting only the data from the smallest size channel (Channel 0) on CIP15 (dark red) shows that these particles are the ones smaller than 15 microns, and are almost certainly part of the liquid cloud population. CIP100 does not observe this mode. Channel 1 only from CIP15 (30 micron particles) (red) and channel 2 (45 micron particles) (pink) do not observe this mode. The total concentration of particles larger than a particular size is shown in the blue colours, with particles larger than 30, 45, 75, 135 and 255 microns shown in ever lighter shades of blue. These distributions show only the ice population and look similar to the observations from CIP100. We took this to show that in this situation with

C4

the observed liquid cloud population on the day that the CIP15 only responded to the liquid cloud in the smallest size channel, bin 0 (less than 15 microns). We therefore have confidence that bins larger than this are responding only to ice, but leave a buffer of two adjacent bins.

Any contamination by small particles would not impact the IWC calculation significantly. We do not use the CIP 15 for effective radius calculations, instead we rely on CIP100.

Bottom of page 6, line 187: the impact of wind-shear across the inversion and resultant contamination of the filter should only occur near the inversion, correct? If that is the case, then why substitute w in place of u and v throughout the depth of the profiles. I think you should only do that near cloud top.

In figure 5 we have added a trace showing the TKE estimate when considering all three wind components. This shows the good performance of just considering w' when above -100 m, and that the estimate only using w' is biased high compared to the three wind component estimate, when below -100 m. This is discussed in the text. Below this altitude the estimate of TKE using only w' is biased high as compared to the full three dimensional estimate.

A correction to Figure 5 Panel (b), now shows the mean values trace as green, rather than the previous incorrect magenta.

The new Figure 5 caption reads: **Figure 5. 2 dimensional histograms for 9 km filtered data, against distance from inversion altitude, for (a) vertical velocity fluctuations, w' and (b) TKE, with PDF (Probability Distribution Function) of 32 Hz data (grey-scale - legend in lower panel) and percentiles (yellow). (a) Mean value of w' (green) and standard deviation of w' (magenta) and Skewness of w' (cyan). (b) Mean value of TKE from three component winds (solid green) and with only the vertical component (see text for details) dashed green).**

Around Line 280, discussing estimate of INC based on aerosol larger than 0.5 μm .

C5

The authors provide an estimate based on DeMott et al (2015) and then appear to adjust (increase) that based on a correction factor of 3. But DeMott et al's equation (2) includes CF (calibration factor as they call it). Thus to be consistent, authors should only report the larger value; the lower value of 0.3 L^{-1} doesn't make any sense, or at least is NOT CONSISTENT with DeMott et al (2015).

We agree with the reviewer. Reference to uncorrected INP concentration is removed from the figure and the text. The formatting for the DeMott 2015 reference is also corrected.

Resulting INP concentrations are $N_{INP} = 1.0 \text{ L}^{-1}$ (DeMott et al., 2015) and 0.60 L^{-1} (Tobo et al., 2013).

2 Technical Comments

Line 4: change Turbulence data to Turbulence measurements

Changed Turbulence data to Turbulence measurements

Turbulence measurements are presented from both the liquid cloud layer and ice virga below

Line 16: . . . mixed-phase conditions began within a few metres. . . began is probably not the right word, it implies time. I suggest something like . . . mixed phase conditions were observed within a few metres of cloud top extending downward through the cloud

We agree, began was not the correct word. Text has been altered.

Carey et al. (2008) observed that mid-latitude altocumulus layer clouds were of mixed-phase composition on more than two-thirds of occasions and that mixed-phase conditions were observed within a few tens of metres of observable cloud top and extended down through the cloud

C6

Line 89: abbreviation of knots should it be kts ??

Abbreviation changed to kts

During sampling the aircraft typically had a nominal Indicated Airspeed kts...

Line 275: Fig. 12@ (b) ?

typo corrected

Fig. 12(b)

A.A. Sinkevich and R.P. Lawson, A Survey of Temperature Measurements in Convective Clouds, J. Appl. Met. 44, 1133-1145, 2005

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-749>, 2019.

C7




Fig. 1. Cloud particle number concentration histograms from CIP15 and CIP100

C8